





51213/B/1

1690

# This Volume

was given to the

Manchester Medical Society

by

J. A. Kelham Esq.

of

Manchester

August 18<sup>th</sup> 1908







07463.


Supp.

pages

Index

at end.

missing





THE  
**LONDON DISPENSATORY:**

CONTAINING

- I. THE ELEMENTS OF PHARMACY;  
II. THE BOTANICAL DESCRIPTION, NATURAL HISTORY, CHEMICAL  
ANALYSIS, AND MEDICINAL PROPERTIES, OF THE  
SUBSTANCES OF THE MATERIA MEDICA;  
III. THE PHARMACEUTICAL PREPARATIONS AND COMPOSITIONS OF  
THE PHARMACOPŒIAS OF THE LONDON, THE EDINBURGH,  
AND THE DUBLIN COLLEGES OF PHYSICIANS.

THE WHOLE FORMING  
A PRACTICAL SYNOPSIS OF  
MATERIA MEDICA, PHARMACY, AND  
THERAPEUTICS:

*ILLUSTRATED WITH MANY USEFUL TABLES,*  
AND WOOD-CUTS OF THE PHARMACEUTICAL APPARATUS.

---

BY  
**ANTHONY TODD THOMSON, M.D. F.L.S.**

MEMBER OF THE ROYAL COLLEGE OF SURGEONS,  
AND OF THE MEDICO-CHIRURGICAL SOCIETY OF LONDON;  
FELLOW OF THE MEDICAL, THE SPECULATIVE, AND THE ROYAL PHYSICAL  
SOCIETIES OF EDINBURGH;  
MEMBER OF THE SOCIÉTÉ D'ÉMULATION DE PARIS, AND OF  
THE SOCIÉTÉ DE MÉDECINE DE MARSEILLES.

---

*FOURTH EDITION.*

---

LONDON:  
PRINTED FOR LONGMAN, REES, ORME, BROWN, AND GREEN,  
PATERNOSTER-ROW.

1826.





# LONDON DISPENSATORY:

CONTAINING

I. THE ELEMENTS OF PHARMACY;  
II. THE BOTANICAL DESCRIPTION, NATURAL HISTORY, CHEMICAL  
ANALYSIS, AND MEDICINAL PROPERTIES, OF THE  
SUBSTANCES OF THE MATERIA MEDICA;  
III. THE PHARMACEUTICAL PREPARATIONS AND COMPOSITIONS OF  
THE PHARMACOPOEIA OF THE LONDON, THE EDINBURGH,  
AND THE DUBLIN COLLEGES OF PHYSICIANS.

THE WHOLE FORMING

## A PRACTICAL SYNOPSIS OF MATERIA MEDICA, PHARMACY, AND THERAPEUTICS:

ILLUSTRATED WITH MANY ENGRAVED TABLES  
AND WOOD-CUTS OF THE PHARMACEUTICAL APPARATUS.

BY  
ANTHONY TODD THOMSON, M.D. F.R.S.

MEMBER OF THE ROYAL COLLEGE OF SURGEONS,  
AND OF THE MEDICO-CHIRURGICAL SOCIETY OF LONDON;  
FELLOW OF THE MEDICAL, THE SURGEON, AND THE ROYAL PHYSICAL  
SOCIETIES OF LONDON;  
MEMBER OF THE SOCIETY OF PHYSICIANS IN PARIS, AND OF  
THE SOCIETY OF MEDICAL PHYSICIANS.



LONDON:  
Printed by A. & R. Spottiswoode,  
New-Street-Square.



TO  
SIR JAMES MACGREGOR, KNT. M.D.F.R.S.

FELLOW OF THE ROYAL COLLEGES OF PHYSICIANS  
OF LONDON AND EDINBURGH;

PHYSICIAN EXTRAORDINARY TO THE KING;

AND  
DIRECTOR GENERAL OF THE ARMY MEDICAL DEPARTMENT,  
&c. &c. &c.

AS  
A TESTIMONY  
OF RESPECT AND ADMIRATION,  
FOR  
HIS ARDUOUS AND WELL DIRECTED EFFORTS  
TO PROMOTE  
THE CULTIVATION OF GENERAL SCIENCE  
IN  
THE PUBLIC DEPARTMENT UNDER HIS CONTROL,  
AND CONSEQUENTLY,  
TO ELEVATE THE CHARACTER OF THE  
MEDICAL PROFESSION,

*THIS VOLUME IS INSCRIBED,*

BY  
HIS FRIEND,  
THE AUTHOR.



Digitized by the Internet Archive  
in 2018 with funding from  
Wellcome Library



## ADVERTISEMENT.

---

THE progressive advancement of Chemical Science and many new discoveries in Natural History, have contributed, within the space of a very few years, to alter materially the character of Pharmacy; and consequently to give to each new Edition of the Dispensatory the aspect, almost, of an original work. These circumstances have augmented the author's desire to make this volume as complete as possible: and, while he has endeavoured to correct some errors which found their way into the former editions, he has endeavoured to add as much new matter, both to the text and the notes of this edition, as the course of his reading has enabled him to collect.

Among the new works which he has consulted, candour obliges him to say, that he has taken advantage of the excellent translation of the Pharmacopœia of the Royal College of Physicians of London, by Mr. R. PHILLIPS, Professor BRANDE'S Manual of Pharmacy, and the Parisian Codex Medicamentarius. He has, also, availed himself of every work on Natural History which has come within his knowledge, as well as the numerous Voyages and Travels which have appeared since the publication of the last Edition of this



work; and he trusts, that the information which he has obtained from these sources, has greatly enriched the field of his exertions on the present occasion. It has been indeed objected that the facts in Natural History which the author has endeavoured to accumulate in this work are of no practical utility, and are out of place. He replies in the words of Pliny, (lib. i. in Proœm.) “*Rerum Naturæ contemplatio quamvis non faciat Medicum; aptiorem tamen medicinæ reddit, atque perfectum.*”

91. *Sloane Street,*  
30th January, 1826.



# PREFACE

## TO THE FIRST EDITION.

---

THE unnecessary multiplication of books on the same subject may be said, with propriety, to be a great evil: and, therefore, in undertaking the compilation of the following Work, which in its arrangement and plan coincides very closely with the Edinburgh New Dispensatory, it has been my endeavour to render it essentially so different from that work, as to prevent it from being placed under this reproach. But the plan of that volume, which is nearly the same as was originally adopted by Dr. Lewis, with some little alterations, has become so familiar to the profession, and is so well adapted for the purposes of a Dispensatory, that I have thought it prudent not to deviate from it. I trust, however, that the alterations and the additions which I have introduced, particularly in the history of the different articles of the *Materia Medica*, will give a legitimate value to the London Dispensatory; and, without a wish to detract from the high character of its precursor, will enable it to excite a new interest in the branch of the science of medicine of which it treats; and especially to turn the attention of the student towards Medical Botany, which has been so unaccountably neglected, as to be almost regarded as unnecessary in the education of a physician. Indeed, although it has always been admitted that a correct knowledge of *Materia Medica* and Pharmacy can be obtained by those only who have a previous knowledge of Botany and Chemistry, yet neither the Dispensatories nor the systems of *Materia Medica* published in this country, have described plants in a scientific manner; or noticed, in their descriptions, those characteristics which botanists have fixed on as the



only means by which a plant, that is not familiar to the reader of a description of it, can with certainty be known, when he wishes to possess it, or is in any doubt regarding it when it is obtained. From the want of this degree of accuracy in the descriptions of plants, many valuable remedies, used by the inhabitants of one part of the world, have been lost to those of another part, where they are, nevertheless, indigenous; or, instead of the proper plants, other species of the same genera, which possess little or no virtue, have been employed; and even plants, not in any respect medicinal, but highly deleterious, have been used as remedies, merely from their bearing names in common or in pharmaceutical language similar to those of some medicinal plants. To prevent this evil, therefore, I have added to the usual account of each vegetable substance the characters of the genus to which the plant belongs, as they are given by Willdenow in his excellent edition of the *Species Plantarum*; and have also given detailed descriptions of each in the language employed by modern botanical writers.

In the performance of the task of compilation, I have endeavoured to bring together as much useful information, regarding each of the substances treated of, as could be crowded into a small space; and to obtain it, have had recourse to every work of reputation to which I could gain access. For the liberality of Sir Joseph Banks, who, through the kind interference of Dr. Gartshore, opened to me the door of his princely collection, I have to return my most grateful thanks; as I obtained information there which I could not otherwise have procured, and some which has never before, I believe, been given to the public in an English garb. Of this nature, in particular, are the observations of Zea on the medicinal species of Cinchona, extracted from the *Anales de Historia Natural*; those of Humboldt on the same subject, and on some other South American plants, from his splendid work entitled *Plantæ Æquinoctiales*; and Willdenow's description of the *Heracleum gummiferum*, which I have translated from the *Hortus Berolinensis*. Information has also been



sought for, and obtained, from other sources besides books; and I have received from individuals engaged in the drug trade some notices regarding the forms in which drugs are imported, and the modes of selecting them; which I hope will prove useful.

The botanical descriptions of the plants have been selected chiefly from *Martyn's edition of Miller's Gardener's Dictionary*; the last edition of *Woodville's Medical Botany*; *Smith's Flora Britannica*; *Sowerby's English Botany*; the *Flora Peruviana*; *Rheede's Hortus Malabaricus*; and the *Flora Danica*; but in every instance when the living plants could be obtained,—which was the case with the greater number of the indigenous plants, and many exotics also,—the descriptions have been either drawn up from nature; or those adopted have been carefully compared with the plants themselves, and any errors corrected. For other information on this part of the subject, the excellent work of *Gærtner de Fructibus*; *Bergius' Materia Medica a Regno Vegetabili*; *Murray's Apparatus Medicaminum*; *Alston's Materia Medica*; and the *Linnean*, and the *Philosophical Transactions*, with the best books of travels, have been consulted.

For the chemical part I have principally consulted the last editions of the *System of Chemistry of Doctor Thomson*, and that of my very ingenious friend *Dr. Murray*; the *Annales de Chimie*; the *Journal de Physique*; the *Philosophical Transactions*; and the best chemical papers inserted in the periodical publications; nor have I refrained from taking advantage of the observations of *Dr. Duncan* contained in the *Edinburgh New Dispensatory*.<sup>1</sup> As this is undoubtedly the most import-

---

<sup>1</sup> The following are the more important of the additional works, which have also been consulted in compiling the three former and the present edition:—

*Flore, du Dictionnaire des Sciences Médicales, décrites par F. P. Chaumeton, Chamberet et Poiret*, 8vo. Paris.

*Histoire Nat. et Med. des Cases, &c. &c.*: L. T. Calloden de Geneve, M. D. 4to. Montpelier, 1816.

*Materia Medica of Hindostan, &c. &c.* By Whitelaw Ainslie, M. D. 4to. Madras, 1813.

*A Treatise on the Medicinal Leech, &c. &c.* By T. Johnson, M. D. 8vo. London.

*Mem. de la Société Imperiale des Nat. de Moscow*, 4to. Moscow, 1809.

*Asiatic Researches*, 4to. Calcutta.



ant part of the work, I have given it all the attention I am possessed of; and although it would be almost impossible, and altogether unnecessary, to prove by actual experiment the correctness of all the formulæ for the preparations and compositions ordered in the Pharmacopœias, yet the greater part of those ordered in the London Pharmacopœia, which I have chosen for my text-book, have been repeated, and an unrestrained opinion regarding them delivered. Those chemical theories only, however, which are fully established have been adopted; and I have studiously avoided ranking myself on either side, on the important questions now at issue, suggested by the late investigations of Mr. (now Sir H.) Davy.<sup>1</sup>

With regard to the reformed nomenclature of the Pharmacopœias, the entire principle of which has been strongly objected to by Dr. Bostock, in his *Remarks on the Nomenclature of the London Pharmacopœia*, I conceive it to be my duty to state here, that although I have not scrupled freely to criticize such names as appear to me to be improper in the work of the London College, and incompatible with the views of the subject which it professes to have taken, yet, that I do not

*Tableaux Chimiques du Règne Animal, &c. par Jean-Frédéric John, &c. traduit de l'Allemand, par Stéphane Robinet, 4to. Paris, 1818.*

*Annales de Chimie et de Physique, 8vo. Paris.*

*Henry's Elements of Experimental Chemistry, 7th edit. 8vo. London.*

*Elementi di Chimica Farmaceutica e d'Istoria Naturale, &c. 8vo. Torino.*

*Nouveaux Elémens de Therapeutique et de Matière Médicale, &c. par I. L. Alibert, 8vo. 4me edit. Paris, 1817.*

*Farmacopea Generale ossia Dizionario delle Preparazioni Farmaceutico-Mediche, &c. di L. V. Brugnatelli, 8vo. Pavia, 1814.*

*Traité des Poisons, &c. &c. par M. P. Orfila, 8vo. Paris.*

*Philips' Remarks on the Editio Altera of the Pharm. Lond. 8vo. London.*

*Journal de Pharmacie et des Sciences Accessoires, &c. 8vo. Paris.*

*Journal of Science and the Arts, 8vo. London.*

*The Annals of Philosophy, 8vo. London.*

*Medico-Chirurgical Transactions, 8vo. London.*

*The London Medical Repository, 8vo. London.*

*Bancroft's Philosophy of Permanent Colours, 2d edit. 8vo. London.*

*Lambert's Illustrations of the Genus Cinchona, 4to. Lond. 1821.*

*Burckhardt's Travels in Nubia, 4to. Lond. 1820.*

*Davy's Account of the Interior of Ceylon, 4to. Lond. 1821.*

*Humboldt's Personal Narrative, 8vo. Lond. 1821.*

*Opium Historice, Chemice atque Pharmacologicæ Investigatum, per C. A. Christen, 8vo. Vind. 1820.*

*Pharmacologia, &c. &c. ; 3d. edit. 8vo. Lond. 1820.*

*Pharmacopœa Batava et Notis, &c. Lipsiæ, 1824.*

<sup>1</sup> The establishment of Sir H. Davy's theories has rendered the reserve implied in this sentence no longer necessary.



concur with that gentleman in thinking that the reform was unnecessary, or now, when it is accomplished, that it is not likely to be generally adopted. It is, indeed, deeply to be lamented, that a work issuing from so respectable and learned a body as the London College is acknowledged to be, is not more perfect in its execution: but, nevertheless, its nomenclature is much superior to that of its predecessor; and within the scope of my own observation, as well as from information which I have received from others, it is now coming into very general use. Many of the older physicians already write their prescriptions in the reformed language; and it is so congenial with the habits of thinking, and the ordinary language and mode of writing of the younger practitioners, that there is no doubt of its being readily adopted by them.<sup>1</sup>

The best authors have been consulted regarding the medicinal properties and uses of the various substances treated of; but I must confess that less attention has been given to this, as it is not intended that the practice of medicine should be taught by a Dispensatory: and surely little is to be expected from those who would attempt to acquire it from such a source. As memorandums, however, I have been anxious to make the medical notices as correct as possible; and have, in most instances, given references to the works from which the best information on this part of the subject may be obtained.

After all my efforts, however, to avoid errors, I am sensible that the work may, nevertheless, contain many; I have, therefore, determined to lend a willing ear to the remarks of candid criticism; and, in future editions, to take advantage of every suggestion that may tend to bring it nearer to that state of perfection, the attainment of which must afford more satisfaction to an author than all the indiscriminate praise which can be lavished upon his labours. I might, indeed, plead as an apology for its imperfections, that the work has not been prepared in the repose of retirement, and with an abundance of leisure; but on the contrary, that I have been

---

<sup>1</sup> The experience of ten years has fully confirmed these anticipations.



able to bestow on it the few leisure hours only which I could snatch from full employment in a laborious branch of the profession, and during the wasting of the midnight oil, amidst inconvenience and anxieties; and with the perpetual annoyance of professional interruptions. I am sensible, however, of the futility of such an apology; and as the book is now in the hands of the profession, it must fall or rise according to its merits: I can only hope these will be justly appreciated by the Public, that tribunal, by which every production, whether of intellect or of labour, must be ultimately judged.

ANTHONY TODD THOMSON.

91. *Sloane Street,*  
*April, 1811.*

### Explanation of the Abbreviations employed in the Synonyma.

( <i>F.</i> ) French.	( <i>San.</i> ) Sanscrit.
( <i>G.</i> ) German.	( <i>Tam.</i> ) Tamool.
( <i>I.</i> ) Italian.	( <i>A.</i> ) Arabic.
( <i>S.</i> ) Spanish.	( <i>Pers.</i> ) Persian.
( <i>Dutch.</i> )	( <i>Beng.</i> ) Bengalese.
( <i>Swed.</i> ) Swedish.	( <i>Duk.</i> ) Dukhanie.
( <i>Dan.</i> ) Danish.	( <i>Malay.</i> )
( <i>Pol.</i> ) Polish.	( <i>Cyng.</i> ) Cynghalese.
( <i>Port.</i> ) Portuguese.	( <i>Celebes.</i> )
( <i>Russian.</i> )	( <i>Jav.</i> ) Javanese.
( <i>H.</i> ) Hindoostanie.	( <i>Esqui.</i> ) Esquimaux



# CONTENTS.

---

## PART I. ELEMENTS OF PHARMACY.

### SECTION I.

#### I. ATTRACTION.

*a.* OF AGGREGATION.

*b.* OF AFFINITY.

#### II. REPULSION, AND THE POWERS PRODUCING IT.

1. OF CALORIC.

2. OF LIGHT.

3. OF ELECTRICITY and GALVANISM.

### SECTION II.

#### CONSTITUTION AND COMBINATION OF SUBSTANCES.

1. OF SOLIDS.

2. OF LIQUIDS.

3. OF GASES.

### SECTION III.

#### PHARMACEUTICAL OPERATIONS—APPARATUS.

---

## PART II. MATERIA MEDICA.

---

## PART. III. PREPARATIONS AND COMPOUNDS.

ACIDS.

ALKALIES AND NEUTRAL SALTS.

EARTHS AND EARTHY SALTS.

METALLIC PREPARATIONS.

PREPARATIONS OF SULPHUR.

PREPARATION OF VEGETABLES.



EXPRESSED OILS.  
DISTILLED OR VOLATILE OILS.  
DISTILLED WATERS.  
DECOCTIONS.  
MUCILAGES.  
INFUSIONS.  
EXTRACTS.  
MIXTURES.  
SPIRITS.  
TINCTURES.  
PREPARATIONS OF ETHER.  
WINES.  
PREPARATIONS OF VINEGAR.  
PREPARATIONS OF HONEY.  
SYRUPS.  
CONFECTIONS.  
POWDERS.  
PILLS.  
TROCHES.  
PREPARATIONS FROM ANIMALS.  
PLASTERS.  
CERATES.  
OINTMENTS.  
LINIMENTS.  
CATAPLASMS.

TABLE *of the Proportions of* OPIUM, &c. *in certain Compounds.*  
TABLES *of* NEW NAMES.

## APPENDIX.

Nº I. OF WATERS.  
Nº II. *Method of Analysing* MINERAL WATERS.  
Nº III. OF THE DOSES OF MEDICINES.  
Nº IV. TABLE OF INCOMPATIBLES.  
Nº V. EXAMPLES OF EXTEMPORANEOUS PRESCRIPTIONS.  
Nº VI. TABLE OF PHARMACEUTICAL EQUIVALENTS.  
Nº VII. *Method of obtaining* MECONATE OF MORPHIA.  
Nº VIII. TABLE *of the* SYNONYMA *of the Terms of*  
*London, Edinburgh, and Dublin Pharmacopœias.*  
LATIN INDEX.



THE  
LONDON DISPENSATORY.

---

PART I.

---

ELEMENTS OF PHARMACY.

**P**HARMACY is that branch of the science of chemistry which relates to the combination and mixture of different substances, for the purposes of medicine.

Its practice presupposes a knowledge of the ultimate principles of the substances employed in its operations, and of their chemical agencies; and thence, of the general doctrines of Chemical Science. The elements, therefore, of Pharmacy, properly speaking, are those of Chemistry; and without a knowledge of these, it can neither be theoretically understood, nor advantageously practised as an art.

As, however, it would be impossible in this place to give more than an outline or epitome of the elements of Chemistry; and as the second part of this work is intended to contain the analysis as well as the history and an account of the uses of the different articles of the *Materia Medica* which constitute the subjects of Pharmacy, I shall confine the term Elements of Pharmacy to comprehend those general principles of chemical action which enable us to reason on, and perceive the proximate causes of the results of pharmaceutical combinations; and to the explanations of the operations of Pharmacy, with a description of the apparatus.



## TABLE OF ARRANGEMENT.

## SECTION I.

OF THE MORE GENERAL AGENTS INFLUENCING  
PHARMACEUTICAL COMBINATIONS.

## I. ATTRACTION.

- a. *Attraction of Aggregation.*
- b. *Chemical Attraction, or Affinity.*

## II. REPULSION.

*Powers by which it is produced.*

1. CALORIC.
  2. LIGHT.
  3. ELECTRICITY and GALVANISM.
- 

## SECTION II.

OF THE CONSTITUTIONS AND COMBINATIONS OF  
SUBSTANCES.

1. OF SOLIDS.
  2. OF FLUIDS.
  3. OF AËRIFORM SUBSTANCES OR GASES.
- 

## SECTION III.

OF PHARMACEUTICAL OPERATIONS AND THE  
DESCRIPTION OF THE APPARATUS.

## SECTION I.

THE agents which more generally influence chemical and, thence, pharmaceutical combinations, are *Attraction and Repulsion*.

## I. ATTRACTION.

ATTRACTION is the term employed to denote that power which causes bodies to approach towards each other, and which preserves them in a state of union after they come into contact. We are ignorant of the cause of this power, but some of the laws respecting it are sufficiently evident; and from observing the different phenomena to which these give rise, we are inclined to believe that there are different species of attractions, although, perhaps, the difference is more in degree than in kind.

When this power is exerted on masses of matter, at sensible distances, in the direct ratio of the quantity of the matter, and



the inverse ratio of the square of the distance, it is named *Gravitation*; but when its operation is confined to the minute atoms of bodies, and is exerted only when these are near to each other, or in apparent contact, it is denominated *Contiguous Attraction*. The former causes bodies on this globe to fall in a line perpendicular to its surface, preserves the planets in their orbits, and sustains in their places all the parts of the magnificent frame of the universe: the second is the cause of the regular figures of natural bodies, and of the various combinations of matter, which take place in and upon the surface of our globe. It is this species which we are here to examine.

CONTIGUOUS ATTRACTION, operating on particles of the same kind, forms an aggregate or mass; and the power, in this instance, is named the *attraction of aggregation* or *cohesion*: but, acting on dissimilar particles, and producing bodies possessed of new properties, different from those of their components, it constitutes *chemical attraction* or *affinity*.

a. OF COHESION.

The Attraction of *Cohesion* is that force which retains together the particles of bodies at insensible distances. According to the degree of force which it exerts, substances assume the solid, the fluid, or the aëriform state.

1. In *solid* bodies this force is sufficiently powerful to prevent their component particles from being moved with regard to one another, except in a very small degree; and to oppose a considerable resistance to any mechanical power applied to separate them. In the same kinds of bodies, all the circumstances being equal, it is always the same; but in dissimilar bodies it is exceedingly various: from which, and the peculiar arrangement of the particles, arise the different qualities of solids, denominated hardness, softness, malleability, ductility, and elasticity.

The attraction of cohesion in solids is exerted at insensible distances only, and may be weakened or altogether overcome by caloric, or that matter which produces the sensation of heat. If a piece of ice, for example, be brought near a fire, the cohesion of its particles is weakened as the caloric flows into it, till it is changed from the solid state to the fluid, or water; and by continuing and increasing the heat, the particles are still further separated from one another, until the fluid passes into the gaseous form, or becomes steam. This power is also weakened by chemical affinity; for, when a solid body is put into a fluid, the affinity between the particles of the fluid and those of the solid is often sufficient to overcome the aggregation of the solid; and its detached particles, being uniformly diffused through the fluid, now form a part of it, without altering



either its fluidity or its transparency. This constitutes the ordinary chemical or pharmaceutical process of *solution*, which is always favoured by the application of heat, owing to the assistance it affords in overcoming the cohesive attraction, as has been already noticed.

2. In *liquid* bodies this force also operates, but in a less degree than in solids, their particles being at greater relative distances, and moveable with regard to each other by a very small force; but, as their mobility does not change their relative distances, they remain within the sphere of this attraction, and are kept together. The exertion of this power varies in different liquids: it is greater in mercury than in water, and in this than in alcohol. It offers, however, scarcely any resistance to the combination of fluids with other bodies: and thence, the mutual affinity of two bodies is always favoured, when one of them is in the liquid state.

3. This attraction is not exerted over *aëriform* substances; for while these remain at the temperature necessary for the preservation of their aërial state, their particles mutually repel each other, and would recede to an indefinite distance, were they not prevented by the pressure of the surrounding bodies. Thus, a portion of air which can be contained in a vessel of 1 cubical inch of capacity, will fill a vessel of 100 cubical inches of capacity, if the pressure which confines it within the smaller vessel be removed.

One of the most important results of this variety of contiguous attraction, in a pharmaceutical point of view, is the formation of crystals, or the regular and determinate figures assumed by many bodies in passing from the fluid to the solid state, when nothing opposes the union of their particles according to the laws of aggregation.

The process of *crystallization* requires that the particles of the substance to be crystallized become moveable; and, consequently, in order to obtain any body in a crystalline state, it must first be rendered fluid, either by solution in a liquid, or by fusion.

The crystallization of salts is usually effected in the first method. When a salt is much more soluble in hot water than in cold, as is the case, for example, with sulphate of soda, nothing more is required for its crystallization than to saturate boiling water with the salt, and set the solution aside to cool. As the caloric is dissipated, the saline particles gradually approach one another, and uniting, form solids of that regular shape which characterizes the crystals of this peculiar salt. But, when the salt is one which is almost equally soluble in hot and in cold water, as muriate of soda, for instance, its



crystallization can be effected only by evaporating a part of the fluid ; and the more slowly the evaporation proceeds, the mutual attraction of the particles is more regularly effected, and the more regular is the shape of the crystals which are obtained. In both cases, however, the affinity of the saline particles at length ceases to act, while the fluid still retains as much saline matter as it can hold dissolved at the temperature of the atmosphere, or is a saturated solution ; but, by a greater reduction of temperature in the one case, and a further evaporation in the other, it will again yield crystals.

By *fusion*, bodies which are not soluble in water, as glass, sulphur, &c. are enabled to assume the crystalline form. In this case the body is, as it were, dissolved in caloric : and the particles being separated from one another, these, when the cooling is gradual, assume, in aggregating again, the regular arrangements which take place in crystallization. This mode of crystallizing substances is never used for pharmaceutical purposes.

Crystallization is promoted or retarded by various circumstances, to be afterwards noticed. (See *Section iii.*) Its theory is still obscure ; but some light has been thrown upon it by the experiments of Haüy. He found that crystals may be mechanically divided, and reduced to certain primitive forms, which are always the same in the same kind of substances, and depend upon the figure and the mode of combination of the integrant particles composing the crystals. The varieties of figure of these particles, notwithstanding the great diversity of crystalline forms, are reducible to three : namely, 1. the parallelopiped, the faces of which are six, parallel two and two ; 2. the triangular prism ; and 3. the tetrahedron, or four-sided pyramid : and these particles, therefore, according to the mode in which they unite, which may be either by their faces or their edges, form primitive crystals, which are the nuclei of the secondary crystals. The forms of primitive crystals may be reduced to the following six : 1. the parallelopiped, which includes the cube, the rhomboid, and all solids terminated by six faces, parallel two and two ; 2. the regular tetrahedron ; 3. the octahedron with triangular faces ; 4. the six-sided prism ; 5. the dodecahedron, terminated by rhombs ; and 6. the dodecahedron, with isosceles triangular faces. The variations of the forms of secondary crystals are considerable in the same salt, and depend, in general, either on variations in the proportions of the ingredients which compose the integrant particles, or on the properties of the solvent in which the crystals are formed. Thus, alum crystallizes in octahedrons, but the addition of a little alumina produces cubes ; and an excess of this earth prevents crystallization altogether :



thus also, muriate of soda, which crystallizes in cubes, when dissolved in water, assumes the regular octahedral form when it is crystallized in urine. Independent, however, of these causes, a variety of secondary forms make their appearance; which the theory of Haüy explains, by supposing that, as the matter which envelopes the primitive nucleus to form a secondary crystal is attracted in thin layers, each layer decreasing in size in consequence of one or more rows of integrant particles being abstracted from its primitive edges or angles. The decrements may be on the *edges* of the slices, which correspond with the edges of the primitive nucleus; or on the *angles*, that is, parallel to the diagonals of the faces of the primitive nucleus; or the decrements may be *intermediate*, parallel to lines situated obliquely between the diagonals and edges of the faces of the primitive nucleus. It would be impossible, however, to give a satisfactory view of this ingenious theory in the narrow compass of this epitome; and therefore I must refer the reader to Haüy's *Traité de Minéralogie*, tomes 1 and 2; to the *Annales de Chimie*, tom. 17; and the third volume of the fifth edition of Thomson's *System of Chemistry*.

Such is the Attraction of Aggregation, and its general effects. It is frequently concerned in modifying pharmaceutical results; but it is a power of much less importance than the next variety of *contiguous attraction*.

b. OF CHEMICAL ATTRACTION, OR AFFINITY.<sup>1</sup>

*Chemical attraction*, or *affinity*, is that power by which dissimilar substances placed under certain circumstances are enabled to unite, and form new aggregates, in which the properties of the component particles are lost or changed. Its action is confined to the minute atoms or particles of bodies, and is exerted only at insensible distances: not indifferently, however, but electively, between the particles of all bodies. The result of its operation is a *combination* of the constituent particles of the substances, so intimate that the components cannot be recognised, nor separated by any mechanical force. Thus, lime acts as a powerful caustic when applied to animal matter, and is partially soluble in water; phosphoric acid has an acid taste, and is very soluble in water; but phosphate of lime, the compound produced by the chemical combination of these substances, is inert when applied to animal matter, insipid, and insoluble in water; and cannot be again resolved into lime and phosphoric acid by any mechanical power.

*Chemical combination*, therefore, is the result of the affinity

---

<sup>1</sup> For many of the following observations on affinity, I am indebted to the remarks of my late respected friend, Dr. Murray: see his *System of Chemistry*, vol. i.



of two or more substances for each other. It differs from *mixture*, in which the substances are only blended without acquiring any new properties, and in which the dissimilar parts are easily discovered, and may be separated by mechanical powers. Chemical compounds can, however, be decomposed, either by exposure to a high temperature, which weakens the force of attraction existing between their principles; or, by mixture, under favourable circumstances, with some other chemical agent, which has a more powerful affinity for one of the components than these have for each other: and by such means, which constitute *chemical analysis*, the principles of a compound may be ascertained.

As *Analysis* separates compounds into their constituent principles, so *Synthesis* may reproduce them by recombining these principles; and when this can be effected, it is the surest proof of the accuracy of any analysis. In many instances, however, this is impossible: and the evidence of the truth of an analysis is to be drawn from other sources.

It is an acknowledged law of chemical affinity, that a compound “does not possess properties merely intermediate between those of its component parts, but has acquired others more or less new.” One of the most general changes is that of form. The combination of two gases, for example, *muriatic gas* and *ammoniacal gas* form a fluid: that of two fluids, again, may form a solid, as is demonstrated by dropping into a saturated solution of *muriate of lime*, some concentrated *sulphuric acid*; and the common process of *solution* presents to us the fact, that, by the combination of a solid with a fluid, the solid assumes the fluid form. In the last-mentioned instance the fluid is generally regarded as the active substance; but, nevertheless, the attraction of affinity is reciprocal; and thence, the general mode of expressing the fact, that the fluid dissolves the solid, or is the *solvent* or *menstruum*, is, in strict language, erroneous. These terms, however, are more correctly applied, when the properties of the solid, except form, are scarcely sensibly altered; as, for example, when common salt, *muriate of soda*, is dissolved in water.

Chemical combination produces an alteration of *density* — that of the compound not being the mean of the components, but often different. In the greater number of cases the density is increased; and there is a diminution of volume, owing, probably, to the compound atom being of a form admitting a more compact aggregation than the component atoms in their separate state; but the specific gravity of a compound cannot be determined by calculation from the specific gravity of its ingredients. There are cases of combination, however, in which



the density is diminished, and there is an increase of volume in the resulting compound: for instance, when a solid is dissolved by a liquid, the increase of volume acquired by the solid, in passing into the fluid state, may be greater than the condensation resulting from its union in that state with the liquid; and this happens from the solution of a considerable number of the salts in water.

“The exertion of chemical attraction is accompanied by a *change of temperature*.” Thus, if four parts of sulphuric acid and one part of water, both at the temperature of  $32^{\circ}$ , be mixed together, the temperature of the mixture rises to  $300^{\circ}$ ; and the density of the compound is much greater than the mean of the densities of the components. The heat also which is evolved by combustion, and in fermentation, is the direct consequence of chemical combination. In all cases the increase of temperature is accompanied with an increase of density, to which, and the change of form suffered by one or both of the components, its production may be ascribed. Thus water, which is solidified by being mixed with quicklime, parts with a large portion of caloric. The contrary effect, however, or an absorption of caloric, is also produced by chemical combination, when the density of the compound is less than the mean; as, for instance, when, by solutions of salts in water, or in some other fluids, very intense colds, greater than any natural cold, are artificially produced.<sup>1</sup>

The exertion of chemical affinity is influenced by various circumstances: these, according to Berthollet, are mass, cohesion, insolubility, specific gravity, elasticity, and inflorescence.

1. That *mass* has a considerable share in influencing chemical affinity was first suggested by Berthollet, who states it as a canon, that combinations do not depend altogether on the attraction of affinity, but on the proportions also of the substances brought into action. Thus, if A and B form a compound, and C be a substance which has a stronger affinity for A than B has, it should be able when mixed with a compound, to withdraw A altogether from B, if combination was regulated by affinity only: but this, he affirms, is not the case in fact; for C does not entirely combine with A, but is shared between it and B, according to the force of the affinity, and the bulk of each. This view of the subject affords a reason why, in pharmaceutical compositions, a small quantity of a substance may be added to a compound, without producing any sensible effect, although if added in large quantity, decomposition would directly ensue. It follows, also, from it, 1st, that “the chemical action of one substance on another, must di-

---

<sup>1</sup> See Appendix to Part I. No. I.



minish as it advances to saturation:" and, 2dly, that a decomposing substance "must oppose a stronger resistance to the decomposing agent, in proportion as the decomposition proceeds, from the increase in the relative quantity of one of its ingredients to the other, which is abstracted:" and, lastly, "that in estimating the relative forces of affinity in bodies, the quantities of them must be taken into account, and ought to be equal." Objections of considerable weight have been advanced to the opinions of Berthollet on this subject, by Pfaff, Sir H. Davy, and others: but it is unnecessary to enter into an examination of them at the present moment; and we may only observe, that the theory of Berthollet, however plausible, is not unobjectionable: so that it is, perhaps, better for the student to regard affinity as truly *elective*, and that the constituents of all compound bodies are constant.

2. *Cohesion* has an evident influence in opposing chemical action, and counteracting the exertion of chemical affinity. Thus all aggregates are more slowly acted on by liquids, in which they are soluble, than when their parts are mechanically divided; and this does not happen altogether from the mere circumstance of a larger surface being presented to the fluid; for native oxide of tin, which, in the aggregate, resists completely the action of any acid, becomes soluble when its aggregation is overcome by mechanical operations; and some other substances are similarly affected. Hence *trituration*, *levigation*, and *granulation*, are ranked among pharmaceutical operations, and are of importance "in facilitating chemical action, partly by diminishing aggregation, and partly by increasing the surface on which affinity is exerted." In some instances mechanical division is not sufficient, and recourse must be had to precipitation. Thus liquid potash will not dissolve silica in powder, as it can be obtained by trituration; but when the silica is precipitated from a state of chemical solution, it is readily dissolved in liquid potash.

Owing to the force of cohesion, also, solid bodies seldom act chemically on solids; while fluids readily combine with fluids, and likewise act with energy on solids for which they have an affinity. Fluidity, however, is not indispensable to chemical action; there being many cases in which two solids, in a state of minute mechanical division, act chemically on one another.<sup>1</sup> (See *Section iii.*) When, however, the specific gravities of even two fluids are very materially different, their chemical combination is opposed, to a certain extent, by the force of cohesion of the heavier fluid; and thence, *agitation* is frequently necessary for aiding the operation of affinity.

---

<sup>1</sup> Thence the axiom *Corpora non agunt nisi sint soluta*, which was formerly established in chemistry, is not generally true.



Cohesion has sometimes a considerable influence in determining the proportion of combinations formed in consequence of new affinities. Thus, if its intensity be sufficient to counterbalance the affinity of the fluid in which the integrant particles resulting from a new combination are formed, it will combine these, and produce crystallizations or precipitations, which, withdrawing the substance thus formed in part from the sphere of action, and opposing a resistance to any further exertion of chemical power, will, consequently, determine the proportions of the combination.

3. *Insolubility* must necessarily modify chemical action. If an insoluble compound substance be acted on by any substance tending to combine with one of its principles, this is protected, in some degree, by the insolubility of the compound withdrawing it from the action of the decomposing substance; and if a compound, which is produced in the progress of combination, be insoluble, it will be directly precipitated, and thus fixed in its proportions. In decomposition this is extremely useful; for the insoluble product being immediately separated, cannot oppose the further action of the decomposing substance, which would be the case were it to remain in solution.

4. *Specific gravity* influences considerably the exertion of affinity, particularly if the substance be of little solubility, by withdrawing it from the sphere of action, and hence retarding its combinations; and in many instances this can be but imperfectly counteracted by agitation.

5. Chemical attraction, as far as the aëriform substances are concerned, is opposed by *elasticity*. Thus, when two gases having mutual affinities are mixed together, they very seldom combine, which is ascribed to the distances between the particles of substances existing in the gaseous state: for, as chemical attraction is exerted at insensible distances only, the particles of the two gases, although mingled together, are yet without the sphere of attraction. That this is owing to elasticity, is evident from the circumstance that the vapours which are not elastic more readily combine. Hence, whatever gives density to highly elastic substances, as, for example, mechanical pressure, or cold to a certain degree, must favour their chemical combination.

6. *Efflorescence* may also influence chemical affinity; a fact which was first observed by Scheele, who ascertained, that if in a paste composed of several saline substances decomposition is going on, one of the resulting compounds often rises through the mass, and forms an efflorescence on its surface; and its being thus withdrawn from the sphere of action contributes towards forwarding the decomposition.



7. The influence of *temperature* in modifying chemical action is very considerable. An increased temperature by promoting fusion, and in other respects weakening the attraction of cohesion in solids, favours combination; but opposes it in some cases, as much as it augments elasticity. In both instances its effects are much modified by the degree of its intensity; combinations effected at a lower, being often dissolved at a higher temperature, owing to one or more of the components having its affinity weakened by an increased elasticity. Thus mercury exposed to air for some time at a temperature equal to its boiling point, combines with the oxygen of the air, and is converted into red oxide of mercury; but, if the fire be raised so as to make the retort red-hot, this oxide is again decomposed, and running mercury and oxygen gas are obtained.

From the influence of the above circumstances on chemical combination, the utility of these pharmaceutical and chemical operations, which diminish aggregation, overcome the effect of specific gravity, diminish elasticity, and regulate temperature, such as pulverization, trituration, granulation, agitation, and compression, with the proper management of furnaces, is sufficiently obvious.

In that department of pharmacy, also, which regards extemporaneous compositions, it is of importance to attend to the slowness with which chemical action is in many instances produced; for substances, which have mutual affinities for each other, may give no indication of any change when newly mixed, but yet, after some time, may act, and produce even complete changes. Such compounds, therefore, when they are intended to act *medicinally*, should be exhibited as soon as possible after they are made.

Chemical attraction may be exerted between more than two bodies, so as to bring three or four into one combination; and such compounds are named *ternary*, *quaternary*, &c. according to the number of their components. Several examples of these are to be found among the saline preparations, (*Part iii.*); and almost all the vegetable substances are compounds of three or more principles.

---

The forces with which chemical attraction is exerted are different in different bodies. In cases where this attraction is exerted in a superior degree, by a third body, to that of either of the components of a compound of two bodies, so as to decompose it, and form a new compound, while, at the same time, one of the components of the previous compound is set free, the affinity thus exerted has been termed *single elective attraction*. To represent the relative forces of affinity, tables were first constructed by Geoffroy; and afterwards much




improved and extended by other chemists, particularly Bergman. The remarks of Berthollet may have tended to lessen their value in the opinions of some; but their utility to a certain extent must undoubtedly be acknowledged.<sup>1</sup> When the elective attractions are more complicated, or when two elective affinities are exerted, and two new compounds formed, this is termed *double elective attraction*. In such cases, Mr. Kirwan denominated the attractions which tend to preserve a compound in its original state *quiescent*; while the others, which tend to separate the principles of a compound from each other, he termed *divellent attractions*. As an example of double elective attraction, let it be supposed that two compounds, one consisting of potash and sulphuric acid, or *sulphate of potash*, and the other consisting of muriatic acid and lime, or *muriate of lime*, be mixed together, a double decomposition will take place, and two new compounds, *sulphate of lime*, and *muriate of potash* will be formed. In this case, if the attraction between potash and sulphuric acid be 62, and that between lime and muriatic acid be 20, the sum of the quiescent attractions will be 82; but if the attraction between potash and muriatic acid be 32, and that between sulphuric acid and lime be 54, the sum of the divellent attractions will be 86: which, exceeding the former sum of the quiescent, will operate and produce the above-stated decompositions and resulting compounds.<sup>2</sup>

According to the opinions of Bergman, the relative force of the affinities which produce these effects is capable of being measured, and the changes are altogether to be ascribed to the predominance of the affinities of one set of substances over those of another.

But the changes produced by the predominance of certain affinities over others, are ascribed by Berthollet to those circumstances which influence attraction, and limit combination. If four substances, for example, be presented to each other, two of which have a greater tendency to cohesion than the other two have, so as to form by their union an insoluble compound, instead of one compound being formed by the union of the four, in which the affinities are balanced, this will be

<sup>1</sup> See Appendix to Part I. No. II.

<sup>2</sup> To represent this effect of double elective affinity more clearly, diagrams are used, the idea of which first occurred to Dr. Cullen. Thus the above operations would be represented in the following manner: The inverted triangle in the centre denotes water or that the decomposition was effected in the humid way.





averted by the force of cohesion, and the two which form the insoluble compound will unite, and be separated by precipitation or crystallization, leaving the other two in combination in the fluid which has been the medium of action. "If even these four substances were previously in the reverse binary combinations, on presenting them to each other, the affinities within the sphere of action must be reciprocally exerted; and the same extraneous forces will cause an exchange of principles, or the phenomena which have been ascribed to elective affinities will be produced." To avoid the term elective attraction, Berthollet denominates cases of this kind *complex affinity*. The explanation of single elective attraction, or where three substances are presented to each other, is precisely the same; the union which takes place between two of them being determined by the tendency to cohesion, or the disposition of the combination of two of them to form a compound of little solubility.

Elasticity, likewise, has a considerable influence in determining decompositions where the application of heat is necessary; and, according to Berthollet, the decomposition of a compound body of which one of the ingredients has a great tendency to assume the elastic form, is to be ascribed to the disposition it has to escape from its combination, when aided by the intervention of even a weaker affinity.

In complex affinities the same cause determines the union of substances disposed to assume the elastic form, and separates them as a volatile compound. "If therefore," says he, "it be desired to know the result of the exposure of two salts to the action of heat, it is only necessary to consider which of the two bases and which of the two acids have the greater volatility, if there be a difference: for the more volatile base and acid will escape and enter into combination, and the fixed base and fixed acid will remain behind, and combine with one another."<sup>1</sup> Tables representing the forces of affinity have been constructed; but, as Dr. Henry has justly remarked, "one great obstacle to the construction of such tables, is the difficulty of ascertaining, with precision, the quantities of bodies required for neutralization."<sup>2</sup>

A knowledge of the doctrines of affinity is of the utmost importance in pharmacy; and, as the foregoing sketch presents little more than an outline, I must refer those who would wish to investigate the subject to Thomson's and Murray's Systems of Chemistry, Bergman's Dissertation on Elective Attraction, Berthollet's Researches into the Laws of Chemical

---

<sup>1</sup> *Researches*, p. 3. quoted by Murray, *System of Chemistry*, i. 120.  
*Henry's Elements of Experimental Chemistry*, 7th ed. vol. i. p. 57.



Affinity, Richter's Foundation of Stochiometry, and Sir Humphrey Davy's Elements of Chemical Philosophy.

## II. REPULSION.

REPULSION is that force which separates the particles of bodies from one another, and consequently counteracts or modifies the attractions by which they are combined and preserved together in masses. It is supposed to depend on the operation of one or more of the three following powers; *Caloric, Light, Electricity*.

### a. CALORIC.

The cause of the sensation of heat is denominated *Caloric*. Philosophers are not completely agreed whether it is a property only of bodies, such as a vibration of their particles,<sup>1</sup> or a peculiar substance; but the latter opinion is the one more generally adopted.

Under this opinion caloric is regarded as a very subtile, elastic fluid, which penetrates more or less all bodies, passing readily from one to another; and is every where diffused. Its particles are supposed mutually to repel each other; and bodies into which it enters in any sensible quantity are increased in bulk, and undergo other changes of form, while their density is diminished. It is radiated in the same manner as light, and in this state forms a part of the solar ray.<sup>2</sup> The rays are refrangible, and capable of reflection, and of polarization, like those of light. It has no ascertainable gravity; and neither the addition nor the abstraction of it alters sensibly the weight of bodies.<sup>3</sup> It exists in two different states; in a *free* state, in which it is merely loosely mingled, as it were, with other bodies; and, in a *latent* state, or one of intimate combination.

Regarding it as matter, the sources whence it may be obtained, the laws which regulate its motion and distribution, and its effects, require to be noticed.

### *Sources of Caloric.*

The known sources of caloric are the *sun, combustion, percussion, friction* and *mixture*.

a. The *sun* is an apparent source of caloric; but the direct action of the sun's rays upon bodies seldom produces a temperature exceeding 160°. When these, however, are concentrated by means of a concave mirror, or a lens; or when means are taken to prevent the communicated heat from being carried off by the surrounding bodies, a much higher temperature can be produced. This source of caloric is not resorted to for pharmaceutical purposes.

---

<sup>1</sup> The idea of caloric being motion or vibration originated with Lord Bacon.

<sup>2</sup> *Philosophical Trans.* 1807.

<sup>3</sup> *Ib.* 1799, p. 179.



*b. Combustion* is a source of caloric highly interesting on account of its utility.

When a combustible is heated to a certain degree, it becomes still hotter of itself, and is consumed, emitting rapidly light and caloric, until the whole substance has suffered a change of properties.

The true nature of this process was first explained by Lavoisier, who laid it down as a chemical axiom, that "in every case of combustion, oxygen combines with the burning body." His explanation of combustion depends on two laws: 1st. That when a combustible body is heated to a certain temperature, it immediately begins to attract and combine with the oxygen of the atmospheric air. 2d. This oxygen being in a state of gas and combined with light and caloric, is decomposed during its union with the combustible, and its caloric and light are set free in a sensible form, while the oxygen itself remains combined with the combustible. The truth of this theory is generally supposed to be proved by the following facts: 1. combustion does not go on unless oxygen be present; and it is more brilliant in oxygen gas than in common air: 2. the products of combustion are always heavier than the body consumed; and, 3. this increase of weight is exactly equal to the quantity of oxygen which the air loses. Every combination of oxygen, however, with bodies does not produce the phenomena of combustion. Brugnatelli has endeavoured to explain this by supposing that oxygen combines with bodies in two states: "1. Retaining the greater part of the caloric and light with which it is combined when in the state of gas; and, 2. After having let go all the caloric and light with which it was combined."

The above theory of combustion is, however, liable to some objections; for instance, the emission of caloric and light is not proportional to the quantity of oxygen that combines with the combustible; and the quantity of light that appears depends, altogether, upon the combustible. Under the supposition, therefore, that the caloric is obtained from the oxygen of the substances supporting combustion, while the light is derived from the combustibles, the process has been regarded as a case of "double decomposition; the oxygen and combustible dividing themselves into two portions, which combine in pairs; the one compound is the *product*," or the combustible base united with oxygen; "the other is the *fire*," or the caloric and light "which escapes."<sup>1</sup>

The caloric set free by the burning or combustion of coal, charcoal, carbonated hydrogen gas, oil, wax, and tallow, is applied to the purposes of life, and is of the first importance in the practice of pharmacy: thence, endeavours have been

---

<sup>2</sup> Thomson's Chemistry, 4th edit. i. 607.



made to ascertain the quantity of caloric evolved during the burning of different combustibles, and several experiments have been instituted, by the most able chemists, at different times, for this purpose. The following table exhibits the quantity of caloric evolved by the combustion of different substances, when all the circumstances are equal; the estimate being formed from the quantity of ice melted during the burning of one pound of each of the substances.<sup>1</sup>

Substances burnt, 1 lb.	Oxygen consumed in lbs.	Ice melted in lbs.			
		Lavoisier.	Crawford	Dalton.	Rumford.
Hydrogen - - - -	6	295	480	320	
Carburetted hydrogen	4	—	—	85	
Olefiant gas - - -	3.5	—	—	88	
Carbonic oxide - -	0.58	—	—	25	
Olive oil - - - -	3.5	148	89	104	93.073
Rape oil - - - -	—	—	—	—	124.097
Wax - - - -	3.5	133	97	104	126.242
Tallow - - - -	3.5	—	—	104	111.582
Oil of turpentine -	—	—	—	60	
Alcohol - - - -	—	—	—	58	67.470
Sulphuric æther - -	3.	—	—	62	107.027
Naphtha - - - -	—	—	—	—	97.834
Phosphorus - - -	1.5	100	—	60	
Charcoal - - - -	2.8	96.5	69	40	
Sulphur - - - -	1.36	—	—	20	
Camphor - - - -	—	—	—	70	
Caoutchouc - - -	—	—	—	42	

From this table it appears, that hydrogen gas would form the best fuel, where a high temperature is required.

*c. Percussion*, as far as it applies to solid bodies, is another source of caloric. Smiths, for instance, are in the habit of kindling their fires by means of an iron-rod, which is smartly and quickly hammered until it become red-hot; and sparks are produced by the collision of hard bodies, particularly of flint with steel. This effect appears to arise from condensation or forcing the integrant particles of the bodies closer together, so as to dislodge the latent caloric they contain, and give it out in the form of sensible caloric. The specific gravity of iron is increased .052 by being hammered; and it becomes so hard and brittle that it cannot again be heated by percussion until it has been exposed for some time to a red heat in the forge. By the collision of flint and steel the oxidizement

<sup>1</sup> Thomson's Chemistry, 4th edit. i. 610.



of the steel is also effected, the sparks being small incandescent pieces of oxidized iron.

*d. Friction* is also a source of caloric. It is a well-known fact, that a considerable quantity of free caloric is disengaged when two substances are smartly rubbed together; but the real source of the caloric thus evolved still remains undetermined.

*e. Finally Mixture*, or the chemical union of two substances, in many cases, evolves caloric. This always takes place when the density or specific gravity of the mixture is greater than the mean of the substances mixed; as in the mixture of alcohol and water, or of sulphuric acid and water: and much caloric is, also, evolved when water is thrown upon quicklime, owing to the solidification of the water when it unites with the lime. The caloric which is evolved in these, and other instances of mixture, is the latent caloric, which is the cause of the fluidity of the components; for, as the compound is less fluid, and consequently requires the presence of a smaller quantity of combined caloric, the superabundance which the more fluid components contained must be necessarily set free.

Such are the sources from which caloric is obtained. Combustion is the most important of these; and the knowledge of the laws by which it is regulated, and of the modes of conducting it, is of the first consequence in the practice of pharmacy. (See *Furnaces*.)

*Distribution and Effects of free Caloric.*

From whatever source caloric is obtained, it passes from bodies in which it is accumulated in a free state into bodies which contain less of it, until both are brought to an equilibrium. "The state of a body, with regard to its power of producing the different effects arising from the presence of caloric, is termed its *temperature*:" and this depends on the quantity of sensible caloric contained in it. Thus, when a vessel containing water is placed on the fire, a quantity of caloric passing from the fire into the water, the temperature of the water is raised, or it is made sensibly hotter; and if the water, thus heated, be taken from the fire and placed in a cold place, the sensible caloric accumulated in it passes from it into the air and surrounding bodies, until it become as cold as they are, or until its temperature be lowered to an equilibrium with theirs. The caloric lost by hot bodies during their cooling is carried off; 1st, by the conducting power of the surrounding medium, which "diminishes as the temperature of the hot body approaches to that of the medium;" 2dly, by radiation; 3dly, by currents, or the repeated changes of the portion of medium immediately in contact with the hot body; produced by the change of density occasioned by the caloric they receive from the hot body, enabling them to rise and give place



to a new portion, which, being heated, is also displaced in its turn, and so on till the temperature of the hot body approaches to that of the medium. By accelerating these changes the rate of cooling is proportionably quickened; and thence the cooling effect of winds, and artificial currents of air.

The temperatures of bodies can be comparatively ascertained, to a certain extent, by the sensations they induce. Thus, a body containing much sensible caloric feels warm or hot to the touch, owing to its caloric flowing into the hand; and one containing less than the human body gives the sensation of cold, owing to the abstraction of caloric from the hand. But this mode of judging of temperature is very limited, and depends on the state of the sentient organ, the conducting power of the body which is touched, and many other external circumstances, which prevent confidence from being placed on it as a comparative measure of temperature: and, therefore, instruments have been invented to measure the degrees of temperature of different bodies; the properties of which depend on the *expansion* or increase of bulk which bodies suffer when caloric enters into them.

The thermometer is a most useful and important instrument of this kind. It is a hollow glass tube, having at one end a hollow globe or bulb, the hollow of the tube being perfectly cylindrical, and of a small bore, and the bulb of a proportional size. The bulb, and a portion of the tube, after the air is expelled, by holding the bulb over the flame of a lamp, are filled with mercury or coloured alcohol, by immersing the open end of the tube in either of these fluids, and the tube is then hermetically sealed at the extremity. When the bulb of this instrument is applied to a hot body, the mercury, or the fluid it contains, rises in the tube, and continues to do so until the thermometer acquire the same degree of temperature as the hot body, when the mercury becomes stationary, and the point to which it rises indicates the relative temperature of the hot body. In the same manner, when the bulb is applied to a cold body, the mercury contracts and falls in the tube. The quantity, which thus rises or falls, indicating the proportion of increase or diminution of temperature, is ascertained by a scale, which divides the tube into a number of equal parts or degrees.<sup>1</sup> As the instrument may be occasionally plunged into corrosive fluids, the stem and bulb should project beyond the scale; or the scale may turn up with a hinge.



<sup>1</sup> Thermometers of great accuracy may be purchased. Persons who may wish to construct them for themselves, will find ample instructions for their guidance, in the third chapter of *Henry's Elements of Experimental Chemistry*.



For ordinary purposes, mercury is the fluid best adapted for thermometers, its expansion being most equable; but alcohol is used when great degrees of cold are to be measured.

The thermometer commonly employed in this country is that of Fahrenheit<sup>1</sup>: but as three other thermometers are used on the continent, it may be proper to notice all of them, and point out the circumstances in which their scales differ.

*Fahrenheit*, in forming his thermometer, began the scale at the temperature produced by a mixture of snow and sea-salt acting on each other; and divided the space between this and the point indicated by the temperature of boiling water into 212 equal parts or degrees: 212° being marked as the boiling point. The part of the scale indicated by the freezing of water, he found to be 32 degrees from its beginning; therefore 32° is marked as the freezing point: and the space between it and the boiling point is equal to 180 degrees. The scale may be extended above this point and, also, below the commencement of the scale, the descending degrees being marked inversely with the same numbers as the ascending.

The scale of the thermometer of *Celsius*, which has been used in France since the Revolution, begins at the freezing point of water, which is consequently marked 0, and the space between that and the boiling point is divided into 100 equal degrees; hence it has been named the *Centigrade Thermometer*. Each degree of this scale is  $\frac{4}{5}$ ths more than a degree of Fahrenheit's, or one of the latter is equal to  $\frac{5}{9}$ ths of a degree of the centigrade scale. To find, therefore, the degrees of Fahrenheit's scale, corresponding to those of the centigrade, the given number of the latter must be multiplied by 9, and divided by 5, adding 32 to the quotient. The sum expresses the degree on the scale of Fahrenheit.

*Reaumur's* thermometer, which is still used in Italy and Spain, also commences at the freezing point, which is marked 0; and between this and the boiling point it is divided into 80 degrees. Each degree is, therefore,  $\frac{4}{9}$ ths more than one of Fahrenheit's; and to reduce the scale of Reaumur to that of Fahrenheit, the given number of degrees of the former must be multiplied by 9, and divided by 4, adding 32 to the quotient.<sup>2</sup>

In *De Lisle's* thermometer, which is used only in Russia, the space between the boiling and freezing points is divided into 150 degrees, the gradation beginning at the boiling point, which is marked 0; and increasing inversely to the freezing point, which is marked 150°. It is seldom mentioned by authors.

These instruments are well adapted for determining the variations of temperature which bodies undergo: but a certain

<sup>1</sup> Fahrenheit was an artist of Amsterdam.

<sup>2</sup> See Note on the next page.



degree of fallacy attends the observations made with them, chiefly owing to the expansion of mercury increasing with the temperature. Thus, the medium degree of heat between the freezing and boiling points, although marked on the scale 122°, yet is actually 118·8° 1' only ; the temperature which is equal to raise the mercury in the tube 86 degrees in the first instance being sufficient, by increased expansion, to raise it 94 in the second.

*Expansion*, or increase of bulk, is the most general effect of caloric, and, with very few exceptions, may be regarded as the general law of its operation.

When caloric flows into a body, it separates its integrant particles from one another, and hence augments its volume. This change is smallest in solids, more considerable in liquids, and most considerable in gaseous bodies ; or the expansibility is greater in the reverse ratio of the force of aggregation. Thus, the expansion of air is 8 times greater than that of water ; and the expansion of this 45 times greater than that of iron.

TABLE shewing the degrees of Reaumur's and Fahrenheit's thermometers corresponding with those of the centigrade thermometer.

Cent.	Reau.	Fahr.	Cent.	Reau.	Fahr.	Cent.	Reau.	Fahr.	Cent.	Reau.	Fahr.
100	80·	212·	68	54·4	154·4	36	28·8	96·8	5	4·	41·
99	79·2	210·2	67	53·6	152·6	35	28·	95·	4	3·2	39·2
98	78·4	208·4	66	52·8	150·8	34	27·2	93·2	3	2·4	37·4
97	77·6	206·6	65	52·	149·	33	26·4	91·4	2	1·6	35·6
96	76·8	204·8	64	51·2	147·2	32	25·6	89·6	1	0·8	33·8
95	76·	203·	63	50·4	145·4	31	24·8	87·8	0	0·	32·
94	75·2	201·2	62	49·6	143·6	30	24·	86·	1	0·8	30·2
93	74·4	199·4	61	48·8	141·8	29	23·2	84·2	2	1·6	28·4
92	73·6	197·6	60	48·	140·	28	22·4	82·4	3	2·4	26·6
91	72·8	195·8	59	47·2	138·2	27	21·6	80·6	4	3·2	24·8
90	72·	194·	58	46·4	136·4	26	20·8	78·8	5	4·	23·
89	71·2	192·2	57	45·6	134·6	25	20·	77·	6	4·8	21·2
88	70·4	190·4	56	44·8	132·8	24	19·2	75·2	7	5·6	19·4
87	69·6	188·6	55	44·	131·	23	18·4	73·4	8	6·4	17·6
86	68·8	186·8	54	43·2	129·2	22	17·6	71·6	9	7·2	15·8
85	68·	185·	53	42·4	127·4	21	16·8	69·8	10	8·	14·
84	67·2	183·2	52	41·6	125·6	20	16·	68·	11	8·8	12·2
83	66·4	181·4	51	40·8	123·8	19	15·2	66·2	12	9·6	10·4
82	65·6	179·6	50	40·	122·	18	14·4	64·4	13	10·4	8·6
81	64·8	177·8	49	39·2	120·2	17	13·6	62·6	14	11·2	6·8
80	64·	176·	48	38·4	118·4	16	12·8	60·8	15	12·	5·
79	63·2	174·2	47	37·6	116·6	15	12·	59·	16	12·8	3·2
78	62·4	172·4	46	36·8	114·8	14	11·2	57·2	17	13·6	1·4
77	61·6	170·6	45	36·	113·	13	10·4	55·4	18	14·4	0·4
76	60·8	168·8	44	35·2	111·2	12	9·6	53·6	19	15·2	2·2
75	60·	167·	43	34·4	109·4	11	8·8	51·8	20	16·	4·
74	59·2	165·2	42	33·6	107·6	10	8·	50·	21	16·8	5·8
73	58·4	163·4	41	32·8	105·8	9	7·2	48·2	22	17·6	7·6
72	57·6	161·6	40	32·	104·	8	6·4	46·4	23	18·4	9·4
71	56·8	159·8	39	31·2	102·2	7	5·6	44·6	24	19·2	11·2
70	56·	158·	38	30·4	100·4	6	4·8	42·8	25	20·	13·
69	55·2	156·2	37	29·6	98·6						



The expansion of solid bodies is, in general, so very inconsiderable as not to be easily ascertained by measurement ; but, as far as it can be known, it is nearly equable. The degree of expansion, however, is not the same in all solids ; thus, for example, the metals expand in the following order, commencing with the least expansible ; platina, antimony, iron, bismuth, copper, tin, lead, zinc. Argil is an exception to the law of expansion in solids ; for the bulk of pure clay diminishes, when heated, in the ratio of the intensity of the heat to which it is exposed.<sup>1</sup> The cause of this anomaly has not been discovered. The expansion of liquids is more evident than that of solids, but not at all uniform ; the differences apparently depending on the fixity or volatility of the components of the liquids ; those expanding the most the boiling point of which is lowest ; and

*TABLE exhibiting the degrees of the centigrade and Fahrenheit's thermometers corresponding to those of Reaumur's thermometer.*

Reau.	Cent.	Fahr.	Reau.	Cent.	Fahr.	Reau.	Cent.	Fahr.
80	100°	212°	46	57°5	135°5	12	15°	59°
79	98°75	209°75	45	56°25	133°25	11	13°75	56°75
78	97°5	207°5	44	55°	131°	10	12°5	54°5
77	96°25	205°25	43	53°75	128°75	9	11°25	52°25
76	95°	203°	42	52°5	126°5	8	10°	50°
75	93°75	200°75	41	51°25	124°25	7	8°75	47°75
74	92°5	198°5	40	50°	122°	6	7°5	45°5
73	91°25	196°25	39	48°75	119°75	5	6°25	43°25
72	90°	194°	38	47°5	117°5	4	5°	41°
71	88°75	191°75	37	46°25	115°25	3	3°75	38°75
70	87°5	189°5	36	45°	113°	2	2°5	36°5
69	86°25	187°25	35	43°75	110°75	1	1°25	34°25
68	85°	185°	34	42°5	108°5	0	0°	32°
67	83°75	182°75	33	41°25	106°25	1	1°25	29°75
66	82°5	180°5	32	40°	104°	2	2°5	27°5
65	81°25	178°25	31	38°75	101°75	3	3°75	25°25
64	80°	176°	30	37°5	99°5	4	5°5	23°
63	78°75	173°75	29	36°25	97°25	5	6°25	20°75
62	77°5	171°5	28	35°	95°	6	7°5	18°5
61	76°25	169°25	27	33°75	92°75	7	8°75	16°25
60	75°	167°	26	32°5	90°5	8	10°	14°
59	73°75	164°75	25	31°25	88°25	9	11°25	11°75
58	72°5	162°5	24	30°	86°	10	12°5	9°5
57	71°25	160°25	23	28°75	83°75	11	13°75	7°25
56	70°	158°	22	27°5	81°5	12	15°	5°
55	68°75	155°75	21	26°25	79°25	13	16°25	2°75
54	67°5	153°5	20	25°	77°	14	17°5	0°5
53	66°25	151°25	19	23°75	74°75	15	18°75	1°75
52	65°	149°	18	22°5	72°5	16	20°	4°
51	63°75	146°75	17	21°25	70°25	17	21°25	6°25
50	62°5	144°5	16	20°	68°	18	22°5	8°5
49	61°25	142°25	15	18°75	65°75	19	23°75	10°75
48	60°	140°	14	17°5	63°5	20	25°	13°
47	58°75	137°75	13	16°25	61°25			

<sup>1</sup> For measuring higher temperatures than the thermometer can be subjected to, instruments named *pyrometers* have been employed ; the best of which is that in-



which, consequently, most readily assume the gaseous form. The degree of their expansion, also, increases, with the augmentation of the temperature; or, the nearer a liquid approaches to the boiling point, the greater is the expansion produced by a degree of caloric; and, the further it is from this point, the more equable is the expansion. Liquids, in the same manner as solids, suffer a difference of expansion from a given change of temperature. The following table, by Mr. Dalton, shews the expansion of the more common liquids, from  $32^{\circ}$  to  $212^{\circ}$  of Fahrenheit, the volume at  $32^{\circ}$  being denoted by 1.

Mercury.	Water.	Water saturated with salt.	Sulphuric acid, sp. gr. 1.85.	Muriatic acid, sp. gr. 1.137.	Oil of Turpentine.	Ether.	Fixed oils.	Alcohol.	Nitric acid, sp. gr. 1.40.
.0200 =	.0466 = $\frac{1}{25}$	.0500 = $\frac{1}{20}$	.0600 = $\frac{1}{17}$	.0600 = $\frac{1}{17}$	.0700 = $\frac{1}{14}$	.0700 = $\frac{1}{14}$	.0800 = $\frac{1}{12}$	.0110 = $\frac{1}{9}$	.0110 = $\frac{1}{9}$

To the general law of the expansion of liquids by heat, water furnishes an exception. Thus, from the lowest temperature at which water can remain liquid, to  $40^{\circ}$ , heat diminishes the bulk of water, instead of expanding it; but, above  $40^{\circ}$  to  $212^{\circ}$ , expands it.

All gaseous bodies suffer the same expansion by the same additions of caloric, supposing the circumstances to be equal. Their expansion is almost perfectly equable, or the same augmentation takes place by the same addition of caloric at every degree of temperature between the freezing and the boiling

---

vented by Mr. Wedgwood. It depends on the degrees of contraction which pure argil suffers when exposed to high temperatures; and for this purpose small cylinders of pure clay are made in a mould, flattened on one side, and fitted exactly to the wider end of a gauge, consisting of two straight pieces of brass, 24 inches long, fixed on a brass plate so as to converge, and divided into inches and tenths. The length to which the pyrometrical pieces can be slid in the converging groove, indicates the heat to which they have been previously exposed; and, as they do not expand again when cold, no fallacy can result from the action of heat on the gauge. Each degree of this scale is equal to  $130^{\circ}$  of Fahrenheit; and the 0, or commencement of it, corresponds with  $1077\frac{1}{2}^{\circ}$  of Fahrenheit's scale. The highest temperature that has been measured by it is  $160^{\circ}$  or  $21.877^{\circ}$  of Fahrenheit, which is  $30^{\circ}$  above the point at which cast iron melts. But, as much higher temperatures than this must exist, so also, there are temperatures much lower than can be measured by any thermometer.



point of Fahrenheit's thermometer. By the experiments of Gay Lussac, 100 parts of atmospheric air, heated from  $32^{\circ}$  to  $212^{\circ}$  expand 137.5 parts, or  $\frac{1}{480}$ th for every degree of the thermometer: and the other gases, the steam of water, and the vapour of ether, undergo the same expansions by the same augmentations of temperature. The cause of the equable expansion of gaseous bodies appears to be the absence of cohesion; so that, at a low temperature, there is no more resistance made to the expansive power of the caloric thrown into the gas, than at a high temperature.

But, besides the change in bulk produced by the introduction of caloric into substances in different quantities, they are changed in state, assuming the *fluid form* and that of *vapour*; or, they are *ignited*.

*Fluidity* is an effect of caloric, arising from the repulsive force of the caloric which enters into any substance fitted to take on the fluid form, separating the particles from one another to such a distance as to render them easily moveable on one another in every direction. All solids, with a very few exceptions, are susceptible of the fluid form, when exposed to a sufficient degree of heat; and all liquids, with the exception of alcohol and ether, become solid when exposed to very low temperatures. The particular temperatures necessary for the production of these changes, however, are exceedingly various, but for the same bodies they are always the same.<sup>1</sup> In some cases, the change is sudden, or the body instantly passes from the solid to the liquid state; in other cases, it passes through several degrees of softness before it be perfectly liquefied; the conversion of ice into water is an example of the first; the melting of glass, of wax, and other unctuous matters, are instances of the second. There are some bodies, nevertheless, which cannot be melted or *fused*, owing to their suffering chemical decomposition at a lower temperature than is required for their *fusion* under the ordinary pressure of the atmosphere: — a piece of wood, for instance, cannot be melted by the application of any degree of heat.

Although the melting point, in most cases, is always the same in the same bodies, yet circumstances may vary it; and the admixture of other substances may alter it very consider-

<sup>1</sup> Table, showing the degree of temperature, according to Fahrenheit's thermometer, at which several solid bodies melt.

Lead - 612 <sup>o</sup>	Copper - - - 4587 <sup>o</sup> , Fahr. 28, Wedg.	Spermaceti 112 <sup>o</sup>
Bismuth - 476	Silver - - - 4717	Phosphorus 100
Tin - 442	Iron - - - 21637	Tallow - 92
Zinc - 703	Sulphur - - - 218	Oil of anise 50
Antimony 809	Bees-wax - - - 142	Camphor - 303
Mercury - 39	Lard - - - 97	Ice - - - 32



ably. Thus, the melting point of ice, or, what is the same thing, the freezing point of water, is  $32^{\circ}$ ; but by exposing water slowly to the action of freezing mixtures, it may be cooled down to  $22^{\circ}$  before it freezes. The addition of salts renders this point still lower, as may be seen by the following table.<sup>1</sup>

Names of Salts.	Proportion by weight dissolved in 100 parts of water.	Freezing point.
Common salt.....	25.	4.
Sal ammoniac.....	20.	8.
Rochelle salt.....	50.	21.
Sulphate of magnesia...	41.6	25.5
Nitre.....	12.5	26.
Sulphate of iron.....	41.6	28.
———— zinc.....	33.3	28.6

When solids pass to the liquid state they receive an additional quantity of caloric, which combines with them, but does not sensibly elevate their temperature; and this *caloric of fluidity*, or *latent heat*, as it has been named, is again given out in a sensible form, when the body returns to a solid state. If water, for example, be exposed to a great degree of cold, and kept free from agitation, it may be cooled several degrees below the freezing point, and yet remain fluid; but if it be then agitated, it instantly congeals, and at the moment its temperature rises to  $32^{\circ}$ . All fluids, therefore, are combinations of solids and certain doses of caloric. Thus, if snow at  $32^{\circ}$  be mixed with an equal weight of water at  $172^{\circ}$ , the snow instantly melts, but the temperature of the mixture is only  $32^{\circ}$ ; so that  $140^{\circ}$  of caloric have disappeared, or rather have entered the ice, and as the caloric is no longer sensible to the thermometer, it is justly said to have become latent: thence the quantity of caloric necessary to give fluidity to ice is  $140^{\circ}$ . These facts were first ascertained by Dr. Black in 1762; and fluidity in general has been proved to depend on a similar cause. Softness, plasticity, malleability, and ductility, probably depend also upon the repulsive force of the latent heat which combines with bodies.

*Vapour*, which is another effect of caloric, is that state into which all fluids and some solids pass when their temperature is raised to a certain point, or caloric is thrown into them in sufficient quantity to separate their integrant particles to distances beyond the sphere of the attraction of cohesion. The fluid passes to the state of vapour, becoming invisible and elastic, and possessing the other mechanical properties of air.

<sup>1</sup> *Phil. Trans.* 1788, 27, quoted by Dr. Thompson.—*Syst. Chemistry*, 4th edit. i. 520.



Evaporation, however, is also spontaneously produced, partly by the agency of caloric alone, partly by the solvent power of atmospheric air, forming a solution of the body in the aërial fluid. By *spontaneous evaporation*, the fluid is gradually converted into the aëriform state at every temperature. Water, alcohol, ether, and volatile oils are susceptible of spontaneous evaporation, so that a portion of any of them exposed to the air, in a flat vessel, soon altogether disappears: “but sulphuric acid and the fixed oils never assume the form of vapour till they are raised to a certain temperature.”

All fluids have a fixed point of temperature at which their *vaporization*, or conversion into vapour, commences, which is denominated their *boiling point*; and beyond this point fluids cannot be heated, if freely exposed to the air so as to allow the vapour to escape as it forms. Thus, water at  $212^{\circ}$  boils, and is progressively converted into steam at the bottom of the vessel, which, rising in bubbles through the water, produces the ebullition that characterizes boiling; but although the fire be raised ever so much, yet the temperature of the water never exceeds  $212^{\circ}$ , the vapour carrying off every additional increment of heat in a latent form. The *boiling point*, as this degree of temperature is termed, varies in different bodies; and in the same body also, if it be placed under different circumstances, particularly with regard to pressure. Thus the boiling point of ether is  $96^{\circ}$ , of alcohol  $173^{\circ}$ , of pure water  $212^{\circ}$ , of water saturated with common salt  $225^{\circ}$ , of mercury  $656^{\circ}$ , and so on. In a vacuum all liquids boil at a temperature  $124^{\circ}$  lower than in the open air; and in Papin's digester, in which water can be heated under a great pressure, the temperature may be raised to  $300^{\circ}$  without ebullition. Owing to this circumstance, highly volatile substances, as Ammonia and Ether, cannot be easily manufactured in elevated situations.

The elasticity of the vapour of liquids boiled in the open air is equal to that of the circumambient atmosphere; but under such a pressure that the temperature of the vapour may be much augmented, the elasticity increases with the temperature. At low temperatures, on the contrary, vapours lose their elasticity, are condensed, and return to their fluid state. The conversion, therefore, of liquids into elastic fluids depends on the same cause as the conversion of solids into fluids; namely, “to the combination of a certain dose of caloric with the liquid, without any increase of temperature.”<sup>1</sup> The vapour carries off all the caloric which enters a fluid after it arrives at its boiling point; and retains it in a latent form: for the vapour is not sensibly hotter than the boiling liquid: thus, steam, the temperature of which is indicated by the thermometer to

---

<sup>1</sup> The important discovery of the causes which produce the changes of bodies from the solid to the liquid and aëriform state was made by Dr. Black in 1760.



be  $212^{\circ}$ , is water combined with  $940^{\circ}$  of caloric, which remain latent as long as the temperature of the steam is maintained at  $212^{\circ}$ , but is again given out when a lower temperature changes that vapour to the state of a liquid.

*Gases* resemble vapours in their constitution, but differ from them in the greater reduction of temperature which is required for their condensation, some of them not being reducible by ordinary pressure, or by any known reduced temperature, to the fluid or solid state. Are they compounds of solid or of liquid substances and caloric? Ammoniacal gas condenses into a fluid at  $45^{\circ}$ ; and many of the other gases have been condensed into fluids by Mr. Faraday; but the question may be regarded as still *sub judice*.

*Ignition* is another effect of caloric, but differing altogether from expansion, fluidity, and evaporation, which may in some measure be regarded as different degrees of one general effect. It implies an emission of light from bodies which are much heated, or combined with a large portion of caloric, without their suffering any change of composition. It is totally independent of the presence of air, and is a simple effect of caloric. Aëriiform substances are not susceptible of ignition.

The degree of temperature at which all bodies capable of ignition begin to be ignited, or become red-hot, is nearly the same, — about the 800th degree of Fahrenheit; and by raising the temperature the illumination increases, until a perfectly white light is produced, which is the highest point of ignition. Ignition is supposed to arise from the extrication of the light, which is regarded as a constituent of the ignited body, by the repulsive agency of the additional caloric: but this explanation of the phenomenon is liable to some objections, and the real cause remains still undetermined.

As a pharmaceutical agent, *Caloric* is of the first importance. In the majority of cases it produces decomposition; but in some it favours combination. The decomposition most easily effected by it, is the separation of the more volatile from the more fixed ingredients of compounds. Thus, in the process of distillation (see *Operations*), when weak spirits are heated, the alcohol separates from the water, owing to its superior volatility, and, by condensation in a different vessel, is obtained as a distinct substance. Almost all compounds into which oxygen has entered without having occasioned combustion, as nitric acid, and some metallic oxides, suffer likewise decomposition by caloric. All compound bodies containing combustibles are also decomposable by it; as are also compounds consisting of two or more combustible ingredients, in combination with oxygen, as almost all animal and vegetable matters. On the contrary, the compounds which are little or not at all affected by caloric, as far as regards their composition, are those which



have been formed by combustion; such as water, phosphoric acid, and carbonic acid.<sup>1</sup> The proper application of Caloric for the purpose of obtaining new combinations by lessening the force of aggregation, and thus favouring the attraction of affinity; or for producing decompositions by weakening or destroying altogether the force of these attractions so as to obtain the principles of bodies in a distinct state, constitutes the most important feature of operative pharmacy. (See *Operations*.)

## 2. LIGHT.

*Light* is a substance consisting of very subtile particles, which are constantly emanating in straight lines from luminous bodies. Some philosophers, however, following Hooke and Huygens, believe, that light is merely the state of undulation of a highly rare, elastic substance, which it is supposed fills the whole of the universe: but this hypothesis is inconsistent with the fact, that a ray of light which has suffered ordinary refraction in passing through a crystal in one direction, should suffer ordinary refraction in passing through it in another direction; whilst this fact perfectly accords with the idea that light consists of particles endowed with rectilineal motion. The size of the particles of light is too minute to be appreciated; but their velocity is estimated to be at the rate of 200,000 miles in a second. They appear to repel each other like the particles of caloric.

A ray of light falling obliquely upon a polished surface is *reflected* from it at an angle equal to the angle of its incidence. But when a ray of light falls at an angle of  $35^{\circ} 25'$  on a polished plate of glass; and, in being reflected from it, falls upon another plate of glass, so placed that its angle of incidence is also  $35^{\circ} 25'$ ; the second plate may be turned round its axis without varying the angle which it makes with the ray that falls upon it. If the two planes of reflection be parallel to each other, the ray of light is reflected in the same manner from both plates of glass; but if the second plate be turned round a quadrant of a circle, so as to make the plane of reflection perpendicular, the whole ray will pass through it, and none of it be reflected; yet, if this plate be turned round another quadrant of a circle, so as to make the reflecting planes again parallel, the ray will be now reflected by it as at first. The light can penetrate through the glass only when the reflecting planes are perpendicular, but is reflected when they are parallel. This property of light has been termed by Malus, by whom it was first discovered, its *polarization*.

---

<sup>1</sup> Thomson's *Chemistry*, 4th edit. i. 546.



When a ray of light, moving in a straight line, passes within a certain distance of a body parallel to its direction, it bends towards the body, or is *inflected*; but when the body parallel to its course is at a greater distance, the ray is bent from it, or *deflected*. When it passes obliquely from one medium to another of a different density, it is bent a little from the line of its former direction, and assumes a new one, or is *refracted*. In passing into a denser medium it is refracted towards the perpendicular; but is refracted from the perpendicular when passing into a rarer medium. The refraction is proportional to the density of the medium; but, in that of a combustible, the refraction is greater than the ratio of its density; and when a liquid is converted into vapour, its refractive power diminishes at a greater ratio than its density diminishes.<sup>1</sup>

Every ray of light is resolvable into seven other distinct rays, each possessing a different degree of refrangibility; and consequently divisible from each other by the prism. The ultimate or component rays are distinguishable by the impression of colours they excite on the eye; and are arranged in the following order, *red, orange, yellow, green, blue, indigo, violet*. The red is the least refrangible, reflexible, and inflexible; the violet the most; and that of the others follow the order in which they are placed. The colour of bodies depends on their transmitting or reflecting those rays only which excite the impressions of their colour. The reflection of the whole prismatic rays constitutes white; the absorption or suffocation of all, or the greater part, of these occasions black, which is the total absence of light.

The illuminating power of the rays of light differs. Those towards the middle of the prismatic spectrum, as above arranged, possess the greatest illuminating power; but this diminishes as the rays approach towards the extremities.

Light enters into combination with bodies; and, in some cases, is again extricated without any change being produced; as in pyrophori, or substances which absorb light, and emit it again when carried into a dark place. In some cases, however, the absorption of light by bodies occasions very sensible changes in them: the colour of plants, for example, their taste and odour, and the quantity of combustible matter they contain, depend on light: for a plant reared in the dark is nearly colourless, insipid, inodorous, and contains a very small proportion of combustible matter.

The natural sources of light are the sun and fixed stars; but it is, also, artificially produced by combustion, chemical combination, heat, and percussion. The sun's rays, the greatest source of light, have been found to be composed of

<sup>1</sup> See the experiments of Arago and Petit, in the *Ann. de Chem. et Phys.* tom. i. 1.



three different species of rays : 1. rays which produce light and colour ; 2. rays of mere heat ; and 3. rays which produce neither light nor colour, nor affect the thermometer ; but which have the power of deoxidizing. Thus constituted, it produces very important chemical effects.

Light partially deoxidizes metallic oxides and salts. Thus it blackens muriate of silver ; and as this takes place when the salt is placed beyond the violet ray, or out of the prismatic spectrum, the effect is apparently to be attributed to the action of the third species of rays. It also reduces the nitro-muriatic solution of gold, when it is placed in contact with charcoal, or any other vegetable, or any animal matter ; and the red oxides of mercury and of lead become much paler when exposed to the sun. The rays that produce these effects are the least refrangible. Dr. Wollaston, however, has pointed out one exception to this effect of these rays, in guaiacum, which becomes green or oxidized in the least refrangible rays ; and is again changed to yellow, or deoxidized in the most refrangible.

Light has a powerful tendency to decompose nitric acid, which it renders red and fuming, even when it is contained in vessels accurately closed. Almost all the vegetable and animal colouring matters have their brilliancy and colour much impaired by long exposure to the sun's rays : and the colour and the properties of vegetable powders kept in clear glass bottles are, also, affected by them. Light even seems to have a strong influence on the process of crystallization ; for, if light be only partially admitted to a crystallizing solution, the crystals will be larger and more numerous on the enlightened side ; and often the whole mass will radiate towards this point. Chaptal<sup>1</sup> found that by using a solution of a metallic salt, and shading the greater part of the vessel, capillary crystals shoot up the uncovered side, and the extent of the exposed part is distinctly marked by the limit of the crystallization.

Such are some of the properties of light, the chemical effects of the operation of which seem to be perfectly independent of its heating power ; and there is even reason to believe, that the greatest chemical changes are produced by the invisible rays : for Ritter<sup>2</sup> affirms that, by transmitting the coloured rays through different prisms, he has separated them from the invisible or chemical rays, and obtained a coloured spectrum devoid of any chemical power.

### 3. ELECTRICITY and GALVANISM.

The phenomena of electricity depend on a very subtile fluid, which is a powerful chemical agent, capable of producing immediate decompositions and new combinations. Galvanism appears to be essentially the same as electricity, differing,

<sup>1</sup> *Journal de Physique*, xxxiii. 297.

<sup>2</sup> *Nicholson's Journ.* viii. 216.



however, in some degree in its effects and the mode of its production. Both are to be regarded as repulsive powers.

*a. Electricity* may be communicated to all substances: by some it is transmitted without any perceivable obstruction, but by others with much difficulty: thence bodies, in their relation to electricity, are distinguished into two classes, *conductors* and *non-conductors*: and as it can be accumulated in the latter by friction and other means, these are also denominated *electrics*; while the former are named *non-electrics*, to indicate their incapability of being excited.

Metals, plumbago, charcoal, and most liquids are conductors: all other substances are non-conductors; although many of these, when made very hot, become conductors. All electrics or non-conductors when rubbed, as, for instance, a glass rod with a piece of woollen cloth, attract light substances; and, when a conductor is approached to them, exhibit an appearance of light, attended with a peculiar sound and smell. Some electrics can be excited by simple heating or cooling. It is necessary, however, for obtaining any considerable excitation, that the rubber have some communication with the earth; from which it appears, that the great source of electricity is in the earth; and, that excitation consists in the mere transferring of the electrical fluid from one substance to another. By rubbing electrics on one another, the distribution of the electric fluid they contain is altered; and on separating them, more than the natural quantity remains with the one, and less with the other. The one is then said to be electrified *plus*, and the other *minus*, or *positively* and *negatively*. When two bodies are both electrified positively, or both negatively, they repel one another; but if one of them be electrified positively and the other negatively, they attract one another. Instead of this distribution of the same fluid, the existence of two fluids has been assumed, each of which repels its own particles, but attracts those of the other; and this assumption is more favourable for the explanation of the chemical agency of this fluid.


The chemical effects of electricity seem to depend, chiefly, on its power of producing a sudden high temperature; and this appears to be proportioned to the resistance opposed to its transmission. It often favours chemical combinations; as for instance, that of oxygen with the metals, and promotes the instantaneous chemical union of gaseous bodies. It, also, effects chemical decompositions; as for example, those of water, ammonia, alcohol, and metallic oxides. But for neither purpose is it employed as a pharmaceutical agent.

*b. Galvanism* may be regarded as a modification of electricity, in which the fluid is evolved during certain chemical actions. It is transmitted through those substances which are conductors of common electricity, and with the same degrees of facility and rapidity. The metals, Charcoal to a certain ex-



tent, Plumbago, Water, the mineral acids, and saline solutions, are perfect conductors; Alcohol, Ether, Sulphur, oils, resins, and metallic oxides, are imperfect conductors: but Glass, dried and baked woods, the dry animal cuticle, and dry gases, are non-conductors of the galvanic fluid.

Galvanism is generally excited by arranging two different metals, as, for instance, copper and zinc, and a fluid, as muriatic acid, diluted with twenty or thirty parts of water, in such a manner that the metals touch each other in one part, and have the fluid interposed between them in another.<sup>1</sup> The metals soldered together in pairs, are placed transversely in the grooves of a well-seasoned wooden trough, or one of earthenware, and fixed in with a cement of resin and wax, to prevent any liquid from passing through; after which, the diluted acid is poured between the pairs, so that it touches the zinc of one pair, and the copper of another, alternately, the copper side of every double plate looking towards one extremity of the trough throughout the arrangement, and the zinc side to the other extremity. This apparatus is named the Galvanic battery; and the distances between the pairs of soldered plates should be from one-fourth to three-fourths of an inch each, according to the width of the trough. The intensity of action of this apparatus, as far as the production of heat is concerned, seems to depend on the size of the plates, or extent of their surfaces, but, for producing chemical decomposition, on the number of plates. The more improved apparatus now used, is a trough of earthenware, *a*, divided in its length by numerous partitions of the same material. Into each of the cells thus formed, and filled with the diluted acid, a plate of zinc and of copper, *b*, are placed, but not so as to touch each other; and a communication is made by a metallic arc between the zinc in one cell, and the copper in the next. The whole of these plates are attached to a wooden bar, so that they can be removed at the pleasure of the operator, and supported on hooks fixed in the cross bars of the props *e*, *f*, attached to the side of the trough.



<sup>1</sup> The pile of Volta, which is not now employed, consists of plates of zinc and silver, and pieces of moistened woollen cloth, piled in the order of zinc, silver, cloth; zinc, silver, cloth, for twenty or more repetitions.



As a chemical agent, Galvanism is the most powerful of all the repulsive forces, and is capable of producing decompositions which could not otherwise be effected. By its means the chemical constitution of the alkalies and the earths has been established, and their bases discovered to be substances hitherto unknown, which have been added to the list of metals.

By placing any body, one, for instance, composed of oxygen and an inflammable substance, in connexion with the metallic wires proceeding from each end of a galvanic battery, the oxygen is attracted by the wire which is in the positive state, and repelled by that which is in the negative; while, at the same time, the inflammable is attracted by the negative wire, and repelled by the positive. Hence the components are separated, and obtained in a distinct state. In the decompositions, thus effected, substances can be conveyed to a distance, and even through interposed ponderable matter, by the galvanic influence; a result which, however singular, is well ascertained.

Galvanism, like Electricity, acts as a stimulus to the living system. Its effects on the animal body are a sensation of light to the eye; a sensation of acidity on the tongue, and of pain on the muscles; and the excitement of strong muscular action.

It would be entering too much into hypothetical discussion to attempt an explanation of the phenomena of Galvanism. It has not yet been employed as a pharmaceutical agent, except for detecting the presence of oxymuriate, bichloride of mercury, when exhibited as a poison. (See PART 3.)

---

On the forces of attraction and of repulsion every chemical, and consequently every pharmaceutical effect, more or less depends. A knowledge, therefore, of the laws which regulate these powers is of the greatest importance, and forms the basis of all chemical science.

---

## SECTION II.<sup>1</sup>

EVERY substance, whether it be regarded generally as forming a part of the mass of this globe, or particularly as an object of science, may be arranged in one or other of the three following classes; *Solids*, *Fluids*, and *Gaseous Bodies*. We shall now examine each of these classes separately, and endeavour to describe the constitutions and combinations of the substances composing them, which are objects of pharmacy.

### I. SOLIDS.

SOLID bodies are masses of homogeneous particles combined

---

<sup>1</sup> In drawing up this section, I have borrowed very freely from the third book of Thomson's *System of Chemistry*.



and held together by the attraction of aggregation or cohesion. The arrangement of the particles with regard to one another is, often, such as to produce regular figures, in which case the solids are said to be crystallized. Cohesion and crystallization have been already considered.

## CONSTITUTION OF SOLIDS.

*Arrangement of the principal Solids according to their Composition.*

### I. SIMPLE, or UNDECOMPOUNDED.

<i>Sulphur.</i>	<i>Carbon.</i>	<i>Metals.</i>
<i>Phosphorus.</i>	<i>Boron.</i>	

### II. COMPOUNDS.

*Sulphurets of Metals.*

————— *fixed Alkalies.*

————— *Earths.*

*Oxide of Phosphorus.*

*Phosphurets of Carbon.*

————— *Metals.*

————— *Earths.*

*Charcoal.*

*Metallic Oxides.*

————— *with Alkalies.*

*Carburets of Iron.*

*Alloys.*

*Earths.*

————— *with Earths.*

————— *metallic Oxides.*

————— *fixed Alkalies.*

*Solid Acids.*

*Salts and Hydrosulphurets.*

*Bitumens, solid oils.*

*Soaps.*

*Most vegetable Substances.*

*Many animal Substances.*

### SIMPLE SOLIDS.

- a. **SULPHUR.** Although the experiments of Sir H. Davy<sup>1</sup> have unequivocally proved that hydrogen is always present in sulphur as we obtain it; yet, they afford no positive conclusion that it is an essential ingredient of sulphur. I have therefore followed Dr. Thomson in regarding it as a simple solid. For its properties, see PART II.

<sup>1</sup> *Elements of Chem. Phil.* p. 283.



- b.* **PHOSPHORUS** is semitransparent, of a yellowish red or amber colour, and a waxy consistence; but, when perfectly pure, it is colourless and transparent. Its specific gravity is 1.770. It is nearly insoluble in water, and is so soft that it may be cut with a knife. It is brittle at a temperature under 32°. Above that point it softens, and under 90° is very ductile. It melts at 90°; in close vessels is volatilized at 219°; and boils at 554°. When heated in the air to 148° it inflames, and emits a white smoke, which is luminous in the dark, has an alliaceous odour, and condenses to an acid. It is obtained from urine, bones, and other animal matters. From Sir H. Davy's experiments, there appear to be circumstances which favour the idea that it is a compound; but it is, still, generally regarded as an undecomposed body.<sup>1</sup>
- c.* **CARBON.** The diamond is this substance in a state of purity; and although charcoal, in its ordinary state, almost invariably contains either hydrogen or water, yet, from the experiments of Messrs. Allen and Pepys, it is probable that charcoal is essentially as pure carbon as the diamond, and that the hydrogen it evolves depends on water, which it always absorbs on the shortest exposure to the air. Carbon is a constituent of almost all vegetable and animal substances.
- d.* **METALS** are simple inflammable bodies, of great specific gravity, density, and opacity; and, as the result of these qualities, possess great brilliancy or lustre from their power of reflecting almost all the light which falls upon their surface. Their colours are, generally, shades of white, gray, or yellow: their hardness is considerable; and, according to its degree, they are more or less elastic. One metal only, Mercury, is in a fluid state at the ordinary heat of the air. Many Metals possess considerable tenacity, and are consequently malleable and ductile; but some are extremely brittle. They are sapid and odorous when heated or rubbed; their fracture is generally hackly; their texture fibrous or foliated; and many of them are sonorous. They are excellent conductors of caloric, electricity, and galvanism. When exposed to the action of caloric, they expand and are melted; but differ greatly with regard to fusibility. Some of them are volatilized at known temperatures. When fused, their surface is convex and globular; and, in cooling, they generally crystallize. They are very susceptible of oxidizement.

---

<sup>1</sup> For experiment, the simplest mode of obtaining phosphorus is to mix a solution of phosphate of soda with a solution of acetate of lead, in the proportion of one part of the former salt to one quarter of the latter. This will yield a precipitate of phosphate of lead, from which phosphorus may be obtained by distillation.  
— *Henry's Elements of Chemistry*, vol. ii. p. 7.



The following metals or their compounds are used as pharmaceutical agents.

### I. MALLEABLE.

1. Silver. (*See Part ii.*)
2. Mercury. (*Ibid.*)
3. Copper. (*Ibid.*)
4. Iron. (*Ibid.*)
5. Tin. (*Part ii.*)
6. Lead. (*Ibid.*)
7. Zinc. (*Ibid.*)
8. Potassium, obtained by decomposing potass, is white, has the metallic lustre of silver, is soft and malleable at 50°, but at 32° is hard and brittle. Its sp. gr. at 60° is 0·86507. It becomes fluid at 136½°, and at a red heat is converted into vapour. It combines rapidly with oxygen and forms *potass*.
9. Sodium, obtained by decomposing soda, is white, has the lustre of lead, is very soft and malleable at 60°, and retains these properties at 32°. It sp. gr. is 0·97223. It melts at 194°; and requires a higher temperature to be volatilized than potassium. It combines rapidly with oxygen, and forms *soda*.
10. Calcium, obtained from lime, is yet imperfectly known. Combined with oxygen it forms *lime*.
11. Barium, obtained by decomposed barytes, is a metal of the colour of silver, melts at a low heat, but is not volatilized in a heat capable of melting plate glass. Combines rapidly with oxygen, and forms *barytes*.
12. Magnesium is a white solid metal, having the colour of silver. It is heavier than water, which it decomposes, although less rapidly than potassium and the other alkaline metals, and is converted into *magnesia*.

### II. BRITTLE AND EASILY FUSED.

1. Bismuth. (*Part ii.*)
2. Antimony. (*Ibid.*)
3. Arsenic. (*Ibid.*)

### III. BRITTLE AND DIFFICULTLY FUSED.

1. Manganese. (*Part ii.*)
- e. BORON is obtained from the decomposition of boracic acid, in the form of powder, of a deep olive colour, insipid, inodorous, and insoluble in water. It undergoes no change at any temperature, when heated in close vessels; but takes fire and burns like charcoal, when heated in the air to about 600°. During combustion it attracts oxygen, and is converted into boracic acid. It is also converted into that acid by decomposing sulphuric and nitric acids when it is treated with them. It combines with the alkalies and with sulphur.



## COMPOUNDS.

## a. COMPOUNDS OF SULPHUR.

1. *Sulphurets of metals* are compounds of sulphur and metals, inodorous, insipid, often possessed of metallic brilliancy, and brittle. They are conductors of electricity. Three of them only are officinal.

Table of Officinal Sulphurets of Metals.

Sulphurets.	Colours of the Sulphurets.	Specific Gravity.	Sulphur united to 100 metal.	Weight of an atom of the sulphur .
Mercury {	1. Black - - }	8.16 {	8	27
	2. Red - - }		16	29
Iron - - {	1. Yellow - }	4.518	57.1	5.5
	2. Yellow - }	4.830	114.2	7.5
Antimony	Leaden gray	4.368	37.25	

2. *Sulphurets of fixed alkalies* are opaque, solid bodies, of a brownish red or liver colour; decomposable by caloric, water, and acids; and, by exposure to the air, convertible into hydrogureted sulphurets.

*Officinal.* Sulphuret of potash. (*Part ii.*)

3. *Sulphurets of earths* resemble the alkaline sulphurets in their properties.
4. *Hydrosulphurets* are compounds of sulphureted hydrogen with alkalies and earths. They are soluble in water, crystallizable, and are decomposed by the atmosphere and acids.

## b. COMPOUNDS OF PHOSPHORUS.

1. *Oxide of Phosphorus*, produced by heating phosphorus in highly rarefied air, assumes the form of red flakes, which take fire when slightly heated, and burn with a very vivid flame. By further exposure to the air they attract moisture, and are converted into an acid.
2. *Phosphuret of carbon* (the substance which remains in the leather through which new-made phosphorus is strained, purified by exposure to heat in a retort,) is a light flocculent powder, of a bright orange, or rather red colour, insipid and inodorous. It burns rapidly when heated in the air, and leaves charcoal behind.
3. *Phosphurets of metals* are generally brittle, and have a metallic lustre. Vide *Annales de Chimie*, tom. i. & xiii.)
4. *Phosphurets of earths* have a brown colour, are generally luminous in the dark; insoluble in water, but decompose that fluid, furnishing phosphureted hydrogen gas.

c. CHARCOAL? (*Part ii.*)



d. METALLIC COMPOUNDS.

1. *Alloys* are compounds of two or more metals. They have generally lustre, hardness, tenacity, ductility, and other properties of the metals; but these properties in alloys differ from those of the metals from which they are formed. The compounds of mercury with other metals are named *amalgams*.
2. *Metallic oxides* are generally in the form of powders, or friable fragments not at all resembling the metals; some times laminated and crystallized; of various colours, determinate with regard to the metals and their treatment; heavier than the metals; and refractory, or fusible into glass. Some are insipid, others acrid and styptic. In general they are insoluble in water, and combine with acids, or with alkalies, or with both at the same time. They are reducible by light, caloric, hydrogen, carbon, oils, &c.<sup>1</sup>

Table of officinal metallic oxides, shewing the quantity of oxygen united to 100 of metal by weight in each.

Metals.	Colour of Oxides.	Oxygen in 100 parts.	Metals.	Colour of Oxides.	Oxygen in 100 parts.
Potassium	White	20	Copper	Red	12.5
Sodium - -	White	33.3		Black	25
Calcium - -	White	38.39	Zinc - - -	White	24.24
Barium - -	White	11.42	Bismuth -	White	11.28
Magnesium	White	66.6	Antimony	Yellow	18.60
Silver - -	Olive	7.272	Arsenic	White	34.93
Mercury	Black	4		White	52.4
	Red	8	Manga- nese	Green Black	28.75 57.5
Iron - -	Black	28.75			
	Red	48.12			
Lead - -	Yellow	7.692			
	Red	11.53			
	Brown	15.384			

3. *Carburet of iron (plumbago)* is of a dark blue or gray colour; has some degree of metallic lustre, a greasy feel, is soft, and blackens the fingers.<sup>2</sup> It is a conductor of

<sup>1</sup> As the same metal is susceptible of different degrees of oxidizement, it has been found necessary to designate the oxides thus formed by distinct appellations, indicative of the comparative quantity of oxygen with which the metals thus formed are combined. The terms *protoxide*, *deutoxide*, *tritoxide* and *peroxide*, imply that a metal is in its first, second, third, and ultimate stage of oxidizement. — These terms were proposed by Dr. Thomson.

<sup>2</sup> Pencils made of plumbago were used so early as 1565, being mentioned by Conrade Gesner, in his book on fossils, printed at Zurich in that year. “Stylus inferiorius depictus, ad scribendum factus est, plumbi cujusdam (factitii puto, quod ali-



electricity; and is not altered by water or by air, nor affected by the most violent heat, if air be excluded. The acids do not affect it. It detonates with nitrate of potash, and reduces the metallic oxides. 100 parts appear to consist of 5 of iron, and 95 of carbon, with, perhaps, a minute proportion of hydrogen.

4. *Metallic phosphurets.*

5. *Metallic sulphurets.* Metals combine with different quantities of sulphur. The sulphurets are brittle, generally destitute of lusture, and differing in colour from the metals that form their basis.

6. *Metallic oxides with alkalies.*

7. *Earths. (Part iii.)*

*Earths with earths.*

—————*metallic oxides.*

—————*alkalies.*

e. **SOLID ACIDS and SALTS**, in a crystallized state, contain a portion of water in their composition. The following are objects of pharmacy, or officinal:

1. *Phosphoric acid* is solid, colourless, transparent, and resembles glass in appearance, but, more usually, it is somewhat opaque, or resembles enamel; is inodorous, very acid, reddens vegetable blues, and deliquesces when exposed to the air. Its specific gravity in a state of dryness is 2.687. It is very soluble in water, dissolving with a hissing noise; but, nevertheless, much heat is not evolved. It is decomposed at a high temperature by hydrogen, charcoal, and several of the metals. The composition of phosphoric acid is 100 of phosphorus, and 200 of oxygen.

*Phosphate of lime* is found very abundantly in the native state. It constitutes the basis of bones, from which it is procured in the state of a white powder, by calcination and solution in muriatic acid, from which it is precipitated by ammonia. It is inodorous, insipid, and insoluble in water, but is decomposed by several of the acids. Exposed to a heat of 378° Wedgwood, it softens, and changes to an enamel. 100 parts contain about 50.8 of acid, and 40.2 of lime.

*Subspecies. 2. Bi-phosphate of lime.*--Acid 100, lime 40.27.<sup>1</sup>

*Subspecies 3. Quadri-phosphate of lime.*

*Subspecies. 4. Sub-phosphate of lime.*--Acid 100, lime 118.5.<sup>2</sup>

*Phosphate of soda. (Part iii.)*

quis stimmi anglicum vocare audio) genere, in mucronem derasi, in manubrium ligneum inserti."—*Dererum fossilium figuris*, p. 104, quoted by Beckman, *History of Inventions*, vol. iv. p. 350.

<sup>1</sup> Rose.

<sup>2</sup> Thomson's *Syst. of Chem.* 5th edit. vol. ii. p. 460; quoted from Vauquelin, *Journ. de Min.* No. 27. p. 2.



2. *Boracic acid* is obtained in the form of white, thin, irregularly hexagonal scales, greasy to the feel. It reddens vegetable blues, is inodorous, but has a sour bitterish taste, leaving a cooling sweet impression in the mouth. Its specific gravity is 1.479. It swells up when exposed to the fire, and melts into a hard, transparent glass. Boiling water dissolves only two parts of boracic acid: alcohol dissolves it, and the solution burns with a green flame; and oils also dissolve it with the assistance of heat. It oxidizes only iron, zinc, and perhaps copper. Its components are, according to the experiments of Sir H. Davy, one part by weight of boron, and two of oxygen: but, according to Gay Lussac, and Thenard, the proportions are, two of boron, and one of oxygen.

*Sub-borate of soda.* (Part iii.)

3. *Benzoic acid.* (Ibid.)

4. *Succinic acid.* (Ibid.)

5. *Oxalic acid*, generally obtained by treating sugar with nitric acid, is in the form of white, transparent, shining, four-sided prisms, which have a very acid taste, redden vegetable blues, and are soluble in their own weight of boiling water. The solution of one part of the acid in 3600 parts, by weight, of water, is perceptibly acid to the taste. Exposed to heat in open vessels, it is decomposed. According to Berzelius, 100 parts contain fifty-two of real acid, and forty-eight of water; and the dry acid is a compound of hydrogen, 0.244.; carbon, 33.222.; and oxygen, 66.534.<sup>1</sup> Oxalic acid has been swallowed by mistake for Epsom salts, and has produced fatal effects in many instances. I was the first who ascertained, by experiments on dogs and rabbits, that it acts on the stomach like other corrosive poisons; and that a mixture of chalk and water is the best antidote.<sup>2</sup>

6. *Tartaric acid.* (Part iii.)

*Tartrate of Potash.* (Ibid.)

Variety. *Supertartrate of potash.* (Part ii.)

*Tartrate of potash and soda.* (Part iii.)

7. *Citric acid.* (Ibid.)

f. BITUMENS. (Part ii.)

g. SOAPS. The true nature of these compounds has been illustrated by Chevrueil. He found that the fatty matter they contain consists of two distinct substances, which he has named *Steärin* and *Eläin*. These, by the action of salifiable bases, are converted into two acids, the *margaritic* and the *oleïc*; and soaps are merely combinations of one or both of these acids, with alkaline, earthy, or metallic bases. The

<sup>1</sup> *Annals of Phil.* vol. v. p. 99. <sup>2</sup> See *London Med. Repository*, vol. iii. p. 382.

<sup>3</sup> Berzelius.



alkaline soaps have a peculiar unpleasant odour and taste, form a milky solution with water, and a transparent one with alcohol; are powerfully detergent, and are decomposed by the earthy and the metallic salts. 2. The earthy soaps are insoluble in water, and not detergent. 3. Metallic soaps are likewise insoluble in water, but some of them are soluble in alcohol, and others in oil.

1. *Hard soap.* (*Part ii.*)

2. *Soft soap.* (*Ibid.*)

Variety. *Liniment of ammonia.* (*Part iii.*)

3. *Liniment of lime-water.* (*Ibid.*)

h. SOLID VEGETABLE SUBSTANCES. It is necessary to notice in this place the solid proximate principles only of the vegetable substances which are officinal, or employed as pharmaceutical agents. The constituents of the whole of them are—carbon, hydrogen, oxygen, and azote, in different proportions.

1. *Sugar.* (*Part ii.*) It is soluble in nitric acid, and yields oxalic acid. When treated with chlorine, sugar is changed into malic acid.

Varieties. a. *Sugar of figs*,—found concrete on the outside of the dried fruit.

b. *Sugar of grapes*,—found crystallized in some kinds of raisins. When artificially extracted from the juice of the grape, it is white and crystallized, but less sweet than common sugar.

c. *Sugar from starch*:—its crystals are small spherical granules. When dissolved in water, it ferments with the addition of yeast.

d. *Mushroom sugar*:—its crystals are four-sided prisms, with square bases; very crystallizable; digested with nitric acid it yields abundance of oxalic acid.

e. *Manna?* besides common sugar, contains mucilage, and a nauseous substance to which it owes its purgative properties. When digested with nitric acid, it yields saccharic as well as oxalic acid. It does not ferment like sugar.

2. *Sarcocoll* is usually in oblong, semi-transparent, yellow globules, which have a bitter sweet taste, and an odour resembling in some degree that of anise seed. It does not crystallize. Soluble in water and alcohol. Treated with nitric acid, yields oxalic acid.

Variety. *Liquorice?* dissolves in nitric acid, and forms tannin; and, when treated with sulphuric acid, yields about one-fourth of its weight of charcoal. It is not susceptible of fermentation.

3. *Gum.* (*Part ii.*)

*Officinal.* Acacia Gum.

Varieties. a. *Gum Senegal*,—in larger masses than gum arabic, and darker coloured. It is often mixed with gum arabic.

b. *Gum kuteera*,—in loose wrinkled transparent drops, inodor-



ous, and insipid; scarcely soluble in cold water, but completely so in boiling water.

- c. *Mucus*—Inodorous, insipid, soluble in water, insoluble in alcohol; not precipitated by silicated potash, but precipitated by alcohol in a fibrous state.

*Officinal.* Linseed mucus, quince seed mucus, marsh mallow mucus.

4. *Cerasin*. This substance, which was formerly confounded with gum, is the produce of the *Astragalus tragacantha*. For its physical properties, see (*Part ii.*) With nitric acid it yields sacclactic, malic, and oxalic acids.

*Officinal.* Tragacanth.

Varieties. a. *Cherry-tree gum*,—partially soluble only in water. When treated with nitric acid, it yields sacclactic acid.

b. *Congo gum*?

c. *Dominica gum*,—in large masses like stalactites, brittle, light yellowish brown, translucent. It contains three parts of cerasin and one of gum.

5. *Ulm*<sup>1</sup>. This is a spontaneous exudation from the elm; but, as Berzelius has hinted, is probably a component of every bark. It is solid black, hard, shining, and insipid. Soluble in water, but does not form mucilage; insoluble in alcohol; precipitated by nitric and oxymuriatic acids in the state of resin.

6. *Inulin* is obtained from the *Inula helenium* in the form of a white powder, which is insoluble in cold water and in alcohol. It is soluble in boiling water, forming a mucilaginous solution; but precipitates as the solution cools. Treated with nitric acid it yields malic and oxalic acids.

*Officinal.* Elecampane root.

7. *Starch*. (*Part ii.*)

Varieties. a. *Potatoe starch* has a perceptibly crystallized aspect, is more friable and heavier than common starch.

b. *Indian arrow root* has all the properties of common starch.

c. *Sago*, in grey granules. It is the prepared pith of a palm, the *Cycas circinades*.

d. *Casava*, prepared from the roots of the *Iatropa Manihot*.

e. *Salep*? from the bulb of the orchis mascula.

f. *Tapioca*, in granules.

All these varieties of starch are very nutritive, and excellent food for the sick and convalescent.

8. *Gluten* is of a gray colour, nearly insipid, and inodorous, very tenacious, ductile, and elastic; partially soluble in water, and soluble in acetic and muriatic acid. Insoluble in alcohol and ether. M. Taddei has ascertained that it may be decomposed into two principles — which he has named *Gliadine* and *Zimome*. When gluten is rubbed with powdered guaiac, a blue colour is evolved. When treated with nitric acid it yields oxalic acid.

<sup>1</sup> So named by Dr. Thomson.



9. *Fibrin* is tasteless, fibrous, elastic, and resembles gluten. It is insoluble in water, alcohol, and diluted alkali; but is soluble in acids, particularly nitric acid. It soon putrifies.

10. *Extractive* is obtained by evaporating aqueous vegetable infusions to dryness. It has a strong taste; and is soluble in water; but insoluble in alcohol and ether, unless when the extractive is united with resin. It is precipitated from its solutions by oxymuriatic acid, muriate of tin, and muriate of alumina; but not by gelatin.

*Officinal.* Catechu, most barks, &c.

11. *Tannin* has a bitterish astringent taste, is soluble in water, and in alcohol of 0·810. It is precipitated by the muriates of tin and of alumina, and by gelatin.

*Officinal.* Rhatany root, Galls, Uva ursi, Tormentil, Rhubarb, Cinchona barks, Swietenia, Simarouba, Kino, Catechu, Willow bark.

12. *Wax* is a fixed oil, saturated with oxygen. (*Part ii.*)

13. *Camphor*. The ultimate components are probably carbon and hydrogen, the proportion of carbon being greater than in oils. (*Part ii.*)

14. *Scillitin*. This substance is procured from the bulb of the Squill. It is white, transparent, and pulverulent: has an intensely bitter taste, leaving a sweetish impression on the palate. It is soluble in water and alcohol, and when heated swells and exhales the odour of caromel.

*Officinal.* The bulb of the squill.

15. *Nicotin* is obtained from the leaves of several species of *Nicotiana*. It is colourless, acrid, and has the odour of tobacco; is soluble in water and alcohol; and approaches the volatile oils in its properties. It is extremely poisonous.

*Officinal.* Tobacco leaves.

16. *Cathartine* is the active purgative principle of Senna. It is uncrystallizable, of a reddish yellow colour; has a bitter nauseous taste, a peculiar odour; and becomes moist in the air. It is insoluble in ether but soluble in alcohol and water in all proportions.

*Officinal.* Senna leaves.

17. *Resins* are brittle, semi-transparent, yellowish substances, inodorous, and having an acrid taste. Their specific gravity varies from 1·0452 to 1·2289. They melt when heated, inflame in a higher temperature, and burn with a strong yellow flame, emitting much smoke. They are insoluble in water; but soluble in alcohol, ether, alkalies, and acetic acid. Nitric acid converts them into artificial tannin.

*Officinal.* Amber, copal, pine resins, mastiche.

18. *Guaiacum* differs from resins in being soluble in nitric acid, and when treated with it, in yielding oxalic acid, and no tannin. (See *Part ii.*)



19. *Hematin* is the colouring matter of Logwood. It is procured in small brilliant crystals of a reddish white colour. Their taste is bitter, acrid, and slightly astringent. They form an orange red solution with boiling water; alkalies in excess change them first to purple, then to violet, and, lastly, to brown; and the hematin is decomposed. (See *Part ii.*)

*Officinal.* Logwood and its extract.

20. *Emetina*. This substance is obtained from *Ipecacuanha* and some other roots, which have an emetic property. It is, when pure, in the form of white scales, inodorous, but having a bitter and slightly acrid taste. It is not very soluble in water and alcohol, but is soluble in acetic acid, and all the other acids forming salts. It is insoluble in ether. Nitric acid converts it into oxalic acid. Half a grain, when swallowed, excites full vomiting.<sup>1</sup>

*Officinal.* *Ipecacuanha* root.

21. *Cinchonia*, an alkaline salt obtained from the bark of *Cinchona lancifolia*. It is inodorous, has little taste unless it be dissolved in alcohol; is white, translucent, and crystalline. It is scarcely soluble in water, ether, or oils, but very soluble in alcohol. It unites with all the acids forming soluble salts.

22. *Quina*, an alkaline salt procured from the bark of *Cinchona cordifolia*. Inodorous, very bitter, white, little soluble in water, very soluble in ether. Combines with acids forming soluble salts.

23. *Picrotoxa*, is procured from the fruit of *Menispermum cocculus*, in white, four-sided prismatic crystals; of an intensely bitter taste, soluble in twenty-five times their weight of water, and in three times their weight of alcohol and of ether. It is insoluble in oils. Nitric acid converts it into oxalic acid. It is intoxicating, and poisonous when swallowed.

*Officinal.* *Cocculus Indicus*.

24. *Morphia*. This is an alkaline substance procured from opium. It is in white pyramidal crystals, scarcely soluble in boiling water, but very soluble in alcohol and ether. It combines with the acids, forming neutral salts.

---

<sup>1</sup> To obtain *Emetin*, treat powder of *Ipecacuanha* with ether at 63°, as long as it acts upon it: then boil the powder several times with fresh quantities of alcohol at 40°, and filter the boiling solutions, which will throw down as they cool a white flocculent precipitate: filter again, and evaporate the clear solution in a water-bath. Dissolve the reddish residue which will be obtained in cold water, and add magnesia, which will precipitate the *Emetin*, that may be now obtained in a state of purity by dissolving the precipitate in alcohol, and evaporating the solution to dryness.



It acts with great energy on the animal economy. (See *Part ii.*)

*Officinal.* Opium, lactucarium.

25. *Aconita*, an alkaline substance procured from the *Aconitum Napellus*, on which the poisonous qualities of that plant are supposed to depend. (See *Part ii.*)

*Officinal.* *Aconitum Napellus* and its extract.

26. *Atropia*, an alkaline substance, on which the poisonous and active properties of *Belladonna* depend. (See *Part ii.*)

*Officinal.* *Atropa Belladonna* and its extract.

27. *Delphia*, an alkali, the active principle of *Delphinium Stavisagria*. (See *Part ii.*)

28. *Hyosciama*, an alkali, the active principle of *Hyosciamus niger*. (See *Part ii.*)

29. *Veratria*, an alkali, the active principle of *Veratrum album*, and *Colchicum autumnale*. (See *Part ii.*)

30. *Strychnia*, an alkaline substance, the active principle of *nux vomica*. It is white, granular, inodorous, extremely bitter, insoluble in ether, slightly soluble in cold water, but very soluble in alcohol. Forms a class of salts with acids; which are most virulent poisons.

31. *Balsams* resemble resins in their appearance; have a strong aromatic odour; yield benzoic acid when heated, or dissolved in sulphuric acid; and when treated with nitric acid yield artificial tannin.

*Officinal.* Balsams of tolu, benzoin, storax,

32. *Gum resins* resemble resins in their appearance; but they are odorous, and form milky solutions with water, and transparent solutions with alcohol. They are soluble in alkalies; and are converted into tannin by nitric acid.

*Officinal.* Ammoniacum, galbanum, scammony, assafoetida, myrrh, sagapenum, gamboge, aloes.

33. *Wood*, which forms the support of all vegetables, is composed of tasteless fibres, insoluble in water and alcohol, but soluble in weak alkaline ley, and in nitric acid yielding oxalic acid. When distilled *per se*, at a red heat, it leaves much charcoal.

#### i. SOLID ANIMAL MATTERS.

1. *Gelatin*. (*Part ii.*)

2. *Albumen* when dried is a brittle, transparent, glassy substance, resembling gum in appearance. It is soluble in cold water; and, when the solution consists of one part of dry albumen and 9 of water, heat coagulates it into a firm white solid mass. Alcohol, ether, the strong acids, many metallic oxides, and tannin, also coagulate the solution.

*Officinal.* White of egg, hartshorn shavings.



3. *Solid oils* are composed, like the other fixed oils, of a solid and a fluid substance, which Chevreul has named *Steärin* and *Eläin*.

Varieties. a. *Spermaceti*. (*Part ii.*)

b. *Fat* is an odorous, insipid, white crystal line substance; greasy to the touch; melts at  $140^{\circ}$ ; vaporized at  $400^{\circ}$ , the vapour being inflammable. Insoluble in water, alcohol, and ether; combines with alkalies, and forms soap; and is decomposed by strong acids.

*Officinal*. Lard, mutton, suet, fat.

4. *Cantharidin* is the active principle of Spanish flies.  
(See *Part ii.*)

*Officinal*. *Lyttae*.

5. *Cochinilin* is the colouring principle of the *Cocus Cacti*.  
(See *Part ii.*)

6. *Castor*. (*Ibid.*)

7. *Musk*. (*Ibid.*)

8. *Bones and Shells*. (*Ibid.*)

9. *Horn*. (*Ibid.*)

#### COMBINATION OF SOLIDS WITH SOLIDS.

Although solid bodies may be made to enter into combination with one another, yet, all do not combine in the same manner, and under similar circumstances. Thus, some unite in any proportion, and some in certain determinate proportions only; while others will not combine with each other under any circumstances.

1. TABLE of the principal Solids which have been ascertained to be capable of uniting in any Proportion.

Sulphur with phosphorus.

Carbon with iron?

Metals with most metals.

Protoxide of antimony with sulphuret of antimony.

Earths with earths.

Earths with some metallic oxides.

Some earths with fixed alkalies.

Solid oils with each other, and with bitumen.

All the products are solids, except those resulting from the union of sulphur and phosphorus, which are liquid.

None of these solids combine spontaneously, even although placed in contact; but require to be mixed, and exposed to a degree of heat capable of melting one or both of them; in which case, the caloric breaking the force of the cohesive attraction which retains the particles of the solids in the aggregate state, the atoms of the one substance are brought into immediate contact with those of the other, or within the sphere of the attraction of affinity, which consequently acts and produces



the new compound. The compounds do not very materially differ in their properties from their constituents, except the compounds of iron with carbon, and some of the earths with each other. The combination is generally accompanied with a change of density.

2. TABLE of the principal Solids which have been observed to unite only in determinate Proportions.

Sulphur with metals.

———— some metallic oxides.

———— earths.

———— fixed alkalies?

Phosphorus with carbon.

———— metals.

———— some earths.

Acid with alkalies.

———— earths.

———— metallic oxides, &c.

These enter into more intimate union than the preceding. They, however, do not unite when both bodies remain in the solid state; “except sulphur and the fixed alkaline hydrates<sup>1</sup>, some acids, and a few hydrates of metallic oxides:” hence they are brought into union, either by *fusion*, or by *solution in water*, or some other liquid menstruum. By the first mode, “sulphur is made to combine with metals, earths, and fixed alkalies, and phosphorus with metals:” by the second, the acids are combined with the alkalies, the earths, and the metallic oxides. The mode of union resembles that of liquids with solids in every respect.

It is important to ascertain the *proportions* in which these bodies unite, and their *change of density*. Berthollet is of opinion, that sulphur may unite indefinitely with the metals, the proportion of sulphur varying indefinitely in many native sulphurets; but Dr. Thomson<sup>2</sup> maintains the contrary opinion, owing to the circumstance, “that when sulphur and a metal are fused together, we obtain always the two bodies combined in determinate proportions.”

The following Table exhibits the composition of the sulphurets of the officinal metals. The first column gives the specific gravity of the sulphuret; the second the weight of sulphur united to 100 parts of the metal; the third the colour of the sulphuret.

<sup>1</sup> These are alkalies in the crystalline form, or containing water solidified.

<sup>2</sup> *System of Chemistry*, 4th edit. iii. 136.



Metals.	Specific gravity.	Weight of sulphur combined.	Colour.
Silver - -	7.215	14.544	black
Bismuth	3.3384	22.52	{ blue
Arsenic -		45.	{ blue
Copper -		71.42	red
Mercury	8.16	25.	black
		8.2	{ black
		16.	{ red
Tin - - -	7.602	27.1	{ blue
		54.2	{ yellow
		15.384	{ white
Lead - -	4.368	30.768	{ white
Antimony		33.333	lead-en-grey
Iron - -		57.1	{ bronze
	4.5	114.2	{ yellow

The metallic sulphurets are rarer than the mean of their components, owing to the substances expanding during their union sometimes more than one-fifth of the whole. Pyrites, however, is an exception, its specific gravity being greater than the mean.

Nothing precise is known of the other combinations of sulphur, or of those of phosphorus with solid bodies.

The combinations of the acids with alkalies, earths, and metallic oxides are well understood. When an acid and an alkali are mixed together, we find that, after several small additions of the alkali to the acid, diluted with a little water, the mixture still retains acid properties; but by continuing to add the alkali these disappear, and alkaline properties are acquired by the next addition that is made; the acid or the alkaline properties of the compound, therefore, predominate according to the proportions of each; but there are certain proportions, according to which they destroy, by their union, the properties of each other, so that neither predominates. In this case they are said to *neutralize* each other; the products are named *neutral salts*, and the proportions in which the acids and alkalies unite to form neutrals are fixed and determinate.

All salts, however, are not neutrals; but, in some, the proportions of the acid, in others, that of the base, predominate. The former, which are named *super-salts*, are supposed to be compounds of two or more atoms of the acid with one of the base; and the latter, which are named *sub-salts*, of two atoms

<sup>1</sup> Vauquelin,



of the base with one of the acid. <sup>1</sup>Thus supertartrate of potass consists of one atom of potass united to two of tartaric acid, or by weight, of 5.23 parts of base, and 100 of acid <sup>2</sup>; while carbonate of potass consists of two atoms of potass and one of carbonic acid, or by weight, of 2.75 of acid and 6.00 of base. *Triple salts*, are salts composed of one acid united to two bases at the same time, — as the tartaric acid, for instance, with potass and soda, to form the tartrate of potass and soda.

The metalline salts are seldom neutral, having generally an excess either of acid or of base.

## II. LIQUIDS.

It has been already observed, that by throwing caloric into a solid body, or in other words, heating it, the force of the attraction of cohesion which preserved it in the solid state is gradually weakened, and finally overcome. When the particles of a body, which were at a low temperature immoveable relatively to each other, are separated by interposed caloric, so as to move easily upon each, but are yet within the limits of the sphere of the attraction of aggregation, the body is denominated a *liquid*, provided it remain in this state under the medium temperature of the atmosphere. Thus *Ice*, when brought into a place the temperature of which exceeds 32°, loses its solidity and becomes a liquid or water, which form it retains in every degree of temperature between 32° and 212° on the scale of Fahrenheit's thermometer. Let us now examine the constitution of liquids; their combinations with other liquids; and their combination with solids.

### 1. OF THE CONSTITUTION OF LIQUIDS.

*Liquids*, as we have explained, are compounds of caloric with a solid base. Their parts move easily upon each other, and yield to the smallest impression; but they are not sensibly elastic. The greater or smaller degree of liquidity of different substances, depends upon a difference of the force of cohesion exerted between their particles, which may be regarded as placed in the limit between attraction and repulsion: thus the cohesion of mercury is greater than that of water. Liquids differ very much in specific gravity; and the degree of this bears a relation to their density. “The distances of the atoms are so regulated, that the attraction and repulsion by which they are at once actuated just balance one another; while their form, which is supposed to be spherical, is such, that they can move freely among each other without altering these dis-

---

<sup>1</sup> For an account of the Atomic Theory, I must refer my readers to Mr. Dalton's writings, and Thomson's *Annals of Philosophy*, passim.

<sup>2</sup> Berzelius.



tances. It is this which seems to constitute the real cause of liquidity.”

All liquids may be arranged into two great classes. “The following Table<sup>1</sup> exhibits a list of almost the whole of them, arranged according to their composition: —

I. SIMPLE.

	Sp. grav.
Mercury. ( <i>Part ii.</i> ) .....	13·568

II. COMPOUND.

a. Simple gases combined.

Water .....	1·000
Nitric acid. ( <i>Part iii.</i> ) .....	1·583

b. Gases with a solid base.

Sulphuric acid. ( <i>Part ii.</i> ) .....	1·898
Alcohol. ( <i>Part iii.</i> ) .....	0·794
Ethers. ( <i>Ibid</i> ) .....	0·632 to 0·900
Volatile oils. ( <i>Ibid.</i> ) .....	0·792 to 1·094
Fixed oils. ( <i>Ibid</i> ) .....	0·913 to 0·968
Petroleum. ( <i>Part ii.</i> ) .....	0·730 to 0·678
Supersulphureted hydrogen .....	—
Oxymuriate, or Bichloride of tin .....	—

c. Solids combined.

Phosphuret of sulphur .....	—
Sulphuret of carbon .....	1·272

If mercury be excepted, all the known liquids are compounds, and the greater number of them contain water as an ingredient.

*Water.* The ordinary appearances and properties of this liquid are too well known to require description. Its maximum of density is at the temperature of 36°. A cubic foot of it, at 30 inches of the barometer, and 55° of the thermometer, weighs 998·74 avoirdupoise ounces of 437·5 grains troy each. Its specific gravity is supposed = 1·000, and it is made the standard of unity in the measurement of the gravity of every other liquid. The gravity of ice is less than that of liquid water. In the form of steam, under a pressure of 28 inches of mercury, it occupies 1800 times the space which it does in the form of water. It is not decomposed by heat alone; nor altered by light: but is decomposed by iron, zinc, antimony, and tin, when assisted by heat. It readily absorbs air and gases<sup>2</sup>, and is a constituent of all gases. It is a compound of oxygen and hydrogen, 100 grains containing 88·286 of oxygen, and 11·714

<sup>1</sup> Thomson’s Chemistry, 5th edit. iii. p. 82.

<sup>2</sup> From Mr. Dalton’s experiments, it appears that water absorbs its own bulk of carbonic acid gas, of sulphureted hydrogen gas, and of nitrous oxide;  $\frac{1}{27}$  of oxygen gas, nitrous gas, and carbureted hydrogen; and  $\frac{1}{8}$  of carbonic oxide, azotic gas, and hydrogen gas.



of hydrogen.<sup>1</sup> It liquefies a great number of solid bodies, its solvent power being increased by diminishing the pressure of the atmosphere; and, as has been already stated, the greater number of liquids contain it as an ingredient.

*Supersulphureted hydrogen* is a transparent colourless liquid when pure, but more frequently has a greenish yellow tinge. It has a strong peculiar odour, and a pungent yet cooling taste. Its specific gravity is 1.3. It burns like spirit of wine, and during the combustion emits a sulphurous odour. It is a compound of 96.75 parts of sulphur, and 3.25 of hydrogen.

*Oxymuriate, or bichloride of tin*, is a transparent liquid, which exhales a very heavy dense smoke when exposed to the air. Twenty-two parts of it united with seven of water condense into a solid mass. It yields by evaporation small crystals, which are deliquescent, and sublime in a moderate heat.

*Phosphuret of sulphur* is of a yellowish white colour crystallized appearance, and exceedingly inflammable. Its components are 100 parts of sulphur, and 75 of phosphorus.

*Sulphuret of carbon* is a transparent colourless liquid, having an acrid pungent, slightly aromatic taste, and foetid odour. It is very inflammable, burning with a blue flame, and emitting fumes of sulphurous acid. It is a compound of 84.83 parts of sulphur, and 15.17 of carbon.

## 2. COMBINATION OF LIQUIDS WITH ONE ANOTHER.

When liquids are mixed together they either unite in all proportions, or in certain determinate proportions only; or they cannot be united, but separate, howsoever carefully they be mixed together; or they decompose one another.

### I. TABLE of liquids which unite when mixed together in all proportions, and do not afterwards spontaneously separate.

Water with alcohol.

—————nitric acid.

—————sulphuric acid.

Alcohol with ether.

Sulphuric acid with nitric acid.

Fixed oils with petroleum.

—————volatile oils.

—————fixed oils.

Volatile oils with petroleum.

—————volatile oils.

When these liquids are mixed together, such a mutual penetration takes place, that every portion of the mixture contains the same proportions of both ingredients; and this is the case, although there may be the greatest difference in the specific

<sup>1</sup> Biot and Arago.



gravity of the individual liquids. Agitation assists the rapidity of this effect very much, but the mixture is never perfect until some time afterwards. If, on the contrary, agitation be not employed, the mixture is always more quickly effected when the denser liquid is added to the rarer; for in the opposite case a long period often elapses before it is completed. A partial muddiness occurs when even transparent liquids of different densities are mixed together, and continues until the mixture be perfect; but when it is completed, the compound is homogeneous, and the liquids do not afterwards separate from one another.

As the density and specific gravity of a compound thus formed are always greater than the mean, caloric is evolved during the mixture. In some cases the quantity is scarcely sensible; but in other cases it is capable of affecting considerably the thermometer. If fixed and volatile oils be mixed, the temperature is not very sensibly raised; but if alcohol and water are mixed, the evolution of heat is very sensible; and if four parts of sulphuric acid and one part of water, both at  $32^{\circ}$ , be mixed together, the temperature rises to  $212^{\circ}$ . When equal parts of sulphuric acid and water are mixed, the density is augmented by 13 per cent.; of nitric acid and water, the increase is equal to  $\frac{1}{12}$ ; and when water and pure alcohol are mixed, it is rather more than  $\frac{1}{28}$  of the whole weight. These mixtures are cases of real chemical combination; the force which holds them combined being that of chemical attraction exerted between the integrant particles of the two liquids.

II. TABLE *exhibiting a list of the liquids that unite with each other only in certain proportions.*

Water with ether.

————— volatile oils.

————— sulphuret of carbon.

Alcohol with volatile oils.

————— petroleum.

————— supersulphureted oxygen?

————— phosphuret of sulphur?

Ether with volatile oils.

————— petroleum.

Volatile oils with petroleum.

Water dissolves rather less than one-tenth of its bulk of sulphuric ether; and the proportion of volatile oil it takes up is also very minute, being scarcely more than is sufficient to communicate the odour of the oil to the water, without any other of its properties. Although alcohol unites readily with the volatile oils, yet the quantity of each is limited: and the proportion of petroleum which alcohol dissolves is very small. The proportions of volatile oils and petroleum which ether dissolves are considerable.



The affinity of the compounds of this table is much weaker than those of the former; which, “with the difference between the cohesion of the particles of the two liquids, limits the combination to certain proportions.” They are also more easily decomposed; for, if a spirituous solution of volatile oil be poured into water, the alcohol leaves the volatile oil to unite with the water, while the greater part of the separated oil swims on the surface of the new compound.

III. TABLE *exhibiting the principal liquids which do not sensibly combine in any proportion.*

Water with	petroleum.
—————	fixed oils.
—————	supersulphureted hydrogen.
Fixed oils with	alcohol.
—————	ether.
Mecury with	water.
—————	alcohol.
—————	ether.
—————	volatile oils.
—————	petroleum.

In these cases the affinity between the two liquids is not sufficient to overcome the cohesion between the particles of each liquid. The spreading of oil, however, upon the surface of water, and adhering to it, is supposed to depend on the exertion of some degree of affinity, although less than is requisite to produce a combination of the two liquids.

If a liquid have an affinity for one of the constituents of another liquid, although not for the liquid itself, it frequently decomposes it, and forms new compounds.

IV. TABLE *of the principal liquids which decompose each other.*

Water is decomposed by phosphuret of sulphur.

Nitric acid by all liquids, except water and sulphuric acid.

Sulphuric acid by all liquids, except nitric acid and water.

During the first case of decomposition, which is facilitated by a high temperature, sulphureted and phosphureted hydrogen exhale, and sulphuric and phosphoric acids are formed.

The combinations of solids reduced to the liquid state, are regulated by the very same laws as those of proper liquids.

3. OF THE COMBINATION OF LIQUIDS WITH SOLIDS.

The principal liquids, the action of which upon solids has been examined, are *water, alcohol, ether, petroleum, volatile oils, fixed oils, mercury*, and the *acids* which have been already noticed.



a. *Water enters into combination with solid bodies in two states. In the first, the proportion of solid matter exceeds that of water, and the liquid becomes a part of the solid body without rendering it liquid; in the second, the solid is much exceeded by the quantity of fluid, which liquefies it, and imposes its peculiar form upon the compound. The products of the first state are denominated *hydrates*? the second constitutes *solutions*.*

I. TABLE of *hydrates or compounds of solid bodies and water, still retaining the solid form.*

1. *Sulphur* is found native in the state of a hydrate; but the hydrate most generally known is precipitated sulphur. (Part iii. p. 389.)
2. *Metallic oxides*, when in the state of hydrates, are powders possessed of very intense colour, having usually a strong taste, and being easily acted upon by acid or alkaline solutions.
3. *Earthy hydrates* are powders, and in some cases crystals.
4. *Alkaline hydrates* are what are commonly termed the crystals of alkalies.
5. *Acid hydrates* are those acids which are generally procured in a solid state, and known under the name of crystallized acids.
6. *Saline hydrates* comprehend the whole class of saline preparations, whether assuming the form of crystals, powders, or solid masses.
7. *Hydrates of hydrosulphurets* are the crystallized hydrosulphurets.
8. *Soaps* are hydrates, water being always present in them as a constituent.
9. *Tannin* and many animal and vegetable solids.

In the two last classes the proportion of combined water does not appear to be determinate, although this is the case with all the others.

*Solution.*—During solution, both bodies, or the solid and the liquid, act mutually upon one another at the same time; and the force exerted by each is equal to its mass. The action goes on at the point of contact only: hence, as far as the mass is concerned, the quantity of liquid has no effect in hastening the solution. When a solid body is plunged into a liquid, if the affinity between them be weak, the combination of the two goes on at long only as the force of the affinity is able to overcome the force of cohesion of the particles of the solid; when it stops, the compound remains solid, and is consequently a *hydrate*. But if the affinity be strong, the cohesion of the solid is gradually destroyed, and its particles, being united with



those of the liquid, are dispersed equally through it, forming a *solution*. By the addition, however, of new portions of the solid, the action of the liquid is gradually weakened; and at length, being unable to overcome the cohesion of the solid, no more of it is dissolved. In this case, the sums of the force of the attraction of affinity exerted between the solid and the liquid, and of the force of the cohesive attraction of the particles of the solid for each other, are balanced; and the liquid is said to be *saturated*. The union of the two bodies is accompanied by the usual phenomena of chemical combination. If a portion of the liquid be now abstracted (as for example by evaporation), the force of the cohesive attraction of the particles of the solid becomes again superior to the force of the affinity which separates them, so that the solid is reproduced. When this is slowly accomplished, it produces crystallization, the phenomena of which have been already noticed.

In the formation of hydrates the increase of density is often very great, and much caloric is evolved. Thus, hydrate of lime is specifically heavier than pure lime. Hydrate of alum, which is simply crystallized alum, has a specific gravity of 1.7065; but when its water is driven off by calcination, the gravity is reduced to 0.4229; and crystallized nitrate of potash, or hydrate of nitre, is of the specific gravity 1.9639; but nitre deprived of its water of crystallization is only 1.7269.

The density of solutions is greater than the mean, when pure solids are employed; but when it is the hydrates which are dissolved, the specific gravity is more generally less than the mean. The following useful tables, drawn up by Hassenfratz, show the specific gravity of saline solutions containing different proportions of salt, at 55°.¹ By consulting them we can readily know the exact quantity of salt contained in any saline solution of a specific gravity corresponding with the numbers marked in the tables; and when the gravity of the solution is not found in the tables, its saline contents can still be found by calculation.

---

¹ The salts were generally in a crystallized state. The column belonging to each salt terminates at the point of saturation at the temperature of 55°.



TABLE OF SALINE SOLUTIONS.

Weight of Salt in 100 parts of the Solution.	Sul- phate of Soda.	Sul- phate of Potash.	Alum.	Weight of Salt in 100 parts of the Solution.	Sul- phate of Mag- nesia.	Sul- phate of Iron.	Sul- phate. of Zinc.	Sul- phate of Copper.
1	1.0039	1.0086	1.0047	2	1.0096	1.0096	1.0080	1.0141
2	1.0078	1.0171	1.0094	4	1.0192	1.0203	1.0165	1.0280
3	1.0116	1.0257	1.0142	6	1.0286	1.0314	1.0255	1.0413
4	1.0154	1.0343	1.0189	8	1.0379	1.0436	1.0345	1.0539
5	1.0192	1.0429	1.0236	10	1.0470	1.0560	1.0440	1.0660
6	1.0230	1.0515		12	1.0555	1.0696	1.0540	1.0795
7	1.0268			14	1.0646	1.0829	1.0665	1.0938
8	1.0306			16	1.0711	1.0961	1.0790	1.1083
9	1.0344			18	1.0771	1.1095	1.0915	1.1230
10	1.0381			20	1.0860	1.1220	1.1040	1.1380
11	1.0418			22	1.0976	1.1358	1.1165	1.1513
12	1.0455			24	1.1092	1.1498	1.1290	1.1747
13	1.0492			26	1.1178	1.1638	1.1420	
14	1.0528			28	1.1324	1.1781	1.1550	
15	1.0564			30	1.1440	1.1920	1.1680	
16	1.0598			32	1.1557	1.2031	1.1820	
				34	1.1675	—	1.1960	
				36	1.1789	—	1.2100	
				38	1.1905	—	1.2240	
				40	1.2122	—	1.2380	
				42	1.2262	—	1.2525	
				44	1.2302	—	1.2680	
				46	1.2432	—	1.2855	
				48	1.2562	—	1.3045	
				50	1.2683	—	1.3310	
				52	1.2833	—	1.3485	
				54	1.2973	—	1.3565	



TABLE OF SALINE SOLUTIONS — *continued.*

Weight of Salt in 100 parts of the Solution.	Muriate of Soda.	Muriate of Potash.	Hyper- oxymuriate of Potash.	Muriate of Ammono- nia.	Muriate of Barytes.	Weight of Salt in 100 parts of the Solution.	Muriate of Mag- nesia.	Muriate of Lime.
1	1·0064	1·0047	1·0055	1·0029	1·0073	2	1·0068	1·0125
2	1·0128	1·0095	1·0105	1·0059	1·0146	4	1·0136	1·0212
3	1·0192	1·0143	1·0150	1·0089	1·0217	6	1·0204	1·0319
4	1·0256	1·0192	1·0193	1·0118	1·0289	8	1·0274	1·0429
5	1·0320	1·0240	1·0220	1·0149	1·0360	10	1·0340	1·0540
6	1·0384	1·0288	1·0301	1·0179	1·0430	12	1·0408	1·0650
7	1·0448	1·0338	1·0376	1·0209	1·0503	14	1·0476	1·0759
8	1·0502	1·0388	1·0461	1·0239	1·0575	16	1·0554	1·0870
9	1·0576	1·0438	1·0567	1·0269	1·0647	18	1·0612	1·0979
10	1·0640	1·0490	—	1·0300	1·0720	20	1·0681	1·1000
12	1·0775	1·0612	—	1·0358	1·0919	22	1·0751	1·1212
14	1·0910	1·0701	—	1·0416	1·1014	24	1·0823	1·1323
16	1·1045	1·0801	—	1·0474	1·1309	26	1·0895	1·1445
18	1·1182	1·0901	—	1·0532	1·1504	28	1·0967	1·1547
20	1·1320	1·1000	—	1·0590	1·1700	30	1·1040	1·1670
22	1·1462	1·1090	—	1·0642	1·1901	32	1·1114	1·1803
24	1·1608	1·1178	—	1·0693	1·2227	34	1·1190	1·1935
26	1·1760	1·1264	—	—	1·2363	36	1·1266	1·2067
28	1·1920	1·1344	—	—	1·2600	38	1·1343	1·2198
30	1·2100	1·1420				40	1·1420	1·2330
						42	1·1507	1·2478
						44	1·1597	1·2528
						46	1·1686	1·2789
						48	1·1777	1·2949
						50	1·1870	1·3120
						52	1·1963	1·3310
						54	1·2068	
						56	1·2164	
						58	1·2261	
						60	1·2380	
						62	1·2507	
						64	1·2646	



TABLE OF SALINE SOLUTIONS—*continued.*

Weight of Salt in 100 parts of the Solution.	Nitrate of Potash.	Acetate of Lead.	Acetate of Iron.	Tartrate of Soda.	Tartrate of Potash.	Phosphate of Soda.	Borax.	Soda of Commerce.	American Potash.
1	1.0063	1.0070	1.0035	1.0034	1.0050	1.0040	1.0040	1.0042	1.0050
2	1.0125	1.0140	1.0075	1.0072	1.0102	1.0081	1.0084	1.0086	1.0102
3	1.0186	1.0211	1.0112	1.0108	1.0153	1.0120	1.0122	1.0130	1.0156
4	1.0244	1.0283	1.0150	1.0148	1.0212	1.0166	—	1.0175	1.0212
5	1.0302	1.0366	1.0188	1.0190	1.0258	1.0200	—	1.0220	1.0269
6	1.0353	1.0430	1.0225	1.0231	1.0311	1.0237	—	1.0264	1.0327
7	1.0408	1.0505	1.0264	1.0272	1.0363	1.0270	—	1.0310	1.0385
8	1.0468	1.0580	1.0302	1.0313	1.0417	1.0300	—	1.0356	1.0443
9	1.0531	1.0655	1.0341	1.0355	1.0470	—	—	1.0403	1.0503
10	1.0595	1.0731	1.0380	1.0397	1.0525	—	—	1.0458	1.0563
12	1.0722	1.0891	1.0458	1.0481	1.0634	—	—	1.0544	1.0684
14	1.0850	1.1055	1.0537	1.0567	1.0744	—	—	1.0640	1.0807
16	1.0984	1.1221	1.0616	1.0655	1.0856	—	—	1.0736	1.0930
18	1.1119	1.1330	1.0697	1.0745	1.0968	—	—	1.0833	1.1053
20	1.1235	1.1560	1.0780	1.0837	1.1080	—	—	1.0930	1.1179
22	1.1389	1.1740	1.0863	1.1032	1.1196	—	—	1.1031	1.1307
24	1.1520	1.1928	1.0948	1.1153	1.1317	—	—	1.1135	1.1438
26	—	—	1.1045	1.1283	1.1447	—	—	1.1241	1.1571
28	—	—	1.1140	1.1436	1.1569	—	—	1.1349	1.1724
30	—	—	1.1224	1.1600	1.1700	—	—	1.1460	1.1840
32	—	—	1.1323	1.1801	1.1838	—	—	—	1.1989
34	—	—	—	—	1.1978	—	—	—	1.2142
36	—	—	—	—	1.2118	—	—	—	1.2304
38	—	—	—	—	1.2259	—	—	—	1.2478
40	—	—	—	—	1.2400	—	—	—	1.2660
42	—	—	—	—	1.2547	—	—	—	1.2882
44	—	—	—	—	1.2696	—	—	—	—
46	—	—	—	—	1.2861	—	—	—	—
48	—	—	—	—	1.3015	—	—	—	—
50	—	—	—	—	1.3180	—	—	—	—
52	—	—	—	—	1.3351	—	—	—	—
54	—	—	—	—	1.3527	—	—	—	—
56	—	—	—	—	1.3707	—	—	—	—
58	—	—	—	—	1.3902	—	—	—	—
60	—	—	—	—	1.4120	—	—	—	—

It is necessary to keep in view, that the solvent powers of water are augmented by an increase of temperature, and that the proportions in the foregoing Tables are such as take place only at a temperature of  $55^{\circ}$ .

If a new substance be added to the saturated aqueous solution of another substance, the result is different according to the nature of the matters employed. Sometimes the second substance is not dissolved: thus a saturated solution of muriate of lime at  $60^{\circ}$  cannot dissolve any common salt. Sometimes the whole, or a part of the new solid, is dissolved without any of the already-dissolved solid being lost or precipitated: thus a saturated solution of nitrate of potash at  $51^{\circ}$  can dissolve more muriate of soda than can be dissolved by pure water, and the same is the case with nitrate of soda; but, in the latter case, a great portion of the nitrate is precipitated. Sometimes the new



solid is dissolved at the expence of the whole of the substance already dissolved, which is consequently precipitated: thus, if a sufficient quantity of muriate of soda be added to a saturated solution of muriate of ammonia at  $61^{\circ}$ , the former salt is dissolved, but the whole of the latter precipitates during its solution. This last result, however, does not take place at every degree of temperature; for, at a boiling heat, muriate of soda is separated by those very salts which it precipitates at a low temperature.

b. Alcohol acts less extensively upon solids than water; and it forms no solid combinations similar to the hydrates.

*Table of the solids which alcohol is capable of dissolving.*

1. Sulphur.
2. Phosphorus, and its compounds.
3. Fixed alkalies.
4. Some of the alkaline earths in minute portions.
5. Most of the solid acids.
6. Many salts.
7. Alkaline sulphurets.
8. Alkaline soaps.
9. Tannin, and many vegetable substances.

A mixture of water and alcohol appears to possess greater energy as a solvent in many cases, than is possessed by either of them in a separate state.

- c. The action of ether upon solids is still more limited than that of alcohol.
- d. The actions of *petroleum*, *volatile oils*, and fixed oils, have been too little investigated to permit any general deduction.
- e. The action of mercury as a liquid is altogether confined to the metals, for many of which it has a considerable affinity, and forms compounds with them, which are denominated *amalgams*. None of these are objects of pharmacy.

### III. GASES.

GASES are aëriiform fluids, possessed of very different properties, but all agreeing in that peculiar kind of elasticity which constitutes aërial bodies.

#### 1. CONSTITUTION OF GASES.

The particles of *gases*, like those of liquids, are moveable upon each other; but gases differ from liquids in possessing elasticity, or that power which allows them to be compressed into a smaller bulk; and by which, however large a portion of any gas contained in a vessel be taken away, the small portion which is left is enabled to expand so as to fill the vessel. The bulk of common air may be thus easily reduced or increased 3000 times; and indeed, there does not appear to be any limit to expansion. These properties of airs depend on the



repulsion which exists between their component particles; the force of which, according to Newton, is always inversely, as the distance of their centres from each other. As gases contain a larger proportion of caloric than any other class of bodies, it is very probable that caloric is the cause of the repulsion which exists between their particles, or of their elasticity; and hence the addition of sensible heat to gases increases their elasticity, while the abstraction of it, or the application of cold, diminishes it. No degree of compression yet attained, nor any abstraction of caloric, can alter the constitution of air; but by compression several of the gases may be reduced to liquids, and by the simple abstraction of caloric all the vapours can be reduced to the liquid, or even the solid state.

*Arrangement of the known gases according to their composition.*

### I. SIMPLE GASES.

- |              |                    |
|--------------|--------------------|
| 1. Oxygen.   | 4. Chlorine.       |
| 2. Hydrogen. | 5. Sulphur vapour. |
| 3. Azote.    | 6. Iodine vapour.  |

### II. COMPOUND GASES.

#### a. Simple gases combined.

- |                           |                        |
|---------------------------|------------------------|
| 7. Hydriodic acid.        | 11. Muriatic acid gas. |
| 8. Protoxide of chlorine. | 12. Ammonia.           |
| 9. Protoxide of azote.    | 13. Steam.             |
| 10. Deutoxide of azote.   |                        |

#### b. Oxygen and a solid base.

- |                        |                            |
|------------------------|----------------------------|
| 14. Carbonic oxide.    | 16. Sulphurous acid gas.   |
| 15. Carbonic acid gas. | 17. Sulphuric acid vapour. |

#### c. Hydrogen and a solid base.

- |                           |                                 |
|---------------------------|---------------------------------|
| 18. Cyanogen.             | 22. Hydroguret of phosphorus.   |
| 19. Olefiant gas.         |                                 |
| 20. Carbureted hydrogen.  | 23. Bihydroguret of phosphorus. |
| 21. Sulphureted hydrogen. |                                 |

#### d. Fluorine, Chlorine, Cyanogen, with a base.

- |                        |  |
|------------------------|--|
| 24. Fluoboric acid.    | 26. Hydrocyanic ( <i>Prussic</i> ) acid. |
| 25. Chlorocyanic acid. | 27. Chlorocarbonic acid.                 |

#### e. Two solid bases.

- 28 Sulphuret of carbon.

#### f. Triple or quadruple compound gases.

- |                      |                               |
|----------------------|-------------------------------|
| 29. Hydriotic ether. | 33. Vapour of alcohol.        |
| 30. Chloric ether.   | 34. ——— of oil of turpentine. |
| 31. Sulphuric ether. |                               |
| 31. Muriatic ether.  |                               |



All these gases are invisible, except chlorine and the protoxide of chlorine, which have a yellowish green colour; but when gases of very different specific gravity are mixed together, they become in a certain degree visible. With respect to the specific gravity of gases, there is a greater difference between them under the same pressure, and at an equal temperature than exists between liquid substances; a circumstance which must depend either on a difference of the repulsive force or of the weight of the atoms in different gases.

*A Table of the density and weight of 100 cubic inches of the gases at the temperature of 60°, and a barometrical pressure of 30°.*

	Specific Gravity.	Weight of 100 cubic Inches.		Specific Gravity.	Weight of 100 cubic Inches.
Air.....	1.000	30.5	Cyanogen .....	1.804	55.028
Oxygen .....	1.111	33.888	Olefiant Gas .....	0.974	29.720
Hydrogen .....	0.0694	2.117	Carbureted hydrogen	0.555	16.99
Azote .....	0.9722	29.652	Hydroguret of } Phosphorus .....	0.9022	27.517
Chlorine.. .....	2.500	76.250	Bihydroguret of } Phosphorus .....	0.9716	29.634
Sulphur vapour.....	1.111	33.888	Fluoboric Acid .....	2.3709	72.312
Vapour of Iodine....	8.678	264.679	Chlorocyanic Acid } vapour .. .....	2.152	65.636
Hydriodic Acid.....	4.375	133.434	Hydrocyanic Acid } vapour .. .....	0.9368	28.572
Protoxide of Chlorine	2.440	74.420	Chlorocarbonic Acid	3.472	105.896
Protoxide of Azote	1.5278	46.598	Hydriodic Ether ...	5.475	166.987
Deutoxide of Azote	1.0416	31.769	Chloric Ether .....	3.474	105.957
Muriatic Acid .....	1.284	39.162	Sulphuric ethervapour	2.586	78.873
Sulphureted hydrogen	1.180	35.890	Muriatic Ether .....	2.219	67.679
Steam.....	0.625	19.062	Alcohol vapour .....	1.6133	49.206
Ammonia .....	0.590	18.000	Oil of Turpentine } vapour .. .....	5.013	152.896
Carbonic Oxide.....	0.9722	29.652			
Carbonic Acid.....	1.527	46.373			
Sulphurous Acid ...	2.222	67.771			
Sulphuric Acid .....	2.777	84.698			
Sulphuret of Car- } bon vapour .....	2.6447	80.663			

Water is a constituent of almost every gaseous body; and the quantity of it contained in each depends upon the bulk, not the density, of the gas. It also appears probable, that the weight of it contained in 100 inches of all gases under the same pressure, and at the same temperature, is very nearly the same. It can be separated, in a great degree, by sulphuric acid, very dry alkalis, lime, and other matters which have a powerful attraction for water; but the whole of the moisture cannot be absorbed by these substances; and it is therefore undetermined whether gases can exist independent of the presence of water. The quantity present in any gas is regulated, in a great degree, by the temperature: for, if this be high, a much larger proportion of moisture can be retained in the elastic form; but in a low temperature it is deposited.



Vapours differ from gases in several particulars. Their “elasticity does not increase as the pressure, like that of gases;” they can be condensed by pressure, and by the abstraction of caloric, into liquids; and even some of those bodies which are regarded as real gases, such as ammonia and chlorine, are reduced by pressure at a low temperature to the liquid form. The elasticity of the majority of vapours is sensible at a high temperature only; but some become sensibly elastic at the common temperature.

#### 1. OF THE MIXTURE OF GASES WITH GASES.

1. Gases may be mixed together in the same manner as liquids, and with nearly similar results. Some never intimately combine, or are merely mechanically mingled, while others unite closely, and form new chemical compounds, possessing properties very different from those of their components.

*TABLE of gases which may be mixed together without any apparent change in their state.*

##### i. Gases that may be mixed, but which do not combine.

Oxygen with fluoboric, fluosilic, and carbonic acid gases.

Hydrogen with muriatic acid, fluoboric acid, and fluosilic acid; carbonic oxide, olefiant gas, carbureted hydrogen, phosphureted hydrogen, sulphureted hydrogen, and ammoniacal gas.

Azote, with almost all the other gases.

Fluoboric acid and fluosilic acid, with almost all the other gases.

##### ii. Gases which mix without any change, but may be made to combine.

Oxygen with chlorine, iodine, hydrogen, azote, carbonic oxide, sulphurous acid, and protoxide of azote.

Hydrogen with oxymuriatic acid, iodine, and azote.

##### iii. Gases which mix without change, but may be made to decompose each other.

Oxygen with carbureted hydrogen, bihydroguret of phosphorus, olefiant gas, sulphureted hydrogen, cyanogen, and ammonia.

Hydrogen with carbonic acid, nitrous gas, nitrous oxide, and sulphurous acid.

Although these gases, when simply mixed, do not chemically combine, or act on each other, yet the mixtures, even independent of agitation, are homogeneous compounds, or the gases do not arrange themselves according to their gravities, but are all equally diffused in the mixture, every portion of it containing exactly the same proportion of each of the mixed gases, and when once mixed they never afterwards separate. The bulk also, after mixture, is exactly equal to the sum of the bulks of the gases which have been mixed; or each gas occu-



pies the same space as when separate; and the specific gravity of the mixture is exactly the mean of that of the gases mixed. Hence the mixture of these gases appears to be a species of combination, similar to that which takes place in mixing together vinegar and water, or similar liquids; and that this is actually the case appears from the experiments of Mr Dalton<sup>1</sup>, who found that two gases of different specific gravity, when merely brought into contact, the lightest being placed uppermost and the heaviest undermost, will mix together spontaneously, if left at rest; and the same effect will take place if the gases be put in separate vessels, communicating with each other by a tube only, as in the experiments of M. Berthollet<sup>2</sup>, who supposes that the gases dissolve one another, while Mr. Dalton conceives that they are merely mechanically combined.

Vapours and gases unite in nearly the same manner as gases and gases; and this combination enables the vapour to sustain the pressure of the incumbent atmosphere, which it could not otherwise support, without being condensed. They are also retained together by a species of affinity, sufficient to cause their intimate and uniform mixture, but not strong enough to produce chemical combination.

2. Of the gaseous bodies which chemically unite when they are mixed, "some combine in all circumstances by mere mixture; others unite only in particular states."

TABLE of some of the gases which unite by simple mixture, and form pharmaceutical products.

Names and proportions of the gases.		Products.
Volumes.		
Oxygen	$\left\{ \begin{array}{l} 100 + 133 \\ 100 + 200 \end{array} \right\}$	Nitrous gas $\left\{ \begin{array}{l} \text{Nitric acid.} \\ \text{Nitrous acid.} \end{array} \right.$
Ammoniacal gas with vapour	.....	Liquid ammonia.
—————100	+ 100 muriatic acid gas	Muriate of ammonia.
—————100	+ 100 carbonic acid ...	Carbonate of ammonia.
—————100	+ 100 sulphureted hydrogen	$\left\{ \begin{array}{l} \text{Hydrosulphuret of} \\ \text{ammonia.} \end{array} \right.$

The two first of these products are vapours, the third is a liquid, and the rest are solid bodies.

a. *Oxygen and nitrous acid* unite in two different proportions; or 100 volumes of oxygen gas is capable of uniting with 133, and also with 200 volumes of nitrous gas. The first proportions produce nitric acid, which, as nitrous gas is a compound of  $66\frac{2}{3}$  volumes of azote, and  $66\frac{2}{3}$  of oxygen, therefore appears to be a compound of  $166\frac{2}{3}$  volumes of oxygen combined with  $66\frac{2}{3}$  of azote; the second produce nitrous acid, or nitric acid saturated with

<sup>1</sup> *Phil. Mag.* xxiv. 8.

<sup>2</sup> *Statique Chimique.* i. 274. 487.



nitrous gas, which appears on the same principles to be a compound of 200 volumes of oxygen united to 100 of azote. It is, however, probable that these gases will combine in different proportions from the above, and produce nitric acid, containing various proportions of nitrous gas in solution. The immediate effect of their combination is the formation of a yellow-coloured vapour.

- b. *Ammoniacal gas* and *aqueous vapour* combine the moment they are brought into contact; and are condensed to a liquid; but the exact proportions are unknown.
- c. *Ammoniacal gas* and *muriatic acid gas* unite when equal volumes of each are brought into contact; and the result of the mixture is a mutual condensation into a white powder, or muriate of ammonia. If 100 cubic inches of muriatic acid gas, therefore, weigh 59·80, and the same bulk of ammoniacal gas 18·67, grains troy, muriate of ammonia must be a compound of 3 parts of muriatic acid by weight, united to 1 part of ammonia.
- d. *Ammoniacal gas* and *carbonic acid* unite in equal volumes, and condense into a solid salt, which is carbonate of ammonia.
- e. The combination of *ammoniacal gas* with sulphureted hydrogen in equal volumes, forms also a solid compound.

TABLE of some of the gases which mix without chemically combining, but may be made to combine; and of the pharmaceutical products formed by the combinations.

Names and proportions of the gases.				Products.
Oxygen	100	+	200 hydrogen	..... Water.
————	50	+	100 carbonic oxide	Carbonic acid.
————	250	+	100 azotic gas	..... Nitric acid.
————	250	+	100 chlorine	..... Chloric acid.
————	50	+	100 sulphurous acid	Sulphuric acid.
————	200	+	100 protoxide of azote	Nitric acid.
Hydrogen	{		100 + 100	chlorine ... { Muriatic acid.
			100 + 100	cyanogen { Hydrocyanic acid.
			100 + 100	iodine ..... { Hydriodic acid.

- a. The two first combinations may be effected by combustion, and the third by electricity. It has been supposed that the heat in these cases acts indirectly only, and produces the combination by forcibly expanding one portion of the gas, and thence producing a sudden compression in the neighbouring portions, so that some or the atoms of the two gases, being brought within the sphere of action of the attraction of affinity, combine; while the caloric evolved by this union occasioning the same expansion to be constantly renewed, the whole gaseous mixture



is by degrees combined. This theory is confirmed by the experiments of Biot, which proved that oxygen and hydrogen gases can be made to combine by simple pressure.

b. *Oxygen gas and sulphurous acid gas* probably combine when simply mixed together, but the fact has not been ascertained in a decisive manner. They undoubtedly combine in a red heat: it is, however, probable that the combination is not direct, but that a portion of the sulphur is first separated, and then enters into combustion.

All these gases suffer condensation when they combine, as displayed in the following table.<sup>1</sup>

Constituents.	Volumes of ditto.	Products.	Volumes of Products.	Volumes condensed.
Hydrogen . . . Oxygen . . . .	1 2	Water	2	1
Oxygen . . . . Carbonic acid .	1 2	Carbonic acid	2	1
Oxygen . . . . Azote . . . . .	2.5 1	Nitric acid	1 ?	2.5 ?
Oxygen . . . . Sulphurous acid	1 2	Sulphuric acid	1.2	1.8
Oxygen . . . . Protoxide of azote	2 1	Nitric acid	1 ?	2 ?
Hydrogen . . . Chlorine . . . .	1 1	Muriatic acid	2	0
Hydrogen . . . Iodine . . . . .	1 1	Hydriodic acid	2	0
Hydrogen . . . Cyanogen . . . .	1 1	Hydrocyanic acid	2	0

TABLE of the principal gases which mutually decompose each other when mixed together.

Oxygen with ..... Phosphureted hydrogen.  
Chlorine with ..... Ammonia.

<sup>1</sup> This Table is copied from that drawn up by Dr. Thomson (*Sys. of Chemistry*, 5th edit. iii. 47.)



Chlorine with .....	Phosphureted hydrogen.
————— .....	Carbureted hydrogen.
————— .....	Olefiant gas.
————— .....	Sulphureted hydrogen.
————— .....	Nitrous gas.
Sulphureted hydrogen with ...	Nitrous gas.
————— ———— ...	Sulphurous acid.

The three first decompositions are attended with combustion.

- a. *Phosphureted hydrogen* with *oxygen*. The first of these gases may be regarded as phosphorus dissolved in hydrogen; and consequently in that state which enables oxygen to act upon it at the ordinary temperature of the atmosphere. When the proportion of phosphorus is considerable, the combustion is extremely brilliant, and the caloric evolved is sufficient to set fire to the hydrogen. The products, when the combustion is incomplete, are water, phosphorous acid, and hydrophosphorous acid; but when it is complete, they are water and phosphoric acid.
- b. *Chlorine* and *ammonia*, when brought into contact, excite spontaneous combustion; one fourth of the ammonia is decomposed; the hydrogen of that portion uniting with the chlorine, and forming muriatic acid, which enters into combination with the remaining ammonia, whilst the azote is dissipated.
- c. *Phosphureted hydrogen gas*, when mixed with chlorine, burns, and exhibits a lively combustion. The products are bichloride of phosphorus and common muriatic acid.
- d. *Chlorine*, when mixed with the heavy inflammable gases, does not occasion combustion, but slow and imperceptible decomposition takes place.
- e. *Chlorine* with nitrous gas has no reciprocal action: it is brought to the state of common muriatic acid; while the nitrous gas is converted into nitric acid. The requisite proportions, according to Humboldt,<sup>1</sup> are equal bulks of each gas.
- f. *Sulphureted hydrogen gas* and *nitrous gas* mixed together in a dry state suffer spontaneous decomposition; sulphur is deposited, and protoxide of azote, ammonia, and water, are produced.

TABLE of gases which mix without spontaneous decomposition, but which may be made to decompose each other by peculiar treatment.

Oxygen with sulphureted hydrogen.
————— carbureted hydrogen.
————— olefiant gas.

<sup>1</sup> *Ann. de Chimie.* tom. xxviii. p. 142.



Oxygen with vapour of ether.

————— alcohol.

Nitrous oxide with hydrogen.

————— phosphureted hydrogen.

————— sulphureted hydrogen.

• ————— carbonic oxide.

————— carbureted hydrogen.

————— olefiant gas.

————— vapour of ether.

————— alcohol.

Nitrous oxide with sulphurous acid.

Nitric acid with hydrogen, and probably all the preceding combustible gases and vapours.

————— sulphurous acid.

Nitrous gas with hydrogen.

————— sulphurous acid.

Hydrogen with sulphurous acid.

————— carbonic acid.

Vapour of water with carbureted hydrogen.

————— olefiant gas.

Some of these decompositions are produced by combustion, and are instantaneous; others take place without combustion, and are, consequently, very slow.

*Oxygen* and *sulphureted hydrogen* gases, when mixed together, do not suffer any change; “but, if the mixture be made to approach an ignited body, combustion immediately takes place, and the products vary according to the proportion of the gases mixed.” In all cases a great proportion of the sulphur is deposited, and some sulphurous acid is formed, owing to the caloric evolved by the combustion of the hydrogen setting fire to a portion of the sulphur.

If *oxygen* gas, in a small proportion, and *carbureted hydrogen*, or *oxygen* and *olefiant* gas, be exploded together, charcoal is deposited, and water and carbonic acid are produced; beside which, an inflammable gas, named by Berthollet oxycarbureted hydrogen, remains, occupying more than double the space of the original gases. But, if the proportion of oxygen be large, that is, twice the bulk of the carbureted hydrogen, and three times that of the olefiant gas, these two gases are completely consumed, and the products are water and carbonic acid.

The *vapours of ether* and of *alcohol* detonate when certain proportions are mixed with common air, or rather with the oxygen it contains; and the products are carbonic acid and water; the quantity of the former being very considerable. when the vapour fired is that of alcohol. If the proportion of the ethereal vapour be one cubic inch, which should weigh 0·7 grains, and that of the oxygen 6·8 inches, weighing 2·3 grains,



the products will be 4·6 inches of carbonic acid, and a portion of water.

The following TABLE shows the quantity of oxygen necessary for decomposing 100 inches of each of the above gases.

100 Measures of	Measures of oxygen.	100 Measures of	Measures of oxygen.
Sulphureted hydrogen .	75	Vapour of ether . . . .	680
Carbureted hydrogen .	200	———— alcohol . .	680 <sup>1</sup>
Olefiant gas . . . . .	200		

When 100 measures of *nitrous oxide* are mixed with 100 of *hydrogen*, and fired by the electric spark, a complete combustion of the hydrogen and decomposition of the nitrous oxide take place, and water and azote are produced. The superior affinity of the hydrogen over azote is, in this case, aided by the caloric which is evolved during the combustion of the former gas.

One measure of *phosphureted hydrogen*, and three measures of nitrous oxide, are completely decomposed when exploded by the electric spark, producing water and phosporic acid, and leaving three measures of pure azote. In this case, the proportion of nitrous oxide is sufficient to combine with both the components of the phosphureted hydrogen gas. In the same manner *sulphureted hydrogen gas* is acted on by *nitrous oxide*; and the products of the detonation are water, sulphuric acid, and azote.

When a mixture of *olefiant gas* and *nitrous oxide* is detonated, the products are modified by the quantity of the latter gas employed; when this is large, the constituents of the inflammable gas are both saturated with oxygen, and water, carbonic acid, and azote, are produced: but when a smaller proportion is used, an inflammable gas remains.

In all cases of the combustion of *nitrous oxide* with inflammable gases, the phenomena are analogous to their combustion in oxygen; and the same is the case when mixtures of nitrous gas and the combustible gases are fired by being passed through a red-hot tube. Nitrous gas, however, does not detonate with, nor decompose, any of the combustible gases which have been just considered; but when moist iron is placed in contact with nitrous gas, the hydrogen evolved by the decomposition of the water for the oxidizement of the iron, decomposes the nitrous gas, converting it into nitrous oxide, and forming ammonia.

The other cases of decomposition enumerated in the table are slowly produced by the continued action of electricity, without any combustion taking place.

<sup>1</sup> Thomson's Chemistry, 4th edit. iii. 469.



The combinations of gases with gases are not immediately effected for the purposes of pharmacy, but several of them occur during many of the operations for the preparation of the saline and metallic compounds; and, therefore, require to be known and understood for comprehending the theory of these operations.

## 2. OF THE COMBINATION OF GASES WITH LIQUIDS.

Water is the only liquid the action of which upon the gases has been accurately examined. In its ordinary state it contains in solution a considerable portion of atmospheric air, which can be separated from it by boiling; and it is then capable of reabsorbing air, and any other gaseous fluid with which it may come in contact. All gases, however, are not equally absorbable; some being taken up in great quantity, and others only in a very small proportion.

*TABLE of gases which are but little absorbable by water, placed in the order of their absorption, beginning with the least absorbable.*

- |                           |                           |
|---------------------------|---------------------------|
| 1. Azotic gas.            | 7. Oxygen gas.            |
| 2. Hydrogen gas.          | 8. Nitrous gas.           |
| 3. Arsenical hydrogen.    | 9. Olefiant gas.          |
| 4. Carbureted hydrogen.   | 10. Nitrous oxide.        |
| 5. Carbonic oxide.        | 11. Carbonic acid.        |
| 6. Phosphureted hydrogen. | 12. Sulphureted hydrogen. |

The quantity of any gas absorbed by water is very much increased by pressure; but by diminishing pressure, the gas again separates in its elastic form. Temperature also regulates the quantity, which diminishes as the temperature increases, owing to every additional increment of caloric augmenting the elasticity of the aëriform fluid. Thus, Dr. Henry found that 100 inches of water at 55° absorbed 108 inches of carbonic acid, while at 85° it absorbed only 84 inches.

When water is pure, and the pressure and the temperature are equal, it then “absorbs a determinate quantity of every individual gas.”



TABLE, exhibiting the bulk of each of the foregoing gases, absorbed by 100 cubic inches of water at 60°, according to the experiments of Mr. Dalton, Dr. Henry, and Mr. Saussure.<sup>1</sup>

Names of Gases.	Bulk absorbed by 100 cubic inches of water, according to		
	DALTON.	HENRY.	SAUSSURE.
Carbonic acid.....	100	108	106
Sulphureted hydrogen..	100	106	253
Nitrous oxide.....	100	86	76
Olefiant gas.....	12·5	—	15·3
Nitrous gas.....	3·7	5·	—
Oxygen gas.....	3·7	3·7	6·5
Phosphureted hydrogen	—	2·14	—
Carbureted hydrogen...	3·7	1·4	5·1
Azotic gas.....	1·56	1·53	4·1
Hydrogen.....	1·56	1·61	4·6
Carbonic oxide.....	1·56	2·01	6·2

From this table it appears that water absorbs its own bulk, or rather more, of the first three gases; one-eighth of its bulk of the fourth; one-twenty-seventh of the fifth, sixth, and seventh; and one-sixty-fourth of the last three; and the absorption is in the direct ratio of the densities of the gases.

With regard to pressure, water of the same temperature always takes up the *same bulk* of each gas, whatever be the density of the gas; and, therefore, by increasing the pressure sufficiently, water may be made to absorb any quantity of gas. Thus, twice its bulk of carbonic acid will be absorbed under an additional pressure of 30 inches of mercury; three times its bulk under a pressure of 60 inches, and so on. A fact which has been applied to practice, in the manufacture of aërated soda-water, on a great scale. From this circumstance it would appear, that the absorbed gas still retains its elasticity; yet it is probable that a chemical attraction is exerted between the particles of the water and those of the gas, and it is taken up until the repulsion between the particles of the absorbed gas just balances the affinity of the water for them. Owing, however, to the weak affinity exerted between the gas and the water, if a quantity of water fully impregnated with any gas, as carbonic acid, for example, be exposed to the atmosphere, or any other gaseous body, the greater part of the absorbed gas escapes from the water and mixes with the superincumbent air; and, therefore, to preserve the impregnation complete, the aërated water must be preserved in well-stopped bottles; or under an atmosphere of the same gas it contains.

<sup>1</sup> Thomson's Chemistry, 5th edit. iii. 58.



Such are the principal circumstances connected with the absorption of the less absorbable bases : those which are more absorbable appear to belong to the class of *acids* and *alkalies*.

TABLE of the very absorbable gases, with the numbers of measures of each absorbed by one measure of pure water<sup>1</sup> ! and the increase of bulk produced on the fluid, supposing the original bulk to be 1.

Names of gases.	Measures absorbed.	Bulk in cubic inches.
Chlorine . . . . .	2 . . . . .	1.002
Cyanogen . . . . .	4½ . . . . .	—
Sulphuris acid . . . . .	43.78 . . . . .	1.040
Fluosilic acid . . . . .	363 . . . . .	—
Muriatic acid . . . . .	516 . . . . .	1.500
Fluoboric acid . . . . .	700 . . . . .	—
Ammoniacal . . . . .	780 . . . . .	1.666

The absorption of these gases is the consequence of the exertion of an affinity between them and water ; but, in every respect, the circumstances attending it are exactly the same as those attending the absorption of the former class of gases ; except that “ most of the gases belonging to the first class experience an expansion when absorbed ; while all those of the second undergo a condensation, their affinity for water being greater than their elasticity.”

With regard to the absorption of gases by other liquids, scarcely any very decisive experiments have been made ; but the experiments of Saussure render it probable, that alcohol and oil absorb a much greater proportion than water.

### 3. OF THE COMBINATION OF GASES WITH SOLIDS.

From the difference which exists between the constitution of gases and solids, their combination appears to be opposed by the elasticity of the former and the cohesion of the latter : but, nevertheless, under proper circumstances, both the simple and the compound gases combine with solids.

The simple gases are *oxygen*, *chlorine*, *hydrogen*, *azote*. Of these oxygen combines with all the known simple solids ; which are *carbon*, *boron*, *silicon*, *phosphorus*, *sulphur*, and the *metals* ; hydrogen requires that the solids be brought into a fluid state before it can combine with them ; and azote combines with one solid only, which is carbon.

*a. Oxygen gas* unites with carbon in two proportions, and forms *carbonic acid*, and *carbonic oxide*, which are gaseous fluids. Experiment has demonstrated that carbonic acid is composed of 27.27 parts of carbon and 72.73 of oxygen : thence, according to the theory of Dalton, an atom of it must consist of two atoms of oxygen and one of carbon. *Carbonic oxide* is composed of 41 parts by weight of oxygen and 28 of carbon ;

<sup>1</sup> Thomson's Chemistry, 5th edit. iii. 68.



or of one atom of oxygen and one of carbon. Oxygen combines with boron in one proportion only, forming boracic acid, which is a compound of one atom of boron and two atoms of oxygen, or 100 parts of boron and 228.57 of oxygen.

With phosphorus, oxygen unites in three proportions, and forms *hypophosphorus acid*, *phosphorus acid*, and *phosphoric acid*, which are all solid substances. The proportions of the constituents of these three compounds are the following: hypophosphorus acid consists of 100 of phosphorus and 66.6 of oxygen, or one atom of phosphorus and one of oxygen: phosphorus acid of 100 of phosphorus and 133.3 of oxygen, or of one atom of phosphorus and two of oxygen; and phosphoric acid of 100 of phosphorus and 200 of oxygen, or one atom of phosphorus and three of oxygen.

Oxygen combines with sulphur in the same manner, and forms *hyposulphurous acid*, *sulphurous acid*, and *sulphuric acid*; the first of which is supposed to be a solid, the second and third are gases. According to Dalton's theory, the constituents of hyposulphurous acid are one atom of sulphur and one of oxygen; or of 100 parts of sulphur and 50 of oxygen; of sulphurous acid, one of sulphur, and two of oxygen, or 100 of sulphur and 100 of oxygen: while sulphuric acid, which is found by experiment to be composed of 150 parts by weight of oxygen and 100 of sulphur, is formed by the union of one atom of sulphur to three atoms of oxygen. The oxygen in these compounds is much more loosely combined than in the preceding; and thence the greater facility with which they are decomposed by combustibles.

Oxygen unites readily with the *metals*, forming solid compounds. In them the oxygen is condensed, while the cohesion of the metallic particles is merely weakened, but not overcome. It has been supposed that the metals combine with determinate proportions only of oxygen, that there are no intermediate combinations, and that, in general, there are only two degrees of metallic oxidizement. There is reason, however, to believe with Berthollet, that the proportions are indefinite from the commencement to the highest degree of oxidizement of which any metal is susceptible, or to complete saturation; and in cases where determinate proportions are observed, these are owing to peculiar circumstances, which limit the combination, and "which in general being uniform, give rise to an invariable proportion." Thus, if the oxidizement of a metal takes place at its melting point, or at its vaporific point, these being uniform, the oxide will consequently be so, or the same determinate proportion will be observed in the combination. The compounds formed by the combination of oxygen with the metals have a powerful action on the animal economy, and



are consequently very important objects of pharmacy. In this state metals become, also, capable of combining with acids, and acquire still greater activity; and as the degree of oxidization varies, so the combination of the oxide in these different states with the same acid forms compounds differing from one another, and exerting various degrees of medicinal power.

*b. Chlorine* forms two compounds with phosphorus, one liquid and the other solid; and one compound, which is liquid, with sulphur. It unites with all the metals forming solid compounds.

*c. Hydrogen* has a considerable affinity for the simple combustibles; but they do not combine unless the cohesive force, which keeps together the particles of the solid, be overcome, or the hydrogen be exhibited in a nascent state; and, therefore, it is chiefly by the decomposition of water that these combinations are effected. Owing to the great elasticity of hydrogen gas, all the known combinations of it with the simple combustibles, except one, are gases.

Hydrogen unites with *carbon* in three proportions, constituting, 1. *olefiant gas*, which is composed of one atom of carbon and one of hydrogen; 2. *carbureted hydrogen*, composed of two atoms of hydrogen and one of carbon; and 3. *ether*, composed of 6 atoms of hydrogen, 5 of carbon, and 1 of oxygen. The first is the most intimate compound of the three, and carbureted hydrogen the next; for neither of these is affected by a red heat; whereas ether is decomposed, and converted into olefiant gas, carbureted hydrogen, and charcoal.

“Sulphureted hydrogen is the most intimate of the combinations of *sulphur* and hydrogen. A red heat does not decompose it.” It is commonly formed by the “decomposition of water by the compound agency of an acid and a metal united to sulphur.” In this case no obstacle is raised to the combination of the hydrogen, which is nascent, by the attraction of cohesion, the sulphur being just separated from the metal.

The combination of hydrogen with *phosphorus* is also obtained by the decomposition of water, by boiling the phosphorus in a liquid alkali, which retains the phosphorus in a temperature sufficient for enabling it to effect the decomposition. The oxygen of the water unites with one portion of the phosphorus, and forms phosphoric acid; while, at the same time, the hydrogen unites with the other portion, and forms phosphureted hydrogen.

*d.* The combinations of *azotic gas* with the simple solid combustibles are not yet sufficiently understood, except its combination with carbon to form *cyanogen*<sup>1</sup>, which is a compound of one atom of azote and two atoms of carbon.

---

<sup>1</sup> This name was imposed upon this gaseous compound by Guy Lussac, who discovered it in 1815; and who found, that, in combination with hydrogen, it forms



The compound gases do not enter into many combinations with solids, if the salts, which the acid gases form with alkalies, earths, and metallic oxides, be excepted; and those formed by ammonia with the solid acids. In general they are rather decomposed. Thus, carbon, phosphorus, sulphur, and many metals decompose nitrous gas, nitric acid, and oxymuriatic acid; and sulphurous acid is decomposed by the metals.

---

Such are the effects of the combinations of solids, liquids, and gases. The knowledge of the laws which regulate them, and the results of the combinations, are of the utmost importance; the greater number of the operations of pharmacy consisting of the combination of substances, with a view either of obtaining compounds by their direct chemical union, or the products of chemical action resulting from their mutual decomposition.

---

### SECTION III.

## PHARMACEUTICAL OPERATIONS AND APPARATUS.

THE operations of pharmacy may be arranged under two classes:—

I. *Operations which are purely mechanical.*

II. *Operations which are performed by chemical powers and agents.*

The first are intended for determining the weight and bulk of bodies, diminishing their cohesion, and separating their integral parts: the second are intended for separating the elements of bodies from one another, and for reuniting these elements into new combinations.

#### I. PHARMACEUTICAL OPERATIONS PURELY MECHANICAL.

a. *Of the means of determining the weight and bulk of bodies.*

In pharmaceutical processes it is essential that the quantities of the substances employed be accurately ascertained; and for this purpose beams with scales, and measures, must be provided. Several sets of beams and scales are necessary; one set for large weights, from one pound to one hundred weight or more; another for weights not exceeding five pounds; and a third for small weights under two drachms. A good beam should remain in equilibrium, both by itself, and when the scales are

---

Prussic acid, which he therefore named *hydrocyanic acid*. It may be obtained from exposing dry prussiate of mercury in a small retort to a moderate heat. It is a colourless, transparent, elastic fluid, with a strong, disagreeable odour, solubly in water and alcohol; and highly inflammable.



suspended to each extremity : the largest set should be exact to within half a drachm ; the second should be sensibly affected by two or three grains at most ; and the smallest by the hundredth part of a grain. Apothecaries, however, seldom have beams of such accuracy, and, generally, those that they employ are much injured by exposure to acid fumes, and from want of cleanliness. To preserve the delicacy of beams they should be kept in very close cases, and not left suspended longer than is absolutely necessary ; nor should they be overloaded.

Drugs are bought in the gross by *avoirdupois weight*, which is the standard of most articles of merchandize ; but, for the composition of medicines, *troy weight* is directed to be used by the British Colleges. The following TABLE exhibits the manner in which the pound is divided, and the signs used, in prescriptions, for denoting the different weights.

A pound ( <i>libra</i> ), ℔	} contains {	Twelve ounces.
An ounce ( <i>uncia</i> ), ℥		Eight drachms.
A drachm ( <i>drachma</i> ), ʒ		Three scruples.
A scruple ( <i>scrupulus</i> ), ʒ		Twenty grains.
A grain ( <i>granum</i> ), gr.		_____

The differences between the avoirdupois pound and the troy or apothecaries' pound, and their subdivisions, are exhibited in the following TABLES.

#### Troy or Apothecaries' weight.

Pound.	Ounces.	Drachms.	Scruples.	Grains.
1	= 12	= 96	= 288	= 5760
	1	= 8	= 24	= 480
		1	= 3	= 60
			1	= 20 <sup>1</sup>

#### Avoirdupois weight.

Pound.	Ounces.	Drachms.	Grains.
1	= 16	= 256	= 7000
	1	= 16	= 487.5
		1	= 27.975

The Troy weight has also been adopted by the Edinburgh College for apportioning liquids ; but the London and Dublin Colleges with more propriety order liquids to be measured : and for this purpose the London College employs measures derived from the wine gallon, which is subdivided for medical purposes, in the manner exhibited by the following TABLE, which shows also the symbols used for denoting the several measures.

<sup>1</sup> Tables of the method of reducing the subdivisions of the troy pound into decimals of the troy pound are given in the Appendix to Part I.




A gallon ( <i>congius</i> ), Cong. or c	} contains {	Eight pints.
A pint ( <i>octarius</i> ), o		Sixteen fluid ounces.
A fluid ounce ( <i>fluid uncia</i> ), f 3		Eight fluid drachms.
A fluid drachm ( <i>fluid drachma</i> ), f 3		Sixty minims.
A minim ( <i>minimum</i> ), m		_____

Table of the proportions of the wine gallon.

Gallon.	Pints.	Fluid Ounces.	Fluid Drachms.	Minims.
1 =	8 =	128 =	1024 =	61440
	1 =	16 =	128 =	7680
		1 =	8 =	480
			1 =	60

The London College have introduced the minim measure as a substitute for the drop, the inaccuracy of which had been long experienced; as the fluidity and specific gravity of the liquid, the thickness of the lip of the phial, and even its degree of inclination, were all liable to vary the size of the drop; but by dividing the fluid drachm into sixty equal parts, a measure of bulk is obtained, which is as constant and uniform as the grain weight employed as solids.

For measuring liquids graduated glass measures of different sizes *a*, *b*, are to be preferred: and for quantities under five minims a slender graduated glass tube, *c*, open at both ends, is to be employed. When this tube is used, the graduated end is to be inserted into the liquid to be measured, down to the mark indicative of the quantity required; and the upper end being then closely covered by the finger, it is removed, retaining the proper quantity of liquid, which again drops from it on raising the finger from the upper end. A small glass measure, properly graduated, is now pretty generally used instead of the tube. In extemporaneous prescriptions, the measures of a table-spoonful and a tea-spoonful are used, when great accuracy is not required; the former being supposed to be equal to half a fluid ounce, the latter to a fluid drachm.



Elastic fluids or gases are also measured in glass jars, or tubes hermetically closed at one extremity, and graduated by inches with their decimals; but, in ascertaining the bulk of gases, the temperature of the atmosphere, and its density at the time, as indicated by the thermometer and the barometer, must be attended to; for if the former be above or below 60°, the mean heat of the air, or if the mercury in the barometrical tube be under or above 30 inches, corrections must be made by calculation relative to the degrees of temperature and pressure. For the former, the observed column of air must be divided by 80



and the quotient multiplied by the degree of temperature above or below  $60^{\circ}$ . This correction is negative when the actual temperature is above the standard, and positive when it is below. For making the corrections with regard to pressure, see the table in the Appendix.

The SPECIFIC GRAVITY of bodies is also necessary to be known in many pharmaceutical processes; and as the effects of acids and alcohol depend on the degree of their concentration, a knowledge of their gravity enables this to be correctly ascertained. The specific gravity of any substance is "the quotient of its absolute weight divided by its magnitude, or the weight of a determinate bulk of any body; and as a standard for this purpose, the weight of a determinate magnitude of distilled water has been generally assumed as unity."<sup>1</sup> It is seldom necessary to determine the specific gravity of solids; but for ascertaining that of fluids various means may be employed. If a little ball of rock-crystal, for instance, suspended by a hair or a fine platinum wire, be weighed first in air, and afterwards in distilled water, of the temperature of  $60^{\circ}$ , Fahr., the weight lost by the ball is equal to the weight of an equal bulk of the liquid; so that by repeating this operation in other fluids, and dividing its loss of weight in any other liquid by its loss of weight in water, the quotient is the specific gravity of the particular liquid. The specific gravity of liquids, however, is more generally determined by hydrometers, of which Mr. Nicholson's is by far the most accurate.<sup>2</sup>

The specific gravity of liquids is also very easily determined by the following simple method: Take a small light bottle which stands firmly, *e*, and holds about a fluid ounce or two fluid ounces of water, and stop its neck by a piece of barometer-tube very accurately ground; or a conical stopper with a notch in it, *f*. First weigh the empty bottle and tube, then fill it with distilled water at  $60^{\circ}$ , recently boiled, till the water rises a little into the bore of the tube, and weigh the whole, scratching the weight in grains on the bottle, and also the weight in grains of the empty bottle and tube. For facilitating calculation, the water should be brought to that height in the tube, at which its weight will be 500 grains, or 1000, or 1500 or 2000; and this height must be accurately marked on both sides of the tube with a file. A bottle of this kind is now sold under the name of "*Thousand grain bottle*," with a weight which is an exact counterpoise to it, when it is full of distilled water at  $60^{\circ}$ .



<sup>1</sup> Lavoiser's *Elements of Chemistry*. — Trans. 376.

<sup>2</sup> Nicholson's *Journal*, 4to. ; 110.



By filling this bottle with any any other fluid, and weighing it, the specific gravity of that fluid is ascertained by only calculating how much lighter or heavier it is than the same bulk of water.<sup>1</sup> Thus, if it be filled with sulphuric ether, instead of weighing 1000, it will require 739 grains to be put into the scale with the bottle to restore the equilibrium, and thence, its sp. gr. is 0·739: but, if sulphuric acid be used, 875 grains will be required to be added to the 1000 grains weight in the opposite scale, and consequently its sp. gr. is 1·875.

In ascertaining specific gravity the substances should be brought by calculation to the temperature of 60°, if the thermometer be above or below that point at the time of performing the experiments; and the gravities should always be expressed according to their relation to distilled water. Although this is the method generally employed in philosophical and pharmaceutical operations, yet it is necessary to observe, that the strength of spirits, according to the excise laws in this country, is estimated by the proportion they contain of a standard spirit, termed hydrometer proof, which consists of 40 parts of pure alcohol and 51 of water. Sikes' hydrometer is the one employed by the Excise. The strength of spirits stronger than proof, or *over proof*, is ascertained by the bulk of water required to reduce a given bulk of the spirits to the specific gravity denominated proof, on Sikes' hydrometer; and the strength of weaker spirits, or *under proof*, is estimated by the quantity of water it would be necessary to abstract to bring the spirits up to proof. Thus, if 20 gallons of the spirit require the addition of one gallon of water to bring it to proof, the spirit is said to be *one to twenty over proof*; and if, from the same quantity of spirit, one gallon of water must be abstracted to bring it to proof, it is said to be *one in twenty under proof*; and so on.

b. *Of the mechanical division of bodies.*

The cohesion of solid bodies often opposes an obstacle to their immediate chemical combination with other substances, and their medicinal action in the stomach; and therefore the following preliminary mechanical operations are instituted for overcoming to a certain degree that power, and separating the integrant particles of bodies, or reducing them to the state of powder. These are denominated pulverization, trituration, levigation, and granulation.

1. *Pulverization* is that process by which friable and brittle solid bodies are reduced to powder. It is generally performed in mortars by means of pestles. These are made of various materials, of brass, iron, marble, granite, glass, agate, and


---

1 *Aiken's Dictionary of Chemistry*, ii. — *Appendix*.




porcelain, or of Wedgewood's ware, according to the nature of the substances for the pulverization of which they are intended to be used; it being requisite that the materials of which pestles and mortars are made be such as to resist both mechanical force and the chemical action of the substances they contain. Thus a mixture containing any astringent matter, is turned black if rubbed in an iron mortar.

Mortars are required to be of various sizes. The largest size is usually made of cast-iron, *a*. fitted with a wooden cover, *b*. perforated to admit the pestle, but close enough to prevent the finer and lighter parts of the substances from flying off, and to defend the operator from disagreeable and noxious matters, such as aloes, ipecacuanha, &c. This may be more completely attained by tying closely round the mouth of the mortar, and round the stalk of the pestle, *c*. a



large piece of leather, so pliable as to permit the free motion of the pestle. But, notwithstanding these guards, it is sometimes necessary for the operator to cover his mouth and nostrils with a wet cloth, and to stand with his back to a current of air, that the particles which arise may be carried from him, when very acrid friable matters, as Euphorbium or Spanish-flies, for instance, are to be powdered. To lessen the labour, the pestle is often attached by a cord to the end of a flexible wooden beam, placed horizontally over the mortar, the elasticity of which elevates the pestle to the proper height after each stroke is made. For lighter purposes, brass and bell-metal mortars are sometimes used; but as, in the pulverization of every hard body, the mortar also is worn by the operation, these mortars are improper for pharmaceutical purposes; neither must marble or metallic mortars be used for acid substances. The

most useful mortars for smaller articles are those of porcelain and Wedgewood's ware, *d. e.* as they are smooth, hard, and resist the action of any chemical re-



agent. The handles, *f*. of the pestles are made of wood, and the heads, *g*. of the same material as the mortar.



Of whatever materials mortars are made, they should be internally, at bottom, of the form of a hollow hemisphere, and their sides should have such a degree of inclination as to make the substances fall back to the bottom every time the pestle is lifted. The operation, however, is retarded when too great a portion of the ingredients falls under the pestle; hence a large quantity of any substance should not be put into the mortar at a time, and the finer parts should be from time to time removed.

Vegetable matters require to be dried before they can be pulverized; and wood, roots, and barks should be previously cut, chipped, or rasped. When roots are very fibrous, as those of ginger, for example, it is advisable to cut them diagonally, which prevents the powder from being full of hair-like filaments. Resins and gum resins, which soften in a moderate temperature, or in warm weather, should be powdered in cold weather, and only gently beaten to prevent them from running into a paste instead of forming a powder; and when the powdered substance is intended to be dissolved in any menstruum, except a pure alkali, the pulverization is much facilitated by mixing them with a portion of clean well washed white sand. The pulverization of camphor is assisted by the addition of a few drops of alcohol: sugar is the best addition to aromatic oily substances, as nutmegs, mace, &c.; and to the emulsive seeds some dry powder must be added, without which they cannot be reduced to powder. Metals which are scarcely brittle enough to be powdered, and yet are too soft to be filed, as zinc, for instance, "may be powdered while hot in a heated iron mortar, or metals may be rendered brittle by alloying them with a small quantity of mercury<sup>1</sup>;" but as metals are not required to be reduced to the state of very fine powder for pharmaceutical purposes, these processes are seldom performed.

2. *Trituration* is intended to produce the same effect as pulverization, but in a greater degree. It is performed by a rotatory motion of the pestle, either in the common mortars of glass, agate, or Wedgewood's ware, or in flatter mortars made of the same materials. On a great scale, this operation is performed by means of large rollers of hard stone, which turn upon each other as in corn-mills, or by one vertical roller turning upon a flat stone. The fine powders kept in the shops are generally ground in this manner; but there appears to be an error in reducing vegetable matters to the state of impalpable powder: for in this state, both during the process of grinding and afterwards, the air and light act powerfully upon


---

<sup>1</sup> *Lavoisier's Chemistry.*—Trans. 437.




them, and produce changes, which, although they be not well understood, yet appear to alter the medicinal virtues of the substances.

3. *Levigation* is a process similar to trituration, except that the rubbing is assisted by the addition of a liquid in which the solid under operation is not soluble. Water or spirit of wine is usually employed, and occasionally viscid and fatty matters, as honey and lard. The substance to be levigated is spread on a flat table of porphyry, or some other hard stone, *b.* and is then bruised and rubbed with a muller of the same materials, either of a pyramidal shape, as *a.* or a portion of a large sphere. A thin spatula of ivory, horn, wood, or iron, is employed to bring back the materials from the edges of the table, to which the operation of the muller continually drives them. Earths and some metallic substances are thus prepared.



4. *Granulation* is employed only for the mechanical division of metals and of phosphorus. It is performed by melting the substance, and either stirring it briskly until it is cold, or pouring it, in the melted state, into water, and stirring or agitating it till it cool. For the granulation of phosphorus the latter process must be employed.

Substances are also reduced to the state of coarse powder by rasping and filing; and softer vegetable bodies are converted to a pulp by means of the grater, *c.*



#### MECHANICAL SEPARATION.


The parts of substances, under certain circumstances, may be separated from each other by different mechanical means; as sifting, washing or elutriation, filtration, expression, and despumation.

1. *Sifting*. The particles of the powders obtained by the longest and most accurate pulverization and trituration are still of very unequal degrees of fineness, and therefore require to be separated, the finer from the coarser, by the operation denominated sifting. For this purpose sieves are employed, made of iron-wire, or of hair-cloth, or of gauze. These permit the finer particles to pass through them, and leave the coarser to be again submitted to the pestle; and thus by de-




grees the whole assumes an uniform fineness.

The simple sieve is a broad wooden hoop, with a cloth of one or other of the above textures stretched over it in the manner of the parchment of a drum. The compound sieve, which is more employed, consists of the simple sieve, *b*, with a deeper rim, *a*, a lid covered with leather, and *d*, a receiver, with leather stretched across one end, and made sufficiently wide to admit the lower portion of the sieve to enter and fit tightly within it. When these are put together, the finest powders may be separated by them without any loss or inconvenience to the operator.



2. *Washing*, or *Elutriation*, is intended for separating the finer parts of powders prepared by trituration or levigation, which are not acted upon by water. The powdered substance is mixed with a large quantity of water, and briskly stirred, so as to diffuse it pretty equally through the fluid, which retains the finer particles suspended for a short space of time, and permits the coarser to settle to the bottom. The liquor still containing the finer particles suspended is poured off from the sediment; and by allowing it to remain at rest for a sufficient length of time, it deposits the fine powder, from which the clear water is separated, either by carefully decanting it; or, if the sediment be very light, so as to be easily disturbed, by drawing it off through the glass syphon, *g*, the longer limb of which being plunged into the vessel containing the fluid till it nearly touches the subsided powder, where it is supported by the board, *f*, placed on the mouth of the vessel, *e*, and the air sucked from it by means of the arm, *h*, the whole of the supernatant fluid is drawn off into the vessel, *i*, and the powder left in a fit state to be dried.<sup>1</sup> The coarser particles first separated may be again levigated, and the elutriation repeated. Chalk and some metallic matters are thus prepared; and the process may likewise be employed for separating substances of different degrees of specific gravity, although of the same degree of fineness.




<sup>1</sup> The fluid may also be drawn off by means of wet cotton or worsted threads, having one end in the fluid, nearly touching the precipitate from which it is to be separated, and the other hanging over the side of the vessel, below the plane of the fluid in it.



3. *Filtration* is intended for separating fluids from solid bodies suspended in them. Filters may be regarded as kinds of sieves; and are generally made either of very fine and close flannel, or of linen, or of unsized paper formed into a conical shape, through which the liquid percolates clear, while the solid is collected at the apex of the cone, which is inverted. When the quantity of materials is large, and the solid suspended in the fluid is not in the state of very fine powder, flannel or linen bags are to be preferred, as performing the process more quickly than paper. These are generally made in a conical shape, with the mouth stretched on a hoop or frame supported upon a wooden stand. When the solid residue is the part to be preserved, flannel filters may be used; but when the filtered liquor is the valuable product, linen is preferable, as it absorbs less of the fluid, which is obtained, also, in a more limpid state. The cloth must be well cleaned after every time that it is used, to prevent any thing from remaining to injure subsequent operations. For smaller processes, unsized paper is the best material for forming filters. A square piece of this paper of a size proportionate to the quantity of the substance to be filtered is taken, and first doubled from corner to corner into a triangle, which by a second doubling forms again a smaller triangle, and this when opened constitutes a paper cone, *k*, which is to be supported in a glass funnel, *l*, before the liquor is poured into it.

Funnels are made of tin, or of Wedgewood's ware, or of glass; but only the two latter should be used in the laboratory. Those which are ribbed are preferable, as the paper adheres so closely to the sides of smooth funnels as nearly to prevent the filtration from proceeding, unless pieces of straw, or quills, or thin glass rods, be arranged round the inside, so as to form an unequal surface for the paper to rest upon.

In most instances, the first portions of fluid that pass through a filter are turbid, and therefore require to be poured back again into the filter, sometimes repeatedly, until the pores are sufficiently obstructed to permit the most limpid part only of the liquor to pass. In cases where the solid residue is small, and it is requisite to collect the whole of it, it is useful to have a small glass tube, drawn out to a fine capillary point at one extremity; and blown to a globe in the center; by filling which with distilled water, and putting the larger end into the mouth, the force of the breath can direct a small, but strong stream of water round





the sides of the paper in the funnel, which will wash down to its bottom all the minute particles of solid matter lodged on its sides.

The concentrated acids and alkaline solutions act too powerfully on the materials commonly employed for filters, to be filtered in the ordinary way; and therefore, when it is required that they should be filtered, which is not often the case, they are passed through strata of siliceous matter arranged in a glass funnel, in the following manner: An irregularly shaped pebble is first dropped into the throat of the funnel; then a layer of pieces of quartz, or broken flint glass, is placed over it; and lastly, a thick stratum of coarsely powdered glass, or of well washed white sand, covers the whole. The substance to be filtered is poured gently on the surface of the sand, and soon passes through it and the substrata, leaving the impurities behind.

*Expression* is employed for obtaining the juice of fresh vegetables, and the unctuous vegetable oils. The subject is first bruised or coarsely ground, then inclosed in a hair-cloth bag, and subjected to violent pressure between the plates of a screw press. The bags should be nearly filled; and the pressure should be gentle at first, and gradually increased.

Vegetables in general, intended to be expressed, should be perfectly fresh, and cleansed from all impurities; and should be submitted to the press as soon as they are bruised, as the bruising disposes them more readily to ferment: but subacid fruits yield more juice, and of a finer quality, when the bruised fruit is allowed to stand for some days in an earthen or a wooden vessel. It is necessary to peel oranges and lemons before pressing them, to prevent the essential oil which their rind contains from mixing with the juice; and to some vegetables, which are not very juicy, the addition of a little water is requisite.


For expressing the unctuous seeds, in order to obtain the oil which they contain, iron plates are employed; and the bruised seeds should be previously exposed in a bag to the steam of boiling water.

*Despumation* is employed to clarify fluids which are so thick and clammy as not to be able to penetrate through the substances of which filters are made, without some previous preparation. For this purpose it is, sometimes, required only to heat the liquor, which then throws up a scum that is to be carefully removed; but more frequently it is necessary to clarify it with the white of egg. When the substance is not spirituous, as syrup, for example, the albumen which is mixed with the fluid coagulates when it is boiled, and, entangling



the impurities of the fluid, rises with them to its surface in the form of scum; but spirituous liquors may be clarified with isinglass without the assistance of heat, the alcohol coagulating the isinglass, which forms a scum, and descending to the bottom of the vessel, carries with it all the impurities. Some expressed juices are clarified by the simple addition of any vegetable acid.

Besides the above methods of mechanically separating the parts of substances from one another, fluids of different specific gravities, mixed together, are separated by means of the separatory funnel, *m*. It is chiefly used for separating the essential oils from the water with which they are entangled during their distillation. The funnel is first stopped at the bottom, and then filled with the mixed fluids, the heaviest of which gradually subsides into the narrow part below; and when the cork at the bottom is taken out, and the stopper above a little loosened, it flows out; by which means the lighter is easily obtained in a separate state. Some of the essential oils are heavier, others lighter than water, but both can be thus separated with equal facility.



## II. CHEMICAL OPERATIONS.

The operations of Pharmacy, which are strictly chemical, may be arranged in three classes.

- a.* Operations which produce changes in bodies, separating the constituents, without any obvious decomposition.
- b.* Operations in which changes are produced by the chemical action of one set of bodies upon another, or attended with obvious decomposition.
- c.* Operations in which the oxidizement and the deoxidizements of bodies are effected by means of a very high temperature.

*a. Of the operations which produce changes in bodies, separating the constituents, without any obvious decomposition.*

These changes are effected —

- |                          |     |                |
|--------------------------|-----|----------------|
| 1. By caloric — inducing | . . | Liquefaction.  |
|                          |     | Fusion.        |
|                          |     | Evaporation.   |
|                          |     | Exsiccation.   |
|                          |     | Distillation.  |
|                          |     | Rectification. |
|                          |     | Dephlegmation. |
|                          |     | Sublimation.   |



- |                               |                |
|-------------------------------|----------------|
| 2. By water and other fluids. | Solution.      |
|                               | Lixiviation.   |
|                               | Maceration.    |
|                               | Digestion.     |
|                               | Infusion.      |
|                               | Decoction.     |
|                               | Extraction.    |
| 3. By other chemical agents   | - Coagulation. |

1. *Operations in which the changes are effected by caloric.*

*Liquefaction* is that operation by which certain bodies when exposed to a moderate heat are rendered fluid, after passing through several intermediate states of softness. Fat, lard, wax, resin, and many other similar bodies undergo liquefaction; which is, therefore, employed in pharmacy to facilitate the combination of these bodies in the formation of ointments. The best vessels for conducting the process of liquefaction are earthen ware pans.


*Fusion* differs from liquefaction, in the sudden change from the solid to the fluid state which those bodies, which are liable to it, suffer on exposure to heat. There are no intermediate states of softness; but the fusible body, when heated to a certain point, immediately assumes the fluid form. This point differs very considerably in different solids; but in general simple substances are less fusible than compounds; and some of the simple earths cannot be fused without the addition of some other substances to promote their fusion. These are, generally, saline bodies, and are denominated *fluxes*.

Fusion may take place without changing the nature of the fused matter; but in general this operation is intended as a mean of promoting chemical action and of decomposing bodies. It is, however, generally confined to the metals which are extracted from their ores; and afterwards smelted and alloyed by it. It is a species of operation seldom employed in pharmaceutical processes.

Fusion is usually performed in crucibles, the best of which are made of very pure clay, or potter's earth. Those formed of common clay with calcareous or siliceous earth are easily vitrified, and then melt. The Hessian crucibles are composed of better clay and sand, and when good, stand the fire very well; as do also Wedgwood's crucibles; but they are apt to crack when suddenly heated or cooled, — a circumstance, however, which may be remedied by using a double crucible, and filling the interstice with sand, or by coating the crucible with a paste of clay and sand. Crucibles formed of black lead resist very sudden changes of temperature; but, they are destroyed, if nitre be melted in them, and even a current of air acting upon them,



whilst they are hot, destroys them. Crucibles are made of various forms, three-cornered or round, *b b*, and fitted with covers, *a a*. The lids may be luted on if necessary, with a mixture of clay and borax. Those crucibles, which are of uniform thickness, which have a reddish-brown colour without black spots, and a clear sound when struck, are to be preferred.



In order to expose the lower part of a crucible to the utmost intensity of heat, and to prevent it from cracking by the draught of cold air which would be directed upon it, were it to be placed directly upon the grate of the furnace, it is usually raised upon a small stand *c c*, either solid or hollow, an inch above it, which, according to Dr. Kennedy, is the hottest part of the furnace.

Crucibles are also made of cast-iron, of fine silver, and of platina. The first, however, are destroyed when saline substances are melted in them; and, when made red-hot in a current of air, are apt to suffer oxidation; but in other respects they are durable, and can sustain sudden alternations of heat and cold without cracking. Some of the metallic crucibles combine many of the best qualities necessary for this set of instruments; particularly those of platina, which, however, are too expensive for ordinary use.

*Evaporation* is the dissipation of a liquid by means of heat, and is employed in pharmacy generally with the view of obtaining, in a separate state, any fixed substance which may be combined with water, or some other evaporable fluid. Thus, by exposing an aqueous solution of a salt to a certain degree of heat, the caloric which combines with the water renders it volatile, and disperses it in the form of an elastic aeriform fluid, while the particles of the salt being brought nearer to each other, and within the sphere of their mutual attraction, reunite, and the salt is obtained in its concrete state. This process differs from spontaneous evaporation, in which air is the principal agent, the liquid being diminished in quantity and dissipated in that fluid, independent of the action of caloric; whereas evaporation is not carried on by the air, nor even much accelerated by the exposure of a large surface, but only in proportion to the quantity of caloric which combines with the fluid, or the degree of heat at which the process is conducted. As the fluid which is dissipated is entirely lost, and sacrificed for the sake of the fixed substance with which it was combined, evaporation is only employed where the liquid is of




little value, such as water ; but where a solid is to be recovered from a more valuable liquid, as alcohol, for instance, the process of distillation is employed.

For small processes, very good evaporating dishes are made of the bottoms of broken retorts and matrasses, which may be cut smooth round the edges by means of a hot iron or ring *d*, with a wooden handle *e* ; and, thus, they are converted into semiglobular<sup>1</sup> basins. The best evaporating dishes, however, are those of biscuit porcelain made by Wedgwood, and sold in assortments, the largest of which is capable of holding eight or ten pints. They are flat-bottomed, shallow vessels, *f*, with a lip, *g*, glazed in the inside ; and thin, but of a dense hard texture. They will bear to be heated to the boiling point over a clear hot fire ; but are apt to crack when a flame is allowed to play on them, or when the liquor is boiled to dryness, at the moment the last drop of fluid is expelled, unless the fire be much lowered.

It is preferable, therefore, when glass or earthenware vessels are employed, to apply the heat by the medium of sand ; or, if a still more moderate heat be necessary, by means of boiling water, over which the evaporating dish should be placed. The first is denominated a *sand-bath* ; the second, a *water-bath* ; but for processes on a large scale, shallow iron pots or leaden troughs are used, to which the fire is directly applied.

*Exsiccation* is a variety of evaporation, producing the expulsion of moisture from solid bodies by means of heat. It is generally employed for depriving salts of their water of crystallization. They are exposed to the action of a fire in an iron ladle or pot, or in a glass vessel ; and after dissolving as they are heated, in the water they contain, or undergoing what is termed the *watery fusion*, the water boils ; and, evaporating, leaves the salt in the form of a dry mass. When the substances to be exsiccated are liable to decomposition in a temperature above 212°, as is the case with some of the compound oxides, the process must be conducted by the heat of a water-bath.

*Distillation* differs from evaporation only in the circumstance, that the vapour of volatile matter is elevated to be condensed




<sup>1</sup> The iron ring for this purpose is made red-hot in the fire, then put upon the matrass which is to be cut ; and, when the glass is sufficiently heated, by throwing on it a little cold water, it will generally break exactly at the circle heated by the iron ring.




in close vessels, and preserved. The mode of conducting the operation, and the regulation of the heat, differs according to the nature of the substances operated on.

The simplest distilling apparatus, for smaller processes, is the *Retort* and the *Receiver*. The former consists of a nearly globular body, *h*, with a long, gradually tapering neck, *i*, which is bent nearly at a right angle with the body. This is the simplest kind of retort; and

if the materials to be distilled be liquid, they should be poured into the body of the vessel by means of a very long funnel, *k*, which, by reaching completely into it, prevents any thing from trickling down the sides of the



neck. In withdrawing the funnel, it is necessary to keep it applied to the upper part of the retort, that the drop hanging from it may not touch the inside of the neck. For nicer purposes the tubulated retort, *l*, is to be preferred. The bottom of either kind should be very thin, and of a uniform degree of thickness, so as to bear the sudden application of heat from an Argand lamp, or even from a naked fire. The receiver, *m*, should be larger than the retort, and of a globular form, so as to allow of a large surface for cooling the condensing vapour: and it may be either joined directly to the retort, by the neck of the latter passing into it, or by the intervention of a third piece, *n*, denominated an *adopter*; and




in either case the joinings are usually protected by lutes. When the substance to be condensed is of a very volatile nature, as ether, for instance, the receiver must be artificially cooled; and kept during the whole process at the temperature of the atmosphere, either by surrounding it with ice, or allowing water to trickle slowly over it, brought down from a trough placed above the receiver, by means of worsted threads. The constant evaporation which the water suffers, on the surface of the receiver, keeps it at the requisite degree of temperature for condensing the ether. Both the retort and the receiver may be tubulated.


Sometimes, instead of the retort and receiver, the stone-



ware cucurbit, *o*, with its capital, *p*, and receiver, *q*, is used. It is necessary, occasionally, to coat the retort, and the latter-mentioned vessels, with sand and clay, to enable them to sustain a high temperature, and the sudden alternations of heat and cold to which they are liable in common operations. By these



kinds of apparatus, acids and other substances which arise from chemical decompositions aided by heat, are distilled; and the process is named distillation *per latus*: but if the products be highly volatile, or of a gaseous nature, the pneumato-chemical apparatus, to be afterwards described, is required.



For the preparation of alcohol, and of distilled waters, the common still is employed. It consists of two parts, — the boiler, *b*, and the head or capital, *a*. The boiler, which is the part to which the fire is applied, and contains the materials, is generally of a cylindrical shape, and may be sunk into a furnace, or immersed in a water-bath, *d*, when the temperature requires to be nicely regulated. The *head* or *capital* is a large hollow globe, the upper part of which is drawn




out into a tapering pipe, *c*, bent to a curve or arch, and terminating in the serpentine, or worm *e*. These parts are generally made of copper; but the *worm* is a long pewter pipe, of a decreasing diameter, which winds in a spiral direction obliquely through a deep tub, *B*, filled with cold water. The body, head, and worm require to be luted together; but in general, slips of paper dipped in flour paste, or pieces of wet bladder, are sufficient for this purpose. In this apparatus, the vapours are raised into the *head*, whence they pass into the *worm*, where they are condensed, and issue in drops from the lower end of the pipe, *d*. By degrees the water in the refrigeratory becomes warm, and requires to be renewed; and thence the necessity of the tube being furnished with a stop-cock, by which the heated water may be drawn off without disturbing the apparatus. As in this species of distillation the vapour ascends before it is condensed, it is named distillation *per ascensum*.

In some cases, as in the distillation of several essential oils, the vapour, instead of passing laterally, or ascending, is forced to descend. To produce this effect, a plate of tinned iron is fixed within any convenient vessel, so as to leave a space beneath it; and the materials to be distilled being laid upon this, they are covered by another plate accurately fitted to the sides of the vessel, and strong enough to support the fuel, which is burnt upon it. By this means, the volatilized matter of the materials, under the fire, is forced into the lower cavity of the vessel, and there condensed. This mode of distilling is denominated distillation *per descensum*.

In many processes, a large proportion of the vapours which are extricated is incondensable; and, unless there were some means by which these could escape, the apparatus would be burst in pieces. To prevent accidents, therefore, a small hole was generally left, either in the joinings of the vessels, or in the receiver, which could be kept shut, and occasionally opened when the quantity of confined vapour was supposed to be such as might endanger the rupture of the vessels. By this contrivance, however, much condensable vapour escaped, and a large proportion of the products of the distillation was necessarily lost. This defect of the old apparatus was first attempted to be remedied by Glauber, whose hints were improved by Woulfe, the inventor of the apparatus now commonly employed. It consists of a retort, *b*, generally tubulated, in which the materials are heated by a lamp, *l*; a receiver, *c*, supported by a stand, *m*, to detain any part of the product which is condensable by cold; and a bent tube, proceeding from the receiver to the bottom of a bottle, *e*, with two apertures, and which is about half full of water. Several bottles, however,







are generally employed; and these, being placed side by side, are connected with one another by means of bent tubes, one limb of each proceeding from the top of the bottle immediately preceding, and the other plunging to the bottom of the liquid of the bottle next in order. The joinings of the apparatus are all made air-tight, except the opening of the last bottle furthest from the retort, so that any vapour which escapes must have passed through the liquid in the whole series of bottles, and left all its condensable matter before it can escape. One inconvenience, however, attends this apparatus when it contains no other parts than the above; which is, that after the distillation, as the retort cools, a vacuum is produced in it and the first receiver, which induces a suction or absorption from the other receivers through the bent tubes, and a retrograde motion of the liquid contained in them takes place through the whole apparatus; so that the products are mixed, — unless the operator be on the watch, to separate the retort and receiver, the moment the liquor begins to rise in the bent tube between the receiver and the first bottle. The best contrivance for remedying this defect is the tube of safety, invented by *Welter*, and represented at *d*. It is a bent tube with a bulb blown in that part of it which lies between the upper and lower flexure; and a small funnel at the top. This tube is sometimes used as a stopper to the tubulure of the retort, or to a separate opening in the receiver; or, as is represented in the cut, it is cemented into the tube passing from the receiver to the first bottle. When it is to be used, a little mercury is dropped into the funnel, so as to occupy the space of the tube which lies between the two lower flexures. The mercury excludes the external air during the distillation; but as soon as the vacuum is formed by the cool-



ing of the vessels, the mercury is forced by the pressure of the atmosphere into the bulb; and not being in sufficient quantity to fill it, the external air passes by it in the bulb, and rushes into the apparatus; by which means the vacuum is filled up, and the absorption of the liquid prevented. Various modifications of this have been suggested; but the best is the invention of M. Pepys. It consists of a globular or rather pyriform vessel, *c*, with which the receiver, *b*, is surmounted, and into which it is accurately ground. This is furnished with a glass valve, which allows gas to pass freely into it from the receiver, but prevents the water which it contains from falling into the receiver.



In chemical operations, when the gases which are separated during a process are to be preserved, the pneumatic trough, *h*, with its inverted jar, *i*, under which the curved tube, *k*, proceeding from the bottle *g*, opens, is attached to Woulfe's apparatus. The construction of the trough differs according to the nature of the fluid with which it is filled.




If water be employed, the trough may be of stone ware, or of tinned iron well japanned, and of an oblong or a circular shape. It may be about 18 inches long, 14 broad, and 8 inches deep; with a shelf of the same materials, which should extend entirely across the trough, and have two small holes in it to convey the gas into the inverted jars set upon it; and two larger holes to receive two bottle supporters. This trough should be nearly filled with water, and the jars intended to hold the gas should be, also, filled with the same liquid, and inverted; so that when placed upon the shelf, the water in the trough may ascend about half an inch up their sides, which enables them to retain either water or gas. If mercury be employed, which is essential when the gases to be extri-



cated are absorbable by water, the trough may be made of some hard wood, as mahogany, or of marble. It is not required to be so large as the trough for water, and one part only need be sunk; the shelf should be on each side of this part which is called the well; and it is useful to have an iron or brass stem supporting a semicircular clip fastened into the substance of the trough, to support the jar when it is filled with mercury and inverted. By this apparatus any gases given out during distillation may be collected and preserved; but this is a circumstance, in pharmaceutical operations, which is attended to more with the view of guarding the operator against the effect of noxious gases, than of preserving gases for examination.

*Rectification* is the repeated distillation of any product obtained by distillation, when it is not perfectly pure. This second operation is carried on at a lower temperature, so that the more volatile parts only are raised, and pass over into the receiver, leaving the impurities behind. When the fluid is simply rendered stronger, as in the case of alcohol, by bringing over the spirit, and leaving behind the superfluous water, the operation is named *dephlegmation* or *concentration*. When the liquid is distilled off from any substance, the process is called *abstraction*; and *cohobation*, if the product be re-distilled from the same materials, or from a fresh parcel of the same materials.

*Sublimation* is a species of distillation, in which the product of the volatilization is condensed in a solid form; but as this condensation takes place at a higher temperature than that of watery vapour, a much more simple apparatus is required. The process is conducted sometimes in a crucible, with a cone of paper or another crucible inverted over it, in which the product is condensed; and as in this case it is light and spongy, it was formerly denominated *flowers*. For other matters which are less volatile, a cucurbit and capital, or a flask, or phial, are employed, and sunk about two-thirds in a sand-bath. The aludel, *a, a, b, b, c*, which may be extended to any number of vessels, is generally used in this operation. The product in these cases is solid, and is denominated a *sublimate*.



*b. Of the operations by which chemical changes are produced in the forms of bodies by the action of water, and other fluids.*

When a solid body is thrown into a liquid and disappears, the transparency of the liquid remaining the same, the process is named *solution*; or *solution* is that operation by which the



aggregation of a solid is overcome by a liquid, and a compound produced, which, retaining the fluid form, is transparent, and perfectly homogeneous. The liquid is generally supposed to be the substance exerting the active power, and has therefore been called the *solvent* or *menstruum*. It separates the particles of the solid or *solvend* from each other, and permanently suspends them by the state of combination into which they enter; but the attraction, as was before stated, is reciprocal, both as it regards the solid and the fluid. In general, the solution of every solid in a liquid can be effected in a certain quantity only, or is limited; and when it is carried to its ultimate point, the liquid is said to be saturated. The solvent power, however, is not always limited; there being some instances in which a solid dissolves in a liquid in any proportion: thus gum and sugar dissolve in water in every proportion. The solvent power of a fluid diminishes as it approaches to saturation, and the solution consequently goes on more slowly; but by raising the temperature, it proceeds again more rapidly, and a much larger portion of the solid is taken up than could have been dissolved at a lower temperature. This effect of temperature, however, does not take place in every instance; for muriate of soda, for example, and some other salts, can be dissolved in nearly as great quantity by cold as by hot water. When an increase of temperature increases the solubility of bodies, a portion of the solid, taken up by a heated liquid, is retained in combination as long as the increased temperature exists, but separates again as the solution is cooled down to the temperature of the atmosphere, or lower; and when this is properly conducted, salts are obtained in regular forms, or crystallization takes place.

Although a liquid be saturated with one solid, yet it may be still capable of dissolving a portion of another, and even of a third when saturated with the second; until it be combined with, or hold in solution, three, four, or five different bodies at the same time. The liquid, indeed, in this case does not dissolve so large a portion of any of the substances; but sometimes, from the mutual affinities which the substances exert, the whole proportion of solid matter dissolved is very much increased.

The solution of saline bodies in water requires no particular apparatus; as it can be conducted equally well in phials, or jars, or basins, provided the materials of which they are composed be such as can resist the action of the solvent.

*Lixiviation* is a term applied to solution when the saline body consists of both soluble and insoluble ingredients. On a great scale it is generally performed in large tubs, or vats, having a hole near the bottom containing a wooden spigot and faucet. A layer of straw is placed at the bottom of the tub




over which the substance is spread, and covered by a cloth; after which hot or cold water, according as the salt is more or less soluble, is poured on. The water, which soon takes up some of the soluble parts of the saline body, is after a little while drawn off by the spigot; and a fresh portion of water is successively added and drawn off, until the whole of the soluble matter be dissolved. The straw in this operation acts as a filter; and the cloth prevents the water from making a hollow in the ingredients when it is poured on, by which it might escape without acting on the whole of the ingredients.

In smaller operations lixiviation may be conducted in glass matrasses, and the *ley*, which is the name given to the solution, filtered through paper in a glass funnel.

*Maceration* is that operation by which the soluble parts of substances, chiefly of a vegetable nature, are obtained in solution by keeping them immersed in *cold* water or in spirituous fluids for a sufficient length of time. It is frequently employed as a preparation for infusion and decoction, which are always rendered more effective by the previous maceration of the materials.

*Digestion* is an operation similar to maceration, except that the power of the fluid is aided by a gentle heat. It is usually performed in a glass matrass, *p*, and the evaporation of the liquid impeded by stopping the mouth of the matrass slightly, with a plug of tow, or tying over it a piece of wet bladder perforated with small holes. When the menstruum is valuable, as alcohol, for instance, another matrass, with a smaller mouth, may be inverted over the former, and the joinings secured by a piece of wet bladder; or, what is perhaps preferable, a long open glass tube may be luted to the mouth of the matrass which contains the materials. By these means, any part of the liquor which is resolved into steam by the heat, is condensed, and conveyed back upon the materials. The matrass may be heated either by a common fire, or a lamp, a water-bath, or a sand-bath; and, when either of the latter are used, the matrass should not be sunk deeper in the water, or the sand, than the portion that is filled. The process has been denominated *circulation*, when the condensed vapours are returned upon the ingredients.



*Infusion* is intended principally to extract the volatile and aromatic principles of vegetable substances, which would be dissipated by digestion or decoction; and also those parts of vegetables which are more readily soluble in water, as gum, sugar, extract, tannin, the salts, and part of the resin, from the insoluble parts. The water is poured boiling hot on the materials, sliced, or reduced to a coarse powder, and kept in a closely covered vessel until they are cold; when the infusion or



liquor is decanted off for use. The best infusion-pots are of a globular form in the body, *q*, with the neck, *s*, cylindrical, and having a very large lip or spout, *r*, furnished with a grate, which should incline inwards towards the top, so as to retain the ingredients in decanting off the infusion. Infusions differ according to the length of time the water has stood on the materials, and the heat used. In some instances agitation is necessary. Infusions may be made in the cold; and these are in general more grateful, although weaker.



*Decoction*, or boiling, is intended to answer the same purposes as infusion; but in a more extended degree. The solvent power of the menstruum is increased by the degree of heat: hence the liquor is deeper coloured, and more loaded with the soluble principles of the vegetable. Decoction is employed with advantage to extract the mucilaginous parts of plants, their bitterness, and several other of the vegetable principles. It is generally performed in slightly covered vessels; but when the menstruum is valuable, as alcohol, for instance, a retort and receiver, or the common still, may be used, in the body of which the decoction is prepared, while the vapours that would otherwise escape are condensed and preserved.

Decoction, however, is often a prejudicial mode of preparation, particularly for those vegetables the virtues of which depend wholly or in part on the essential oil, or other volatile principles they contain; and even some fixed principles, such as Extractive, are injured by it.

*Extraction*. If the liquor, which is obtained by either infusion or decoction, be subjected to evaporation, the watery part is dissipated, and the part which was extracted by it is obtained in the solid form, and is denominated an *Extract*. The same objections may be urged against this species of preparation as were stated under *Decoction*.

All the forms of preparation in which water is the agent may be regarded as various modifications of solution. When alcohol or diluted spirits are employed as menstrua, the ingredients subjected to their action are generally macerated, and the filtered fluid, which is the product, is denominated a *Tincture*.

*c. Of the operations by which changes are produced in bodies by chemical agents.*

Under this division we have only to notice *Coagulation*, which is the conversion of a fluid into a solid more or less consistent. The means employed for this purpose are increase of temperature, alcohol, acids, and runnets. The effect appears to arise from a new arrangement of particles produced by the



affinity exerted between the solid particles contained in the fluid and the coagulating substance.


*b. Of the OPERATIONS in which changes are produced by the chemical action of one set of bodies upon another.*

These are — Decomposition,  
Dissolution,  
Precipitation,  
Crystallization,  
Fermentation.

*Decomposition* implies the separation of the component parts of bodies from one another. It is produced, in some cases, by heat, or the introduction of caloric into a body in sufficient quantity to separate the particles from one another to a distance beyond the sphere of the attraction of affinity which held them in combination. It may be effected, also, by electricity or galvanism; but in the greater number of instances it is the result of a superior affinity, which breaks the weaker affinity that holds the principles of the substance about to be decomposed in union, and produces new compounds.

In pharmaceutical operations decomposition frequently occurs; and it is of the utmost importance, in extemporaneous prescription, to be acquainted with the circumstances which occasion it.

*Dissolution* is the appellation given to cases of solution accompanied with decomposition, or some alteration in the nature of the dissolved body. In general, the dissolution of a body is attended with considerable effervescence, owing to the extrication of gases; and therefore the operation requires to be performed in capacious vessels, to prevent the loss of the materials. When the gas is required to be preserved and the operation is not on a large scale, the proof bottle *b* is used. It is furnished with a tubulated orifice *c*, for permitting the introduction of acid or any other decomposing agent; and *a*, a curved tube, which is ground into the opening of the bottle through which the gas escapes. The gas is collected in inverted jars in a pneumatic trough.



*Precipitation* is an operation in which decomposition, also, takes place, a solid substance being thrown down from a liquid in which it was held in solution, by the chemical action of another body which is added to the solution. The substance employed to produce the precipitation is denominated the




*precipitant*; the substance which is separated by its action, the *precipitate*. Thus, if into a solution of sulphate of magnesia a solution of soda be dropped, the magnesia separates from the sulphuric acid, falls to the bottom, and forms the precipitate; while the alkali, which is the precipitant, combining with the acid, thus set free, remains in solution in the state of sulphate of soda. Sometimes, the precipitate is separated by the precipitant having a greater affinity for the liquid, and thence weakening its attraction to the substance which it held in solution. Alcohol, for example, when added to a saturated solution of sulphate of magnesia, precipitates the salt in a crystallized form, and combines with the separated water. In other cases the precipitate is an insoluble compound formed by the union of the added substance with that which was previously held in solution; as, when a solution of barytes is added to a solution of sulphuric acid, sulphate of barytes is formed and precipitated. The mixture of a solution of a compound salt with the solution of another compound salt may produce a precipitate which is an insoluble compound, while a new soluble compound is formed at the same time and remains in solution; in which case the decomposition is produced by double elective attraction. Thus, if a solution of acetate of lead be added to a solution of sulphate of zinc, the oxide of lead leaves the acetic acid, and combining with the sulphuric, forms sulphate of lead, which is insoluble and falls to the bottom; while acetate of zinc, formed by the union of the oxide of zinc with the acetic acid, remains in solution.

When the precipitate is the chief object of the process, it is necessary to wash it, after it is separated by filtration. This operation requires little attention when the substance thrown down is insoluble in water; but when it is in some degree soluble, attention is required to prevent the loss which might result from the use of too much water.

The best precipitating vessel is a very tall glass jar, *t*, narrower at the bottom than at the mouth, so that the precipitate may readily collect by subsidence, and the supernatant liquor be decanted off with more ease.

Precipitation is intended to separate solids from solutions in which they are contained; to produce new combinations, which cannot readily be formed by the direct union of their constituents; and to purify solutions from perceptible impurities. A knowledge of those substances which produce precipitation is, also, of much importance, in extemporaneous prescription; to prevent the virtues





of remedies from being destroyed by improper combinations. The following Tables of Precipitants are extracted from those drawn up by Dr. Thomson.<sup>1</sup> All the substances not employed in Pharmacy are omitted.

1. ALKALIES.	PRECIPITANTS. <sup>2</sup>
Potash.....	Tartaric acid.
Soda.....	0.
Ammonia.....	Fixed alkalies.
2. ALKALINE EARTHS.	
Barytes.....	Sulphuric acid, sulphates.
Lime .....	Oxalic acid, oxalates.
Magnesia.....	Phosphoric acid, phosphate of soda. <sup>3</sup>
3. EARTHS PROPER.	
Alumina.....	Ammonia, hydrosulphuret of potash.
4. METALLIC OXIDES.	
Silver.....	Muriate of Soda.
Mercury.....	Muriate of soda.
Copper .....	Iron.
Iron.....	Succinate of soda, benzoate of soda.
Lead.....	Sulphate of soda.
Zinc.....	0, alkaline carbonates?
Antimony .....	Water, hydrosulphuret of potash.
Arsenic .....	Nitrate of lead.
5. ACIDS.	
Sulphuric.....	Muriate of barytes.
Carbonic.....	Muriate of an alkaline earth.
Boracic .....	Sulphuric acid.
Nitric .....	0.
Acetic .....	0.
Benzoic .....	Muriatic acid.
Succinic .....	Sulphate of iron.
Oxalic .....	Muriate of lime.
Tartaric .....	Potash.
Citric .....	Acetate of lime.

In some cases, when decomposition is effected by the addition of another substance, the separated body is not precipitated, but rises to the surface, and is thence denominated a *cream*: thus, by the addition of any acid to a solution of soap,

<sup>1</sup> *System of Chemistry*, 5th edit. iii. 149. — 154.

<sup>2</sup> In strict language no precipitation takes place, but the fixed alkalies added to solutions containing ammonia render it perceptible by its odour.

<sup>3</sup> The precipitation of magnesia in this case is not direct; but to effect it a solution of carbonate of ammonia must first be added to the solution of muriate of magnesia; no precipitate will appear, but on adding phosphate of soda it falls down in an insoluble state, in combination with the phosphoric acid. Dr. Wollaston suggested this method.



the alkali unites with the acid, while the oil is separated, and swims on the surface of the liquor.

*Crystallization*, although it can scarcely be regarded as a species of precipitation, yet is very nearly allied to it. We have already noticed the theory of the operation, and therefore it only remains to mention in this place the modes in which it is effected for pharmaceutical purposes.

For the crystallization of any substance, it is necessary that it should be either in the aëriform state or a state of fluidity, produced either by the agency of caloric or that of water.

Thus Camphor, Iodine, Arsenic, and Benzoic acid pass at once from the state of vapour into a regular crystalline form; while metals and other bodies, which are capable of being fused, crystallize if they be allowed to cool very slowly, and are left at the same time in a state of rest: but this species of crystallization is never required for pharmaceutical purposes.

Salts are obtained in a crystalline form by a proper management of their watery solutions. When the salt to be crystallized is considerably more soluble in hot than in cold water, as, for example, *sulphate of soda*, it is only necessary to saturate hot water with the salt, and set it aside to cool; but this must be slowly effected, by covering the vessel with a cloth to prevent the access of cold air, and the too rapid consequent formation of a pellicle, which would produce an irregular mass, instead of well formed, distinct crystals. Crystals, thus formed, generally contain, in a state of combination with the salt, a considerable proportion of water, which is thence termed *water of crystallization*. When the salt is not more soluble in hot than in cold water, crystals are obtained by evaporating the solution while hot, until a pellicle form on its surface; when it is set aside to cool, during which the crystals form. After these are separated, the evaporation is repeated, and another crop obtained, until, by a succession of evaporations, the greater part of the salt contained in the solution is separated in the crystalline state. The same effect may be produced by spontaneous evaporation.

The following method of obtaining very large and regular crystals has been pointed out by M. Leblanc.<sup>1</sup> The solution is first evaporated to such a consistency that it shall crystallize on cooling: when it is cold, the liquor is poured off from the mass of crystals, which generally form at the bottom, and is put into a flat-bottomed vessel. In this, solitary crystals gradually form, the largest of which are to be picked out and placed in another flat-bottomed vessel at some distance from each other, and a quantity of liquid, obtained in the same

---

<sup>1</sup> *Journal de Physique*, lv. 300.



way by evaporating a solution of the salt till it crystallizes on cooling, poured over them. The position of each crystal is now to be altered once a-day by means of a glass rod; for, when not turned, the face on which the crystal rests receives no increase of size. When they have gained considerably in magnitude the most regular are to be selected, and each of them put separately into a vessel filled with the same liquid, and turned as already described several times a-day, until they attain the largest size which the species of crystal under treatment is capable of acquiring. It is, however, necessary to observe, that if the crystals be allowed to remain too long in one portion of the solution, the quantity of salt it contains becomes so much diminished, that the liquid re-acts upon the crystal, and partially dissolves it.

If a crystallizable salt be perfectly pure, the whole of its solution may be crystallized: but, if two or more salts exist in the same solution, after crystals have been obtained by several successive evaporations and coolings, the remaining portion of the fluid, although saturated with saline matter, yet refuses to crystallize, and is then denominated *mother water*.

The vessels best adapted for crystallization are large flat dishes of Wedgwood's ware, such as have been already described (p. 87.) as proper for the evaporation of liquids. When the crystallization is to be conducted slowly in the heat of the atmosphere, with the free access of air, deeper vessels are required, that there may be a considerable body of liquid; by which means crystals of considerable size and very regular in figure are procured. Sometimes the crystallization, particularly when it is effected without the aid of heat, is disturbed by minute crystals forming on the edge of the vessel, and the saline matter extending itself over its sides. This is best prevented by smearing the edge of the evaporating dishes with a little oil.

Crystallization is intended to obtain crystallizable substances in a pure state; and to separate them from one another, by taking advantage of their different solubility at different temperatures.

#### FERMENTATIONS.

The constituents of vegetable matter, when separated from the living plant, and placed under certain circumstances, act upon one another, and a spontaneous decomposition takes place, even at the ordinary temperature of the atmosphere. This process has been denominated *Fermentation* by chemists, on account of the intestine motion with which it is accompanied: and, as its phænomena and results vary according to the nature of the vegetable matter subjected to it, and the circumstances



under which it occurs, the general process is divided into three species easily distinguished from one another. The 1st is named the *vinous fermentation* — of which the products are wine, beer, and other vinous fluids : the 2d, the *acetous fermentation*, which produces acetic acid or vinegar : and the 3d, the *putrefactive fermentation*, in which gases, chiefly foetid, are produced, and ammonia.

Each of these is occasionally artificially produced for pharmaceutical purposes, and therefore requires to be described.

*Vinous Fermentation.* All vegetable substances containing saccharine matter, and a peculiar glutinous principle analogous to the gluten of wheat, are susceptible of this fermentation. For its commencement, however, the presence of water, sugar, extract, and a small proportion of vegetable acid, with a certain increased temperature, is requisite. In juices in which these are present, the fermentation is spontaneous ; but as *yeast* contains the peculiar gluten, and the other principles necessary for exciting the vinous fermentation in any sweet vegetable juice or decoction, it is frequently used for this purpose in the formation of beer and wines. Soon after yeast is added to these substances, or to *wort*<sup>1</sup>, or to *must*<sup>2</sup>, an intestine motion commences in the liquor, its temperature rises, it becomes turbid, and carbonic acid gas is extricated : but, after some time, the fermentation again gradually subsides, the scum which was formed during its continuance rises to the surface, or sinks to the bottom ; the liquor becomes lighter, and instead of its sweet taste has acquired that peculiar taste and flavour which is denominated vinous. This process of fermentation is never employed in the laboratory for the preparation of vinous liquors, although these are articles of the *materia medica* : but the cataplasms, which are prepared from carrots and similar vegetables mixed with yeast, derive their virtues from the vinous fermentation into which they enter, extricating a large quantity of carbonic acid gas, which operates as a powerful antiseptic.

*Acetous Fermentation.* All liquors prepared by the vinous fermentation are susceptible of the acetous when kept exposed to the air in a temperature between 70° and 90°. Under these circumstances the liquor gradually becomes thick, its temperature increases, and filaments are seen moving through it in every direction, an intestine motion being excited, accom-

---

<sup>1</sup> Wort is an aqueous infusion of malt. It consists of saccharine matter, starch, gluten, tannin, and mucilage.

<sup>2</sup> Must is the expressed juice of the grape. It contains water, sugar, a peculiar matter, which changes into gluten by contact with the air, mucilage, supertartrate of potash, tartrate of lime, muriate of soda, and sulphate of potash.



panied with a hissing noise : but as this motion subsides, these filaments fall to the bottom, or attach themselves to the sides of the vessel, the liquor becomes clear and transparent, and has acquired a very sharp acid taste. In this state it is denominated vinegar, and contains, besides the acetic acid and water, which are its principal components, mucus, malic acid, supertartrate of potash, and some other vegetable constituents.

Pure alcohol, even when diluted with water, is not susceptible of this fermentation, but it enters into it when united with other fermentable bodies : thence wine and vinous liquors, which contain, besides alcohol, sugar, and some mucilaginous and extractive matter, are employed for making vinegar. In this process the alcohol is supposed to be decomposed, and oxygen absorbed ; carbonic acid is formed, but is retained in the liquor ; and it is probable, as Dr. Murray has suggested, that the ferment affords nitrogen, which it is now ascertained, enters into the composition of acetic acid.<sup>1</sup>

Many vegetable infusions and decoctions undergo this fermentation in warm weather ; and thence the necessity of preparing these every day during summer, as by the decomposition which takes place, their medicinal virtues are completely destroyed, when they are kept.

*Putrefactive Fermentation.* Almost every vegetable product, in a moist place, and in a temperature not under 45°, nor above 70°, undergoes spontaneous decomposition. The solid structure of the body is completely destroyed, and its ultimate principles entering into new combinations, escape in the gaseous or aëriform state, leaving behind a small quantity of earthy and metallic matter only, which it contained.

This process, which is denominated the *putrefactive fermentation*, does not absolutely require the contact of air ; but water, in every case, appears to be essential. Vegetable bodies, which are very soluble in water, most readily undergo it ; the surface of the liquor becomes covered with a mould ; various elastic fluids, in which ammonia and phosphuretted hydrogen gas are often perceptible, with other matters which produce a foetid odour, are extricated, and the vegetable matter is ultimately completely decomposed.

The knowledge of the circumstances which promote this species of spontaneous decomposition points out the necessity of preserving vegetable substances in perfectly dry places ; and, when they have a tendency to attract moisture, of exposing them in a free current of air to dissipate the humidity which they would otherwise absorb.

---

<sup>1</sup> *System of Chemistry*, iv. 465.



*c. Of the OPERATIONS in which oxidizement and deoxidizement are effected by means of a high temperature.*

The degree of temperature at which these operations are conducted cannot generally be obtained from a common fire; and, therefore, before describing the operations themselves, it is necessary to notice the nature of furnaces, which are instruments of the most universal use in pharmaceutical chemistry.

*Furnaces* differ in construction, according to the particular purposes for which they are chiefly intended; but the following essential parts are common to all furnaces. 1st, The body or fire-place for holding the fuel and the vessel containing the materials to be submitted to the action of heat. 2dly, The chimney by which the heated air and the smoke escape. 3dly, The ash-pit, into which the ashes fall, and through holes in the side of which fresh air is admitted to the burning fuel.

In a well-constructed furnace, the whole of the air which enters the ash-pit passes through the body of the furnace, and supports the combustion, after which the residue, being highly rarified, passes off by the chimney; on the due height of which, and the proper regulation of the access of atmospheric air from below, the strength of the combustion, and consequently the heat produced, altogether depend. The access of the air is, generally, regulated by registers; which, in portable and smaller furnaces, are iron plates pierced with many holes of different sizes, which are generally fitted with brass stoppers, so that according to the number of holes opened, a greater or smaller quantity of air is admitted to the burning fuel. The chimney should be narrower than the body of the furnace, and of such a length that it can be heated throughout by the rarefied air which ascends through it; for it is by producing in the chimney a column of air of much less specific gravity than a corresponding column of the external air, that fresh air is constantly forced through the body of the furnace from below, and a strong draught produced. If the chimney be too short, all the advantage to be derived from the above circumstance is not obtained; and if on the other hand it be too long, the air loses its heat before it reaches the summit, and impedes, to a certain extent, the ascent of the rarefied air. According to Macquer, when the internal diameter of the furnace is 12 or 15 inches, and that of the chimney 8 or 9, its height should be 18 or 20 feet.


Of whatever substance furnaces are made, unless they be fixed and built of fire-bricks, they should be coated, to prevent the radiation, and consequent loss of heat; and the best composition for this purpose is clay and sand. It is perhaps better, however, first to put a coating of charcoal and clay next to



the sides of the furnace, as was recommended by Dr. Black, particularly if it be made of plate-iron. The proportions he recommended were one part, by weight, of fine clay, and three parts of charcoal; which, being reduced to powder, is kneaded together with water, and the mass thus formed, divided into balls of a moderate size. These being applied to the sides of the furnace, are to be beat strongly with the face of a broad hammer, until a general coating of about one inch and a half cover the inside of the furnace, and the cavity assume an elliptical form.

A very convenient and useful furnace is that which was contrived by Dr. Black.


It consists of an oval iron case, *a*, about 22 inches in height, 20 inches in diameter in the length of the oval, and 15 inches across; and lined, in the body, with the coating already described. On the top is fixed an iron plate having two apertures; one large, *f*, intended to receive a sand-bath, a still, or similar apparatus; and the other smaller, *e*, to which an iron tube, which acts as a chimney, is to be fixed. At the bottom of the body of the furnace, directly under the larger aperture, the grate is fixed; and under it the ash-pit receives the body, resting on a strong ring that encircles it, at about half an inch deep. The ash-pit is furnished with a door,




*m*, which opens on hinges, intended for removing the ashes; and also a register to regulate the admission of air to the burning fuel. The register is a plate of iron perforated with six apertures, the size of which increases in a geometrical ratio; so that by passing the sliding plate, *l*, over them either one or more may be stopped at a time, and the supply of air, and of course the heat to be excited, can be regulated with great nicety. The fuel is introduced at the top; but there is a door also, *i*, occasionally, in the side of the body of the furnace, through which fuel can be supplied during the conducting of any process; although, unless it be made to



shut very close, this door is a disadvantage, as it prevents the admission of the air from being so precisely regulated. It has also an opening, *k*, with a sliding door, for receiving a muffle: *k* is an earthen or an iron tube, which passes through the body of the furnace, and issues at the opposite side, intended for procuring hydrogen gas; or for inserting the nozzle of a pair of bellows. Any thing may be suspended over the furnace by inserting an iron pillar into the eyes, *g g*, on each side, and stretching a strong bar or wire between them. The sand-bath, *n*, is placed in the opening, *f*, of the furnace when it is required; but at other times that opening is shut by the cover, *p*, which has a moveable plug, *o*, in its center. A pair of tongs, *r*, bent near the points, is a necessary appendage to this furnace. It may be used for a great variety of operations, and may be fitted with a dome for the purpose of throwing down the flame when it is to be used for fusing metals.



For small operations, when a great heat is required, a furnace may be constructed by simply inverting one large black lead, or a Hessian crucible over another which is perforated with six holes in the bottom, which are intended to serve as a grating. This is placed over the portion of a third, cut off so low as to leave the cavity about an inch deep only; and ground smooth above and below. The upper or inverted pot should have a large perforation to permit the heated air to escape; and the portion on which the second pot stands should also be perforated at the side to admit the external air, or the nozzle of a bellows. No luting is required. A heat sufficient to fuse any metal may be obtained in such a furnace. A sufficient heat for a great variety of small operations may be obtained from a lamp, on the principle of Argand's, with a double concentric wick. A standard, *a*, hav-





ing moveable rings, *b, b*, attached to it, is necessary for supporting the retort or matrass at any height above the flame.

With regard to *fuels*, the best are undoubtedly charcoal and coke, or a mixture of these. The advantages of *charcoal* are its kindling readily, burning with a strong clear heat in a small draught, without running into slag, choking the grate, or melting the walls of the furnace; and owing to its containing only matter which is extremely combustible, the flues or chimneys never collect soot or other foulnesses. The chief objection to charcoal is its great expense. *Coke* is much less expensive; but as it contains a mixture of earth and metallic oxides, it is apt in an intense heat to run together into a tough cohesive slag, which adheres to the walls of the furnace, and to the sides of crucibles, choking up the grate, and of course preventing the proper draught of air for carrying on the combustion. These disadvantages, however, are remedied by mixing it with an equal bulk of charcoal; and this mixture forms the best fuel, when an intense heat is required.

The pharmaceutical operations (usually performed by furnaces) are —

Fusion.

Distillation.

Sublimation.

The oxidizement of metals.

The deoxidizement or reduction of metals.

The three first of these have been already described.

*Oxidizement of metals.* This term signifies that process by which metals are converted into oxides, by absorbing oxygen from the air, when exposed to a certain degree of heat. The disengagement of the caloric and the light which oxygen gas contains, by the solidification of the oxygen in the oxide, is scarcely perceptible when the operation is conducted in atmospheric air; but if the oxidizement takes place in oxygen gas, it is rapidly affected, and caloric and light are very evidently extricated. This mode, however, of oxidizing metals is employed in small experiments only; but in all the processes of the laboratory for procuring oxides by the aid of heat, common air yields the oxygen. The metal, if it be not volatile at the temperature required for its oxidizement, is exposed to the heat of the furnace in a flat dish of baked clay called a *roasting test*, and frequently stirred to present fresh surfaces to the air: but if the metal be easily volatilized, as is the case with zinc, it is thrown by pieces, at separate intervals, into a deep crucible, so placed as to admit the air and allow of the additions being made. If mercury be the metal operated on, it is generally put into a flat-bottomed matrass with a very tall narrow neck, the mouth of which is left open. This matrass is placed in a sand-bath, and kept at a degree of heat nearly equal to the



boiling point of the mercury, for several days : but, it is perhaps better to use a retort with the bottom flattened, and the neck only slightly bent, that the globules of mercurial vapour may be condensed, and the metal fall back into the vessel.<sup>1</sup> In this process the atmospheric air furnishes the oxygen, which readily combines with the volatilized mercury, while the form of the apparatus is intended to permit a renewal of it constantly to take place, without allowing the escape of the mercurial vapour.

*Deoxidizement of metals*, or their reduction, is that process in which the oxygen of a metallic oxide is separated, and the metal recovers its metallic form and properties. It is seldom performed on a large scale in pharmacy ; but in cases of metallic oxides having been taken into the stomach, and proving fatal, it is of importance, in ascertaining their nature, to be able to reduce them to the metallic state by means of the blow-pipe and lamp ; an apparatus by which minute substances may be almost instantaneously heated to a great degree, and their nature discovered with much accuracy.

The most common Blow-pipe is a tube of brass or iron, bent near one of its extremities, and drawn out sufficiently fine to keep up a constant stream of air when it is blown into by the mouth applied to the opposite end. This form of blow-pipe is liable to one inconvenience, from the condensation of the moisture of the breath, in the course of blowing ; to remedy which, a hollow ball or bulb, *a*, is made near the small end of the pipe ; and to render it more portable, this is divided through the middle, and fitted with a screw so as to be put together when used. Small separate jet pipes, or caps, are frequently adapted to slip on the small extremity of the blow-pipe, by which means any size of bore may be had recourse to, as a larger and more moderate, or a smaller and more intense flame is required. The flame for blowing through is best obtained from a wax or tallow candle with a very large wick, which must be kept moderately short by snuffing it frequently, and it must also be turned a little aside from the pipe.




In using this apparatus with advantage and ease, a little practice is necessary. As the flame is often required to be kept up for several minutes, the respiration must be carried on through the nostrils without interruption, and the stress of blowing performed merely by the compression of the cheeks upon the air held in the mouth. In subjecting any substance to experiment, it is to be placed either on a piece of charcoal,

<sup>1</sup> *Higgin's Minutes. Aikin's Dictionary, ii. 75.*



or in a platina spoon. When charcoal is employed, a large, compact well-burnt piece should be chosen, and a small shallow hole scooped in it for receiving the substance to be heated. The flame of the candle or lamp is then to be directed upon this by means of the blow-pipe. The charcoal soon kindles round the hole, which is gradually enlarged; and the heat being thereby augmented and kept up uniformly round the substance, the charcoal aids, by its chemical effect, the reduction of the substance or its deoxidizement. Carbonate of lead thus treated is converted into a globule of metallic lead; and the phosphates are partially reduced to phosphurets.

In many operations, much inconvenience arises in using the common blow-pipe from both the hands of the operator being engaged; and therefore a double pair of bellows, which is fixed below the table, and worked by the foot of the artist, has been invented for giving the blast. Means have also been contrived for producing the blast by a stream of oxygen gas, or of mixed gases, as of oxygen and hydrogen, which excite a much more intense heat than can be produced by any other method.<sup>1</sup> A very ingenious blow-pipe is that of Mr. Paul of Geneva, in which the flame is produced by vapour of alcohol.<sup>2</sup>



<sup>1</sup> For descriptions of a blow-pipe invented by Mr. Newman, for this purpose, and improved by Dr. Clarke and others, see *Journ. of Science and the Arts*, vol. ii. p. 104. *Annals of Philosophy*, vol. viii. *passim*. *London Med. Repository*, vol. vi. p. 376. vol. vii. p. 21.

<sup>2</sup> *Mr Paul's alcohol blowpipe:* *a*, A hollow frame of wood, five inches in its longest dimension, supporting the pillar, *d*, and the two lamps, *b*, *c*; the rim, *e*, slips upon the pillar, *d*, as low as the shoulder of the latter will permit; but it may be raised or lowered at pleasure, and kept fast by the screw peg, *f*. The rim supports *g*, the boiler, which is a hollow piece of thick brass, which will hold about fʒj of alcohol, and has four openings; three, *h*, *i*, *k*, at the top, and one at the bottom, to receive the tube *o*. The latter is long enough to reach the level of the outside of the boiler, and consequently the alcohol in the boiler cannot



## COATINGS, CEMENTS, AND LUTES.

In many chemical operations, although the nature of the substances require that glass vessels be used, yet, from the degree of heat to which they are exposed, these must be protected on the outside by a coating; and in all operations where the products are in any degree volatile, it is of importance that the joinings of the parts of the apparatus should be perfectly secured: thence the necessity of coatings and lutes. Cements are requisite for repairing flaws and cracks.

*Coatings* are applied to the insides of furnaces to prevent the too quick dissipation of the heat, and also to protect the iron and materials of which the furnace is made from being destroyed by the action of the fire. The coating used by Doctor Black has been already described; but another nearly as good may be formed, by coarsely grinding fragments of pottery, and mixing the powder with moist clay in sufficient quantity to allow it to be moulded when wet. To render it more tenacious, some fibrous matter is generally added to the mixture, such as chopped cow-dung; the proportion of which, as recommended by Baumé, should be one ounce to every five ounces of the mixture. This is to be applied in the manner already described.

The same kind of coating may be used for glass vessels which are to be exposed to a red heat. The following is the mode of applying it. After kneading the coating material, so as to render it very plastic, let it be spread out on a flat table, and lay the bottom of the retort in the middle of the mass; then turn up the edges of the cake, so as to bring it round the whole of the vessel, pressing it down in every part with the fingers till it apply uniformly and closely. The material may also be applied in the state of thick cream, by dipping the retort repeatedly into it; and drying it after each immersion by turning it before the fire. The different layers of coating may be thus laid on very equally, from the thickness of  $\frac{1}{4}$  to  $\frac{1}{2}$  an inch; so as to make the retort like a strong earthen retort glazed in the inside; and, as the coating agglutinates in a full red heat, it will form an impenetrable covering which cannot be detached from the glass.

---

readily boil over into the tube; and the opening *k*, which corresponds with it, is closely shut by a screw stopper, hollowed out a little beneath, to allow the free passage of the vapour down the tube. By the contiguity of *o* to the lamp *b*, the vapour is prevented from condensing, and as it passes on through the globe *q q*, into the jet tube *r*, it is directly kindled by the flame of the lamp *c*; and the united flames being violently propelled sideways, a long pencil of blue flame is formed, and remains as long as any alcohol is left in the boiler. The boiler is filled at the opening *h*. The central hole, *i*, is nicely fitted with a brass plug, kept down by a thin slip of iron, *l*, which is confined at one end between two flat screws, *m*, *n*, on the top of the upright pillar. This acts as a safety valve, to prevent the vessel from bursting when the vapour cannot escape quick enough at the jet pipe *r*.



*Cements* and *Lutes* are formed of the same materials. They are generally composed of unctuous or resinous substances; mucilaginous or gelatinous substances; or of clay, lime, and similar materials capable of resisting a high degree of heat.

a. *Unctuous and resinous Lutes*. — These should be viscid, plastic, compact, and possess the power of resisting acrid vapours. The following are the best of this class.

1. Melt eight parts of bees' wax with one of turpentine; and, according as it is required to be more or less consistent or pliable, add different proportions of any resinous substance. This lute adheres very closely to the glass, is not easily penetrated by acrid vapours, and is very manageable. It cannot bear a heat higher than 140°. <sup>1</sup>
2. Dissolve spermaceti, and when melted, while it is hot, throw into it bits of caoutchouc. This is an excellent lute where much heat is not required to be employed.
3. Take pure, dry, unbaked clay finely powdered, beat it for several hours with a heavy iron pestle in a brass mortar, dropping in, slowly, some boiled linseed oil; or, some amber varnish, prepared by melting yellow amber in an iron ladle, and mixing it with linseed oil. This lute can sustain a considerable degree of heat, is impenetrable by acids and spirituous liquors, and adheres very strongly to metallic or glass vessels previously rendered perfectly dry. As it softens in some degree, however, by heat, it is necessary to surround the luting with pieces of wet bladder, and to secure the whole by packthread firmly tied round both above and below the joint. <sup>1</sup> This lute improves by age. It should be kept in a covered pan in a cool cellar.
4. Glazier's putty, which is a composition of chalk and drying linseed oil, resembles very much the above lute in its qualities, and may be used as a substitute for it.
5. Take four parts of common resin, one of yellow wax, and one part of fine brick-dust; melt the two former together, and, when they are melted, stir in the brick-dust. This lute adheres with great firmness, and forms also a good cement for stopping cracks in glass vessels.
6. Six parts of clay, one part of iron filings, and enough of linseed oil to form them into a paste, make a good cement for stopping cracks in iron vessels intended to be strongly heated.
7. The following cement is recommended for joining together glass or steel. "Take of mastich five or six bits as big as peas, and dissolve them in as much alcohol as will render them liquid. In another vessel dissolve as much isinglass (previously soaked in water) in brandy or rum as

---

<sup>1</sup> Lavoisier.



will make two fluid ounces of a strong glue; warm it, and incorporate with it by rubbing, two or three small bits of galbanum or ammoniacum, and the mastich solution. Preserve the mixture in a well-stopped bottle, and gently warm it before use.”<sup>1</sup>

8. A solution of shell lac in alcohol, added to a solution of isinglass in proof spirit, forms a cement that will resist moisture.
- b. *Mucilaginous and gelatinous Lutes* are adapted only for operations which do not require a high temperature, and in which very acrid vapours are not extricated. They are easily applied, are sufficiently adhesive, and can be readily removed, by simply moistening them with water.
  1. Under this head may be mentioned the simple application of moistened bladder. To render it very adhesive, it should be soaked in tepid water, until it feel clammy to the touch; after which it contracts considerably as it dries, and adheres with a sufficient degree of force.
  2. Linseed meal kneaded up with water to a sufficient consistence, and applied pretty thick over the joinings of the vessels, or almond meal treated in the same manner, form very convenient lutes, which dry and become firm in a very short time.
  3. Flour paste spread upon slips of moistened paper forms a sufficiently good lute for many purposes.
  4. Smear slips of linen on both sides with white of egg, then apply these neatly to the joinings of the vessels; and, when they have been applied, shake loosely over them some finely powdered quicklime. This lute dries very quickly, is extremely hard, very cohesive, impervious to water, and impenetrable by most kinds of vapours.
  5. Mix powdered plaster of Paris with white of egg, milk, glue, starch, or any mucilage, and apply it immediately.
  6. Mix together equal parts of clay and lime, with about one-third of flour and white of egg.
  7. Mix together equal parts of colcothar and lime, with white of egg.

All the cements containing lime and gelatinous substances become so very hard that they cannot be separated from glass vessels without the aid of a sharp knife and some force; and, therefore, they can scarcely be applied to very thin vessels. They will not confine very corrosive acid vapours for a great length of time; but are excellent lutes for preserving a complicated apparatus steadily united and air-tight; and they will bear nearly a red heat. They are also the most useful kinds of cement for any accidental crack or failure of a lute

---

<sup>1</sup> *Aikin's Dictionary of Chemistry.*



already applied, although a stream of vapour may be bursting through at the time.<sup>1</sup>

c. *Earthy Lutes* are intended for operations which require a high temperature. The following are the best of this class.

1. Mix burnt gypsum, in powder, with water to the consistence of a thick cream, and apply it immediately. This forms a lute which sets as soon as it is applied, and is firm; but a slight blow will easily crack it.
2. Dissolve one ounce of borax in half a pint of boiling water, and add as much slacked lime as will make a paste. By using a smaller portion of lime, this lute forms an excellent glazing for earthen-ware retorts, over which it should be spread with a brush; but when dry, a coating of slacked lime and linseed oil, beaten till the mixture is plastic, should be laid over the whole of the lute.
3. A very valuable fire lute may be made of about one part of glass of borax, five parts of brick-dust, and five parts clay, finely powdered together, and mixed with a little water when used.
4. The same composition which has been already described as a proper coating for the inside of furnaces, is also an excellent earthy lute.
5. Six parts of clay, and one of iron filings formed into a paste by means of linseed oil, form an excellent cement for stopping cracks in iron retorts or boilers.

If the beak of a retort be too small to fit accurately to the neck of a receiver, the vacancy should be filled up, by introducing short pieces of soft wood or of cork; and if the disproportion be very considerable, a cork must be fitted to the neck of the receiver, and a circular hole made in it sufficient to admit the beak of the retort. The curved tube of a Woulfe's apparatus, when not fitted accurately by grinding, may be also fixed by means of corks. After the parts are thus firmly joined, the luting must be neatly and closely applied over the junctures; and the whole covered with slips of wet bladder, or with linen spread with one or other of the above described cements. The application of the lutes, although apparently very simple, yet requires some management, lest the luting of one juncture should disturb another already luted, which is apt to happen when applying the fillets and ligatures. When an operator, therefore, is not pressed for time, he should always allow the luting of one joint to dry before he applies luting to another; indeed, it is preferable not to apply the fillets and ligatures, until after the luting has been applied to all the joints; and is nearly hard.

---

<sup>1</sup> *Aikin's Dictionary of Chemistry.*



## APPENDIX TO PART I.

## No. I.

## TABLE OF FREEZING MIXTURES.

The following Tables were drawn up by Mr. Walker from actual experiments. They show the degree of cold, or the reduction of temperature, which may be obtained by the different combinations mentioned in the first column.

TABLE I.  
FRIGORIFIC MIXTURES — *WITHOUT ICE.*

Mixtures.		Thermometer sinks.	Degree of cold produced.
	Parts.		
Muriate of ammonia	5	From + 50° to + 10°	40
Nitrate of potash	- 5		
Water	- - - 16		
Muriate of ammonia	5	From + 50° to + 4°	46
Nitrate of potash	- 5		
Sulphate of Soda	- 8		
Water	- - - 16		
Nitrate of ammonia	1	From + 50° to + 4°	46
Water	- - - 1		
Nitrate of ammonia	1	From + 50° to - 7°	57
Carbonate of soda	- 1		
Water	- - - 1		
Sulphate of soda	- 3	From + 50° to - 3°	53
Diluted nitric acid	2		
Sulphate of soda	- 6	From + 50° to - 10°	60
Muriate of ammonia	4		
Nitrate of potash	- 2		
Diluted nitric acid	- 4		
Sulphate of soda	- 6	From + 50° to - 14°	64
Nitrate of ammonia	5		
Diluted nitric acid	4		
Phosphate of soda	9	From + 50° to - 12°	62
Diluted nitric acid	4		
Phosphate of soda	9	From + 50° to - 21°	71
Nitrate of ammonia	6		
Diluted nitric acid	4		
Sulphate of soda	- 8	From + 50° to - 0°	50
Muriatic acid	- - 5		
Sulphate of soda	-	From + 50° to - 3°	47
Diluted sulphuric acid			

N. B. If the materials are mixed at a warmer temperature than that expressed in the table, the effect will be proportionably greater; thus, if the most powerful of these mixtures be made when the air is + 85°, it will sink the thermometer to + 2°.



TABLE II.  
FRIGORIFIC MIXTURES—*WITH ICE.*

Mixtures.	Thermometer sinks.	Degree of cold produced.
Parts. Snow, or pounded ice 2 Muriate of soda - 1	From any temperature.	to $-5^{\circ}$ *
Snow, or pounded ice 5 Muriate of Soda - 2 Muriate of ammonia 1		to $-12^{\circ}$ *
Snow, or pounded ice 24 Muriate of soda - 10 Muriate of ammonia 5 Nitrate of potash - 5		to $-18^{\circ}$ *
Snow, or pounded ice 12 Muriate of Soda - 5 Nitrate of ammonia 5		to $-25^{\circ}$ *
Snow - - - - 3 Diluted sulphuric acid 2	From $+32^{\circ}$ to $-23^{\circ}$	55
Snow - - - - 8 Muriatic acid - - 5	From $+32^{\circ}$ to $-27^{\circ}$	59
Snow - - - - 7 Diluted nitric acid - 4	From $+32^{\circ}$ to $-30^{\circ}$	62
Snow - - - - 4 Muriate of lime - 5	From $+32^{\circ}$ to $-40^{\circ}$	72
Snow - - - - 2 Cryst. muriate of lime 3	From $+32^{\circ}$ to $-50^{\circ}$	82
Snow - - - - 3 Potash - - - - 4	From $+32^{\circ}$ to $-51^{\circ}$	83

N. B. The reason for the omissions in the last column of this table, is, the thermometer sinking in the mixtures to the degree mentioned in the preceding column, and never lower, whatever may be the temperature of the materials at mixing.



TABLE III.  
COMBINATIONS OF FRIGORIFIC MIXTURES.

Mixtures.		Thermometer sinks.	Degree of cold produced.
	Parts.		
Phosphate of soda -	5	From 0° to -34°	34
Nitrate of ammonia	3		
Diluted nitric acid -	4		
Phosphate of soda -	3	From -34° to -50°	16
Nitrate of ammonia -	2		
Diluted mixed acids	4		
Snow - - -	8	From -10° to -56°	46
Diluted sulph. acid	3		
Or, Diluted nit. acid	3		
Snow - - -	3	From 0° to -46°	46
Diluted nitric acid -	2		
Snow - - -	1	From 20° to -60°	40
Diluted sulphuric acid	1		
Snow - - -	3	From +20° to -48°	68
Muriate of lime -	4		
Snow - - -	3	From +10° to -54°	64
Muriate of lime -	4		
Snow - - -	2	From +15° to -68°	53
Muriate of lime -	3		
Snow - - -	1	From 0° to -66°	66
Cryst. muriate of lime	2		
Snow - - -	1	From -40° to -73°	33
Cryst. muriate of lime	3		
Snow - - -	1	From -68° to -91°	23
Diluted sulph. acid	10		

N. B. The materials in the first column are to be cooled, previously to mixing, to the temperature required, by mixtures taken from either of the preceding tables.



## No. II.

## TABLES OF SIMPLE AFFINITY.

The following Tables were drawn up by Bergman, and additions made to them by others at different times. The substance, the attractions of which are to be shown, is placed at the head of a column, and the substances to which it has an attraction, placed beneath, in the order of the forces of attraction.

OXYGEN.	Chrome Bismuth Lead Copper Tellurium Platina Mercury Silver Gold	Arsenic Molybdena	Acids :
Carbon			Arsenic
Charcoal			Lactic
Manganese			Benzoic
Zinc		POTASH, SODA, and AMMONIA.	Acetic
Iron			Boracic
Tin		Acids :	Sulphurous
Antimony		Sulphuric	Nitrous
Hydrogen		Nitric	Carbonic
Phosphorus		Muriatic	Prussic
Sulphur	CARBON.	Phosphoric	Sulphur
Arsenic	Oxygen	Fluoric	Phosphorus
Nitrogen	Iron	Oxalic	Water
Nickel	Hydrogen	Tartaric	Fixed oils
Cobalt		Arsenic	
Copper	NITROGEN.	Succinic	STRONTITES.
Bismuth		Citric	
Caloric ?		Lactic	Acids :
Mercury	Oxygen	Benzoic	Sulphuric
Silver	Sulphur	Sulphurous	Phosphoric
Arsenous acid	Phosphorus	Acetic	Oxalic
Nitric oxide	Hydrogen	Mucic	Tartaric
Gold		Boracic	Fluoric
Platina	HYDROGEN.	Nitrous	Nitric
Carbonic oxide	Oxygen	Carbonic	Muriatic
Muriatic acid	Sulphur	Prussic	Succinic
White oxide of manganese	Carbon	Oil	Acetic
White oxide of lead	Phosphorus	Water	Arsenic
	Nitrogen	Sulphur	Boracic
			Carbonic
			Water
OXYGEN. *	SULPHUR.	BARYTES.	
	PHOSPHORUS.	Acids :	LIME.
Titanium		Sulphuric	
Manganese		Oxalic	Acids :
Zinc	Potash	Succinic	Oxalic
Iron	Soda	Fluoric	Sulphuric
Tin	Iron	Phosphoric	Tartaric
Uranium	Copper	Mucic	Succinic
Molybdena	Tin	Nitric	Phosphoric
Tungsten	Lead	Muriatic	Mucic
Cobalt	Silver	Suberic	Nitric
Antimony	Bismuth	Citric	Muriatic
Nickel	Antimony	Tartaric	Suberic
Arsenic	Mercury		

\* Vauquelin's Table of the affinity of the metals for oxygen, according to the difficulty with which their oxides are decomposed by heat.



TABLES OF SIMPLE AFFINITY — *continued.*

<i>Acids :</i> Fluoric Arsenic Lactic Citric Malic Benzoic Acetic Boracic Sulphurous Nitrous Carbonic Prussic Sulphur Phosphorus Water Fixed oil	<i>Acids :</i> Mucic Citric Phosphoric Lactic Benzoic Acetic Boracic Sulphurous Nitrous Carbonic Prussic	<i>Acids :</i> Lactic Succinic Acetic Prussic Carbonic Ammonia	<i>Acids :</i> Prussic Carbonic Fixed oils Ammonia
<hr/> MAGNESIA.	<hr/> SILEX. Fluoric acid Potash	<hr/> OXIDE OF MERCURY. <i>Acids :</i> Gallic Muriatic Oxalic Succinic Arsenic Phosphoric Sulphuric Mucic Tartaric Citric Malic Sulphurous Nitric Fluoric Acetic Benzoic Boracic Prussic Carbonic	<hr/> OXIDE OF COPPER. <i>Acids :</i> Gallic Oxalic Tartaric Muriatic Sulphuric Mucic Nitric Arsenic Phosphoric Succinic Fluoric Citric Lactic Acetic Boracic Prussic Carbonic Fixed alkalies Ammonia Fixed oils
<i>Acids :</i> Oxalic Phosphoric Sulphuric Fluoric Arsenic Mucic Succinic Nitric Muriatic Tartaric Citric Malic Lactic Benzoic Acetic Boracic Sulphurous Nitrous Carbonic Prussic Sulphur	<hr/> OXIDE OF PLATINA. OXIDE OF GOLD.* <i>Acids :</i> Gallic Muriatic Nitric Sulphuric Arsenic Fluoric Tartaric Phosphoric Oxalic Citric Acetic Succinic Prussic Carbonic Ammonia	<hr/> OXIDE OF LEAD. <i>Acids :</i> Gallic Sulphuric Mucic Oxalic Arsenic Tartaric Phosphoric Muriatic Sulphurous Suberic Nitric Fluoric Citric Malic Succinic Lactic Acetic Benzoic Boracic	<hr/> OXIDE OF ARSENIC. <i>Acids :</i> Gallic Muriatic Oxalic Sulphuric Nitric Tartaric Phosphoric Fluoric Succinic Citric Acetic Prussic Fixed alkalies Ammonia Fixed oils Water
<hr/> ALUMINA.	<hr/> OXIDE OF SILVER. <i>Acids :</i> Gallic Muriatic Oxalic Sulphuric Mucic Phosphoric Sulphurous Nitric Arsenic Fluoric Tartaric Citric		<hr/> OXIDE OF IRON. <i>Acids :</i> Gallic

\* Omitting the oxalic, citric, succinic, and carbonic, and adding sulphureted hydrogen after ammonia.



TABLES OF SIMPLE AFFINITY — *continued.*

<i>Acids :</i> Oxalic Tartaric Camphoric Sulphuric Mucic Muriatic Nitric Phosphoric Arsenic Fluoric Succinic Citric Lactic Acetic Boracic Prussic Carbonic	<i>Acids :</i> Tartaric Phosphoric Citric Succinic Fluoric Arsenic Lactic Acetic Boracic Prussic Carbonic Fixed alkalies Ammonia	Glucine Yttria Alumina Zircon Metallic oxides	NITRIC ACID. MURIATIC ACID. § Barytes Potash Soda Strontites Lime Magnesia Ammonia Glucine Alumina Zircon Metallic oxides
		SULPHUROUS ACID. SUCCINIC. † Barytes Lime Potash Soda Strontites Magnesia Ammonia Glucine Alumina Zircon Metallic oxides	FLUORIC ACID. BORACIC ACID.    ARSENIC ACID. ¶ TUNGSTIC ACID. Lime Barytes Strontites Magnesia Potash Soda Ammonia Glucine Alumina Zircon Silex
OXIDE OF TIN. <i>Acids :</i> Gallic Muriatic Benzoic Oxalic Sulphuric Nitric Tartaric Mucic Phosphoric Citric Succinic Fluoric Arsenic Lactic Acetic Boracic Prussic Ammonia	OXIDE OF ANTI-MONY. <i>Acids :</i> Gallic Muriatic Benzoic Oxalic Sulphuric Nitric Tartaric Mucic Phosphoric Citric Succinic Fluoric Arsenic Lactic Acetic Boracic Prussic Fixed alkalies Ammonia	PHOSPHORIC ACID. CARBONIC. ‡ Barytes Strontites Lime Potash Soda Ammonia Magnesia Glucine Alumina Zircon Metallic oxides Silex	ACETIC ACID. LACTIC ACID. SUBERIC ACID. ** Barytes Potash Soda Strontites Lime Ammonia Magnesia †† Metallic oxides Glucine Alumina Zircon
OXIDE OF ZINC. <i>Acids :</i> Gallic Oxalic Sulphuric Muriatic Mucic Nitric	SULPHURIC ACID. PRUSSIC. * Barytes Strontites Potash Soda Lime Magnesia Ammonia	PHOSPHOROUS ACID. Lime Barytes Strontites Potash Soda Ammonia Glucine Alumina Zircon Metallic oxides	

\* With the omission of all after ammonia.

† Ammonia should come before magnesia ; and strontites, glucine, and zircon, be omitted.

‡ Magnesia should stand above ammonia, and alumina and silica be omitted.

§ Ammonia should stand above magnesia.

|| Silex should be omitted, and water and alcohol inserted. ¶ Except silex.

\*\* With the omission of strontites, metallic oxides, glucine, and zircon.

†† Magnesia should stand above ammonia.



TABLES OF SIMPLE AFFINITY—*continued.*

Oxalic Acid.	Benzoic Acid.	Soda	Alcohol.
Tartaric Acid.		Barytes	Water
Citric Acid.*	White oxide of ar- senic	Ammonia	Ether
Lime	Potash	Alumina	Volatile oil
Barytes	Soda	Magnesia	Alkaline sulphurets
Strontites	Ammonia		
Magnesia	Barytes	FIXED OILS.	SULPHURETED
Potash	Lime	Lime	HYDROGEN.
Soda	Magnesia	Barytes	Barytes
Ammonia	Alumina	Potash	Potash
Alumina		Soda	Soda
Metallic oxides	CAMPHORIC ACID.	Magnesia	Lime
Water	Lime	Oxide of mercury	Ammonia
Alcohol	Potash	Metallic oxides	Magnesia
		Alumina	Zircon

## No. III.

TABLES of the specific gravities of substances, which are articles of the *Materia Medica*, at a temperature of 60° of Fahrenheit.

METALS and INFLAMMABLES.		Muriatic	-	-	1.194
Mercury	- - -	Acetic	-	-	1.168
Sulphuret of mercury	10.000	Vinegar	-	1.0135 to	1.0251
Lead	- - - -	Distilled vinegar	1.007 to	1.0095	
Silver	- - - -	Citric acid	-	-	1.0345
Bismuth	- - -	Benzoic	-	-	0.667
Copper	- - -	Arsenious	-	-	3.7
Arsenic	- - -	Potash (hydrate)	-	-	1.7085
Sulphuret of arsenic (red)	3.225	Soda (do)	-	-	1.336
(yellow)	5.315	Ammonia liquid	-	-	0.760
Iron	- - -	Lime	-	-	2.3908
Sulphuret of iron	- - -	Magnesia	-	-	.346
Tin	- - -	Barytes	-	-	4.000
Zinc	- - -	Alumina	-	-	2.000
Manganese	- - -	Sulphate of potash	-	-	2.298
Antimony	- - -	----- soda	-	-	2.246
Sulphuret of antimony	4.368	----- magnesia	-	-	1.6603
Sodium	- - -	Alum	-	-	1.719
Potassium	- - -	Nitrate of potash	-	-	1.933
Phosphorus	- - -	Muriate of soda	-	-	2.120
Sulphur	- - -	----- ammonia	-	-	1.453
Charcoals	- 0.223 to	----- lime	-	-	1.76
Bitumens	- 0.892 to	Muriate of barytes	-	-	2.8257
	1.357	Phosphate of soda	-	-	1.338
SALINE SUBSTANCES.		Acetate of potash	-	-	2.1—?
Sulphuric acid	- - -	Tartrate of potash	-	-	1.5567
Nitric acid	- - -				

\* Zircon after alumina.



Tartrate of potash & soda	1·757	Honey	-	-	1·45
Supertartrate of potash	1·953	Myrrh	-	-	1·3600
Carbonate of potash	2·012	Olibanum	-	-	1·1732
———— soda	1·421	Opium	-	-	1·3359
———— ammonia	0·966	Opoponax	-	-	1·6226
———— lime	2·7	Resin (common)	-	-	1·0727
———— magnesia	0·2941	Sagapenum	-	-	1·2008
———— barytes	4·331	Scammony (Aleppo)	-	-	1·2354
Sub-borate of soda	1·720	———— (Smyrna)	-	-	1·2743

## METALLIC SALTS.

Mercury, corrosive mu- riate of	-	-	5·1398
———— mild muriate	-	-	7·1758
———— subsulphate	-	-	6·444
Copper, sulphate of	-	-	2·1943
———— acetate	-	-	1·779
Iron, sulphate of	-	-	1·880
———— carbonate	-	-	3·333
———— acetate	-	-	1·368
Lead, carbonate of	-	-	7·2357
———— superacetate	-	-	2·345
Zinc, sulphate of	-	-	1·912

## VEGETABLE SUBSTANCES AND PRODUCTIONS.

Cinchona bark	-	-	0·7840
Logwood	-	-	0·9130
Madder root	-	-	0·7650
Mahogany	-	-	1·0630
Red saunders	-	-	1·1280
Sassafras	-	-	0·4820
Gum arabic	-	-	1·5153
Hepatic aloës	-	-	1·3586
Socotorine aloës	-	-	1·3796
Amber (yellow)	-	-	1·0780
Ammoniacum	-	-	1·2071
Assafoetida	-	-	1·3275
Benzoin	-	-	1·0924
Camphor	-	-	0·9887
Catechu	-	-	1·4573
Elemi	-	-	1·0682
Euphorbium	-	-	1·1244
Galbanum	-	-	1·2120
Galipot	-	-	1·0819
Gamboge	-	-	1·2216
Guaiacum	-	-	1·2289

Storax	-	-	1·1098
Sugar (refined)	-	-	1·6060
Tragacanth	-	-	1·8161
Turpentine	-	-	0·991
Wax (yellow)	-	-	0·9648
———— (white)	-	-	0·9686

## FATS AND OILS.

Fat of beef	-	-	0·9232
———— mutton	-	-	0·9235
———— pork	-	-	0·9368
Tallow	-	-	0·9419
Butter	-	-	0·9423
Spermaceti	-	-	0·9433
Oil of linseed	-	-	0·9403
———— olives	-	-	0·9153
———— almonds	-	-	0·9170
Naphtha	-	-	0·8475
Oil of cinnamon	-	-	1·044
———— cloves	-	-	1·036
———— lavender	-	-	1·894
———— mint	-	-	0·8982
———— rosemary	-	-	0·9057
———— chamomile	-	-	0·8943
———— savine	-	-	0·9294
———— carraway	-	-	0·9049
———— anniseed	-	-	0·9867
———— juniper	-	-	0·8577
———— turpentine	-	-	0·8697
———— amber	-	-	0·8867
Sulphuric ether	-	-	0·632
Nitric ether	-	-	0·9088
Alcohol	-	-	0·794
Proof spirit	-	-	0·916

Water distilled	-	-	1·000
-----------------	---	---	-------



## No. IV.

*RULES for reducing the volume of Gases to a mean height of the Barometer, and mean Temperature.<sup>1</sup>*

1. *From the space occupied by any quantity of gas under an observed degree of pressure, to infer what its volume would be under the mean height of the barometer, taking this at 30 inches.*

This is done by the rule of proportion; for, as the mean height is to the observed height, so is the observed volume to the volume required. For example, if we wish to know what space would be filled, under a pressure of 30 inches of mercury, by a quantity of gas, which fills 100 inches, when the barometer is at 29 inches.

$$30 : 29 :: 100 : 96.66$$

The 100 inches would, therefore, be reduced to 96.66.

2. *To estimate what would be the volume of a portion of gas, if brought to the temperature of 60° Fahrenheit.*

Divide the whole quantity of gas by 480; the quotient will show the amount of its expansion or contraction by each degree of Fahrenheit's thermometer. Multiply this by the number of degrees which the gas exceeds, or falls below 60°. If the temperature of the gas be above 60°, subtract, or if below 60°, add the product to the absolute quantity of gas; and the remainder in the first case, or sum in the second, will be the answer. Thus, to find what space 100 cubic inches of gas at 50° would occupy if raised to 60°, divide 100 by 480; the quotient 0.208 multiplied by 10 gives 2.08, which added to 100 gives 102.08, the answer required. If the temperature had been 70°, and we had wished to know the volume, which the gas would have occupied at 60°, the same number 2.08 must have been subtracted from 100, and 97.92 would have been the answer.

3. *In some cases it is necessary to make a double correction, or to bring the gas to a mean both of the barometer and thermometer.*

We must then first correct the temperature, and afterwards the pressure. Thus to know what space 100 inches of gas at 70° Fah., 29 inches barometer, would fill at 60° Fah., and 30 inches barometer, we first reduce 100 inches, by the second process, to 97.92. Then, by the first

$$30 : 29 :: 97.92 :: 94.63.$$

Or 100 inches thus corrected, would be only 94.63.

4. *To ascertain what would be the absolute weight of a given column of gas at a mean temperature, from the known weight of an equal volume at any other temperature.* First find by the second process what would be its bulk at a mean temperature; and then say, as the corrected bulk is to the actual weight, so is the observed bulk to the number required. Thus if we have 100 cubic inches of gas weighing 50 grains at 50° Fah., if the temperature were raised to 60°, they would expand to 102.08. And

$$102.08 : 50 :: 100 : 49.$$

Therefore 100 inches of the same gas at 60° would weigh 49 grains.

5. *To learn the absolute weight of a given volume of gas under a mean pressure, from its known weight under an observed pressure, say, as the observed pressure is to the mean pressure, so is the observed weight to the corrected weight.* For example, having 100 inches of gas which weigh 50 grains under a pressure of 29 inches, to know what 100 inches of the same gas would weigh, the barometer being 30 inches.

$$29 : 30 :: 50 : 51.72.$$

Then 100 inches of the same gas, under 30 inches pressure, would weigh 51.72 grains.

6. In some cases it is necessary to combine the two last calculations. Thus, if 100 inches of gas at 50° Fah., and under 29 inches pressure, weigh 50 grains, to find what would be the weight of 100 inches at 60° Fah., and under 30 inches of the barometer, first correct the temperature, which reduces the weight to 49 grains. Then,

$$29 : 30 :: 49 : 50.7.$$

100 inches, therefore, would weigh 50.7 grains.

---

<sup>1</sup> Vide Henry's *Elements of Experimental Chemistry*, vol. ii. p. 497.



## No. V.

TABLES of the correspondence between measures of weight and capacity: according to the estimations given by Sir George Shuckburgh Evelyn, in vol. 88. of the *Phil. Trans.*, corrected by Mr. Fletcher, in the 4th Vol. of the *Philosophical Journal*.<sup>1</sup>

TABLE I.

For converting Cubic Inches of Water, (at 60 Therm. and 29.5 Bar.) into their equivalents in Troy weight.

<i>Cubic Inch of Water.</i>	<i>Troy grs.</i>	<i>oz.</i>	<i>drachm.</i>	<i>grs.</i>
1 weighs	252.506	= 0	: 4	: 12.506
2 cubic inches weigh	505.012	= 1	: 0	: 25.012
3	757.518	= 1	: 4	: 37.518
4	1010.024	= 2	: 0	: 50.024
5	1262.530	= 2	: 5	: 2.530
6	1515.036	= 3	: 1	: 15.036
7	1767.542	= 3	: 5	: 27.542
8	2020.048	= 4	: 1	: 40.048
9	2272.554	= 4	: 5	: 52.554
1728 (1 cubic foot)	—————	909	: 0	: 10.368

TABLE II.

For converting Troy Grains, Drachms, Ounces, and Pounds of Water into their equivalent Cubic Inches.

<i>Grains.</i>	<i>Cubic Inch.</i>	<i>Drachm.</i>	<i>Cubic Inch.</i>
1 =	.00396	1 =	.237618
2 =	.00792	2 =	.475236
3 =	.01188	3 =	.712854
4 =	.01584	4 =	.950472
5 =	.01980	5 =	1.188090
6 =	.02376	6 =	1.425708
7 =	.02772	7 =	1.663326
8 =	.03168		
9 =	.03564		
<i>Ounce.</i>	<i>Cubic Inch.</i>	<i>Pound.</i>	<i>Cubic Inch.</i>
1 =	1.900945	1 =	22.81134
2 =	3.801890	2 =	45.62268
3 =	5.702835	3 =	68.43402
4 =	7.603780	4 =	91.24536
5 =	9.504725	5 =	114.05670
6 =	11.405670	6 =	136.86804
7 =	13.306615	7 =	159.67938
8 =	15.207560	8 =	182.49072
9 =	17.108505	9 =	205.30206
10 =	19.009450		
11 =	20.910395		

<sup>1</sup> Not having the Fourth Volume of the *Philosophical Journal* by us, we have copied these Tables from the Appendix of Aikin's Dictionary.



TABLE III.

For converting Wine Pints of Water into their equivalent Troy and Avoirdupois Pounds.

<i>Wine Pints.</i>		<i>lbs. Troy.</i>		<i>lbs.</i>	<i>oz.</i>	<i>dr.</i>	<i>grs.</i>		<i>lbs. Avoirdup.</i>
1	=	1.26581783	=	1	:	3	:	1	: 31.1 = 1.04158725
2	=	2.53163566	=	2	:	6	:	3	: 2.2 = 2.08317450
3	=	3.79745349	=	3	:	9	:	4	: 33.3 = 3.12476175
4	=	5.06327132	=	5	:	0	:	6	: 4.4 = 4.16634900
5	=	6.32908915	=	6	:	3	:	7	: 35.5 = 5.20793625
6	=	7.59490698	=	7	:	7	:	1	: 6.6 = 6.24952350
7	=	8.86072481	=	8	:	10	:	2	: 37.7 = 7.29111075
8	=	10.12654264	=	10	:	1	:	4	: 8.8 = 8.33269800
9	=	11.39236047	=	11	:	4	:	5	: 39.9 = 9.37428525

---

TABLE IV.

For converting Troy Pounds of Water into their equivalent Wine Pints.

<i>Troy Pounds.</i>	<i>Wine Pints.</i>	<i>Troy Pounds.</i>	<i>Wine Pints.</i>
1	=	0.7900031	
2	=	1.5800062	
3	=	2.3700093	
4	=	3.1600124	
5	=	3.9500155	
		6	= 4.7400186
		7	= 5.5300217
		8	= 6.3200248
		9	= 7.1100279

---

TABLE V.

For converting Avoirdupois Pounds into their equivalent Troy Pounds.

<i>lbs. Avoird.</i>	<i>lbs. Troy.</i>	<i>lbs. Avoird.</i>	<i>lbs. Troy.</i>
1	=	1.215277	
2	=	2.430555	
3	=	3.645833	
4	=	4.861111	
5	=	6.076388	
		6	= 7.291666
		7	= 8.506944
		8	= 9.722222
		9	= 10.937500

---

TABLE VI.

For converting Troy Pounds into their equivalent Avoirdupois Pounds.

<i>lbs. Troy</i>	<i>lbs. Avoirdup.</i>	<i>lbs. Troy.</i>	<i>lbs. Avoirdup.</i>
1	=	0.82285714	
2	=	1.64571428	
3	=	2.46857142	
4	=	3.29142857	
5	=	4.11428571	
		6	= 4.93714285
		7	= 5.76000000
		8	= 6.58285714
		9	= 7.40571428



TABLE VII.

For converting Ounces, Drachms, and Grains Troy into Decimals of the Troy Pound.

Grains.	lbs. Troy.	Drachms.	lbs. Troy.	Oz.	lbs. Troy.
1 =	·000173611	1 =	·0104166	1 =	·0833
2 =	·000347222	2 =	·0208333	2 =	·1666
3 =	·000520833	3 =	·0312500	3 =	·2500
4 =	·000694444	4 =	·0416666	4 =	·3333
5 =	·000868055	5 =	·0520833	5 =	·4166
6 =	·001041666	6 =	·0625000	6 =	·5000
7 =	·001215277	7 =	·0729166	7 =	·5833
8 =	·001388888			8 =	·6666
9 =	·001562500			9 =	·7500
				10 =	·8333
				11 =	·9166

TABLE VIII.

For converting Decimals of the Troy Pound into Troy Ounces, Drachms, and Grains.

lbs.	oz.	dr.	grs.	lbs.	oz.	dr.	grs.	lbs.	grs.				
·1 =	1	:	1	:	36	·01 =	0	:	0	:	57·6	·001 =	5·76
·2 =	2	:	3	:	12	·02 =	0	:	1	:	55·2	·002 =	11·32
·3 =	3	:	4	:	48	·03 =	0	:	2	:	52·8	·003 =	17·28
·4 =	4	:	6	:	24	·04 =	0	:	3	:	50·4	·004 =	23·04
·5 =	6	:	0	:	0	·05 =	0	:	4	:	48·0	·005 =	28·80
·6 =	7	:	1	:	36	·06 =	0	:	5	:	45·6	·006 =	34·56
·7 =	8	:	3	:	12	·07 =	0	:	6	:	43·2	·007 =	40·32
·8 =	9	:	4	:	48	·08 =	0	:	7	:	40·8	·008 =	46·08
·9 =	10	:	6	:	24	·09 =	0	:	8	:	38·4	·009 =	51·08

No. VI.

TABLE showing the correspondence between the new French and the English Weights and Measures.

I. MEASURES OF LENGTH.

The metre being at 32°, and the foot at 62°.

	English Inches.			English.
Millimetre	=	·03937		
Centimetre	=	·39371		
Decimetre	=	3·93710	Miles.	fur. yds. ft. in.
Metre	=	39·37100	=	0 0 1 0 3·37
Decametre	=	393·71000	=	0 0 10 2 9·7
Hecatometre	=	3937·10000	=	0 0 109 1 1
Kilometre	=	39371·00000	=	0 4 213 1 10·2
Myriametre	=	393710·00000	=	6 1 156 0 6



## II. MEASURES OF CAPACITY.

		<i>Cubic inches.</i>				
Millilitre	=	·06103				
Centilitre	=	·61028				
Decilitre	=	6·10280				
Litre	=	61·02800	=	0	0	0
Decalitre	=	610·28000	=	0	0	2
Hecatolitre	=	6102·80000	=	0	0	26·419
Kilolitre	=	61028·00000	=	1	0	12·19
Myriolitre	=	610280·00000	=	10	1	58·9

## III. MEASURES OF WEIGHT.

		<i>English grains.</i>				
Milligramme	=	·0154				
Centigramme	=	·1544				
Decigramme	=	1·5444				
Gramme	=	15·4440				
Decagramme	=	154·4402	=	0	0	5·65
Hecatogramme	=	1544·4023	=	0	3	8·5
Kilogramme	=	15444·0234	=	2	3	5
Myriogramme	=	154440·2344	=	22	1	2

## No. VII.

TABLES showing the correspondence of English Weights and Measures with those of Holland, Sweden, and Germany.

The English Ale Gallon contains 282 cubic inches; the Wine Gallon 231 cubic inches.

## I. DUTCH.

1 lb. Dutch = 1 lb. 3 oz. 16 dwt. 7 grs. Troy.

787½ lbs. Dutch = 1038 lbs. Troy.

## II. SWEDISH.

1 kanne of water Swedish = 48088·71944 grs. Troy, in weight;  
and 189·9413 English cubic inches.

1 lb. Swedish = 6556 grs. Troy.

## III. GERMAN.

74 lbs. German Apothecaries' Weight = 74 lbs. Troy.

1 oz. Nuremberg medic. weight = 7 dr. 2 dwt. 9 grs. Troy.

1 mark Cologne = 7 oz. 2 dwt. 4 grs. Troy.



## PART II.

---

### MATERIA MEDICA.

**M**ATERIA MEDICA is that department of the science of medicine, that treats of the nature and properties of the substances which are employed as remedies to restore health in diseased bodies.

According to this definition, it should comprehend every remedy, whether it be a simple, the production of nature, or a compound artificially prepared by the pharmacopolist: but the British Colleges of Physicians confine the application of the term, in their pharmacopœias, to those remedies only which are simples, and such compounds as are articles of general commerce, or over the preparation of which they have no control. These pharmacopœias differ also from the works of the generality of systematic writers on Materia Medica, in arranging the substances alphabetically, without any regard to their affinities as natural objects, or their medicinal virtues. This mode, although it be not so scientific, yet is much less liable to objection than many of the other modes that have been occasionally adopted; as the best of these have been, generally, too much modified by the prevailing theoretical doctrines of the day, which, unfortunately for medical science, have hitherto had too slight a foundation on truth to secure their permanence. The plan of the pharmacopœias has consequently been judiciously followed by the compilers of Dispensatories; and the convenience and utility of it is so generally acknowledged, that we the more readily comply with our own opinion of its propriety in adopting it.

This part of our work, therefore, contains the lists of the materia medica of the pharmacopœias issued by the London, the Edinburgh, and the Dublin colleges; and subjoined to the name of each of the substances supplied by the vegetable and the animal kingdoms, a description of the plant or the animal which yields the remedy, is given in the language and after the method of Natural History. The chemical characters, as far as they are known, of these matters are also stated; and the analysis of such remedies as are more immediately the objects of chemical investigation, with the medical properties and uses of all of them, are detailed; so as to afford every useful information regarding them, in a form the most convenient for practical reference.



**AB'IETIS RESINA.** Vide *Pinus Abies*.

**ABSINTHIUM.** Vide *Artemisia Absinthium*.

**ACACIA.** *Spec. Plant Willd.* iv. 1085.

*Cl.* 23. *Ord.* 1. Polygamia Monœcia. *Nat. ord.* Lomentaceæ *Linn.*  
Leguminosæ *Juss.*

*G.* 1902. *Hermaph.* *Calyx* five-toothed. *Corolla* five-cleft, or formed of five petals. *Stamens* 4-100. *Pistil.* 1. *Legume* bivalve.

*Male.* *Cal.* five-toothed. *Cor.* five-cleft, or formed of five petals.  
*Stam.* 4-100.

\*\*\*\* *Leaves bipinnate, stipular thorns or prickles, elongated spikes.*

*Species* 73. *Acacia Catechu*.<sup>1</sup> *Catechu. Med. Bot.* 2d edit. t. 157.

\*\*\*\*\* *Leaves bipinnate, stipular thorns, globular spikes.*

*Species* 87. *Acacia vera.* *Acacia*, or Egyptian Thorn, *Med. Bot.*  
2d. edit. t. 158. *Vesl. Ægypt.* t. 8. bona.

#### 1. ACACIA CATECHU.

*Officinal.* CATECHU EXTRACTUM, *Lond.* ACACIÆ CATECHU EXTRACTUM, *Edin.* CATECHU; EXTRACTUM E LIGNO, *Dub.*  
Extract of Catechu.

*Syn.* Cachou (*F.*), Katechu; Kaschu (*G.*), Cato o Catecu (*I.*), Catecu (*S.*)  
Cato (*Portug.*), Cutt (*Hind.*)

This tree grows pentifully in the mountains of Kanhana in Hindostan; and flowers in June. It seldom exceeds twelve feet in height, and one foot in the diameter of its stem, which is covered with a thick, rough, brown bark, and towards the summit is divided into many close branches. The leaves are placed alternately on the younger branches; and are composed of fifteen or thirty pairs of pinnæ, nearly two inches long, each having about forty pairs of linear leaflets, beset with short hairs; with a small gland on the common petiole between the bases of each pair of the pinnæ. At the base of each leaf are two short, recurved, compressed spines. The flowers are hermaphrodite and male; axillary, and on close spikes four or five inches long; the calyx is tubular, hairy, dividing into five oval-pointed segments; the corolla of one piece, whitish, twice the length of the calyx, and of the same form. The filaments are numerous, double the length of the corolla, crowded with roundish anthers, and adhering at the base of the germen, which is oval, supporting a slender style the length of the filaments, and terminated by a simple stigma. The fruit is a lanceolate, compressed, smooth, brown pod, with an undulated thin margin: and contains six or eight roundish flattened seeds, which emit a disagreeable odour when chewed.

The inner wood of this tree is of a brown colour; and from it, according to Mr. Ker's statement<sup>2</sup>, the *catechu* is prepared. "After felling the trees, the manufacturer carefully cuts off all

<sup>1</sup> Ἀκαυδὸς τῆς Ἀρίας χώρας, Theophrasti. Dioscorides also mentions this species of *Acacia*. The name in Bahar is *Coira* or *Keira*.

<sup>2</sup> *Med. Obs. and Enquir.* vol. v. p. 151.



the exterior white part of the wood. The interior coloured part is cut into chips, with which he fills a narrow-mouthed unglazed earthen pot, pouring water upon them until he sees it among the uppermost chips: and when this is half evaporated by boiling, the decoction, without straining, is poured into a flat earthen pot, boiled to one-third part, and then set in a place to cool for one day. The decoction is afterwards evaporated by the heat of the sun, stirring it several times in the day; and, when it is reduced to a considerable thickness, it is spread upon a mat or cloth, which has previously been covered with the ashes of cow-dung. The mass is lastly divided into square or quadrangular pieces by a string, which are completely dried by turning them in the sun, until they are fit for sale." Before this account was published, catechu was generally supposed to be extracted from the Areca nut: there are, however, two other species which are extracted from that nut; the one named *Cuttacamboo*, the other *Cashcutti*, and both are used by the Indian practitioners.

This extract, when first introduced as a medicine into Europe, was named *Terra Japonica*, from the supposition that it came from Japan, and was an earth. It is named *cutt* by the natives of Hindostan, *cutch* by the English, and by different authors *khaath*, *cate*, *cachou*, *cachore*,<sup>1</sup> and *catechu*.<sup>2</sup> There are two varieties of the true catechu; one brought from Bengal, the other from Bombay. It is imported into Britain in bags, and sometimes in boxes or chests, containing from 3 to 4 cwt. each; and occasionally in small squares, in boxes, which are at all times preferred. Pale and dark-coloured catechu are mixed in the same package.

*Qualities.* — Pale catechu is generally in round balls, or small square cakes of a pale reddish brown colour, light and friable, with a lamellated texture, and rough fracture; has a bitterish and astringent taste, leaving a degree of sweetness on the palate; is inodorous, and has a specific gravity of 1.39. The dark, which is in round masses, has a deep chocolate-colour internally, with the hue of rusty iron on the outside: the texture is uniform, and the fracture resinous and shining. It is heavier than the pale, the specific gravity being 1.45, and has a more austere and bitter taste; but in other respects agrees with it. Both are often much adulterated with sand, and other impurities. Water at 60° dissolves 3 parts in 4 of pale catechu: the alkalies deepen the colour of the solution, until on standing it becomes of a deep mahogany hue. Ac-

<sup>1</sup> Bolduc, *Mem. Acad.* 1709, p. 293.

<sup>2</sup> This name is said to be compounded of two Oriental words, *cate*, which signifies a tree, and *chu*, juice. *Kerr*, l. c.



According to the analysis of Sir H. Davy, there appears to be very little difference between the two varieties. Both are almost entirely soluble in the mouth. 100 grains, macerated in 18 fluid ounces of water at 52°, left 7½ grains only undissolved, and these were chiefly lime, aluminous earth, and sand. The solutions are inodorous, and slightly redden tincture of litmus. Sulphate of iron and gelatine throw down precipitates in them<sup>1</sup>, demonstrating the presence of *gallic acid* and *tannin*: what remains after the action of alcohol is nearly a pure *mucilage*; and, when fine powder of catechu is washed with water until all the tannin and mucilage are dissolved, a pale red *extractive matter*, inodorous, very slightly astringent, sweetish, and soluble in water and in alcohol, is obtained as a residue. The proportions of these constituents, according to Davy's experiments, were as follows: 200 grains of *Bombay catechu* afforded 109 of tannin, 68 of extractive matter, 13 of mucilage, and 10 of earths and other impurities. The same quantity of *Bengal catechu* gave 97 of tannin, 73 of extract, 16 of mucilage, and 14 of impurities.<sup>2</sup>

*Medicinal properties and uses.* — Catechu is one of the most valuable of the vegetable astringents; and as the dark-coloured contains the greater quantity of tannin, on which its astringency depends, it is to be preferred for medicinal use. It is employed with the best effects in dysentery, and diarrhœa, when the use of astringents is admissible; in alvine and uterine hæmorrhages, leucorrhœa, gleet, and in obstinate catarrhal affections. As a local astringent it is used in sponginess of the gums, and aphthous ulcerations of the mouth and fauces: and I have found the slow solution of a small piece of it in the mouth, a certain remedy for the troublesome cough induced by a relaxed uvula, hanging into and irritating the glottis: and on the same principle it is used by public singers to prevent hoarseness. Dr. Paris recommends it as a dentifrice, especially when the gums are spongy: and I can recommend it as an excellent toothpowder when combined with three parts of finely powdered charcoal.<sup>3</sup> In prescribing it, the practitioner should bear in mind, that alkaline salts destroy its astringency; and metallic salts and solution of isinglass form with it insoluble compounds.

An ointment composed of ℥jv of catechu, ℥ix of alum, ℥iv of white resin, and f℥x of olive oil, with a sufficient quantity of water, is in great repute in India as an application to ulcers.

---

Tincture of muriate of iron produces a precipitate, which, on the addition of Lig., Potassæ is redissolved, unless the quantity of the tincture be great, and the solution becomes of a deep purple or port wine colour.

<sup>2</sup> *Philosophical Transactions*, 1803.

<sup>3</sup> *Pharmacologia*, 3d edit.



The dose of catechu may be from grs. x to ʒj.

Officinal Preparations. *Infusum Catechu*. L. E. *Tinctura Catechu*. L. E. *Electuarium Catechu compositum*. E. D.

## 2. ACACIA VERA.<sup>1</sup>

Officinal. ACACIÆ GUMMI, *Lond.* ACACIÆ ARABICÆ GUMMI, *Edin.* GUMMI ARABICUM, *Dub.* Acacia Gum, or Gum arabic.

Syn. Gomme Arabique (*F.*), Arabischen gummi (*G.*), Gomma Arabica (*I.*), Goma Arabiga (*S.*) Tolh (*A.*), Vullām pisin (*Tam.*).

This species of acacia is found in almost every part of Africa; but the tree that yields the gum which is exported from Barbary to Great Britain, grows principally in the Atlas mountains, and at Bled-eljerrede; flowering in July. It has a hard withered aspect, and does not rise many feet in height. The stem is crooked, and covered with a grey bark, which on the branches has a purplish tinge: the leaves are alternate, bipinnate, composed of several pairs of opposite pinnæ, with a small gland on the common petiole, between the base of each pair, and having numerous pairs of narrow elliptical, smooth leaflets. On each side of the base of the leaves are two long diverging white spines. The flowers are hermaphrodite and male, crowded into globose heads, (*capitula*,) rather than spikes, which are supported on slender peduncles, and rise, four or five together, from the axillæ of the leaves: the calyx is small, bell-shaped, and five-toothed; the corolla is divided into five narrow yellowish segments; the filaments are numerous, capillary, bearing roundish yellow anthers; the germen is conical, with a slender style and simple stigma; and the pods, which are three or four inches long, and half an inch broad, contain several flattish brown seeds.<sup>2</sup>

The gum exudes naturally from the bark of the trunk and the branches of the tree, in a soft, nearly fluid state, and hardens in the air without losing its transparency. It is collected about the middle of December. "It appears," Mr. Jackson informs us, "to be the product of disease; for in the hottest seasons, and from the most sickly trees, the greatest quantity is procured. Very little or none is got in a moist, cool, or mild summer. It is gathered in July and August when the weather is hot and parching. It has a faint smell when first stowed in the warehouses, and is heard to crack spontaneously for many weeks. The best gum is procured from Morocco, Rasel-wed in the province of Suse, and Bledhummer in the province of Abda."<sup>3</sup>

<sup>1</sup> Ἀκανθὸς αἰγυπτία. *Hi pocratis*. Ακακία, Dioscoridis. In Barbary it is named *attaleh*. — *Jackson's Morocco*, 4to. p. 33.

<sup>2</sup> From the unripe pods the *acaciæ veræ succus* of the ancients was expressed. *Vide Murray App. Med.* ii. 412. The seeds yield a reddish dye. *Jackson*, l. c.

<sup>3</sup> *Jackson*, p. 83. In 1805, the quantity exported from Mogodor to London was 277,534lbs. *Ib.* l. c.



It is imported from Barbary and Morocco in large casks. Gum Senegal, which was introduced into Europe by the Dutch in the 17th century, and is often mixed with the Barbary gum, is obtained from various trees, but chiefly from two, one called *verreck*, which yields a white gum, the other called *nebuel*, which yields a red gum.<sup>1</sup>

*Qualities.* — Gum is generally in irregularly shaped pieces, hard, brittle, semitransparent, its fracture possessing a considerable degree of lustre; and is neither fusible nor volatile. When pure, it is almost colourless, or of a pale yellowish hue; is insipid, inodorous, and dissolves completely away in the mouth. Its specific gravity varies from 1,3161 to 1,4317. It is often mixed with the gum Senegal, which is nearly as pure, but in larger masses, generally of a darker colour and more clammy and tenacious, and with other gums less pure, particularly a kind brought from the East Indies, which is still darker-coloured and less soluble.<sup>2</sup>

Gum is soluble in water, either cold or hot, and forms a viscid solution; which, if evaporated, becomes very thick and adhesive, and at length the gum is obtained in a concrete form, equally soluble as before. It is also soluble in the vegetable acids; but is insoluble in alcohol, in ether, and in oils: yet, owing to its viscosity, it renders by trituration both the volatile and fixed oils and resins miscible with water, forming a white opaque mixture. Concentrated sulphuric acid blackens, and partially decomposes it, and acetic acid is produced: strong nitric acid converts it into the oxalic, malic, and saccholactic acids; muriatic exerts very little action on it; but the oxymuriatic (*chlorine*) changes it into citric acid. Solutions of the alkalis and alkaline earths dissolve it without producing on it much change. For an account of the action of other agents on it, see *Mucilago acaciæ*.

The chemical analysis of gum, by Gay Lussac and Thenard shows, that its constituents are 42,23 of carbon 6,93 of hydrogen, and 50,82 of oxygen, with a small proportion of nitrogen

<sup>1</sup> Although, by the treaty which delivered up Senegal to France in 1783, the gum trade was reserved to England, and has never been annulled, yet no advantage was taken of this right by the English, until the commencement of the present year (1821); when the trade was again renewed at Portendic, under the protection of Commodore Sir George Collier. — Portendic is in latitude 18° 19' north, by 16° 10' west of Greenwich. The Acacia forest of Sabel, which yields a very pure white gum, is sixty miles, and the forest of El-Hiebar eighty miles from Portendic. The gum is collected and brought to the coast by the Trarzhar Moors, who receive in exchange baft and other English goods: "the usual quantity given for a piece of blue baft, which costs from fifteen to thirty shillings, is from 100 to 133 lbs. weight of gum." *Sierra Leone, Royal Gazette*, May 26th, 1821.

<sup>2</sup> The gum which exudes from the cherry, plum, and other trees of the genus *Prunus*, in this country is cerasin; but the gum alluded to is very similar to gum arabic, and is furnished by the *Acacia Arabica*, or Babul tree of Hindostan. The gum is called *Babuled gund*, by the natives.



and lime; which last element is supposed to render it incapable of undergoing the fermentative process.<sup>1</sup> I have found, however, that it, nevertheless, contains a small proportion of gluten; for when rubbed up with a spiriteous solution of guaiac, a blue colour is evolved.

*Medical properties.* — Gum exerts no action on the living system; but is a simple demulcent, serving to lubricate abraded surfaces, and involve acrid matters in the primæ viæ; and it effects these purposes the more readily from its passing through the bowels, scarcely acted upon by the assimilative functions.<sup>2</sup> In the solid form it is scarcely ever given, unless to sheath the fauces, and allay the tickling irritation which occasions the cough in catarrh and phthisis pulmonalis; in which cases a piece of it is allowed to dissolve slowly in the mouth. It is chiefly used in the state of mucilage. Vide *Mucilago Acaciæ*,

Officinal preparations. *Mucilago Acaciæ*, L.E.D. *Emulsio Acaciæ Arabicæ*, E. *Emulsio Arabica*, D. *Troc. gummosi*, E.

ACETOSÆ FOLIA. Vide *Rumex Acetosa*.

ACETOSELLA. Vide *Oxalis Acetosella*.

ACETUM, *Lond. Edin.* ACETUM VINI, *Dub.* Vinegar.

*Syn.* Vinaigre (*F.*), Essig (*G.*), Aceto (*I.*), Vinagre (*S.*), Vünacdike (*Danish*), Winättika (*Swed.*), Canchiea (*Sans.*), Khull (*Arab.*), Chooca Wynazin (*Malay*), Kadi (*Tam.*), Cirka (*Pers.*), Kudidia (*Cyng.*), (*Belg.*).

This is a well known acid liquor, produced by exciting the acetous fermentation in substances which have undergone, or are susceptible of, the vinous fermentation. Sugar and water, the saccharine vegetable juices, infusions of malt, malt liquors, cyder, and wine<sup>3</sup>, may be converted into vinegar, by adding to them yeast or any other ferment, and exposing them in vessels to which the air has access, in a temperature between 75° and 90°. In wine countries, as France and Italy, vinegar is made from the lees of wine, which are worked up with new wine, then strained, and exposed to the heat of the sun, or placed in stoved rooms, in casks set upright, with a hole cut through the heading, and left open until the whole of the liquor is thoroughly acidified. In this country it is prepared chiefly from malt. An infusion of malt is made, then properly cooled, and put into large and deep fermenting tuns; “where it is mixed with yeast, and kept in fermentation for four or five days.” The liquor is now distributed into smaller vessels, placed in a chamber heated by means of a stove; and kept

<sup>1</sup> *Murray's Chemistry*, vol. iv. 180. The last analysis, which is by Berzelius makes the components in 100 parts to be, hydrogen 6,792, carbon 41,752, and oxygen 51,456.

<sup>2</sup> It is nevertheless sometimes used as food by the Moors: and at Senaar, in Dongola, is much employed in cookery.

<sup>3</sup> New wines are better for this purpose than old, as they contain more extractive matter.



there for about six weeks, or until the whole is soured. This is emptied into common barrels, which are placed in the open air, the bung-hole of each being simply covered with a tile to keep out the wet; and in this situation such a gentle fermentation goes on, that in four or five months, according to the heat of the weather, perfect vinegar is formed. The process is then completed in the following manner: "Large tuns are employed, with a false bottom, on which is put a quantity of the refuse of raisins and other fruit, left by makers of home-made wines, called, technically, *rape*. These rape tuns are worked by pairs; one of them is quite filled with the vinegar from the barrels, and the other only three quarters full, so that the fermentation is excited more easily in the latter than the former, and every day a portion of the vinegar is laded from one to the other till the process is finished."<sup>1</sup>

The theory of the acetous fermentation is not yet fully understood. Air and a moderate temperature are necessary for exciting and keeping it up. The former affinities between the components of the ingredients are broken, and new ones formed; while a quantity of carbon is thrown off, and uniting with the oxygen of the air, produces the carbonic acid gas, which appears during the process. Although alcohol alone cannot be converted into vinegar, yet the strongest wine produces the best vinegar; and hence that made from malt is weaker, less pure, and more liable to spoil, than wine vinegar. The essential part of vinegar is *acetic acid* and *water*, in the proportion of five parts to ninety-five of water<sup>2</sup>: but it also contains colouring matter, some undecomposed *alcohol*, *gluten*, *mucilage*, *sugar*, *extractive matter*, and often some *malic* and *tartaric acids*.

*Qualities*. — Vinegar, when well made, is clear and limpid; has an agreeable penetrating odour, and a pleasant acid taste.

The colour varies from a pale yellow to a deep red; and as it is derived from the extractive matter, malt vinegar is always higher coloured than wine vinegar. When long kept, particularly if it be exposed to the air, vinegar becomes muddy and ropy, acquires an unpleasant smell, loses its acidity, and putrefies. It, however, may be kept good for a much longer time, if it be boiled for a few minutes, so as to coagulate and separate the gluten, on the presence of which the above changes depend; and be preserved in well-corked bottles. It is sometimes adulterated with sulphuric acid, which may be detected by a solution of acetate of baryta, forming, when

---

<sup>1</sup> *Aikin's Dictionary of Chemistry*, art. *Vinegar*.

<sup>2</sup> *Journal of Science and the Arts*, vol. vi. p. 259.



dropped into the suspected vinegar, a white precipitate, which is *insoluble* in nitric acid, after being exposed to a strong heat. If sulphate of baryta be employed, sulphate of potass or of lime will produce a precipitate, if present in the vinegar, in which case a preferable test is chalk short of the quantity necessary to saturate the portion of vinegar; for, by throwing the whole on a filter, and adding distilled water, the acetate of lime is dissolved, but the sulphate which is formed, if sulphuric acid be present, remains undissolved on the filter. Muriatic acid is detected by nitrate of silver forming a precipitate, which is insoluble in nitric acid; but soluble in liquid ammonia. Nitric acid is detected by first mixing some common salt in the vinegar, then saturating with potass and evaporating to dryness. Gold leaf is next to be diffused through equal parts of sulphuric acid and water, and the mixture added to the dry residue and boiled, when, if the vinegar contained nitric acid, the gold leaf will be dissolved.<sup>1</sup> If *grains of paradise*, *spurge flax*, *capsicum* or *Pellitory of Spain*, which are sometimes used to adulterate vinegar, be present, they can be detected by the taste.

The use of vinegar as a condiment, and an antiseptic, for pickling and preserving animal and vegetable matter, is well known.

*Medical properties and uses.* — Vinegar, when taken into the stomach, acts as a refrigerant, promotes diaphoresis and the discharge of urine; and is a powerful anti-narcotic: its external action on the living fibre is moderately stimulant and astringent.

In inflammatory fevers it may be used to acidulate the ordinary beverage. It is given as a remedy in putrid diseases and scurvy; and is the most easily procured, and the best means of counteracting the fatal effects of over doses of narcotic poisons; for which purpose, it should be administered in doses of a table spoonful, frequently repeated, after the stomach has been freely emptied by a proper emetic. It is employed as a glyster in obstinate costiveness; and externally in the form of fomentation, or of lotion, in burns, bruises, sprains, and chronic ophthalmia; and, diluted with water, it is the best lotion for clearing the eye of small particles of lime, when they adhere to any part of the ball or the lids. Its vapour is inhaled in putrid sore throat; and is diffused through sick rooms with the view of neutralizing pestilential effluvia: but as a fumigation it has little efficacy. The dose of vinegar is  $\text{f } 3j$  to  $\text{f } 3iij$ ; and the quantity given in glysters  $\text{f } 3j$  to  $\text{f } 3ij$ .

---

<sup>1</sup> *Brande's Manual of Pharm.* p. 5.



Official preparations. *Acidum aceticum, dilutum*, L.E.D. *Syrupus Aceti*, E.

**ACIDUM ACETICUM** (*impurum?*) FORTIUS, *Lond.*  
Strong (*impure?*) Acetic acid. (*Spec. grav.* 1.046. *Lond.*)

This is the acetic acid distilled from wood. It is procured by decomposing the wood by means of heat in large iron cylinders, during which the acid rises in a gaseous form; but being conducted through pipes kept cool by passing through barrels of water, it ultimately comes over in a liquid state. It is of a deep brown colour, and has a strong empyreumatic odour when it is first obtained; but, as it is allowed to stand undisturbed for a few weeks, until it deposits a peculiar kind of tar, and is then re-distilled, it loses a large portion of its colouring matter, and much of its empyreumatic odour. It is next saturated with lime, so as to form an impure acetate of lime, which is decomposed in a proper apparatus by sulphuric acid. The sulphate of lime, which is formed in this part of the process, remains in the still, and the acetic acid passes over. In some instances the decomposition is effected by means of sulphate of soda, which produces an acetate of soda, and sulphate of lime, the former of which being decomposed by sulphuric acid, the acetic acid and sulphate of soda are obtained. M. Stolze of Halle has discovered a mode of purifying it by means of sulphuric acid, manganese, and muriate of soda, and subsequent distillation.<sup>1</sup>

*Qualities.* — When well prepared, this acid is colourless and limpid like water, has the sharp, pleasant taste and penetrating odour of vinegar; but combined with these, there remain a peculiar flavour, and a slight empyreumatic aroma not unlike that of Westphalia ham. No process has, hitherto, been able to separate this empyreuma; and, therefore, it cannot, strictly speaking, be regarded as pure acetic acid moderately diluted, as its appellation in the Pharmacopœia of the London College imports. When of the specific gravity 1.046, it should contain 28.23 per cent. of real acid, and 71.57 of water, or be six times the strength of common vinegar of the sp. gr. 1.009<sup>2</sup>; and one hundred grains of it should require eighty-seven grains of crystallized subcarbonate of soda to saturate them. It is extremely volatile; and, when held near carbonate of ammonia, its vapour unites with that of the

---

<sup>1</sup> Although this modification of the acetic acid has only lately come into use, yet it is noticed by Glauber under the name "*Vinegar of wood*," and it was known in England before the year 1661, under the name of "*Sour spirit, or vinegar of box-wood*." Vide *Sceptical Chemist*, quoted in *Journal of Science*, vol. viii. p. 368.

<sup>2</sup> At sp. gr. 1.043 the proportions are, acetic acid 23.67, water 76.33, in 100 parts. *Phillip's Trans. of Pharm. Lond.* 1824.



ammonia, forming a white opaque smoke, which is a dry acetate. In every respect, indeed, it possesses the same chemical properties as diluted, pure acetic acid. If pure it will remain colourless when mixed with sulphuric acid, and form a colourless salt when saturated with potass.

*Medical properties and uses.* — This acid is stimulant and antiseptic. When sufficiently diluted, it may be employed for the same purposes as distilled vinegar; but, it must be acknowledged, that, when saturated with ammonia, the solution of the *acetate* is more nauseous than that which is prepared with distilled vinegar. In diluting it for pharmaceutical purposes, care should be taken that distilled water only be employed; for when spring water is used for this purpose, the diluted acid, when added to lotions containing acetate of lead, causes the lotion to become milky, owing to the decomposition of the acetate and the formation of sulphate of lead, which is insoluble. It has been found useful in dressing phagedenic ulcers, the feter of which it corrects; promoting the formation of healthy pus, and softening the hardened edges of the sores. Its antiseptic qualities are so considerable, that it has been used for preserving both animal and vegetable substances; and might be advantageously employed for anatomical preparations.

Official preparations. — *Potasse Acetas*, L. *Plumbi Acetas*, L.

*A'CIDUM CITRICUM*, Lond. *ACIDUM CITRICUM CRYSTALLIS CONCRETUM*, Dub. Vide *Acidum Citricum* among the preparations.

*A'CIDUM SULPHURICUM*, Lond. Edin. Dub. Sulphuric Acid. (*Specific gravity*, 1,850. Lond. 1,845. Edin. Dub.)

*Syn.* Acide sulphurique (*F.*), Vitriolöl schweffelsaure (*G.*), Vitriool oli. — Reines Zwavebzuur (*Dutch*), Vitriol olje (*Dan.*), Vitriolja, Swafwelsyra (*Swed.*), Acido solforico (*I.*), Gundaica Atr. (*H.*), Gunduck Trāvagum (*Tam.*), Arckgowgird (*Pers.*), Rooch (*Arab.*).

This acid is said to be found in a concrete state, in the cavities of some volcanic mountains, and dissolved in some mineral waters: but for the purposes of medicine, and the arts, it is prepared artificially, either by decomposing sulphate of iron<sup>1</sup> by the process of distillation in close vessels; or by the combustion of sulphur. The first mode is the most ancient,

---

<sup>1</sup> Hence the old names, oil of *vitriol* and *vitriolic acid*, which are still the commercial names of this acid, from *green vitriol*, the old name of the sulphate of iron. This method is still practised at Nordhausen in Germany. The acid is disengaged from the oxide of iron by the heat; but, owing to the small quantity of water which it contains, it is of greater specific gravity and more fuming than the acid which is manufactured in this country. The specific gravity of the fuming acid of Nordhausen is 1.896.



and is still employed in several places on the continent; but the second is that generally adopted by the manufacturers in Great Britain, and therefore requires particularly to be described.

Into a chamber lined with sheet-lead, having no opening but a small door placed a few inches from the floor, and made to shut very close, water is poured so as to cover the floor, and rise upon it to the height of one or two inches. A stand is then introduced, on which is placed an earthen pot, containing a mixture of nine parts of refined Sicilian sulphur, and one of nitre, which is kindled by means of a red-hot iron, and the door instantly closed.<sup>1</sup> The oxygen of the air of the chamber, and that of the nitre, keep up the combustion of the sulphur, and form sulphurous acid gas, and nitrous acid, which, uniting, lose their gaseous character, and coming in contact with the water below, the sulphurous acid is changed into sulphuric acid at the expense of the nitrous acid, which being thus robbed of its oxygen, escapes in the form of nitrous gas. This gas now rising to the roof of the chamber, towards the aperture, attracts a fresh supply of oxygen, and becomes again nitrous acid, which descending, owing to its density among the sulphurous acid gas, once more converts it into sulphuric acid, and is again disengaged when it comes in contact with the water, to undergo fresh combinations and decompositions, which go on successively until the whole of the sulphurous acid is changed into sulphuric acid. But repeated charges of the sulphur and nitre are required to be constantly introduced at intervals of six or eight hours, for the space of two or three weeks, before the water is sufficiently acidified. It is then drawn off through a leaden pipe with a stop-cock at the bottom of the chamber; and contains, besides sulphuric acid, some sulphurous acid, and a portion of nitric oxide. The liquor is at first of a brownish colour; but after it is concentrated and purified, first, by evaporation in leaden boilers, and afterwards by boiling in large green glass retorts, or in platina alembics, it becomes a colourless, dense fluid, having, when perfectly pure, a specific gravity not exceeding 1.8485; and is brought to market in large globular glass bottles, surrounded with wickerwork, and sold under the name of *oil of vitriol*.

Sulphuric acid thus prepared is not perfectly pure, but is united with about three or four per cent. of saline matter, which consists of two-thirds of sulphate of potass, (owing to the acid uniting with the potass of the nitrate employed in the combustion of the sulphur,) and one-third of sulphate of lead

---

<sup>1</sup> This process was first used by Dr. Roebuck in 1749. *Edin, Phil. Trans.* vol. iv.



derived from the chambers in which it is manufactured. Both these impurities are precipitated by adding three parts of distilled water to the acid, which can again be concentrated by distillation: thence the addition of water is a good test of the purity of this acid. The lead is thrown down in the form of a white insoluble powder; from which the pure acid can be decanted. The amount of these impurities may be readily ascertained by evaporating a definite weight of the acid in a platina cup, placed on the red cinders of a common fire.<sup>1</sup> It is sometimes adulterated with sulphate of potass, for the purpose of increasing its specific gravity; in which case the best method of detection is to saturate the suspected acid with ammonia, and then to expel the sulphate of ammonia thus formed by a red heat: the sulphate of potass with which the acid was adulterated will remain fixed. According to Dr. Ure, the pure liquid acid, which is, accurately speaking, a hydroacid, consists of about 81.54 parts of real acid, and 18.46 of water; and the elements of the real acid, according to the estimate of Dr. Thomson<sup>2</sup>, which is probably the most accurate on this subject, are 40 of sulphur, and 60 of oxygen, in 100 parts of acid; or one atom of sulphur, and three atoms of oxygen: but with regard to these proportions chemists are not agreed.<sup>3</sup>

*Qualities.* — Liquid sulphuric acid, when pure, is as colourless and transparent as water<sup>4</sup>, inodorous, corrosive, heavy; and has the consistence of oil: its specific gravity at a temperature of 60° Fah. is 1.8485, and one fluid-ounce weighs fourteen drachms. It has all the generic characters of an acid; reddening the vegetable blue colours when diluted with a sufficiency of water; but, when undiluted, it turns vegetable yellow colours brown. Even when largely diluted, it has an intensely acid taste. When rubbed between the fingers it feels at first unctuous, owing to its dissolving the cuticle, and afterwards excites a burning sensation. It freezes at 15° into six-sided prismatic crystals<sup>5</sup>, bevelled at both extremities; and, when of the specific gravity 1.845, boils at 590°.<sup>6</sup> It attracts water so rapidly from the atmosphere, as to increase its weight one-third in twenty-four hours, and to double its

<sup>1</sup> Dr. Ure, who suggested this test, says, "if more than 5 grains of matter remain from 500 of acid, we may pronounce it sophisticated." *Journal of Science and the Arts*, vol. iv. p. 115.

<sup>2</sup> *System of Chem.* 5th ed. vol. i. p. 288.

<sup>3</sup> Bucholz makes the proportions to be 57.5 of sulphur and 42.5 of oxygen.

<sup>4</sup> Sulphuric acid is so easily coloured by contact with either vegetable or animal matters, without being deteriorated, that we cannot regard the slight colour which it acquires from these circumstances as any indication of its impurity.

<sup>5</sup> Macnab, *Hudson's Bay*.

<sup>6</sup> Dalton.



weight in the course of a month, (hence the necessity of keeping the air excluded from it;) and at the moment it unites with water the temperature of the mixture is much raised. It acquires a brown colour when mixed with any vegetable matter<sup>1</sup>, and therefore bottles in which it is kept must be stopped with glass stoppers. When brought to sp. gr. of 1.780, by being diluted with water, it boils at 435°, and freezes at 45°, or 13 degrees above the freezing point of water; a fact important to trading chemists, as in this state it is apt to burst the bottles in which it is kept, by its expansion in the act of freezing.<sup>2</sup> It forms neutral salts, termed *sulphates*, with the alkalies, earths, and metallic oxides, and decomposes the alkaline and earthy sulphurets. The strength of this acid is readily determined by saturating it with dry subcarbonate of soda, 100 grains of which neutralize 92 grains of the pure fluid acid, of the sp. gr. 1.848.

*Medical properties and uses.* — This acid is a valuable tonic, astringent, and antiseptic; but as it is employed internally in a diluted state only, its medicinal powers shall be explained under the article *Acidum Sulphuricum dilutum*. Although it powerfully corrode the skin, yet, on account of its fluidity, it cannot be applied as an escharotic: but when united with sixteen times its weight of lard, it forms an ointment which has been successfully used in the cure of scabies. It is a strong corrosive poison. When it has been taken into the stomach, either accidentally or as a poison, the best antidotes are chalk and water, or magnesia and water.<sup>3</sup>

Officinal preparations. *Acidum sulphuricum dilutum*, L. E. D. *Acid. sulphuricum aromaticum*, E. *Sulphas potassæ*, E. *Ferri sulphas*, L. E. D. *Hydrargyri Oxymurias*, L. E. D. *Sub-sulphas hydrargyri flavus*, E. *Zinci sulphas*, L. E. D.

ACIPENSER. *Syst. Nat. Gmelin*. 1483.

Cl. 4. Ord. 6. Pisces, Chondropterygii.

G. 134. Head obtuse; mouth far beneath the head, without teeth; *cirri* four under the nose, before the mouth. *Branchial apertures* lateral. *Body* elongated, with many series of angular tubercles.

Spec. 2. *Acipenser Ruthenus*. The Sterlet, or small sturgeon.

Spec. 3. *Acipenser Huso*. The Beluga, or great sturgeon.

Officinal. *ICHTHYOCOLLA*, *Dub.* Isinglass.

<sup>1</sup> This is owing to its strong affinity for water, breaking the affinities which exist between the vegetable components, so as to occasion the hydrogen and oxygen to unite and form water, while the carbon is precipitated.

<sup>2</sup> See *Parke's Chemical Essays*, vol. ii.

<sup>3</sup> I witnessed a case in which a fluid ounce of the strong acid was swallowed, with the intention of committing suicide, by a farrier; and, although it remained an hour in the stomach before it was ejected by vomiting, yet, the person recovered. T.



*Syn.* Ichthyocolle (*F.*), Hocusenblase (*G.*) Huisenblas<sup>1</sup> (*Dutch*), Huusblas (*Dan.*), Husblas (*Swed.*), Harukbi (*Pol.*), Colla di Pisce o Ittiocolla (*I.*), Cola de Pescade (*S.*), Cola de Peixe (*Port.*).

The beluga, the sterlet, the kester, the sturgeon, and other species of *Acipenser*, are caught in the Volga, Danube, Ural, Oby, and Irtysh rivers; and the Caspian sea. The sturgeon is oviparous. The body is pentagonal, in length from 5 to 24 feet, of a dirty olive colour on the upper part, which is studded with five rows of bony tubercles, one row on the back and two rows on each side; while the underside of the fish is flat, and of a silvery white colour. There is a single dorsal fin near the tail, one ventral also near the tail, one anal, and two pectoral fins. The tail is bifurcated; and the upper part longer than the under. The swimming bladder is very ample, and communicates by a large aperture with the æsophagus. The flesh of the sturgeon is agreeable to the palate, particularly that of the sterlet; and the roe, prepared in a particular manner, is eaten as a delicacy under the name of Caviar.

Isinglass is prepared in Russia from the air-bladders or sounds of all the species of the sturgeon; and in Lapland it is made from several species of the perch.<sup>2</sup> That however, which is made from the beluga is reckoned the best. The process, which is carried on during summer only, as frost, it is said, impairs its qualities, varies in different places: and on this account the quality of the isinglass made from the same fish at one place differs from that which is made at another. On the Volga, the sound is taken from the fish, slit open, well washed, and freed from the thin membrane which envelopes it; then exposed to stiffen a little in the air, rolled, and pinned, by means of a wooden peg, into a heart shape; in other places it is folded into leaves, or simply dried without any care, or boiled.<sup>3</sup> The first kind is the best. Isinglass is imported from Petersburg in bales. Four sorts are brought; *long staple*, *short staple*, *book* and *leaf*. “The finest is that which has the longest *staple*, as it is called: it is the thinnest and most flexible;” and is perfectly devoid of odour and taste. It should be composed of dry, whitish, nearly transparent, inodorous membranes.

Persons in London subsist by picking the staple into shreds, in which state it is generally sold; but as it admits of adulteration when picked, it should always be purchased in the staple.

*Qualities.*—Isinglass is insipid, and inodorous. When soaked

<sup>1</sup> The English name is a corruption of the Dutch, which means floating bladder; being derived from *hyzen*, to hoist, and *blas*, a bladder.

<sup>2</sup> Isinglass might be made from the sound of the cod-fish, *Gadus merrhua*; hake, *Gadus merluccius*; and ling, *Gadus molva*, which frequent the British seas.

<sup>3</sup> *Philosophical Transactions*, vol. lxxiii. part i.



in cold water it swells, softens, and becomes opalescent; and in one hundred grains of it, rather more than ninety-eight, according to Mr. Hatchett, are soluble in water, and the two parts which are insoluble consist of phosphates of soda and of lime. Three drachms of good isinglass, dissolved in one pint of warm water, produce on cooling a pretty firm, slightly opaline-coloured jelly, which is a compound of pure animal gelatin and water.<sup>1</sup> Gelatin is soluble in the acids and pure alkalies; but is precipitated from its solution by infusions and decoctions of astringent vegetables, the tannin of which forms with it an insoluble compound<sup>2</sup>; carbonate of potass also throws down a precipitate; and alcohol separates it from water when added to its solution in any considerable quantity. The solution putrefies when kept for a few days. According to John, isinglass contains also traces of saline matter.

*Medical properties and uses.* — The solution of ichthyocolla was formerly much given in cases of fluor albus and diarrhœa; but is now rarely used as a medicine. Its nutrient qualities are more obvious. By dissolving it in water, and adding a little sugar and lemon-juice, an excellent nutritious jelly is produced, well adapted for the sick and convalescent. It is employed in the preparation of English court-plaster.

ACONITUM.<sup>3</sup> *Spec. Plant. Willd.* ii. 1235.

*Cl.* 13. *Ord.* 3. Polyandria Trigynia. *Nat. ord.* Multisiliquæ, *Linn.* Ranunculaceæ, *Juss.*

*G.* 1062. *Cal.* none. *Petals* five, the highest arched. *Nectaries* two, peduncled, recurved. *Pods*, three or five.

\*\* *With blue corollas.*

*Sp.* 8. *A. Napellus*. Wolfsbane, Common Monkshood. (*Lond. Edin.*) *Med. Bot.* 2d ed. 461. *t.* 165.

*Sp.* 9. *A. neomontanum*, (Dub.) Jaquin. *Flor. Austr.* *t.* 381.

*Officinal.* ACONITI FOLIA, *Lond.* ACONITI NAPELLI FOLIA, *Edin.* ACONITUM; FOLIA, *Dub.* The leaves of Monkshood.

*Syn.* Aconit, chaperon de Moine (*F.*), Blauer-strumhut (*G.*), Monnikskapen (*Belg.*), Stermhut (*Dan.*), Stermhatt (*Swed.*), Napello (*I.*), Aconito (*S.*).

The species of aconite cited by the London and Edinburgh colleges has been, generally, regarded as the plant originally used by Stoerk: an error which originated with Stoerk himself; but, after a minute comparison of the plant with the de-

<sup>1</sup> As Gelatin is a constituent of many other substances, it is important to know that sulphate of platina will detect its presence, when the quantity is too small to affect astringent infusions. Mr. E. Davy discovered this test. *Journ. of the Sciences*, vol. x. p. 454.

<sup>2</sup> Gelatin is a nice test of the presence of tannin, for which purpose Sir H. Davy recommends 120 grains to be dissolved in 20 ounces of distilled water.

<sup>3</sup> *Labor. chim.* 1808.

“ Quo quia nascuntur dura vivacia caute  
Agrestes Aconita vocant : ”

OVID.

But Theophrastus derives the name from *Acone*, a city of Bithynia, near which it grew in great abundance.



scription, we agree with Willdenow, and the Dublin college, that the *neomontanum* is the species which should be medicinally used. It is a perennial plant, flowering in July: and found native in the alpine forests of Carinthia, Carniola, and the mountainous parts of Germany.

The stem is firm, elongated, rising to the height of five or six feet, leafy, and terminating in a long sparse spike of flowers, racemose and branched below, with single flowered peduncles. The lower leaves are few, alternate on long channelled petioles; palmated or rather pedate, being divided to the base into five broad cuneiform divisions, deeply cleft and toothed; the petioles are shorter, and the leaves less divided the nearer they are to the summit of the stem: the colour of the whole is a deep green on the upper disk, and a pale green on the under; both sides are naked, smooth, and shining. The flowers are of a deep violet colour, and stand alternately on the spikes, on unifloral, erect, axillary peduncles. They have no calyx; but two small, erect, calycinal stipules are placed one on each side of the peduncle within a few lines of the flower; the petals are five, the uppermost helmet-shaped and beaked, covering two singular, peduncled nectaries; the lateral ones broad, and roundish, the lower oblong, elliptical, and divaricating. These four are slightly pubescent. The nectaries are cuculatted, the spur of each being hooked and blunt; the lip lanceolate, revolute, and bifid. The filaments are spread, and white at the base, where they closely cover the germens; but the upper part is filiform, purple, spreading, and bearing whitish anthers. The germens are three, four, or five, with simple reflected stigmas; and become capsules, containing many angular seeds.<sup>1</sup>

For medicinal use the leaves should be gathered when the flowers appear.

*Qualities.* — Aconite leaves when fresh have a faint narcotic odour; and a moderately bitter, acrid taste, leaving a painful sensation of heat in the mouth, when they are much chewed. The whole of the plant is poisonous; but the deleterious qualities are lost in a considerable degree when it is dried, or long kept, and much of its acrimony is dissipated. Its narcotic principle is supposed by M. Brandes to be an alkali, which he has named *aconita*.

*Medical uses and properties.* — Aconite is narcotic, diaphoretic, and in some cases diuretic. In over doses it occasions

---

<sup>1</sup> The flower-spikes of *A. Napellus* are shorter, denser, and the segments of the leaves narrower, and more linear than those of *A. Neomontanum*; the spur of the nectaries is also straight in *Napellus*, and hooked in *Neomontanum*; the lip ascending in *Napellus*, and revolute in *Neomontanum*.



violent nausea, vomiting, hypercatharsis, vertigo, cold sweats, mania, and convulsions which terminate in death; and these effects appear to depend on its action on the nervous system, as dissections of fatal cases have not displayed any particular marks of organic disease.

Stoerk first administered Aconite internally in chronic rheumatism, gout, exostosis, paralysis, and scirrhus; and since the publication of his experiments, in 1702, it has been advantageously employed in similar cases, and also in amaurosis, scrophula, cancer, itch, venereal nodes, and intermittents. Much caution is required in the exhibition of it; and it is absolutely necessary to know the length of time it has been gathered, as its activity varies so very considerably, as to require this to be ascertained before the dose can be apportioned. It is given in the form of powder, extract, and tincture; and may be combined with calomel, antimonials, camphor, and guaiacum. The dose of the powder is one or two grains, gradually increasing it to six or eight.

Officinal preparations. *Extractum Aconiti.* L. E.

ACORUS. *Spec. Plant. Willd.* ii. 199.

Cl. 6. Ord. 1. Hexandria Monogynia. *Nat. ord.* Piperitæ, Linn. Aroideæ, Juss.

G. 663. *Spadix* cylindrical, covered with florets. *Cor.* petals six, naked. *Style* O. *Capsule* three-celled.

*Species* 1. *A. Calamus.* The Sweet-flag. *Med. Bot.* 2d edit. 725. t. 248. *Smith Flor. Brit.* i. 373.

*Officinal.* CALAMI RADIX, *Lond.* ACORI CALAMI RADIX, *Edin.*

ACORUS, (CALAMUS AROMATICUS,) RADIX, *Dub.* Sweet-flag.<sup>1</sup>

*Syn.* Acorus odorant (*F.*), Kalmus wurtzil (*G.*), Kalmus (*Dutch, Dan. Swed.*) Calamo aromatico (*I.*), Acoro calamo (*S.*), Tatarskie Ziele (*Polish*), Vassamboos (*Tam.*), Vadge (*Pers.*), Shiva Buch (*Beng.*), Igir (*Arab.*), Wudda Kuha (*Cyng.*), Bach (*H.*), Vacha (*San.*), Deringgo (*Javanese*).

The sweet-flag is found growing in marshes and rivulets, over the greater part of Europe and Asia, and in Britain, in many parts, producing its flowers in May and June.<sup>2</sup>

The root is perennial, horizontal, long, somewhat flattened, crooked; full of joints or rings, whence the radical fibres spring from the underside, and branched; from half an inch to one inch in thickness; externally, when fresh, of a greenish yellow colour, internally, whitish and spongy. The leaves spring directly from the root; are sword-shaped, about three feet in length, and generally waved along one of the edges; of a bright green colour, and emitting a strong aromatic odour when bruised. The flowers are small, and produced on

<sup>1</sup> Καλάμου ἀρωματικού, Dioscoridis.

<sup>2</sup> In the rivers of Norfolk plentiful. On Hillington-common, Middlesex, and other places about London. *Smith.* l. c.



a very close tessellated conical spike four inches long, pushed out from the side, above the middle, of a naked stalk or scape, two-edged, and terminating like a leaf. They have no calyx. The petals are six: small, erect, regular, with the apex inflected, and of a pale green colour. The filaments are alternate, with the petals, thread-like, supporting double anthers. The germen is elliptical, crowned with a sessile-pointed stigma.

The greater part of the roots in use are now brought from Norfolk; and are equal in quality to those from the Levant.

*Qualities.* — The root of sweet-flag has a pleasant aromatic odour, similar to that of a mixture of cinnamon and allspice; the taste is warm, bitterish, pungent, and aromatic.<sup>1</sup> In the dried state, the cuticle is corrugated, of a yellowish brown colour, with many small, white, elevated circles on the under side, whence the radical fibres issued. It breaks with a short rough fracture; is internally of a pale buff colour, and a spongy texture: both the smell and taste are improved by exsiccation. The aromatic principle is an essential oil, which can be obtained by distillation, and, with the bitter matter, is extracted by infusion in boiling water. Like the root of Florentine iris it contains a considerable quantity of fecula, which is dissolved in the infusion, and copiously precipitated from it by acetate and superacetate of lead.

*Medical properties and uses.* — This root is tonic, and aromatic. It has been employed in medicine since the time of Hippocrates.<sup>2</sup> By the moderns, it is successfully used in intermittent fever, even after bark has failed; and is certainly a very useful addition to Cinchona. It is also a useful adjunct to bitters, and stomachic infusions, in cases of dyspepsia; particularly when vertigo is one of the symptoms. It is too seldom prescribed. The dose in substance is from ℥j. to ʒj.; and of the infusion, made with ʒvi. of the bruised root in f ʒxiij. of boiling water, a cupful three or four times a day.

ADEPS. Vide *Sus Scrofa*.

ÆSCULUS. *Spec. Plant. Willd.* ii. 285.

Cl. 7. Ord. 1. Heptandria Monogynia. *Nat. ord.* Trihilatæ, *Linn.*  
*Acera, Juss.*

G. 717. *Cal.* one-leafed, five-toothed, swelled out. *Cor.* four or five irregularly coloured petals inserted into the calyx. *Cap.* three-celled.

*Species 1. Æ. Hippocastanum.* Common Horse Chesnut. *Med. Bot.*  
*2d edit. t.* 217. *Gærtner De Fructibus*, ii. 135.

<sup>1</sup> Linnæus erroneously considered it the only native aromatic plant of northern climates. — The candied root is employed at Constantinople as a preservative against epidemic diseases.

<sup>2</sup> *Morb. Mul.* ii. 651.



*Officinal.* CORTEX, *Dub.* The bark.

*Syn.* Marronier d'Inde (*F.*), Ross kastanie (*G.*), Paarden kastanie (*Dutch*), Hestekastagne (*Dan.*), Häskastagnier (*Swed.*), Kasztan owdzikich (*Pol.*), Ippocastano (*I. Port.*), Esculo Castano de Aballo (*S.*).

The horse chesnut is a native of the north of Asia, but cultivated in almost every part of Europe.<sup>1</sup> In this country it attains to great perfection, constituting one of the chief ornaments of our parks and avenues. It flowers in May.

This tree rises to a considerable height, branching, from the upper part of the trunk, so as to form a roundish conical mass. The leaves stand on long foot-stalks, are large, and digitated, each consisting of seven leaflets, proceeding from a common centre; of a deep green colour, and spathulate form; ribbed, pointed, and serrated. The middle leaflet of a full-grown leaf is about six inches long, and two broad near the top, the three lateral ones on each side gradually decrease in size, to the smallest, which are only three inches in length and one in breadth. The flowers are peduncled, and disposed in handsome, terminal, erect, conical spikes. The calyx is one-leaved, of a pale green, bell-shaped, divided at the edge into five blunt teeth; the petals are five, ovate, slightly waved at the margin, ciliated, with claws inserted into the calyx, all inclining to one side, so as to leave a gap betwixt the two innermost; white, with an irregular spot of red or yellow directly above the claw. The filaments are awl-shaped, gracefully curved, supporting reddish, upright double anthers: the germen is cylindrical, with a pointed stigma. The fruit is a large, roundish, green-coloured, three-celled, three-valved capsule, having the bark armed with short spines, and generally containing two sub-globular seeds or nuts.

The fruit consists almost entirely of secula. It is eaten with avidity by deer and some other animals; and were the acrimony destroyed by fire, it might be rendered fit food for men in times of scarcity. For medical purposes, the bark is taken from the middle-aged branches.

*Qualities.*—The *bark* is inodorous, and tastes bitter, astringent, and slightly aromatic: both water and proof-spirit extract its virtues. Its infusion has a fawn-brown colour, and a bitter taste; sulphate of iron and sulphate of zinc strike a black colour when added to it, and throw down a dark-coloured precipitate; oxymuriate of mercury and superacetate of lead precipitate it white. Tartar emetic effects no change on it. Gelatin precipitates tannin from it, but much less than from oak bark.

---

<sup>1</sup> The horse chesnut was brought into Europe by Clusius in 1550, and first cultivated in England by John Tradescant in 1633.



From the above detail, those substances which are incompatible in prescriptions with infusion or decoction of this bark are obvious.

*Medical properties and uses.*—The *bark* of the horse chesnut is tonic, and has been successfully exhibited, particularly on the continent, in intermittents, typhus, and other cases in which *Cinchona* is used; externally, its decoction has been employed as a lotion in gangrene. We have had no opportunity of trying this bark; but we doubt much if it can supersede the *Cinchona*, in any case in which it is properly indicated.

The dose of the powder is ʒ ss.; and of the strained decoction, (made with an ounce of the bark and a pint of water,) fʒjss. or fʒij. It may be given every three or four hours.

AGRIMONIA. *Spec. Plant. Willd.* ii. 875.

Cl. 11. Ord. 2. Dodecandria Dygynia. Nat. ord. Senticosæ, *Linn.*  
• Rosaceæ, *Juss.*

G. 957. *Calyx* five-toothed, guarded by another. *Petals*, five.  
*Seeds*, two, in the bottom of the calyx.

*Species* 1. *A. Eupatoria*. Common Agrimony.<sup>1</sup> *Med. Bot.* 2d edit.  
500. t. 180. *Smith, Flor. Brit.* 511. *Flor. Dan.* t. 588.

*Officinal.* AGRIMONIA; HERBA, *Dub.* Agrimony; the herb.

*Syn.* Aigremoine (*F.*), Odermennig (*G.*), Agrimonie (*Belg.*), Agermaane (*Dan.*), Akermünja (*Swed.*), Rzepik (*Polish*), Kleine klette (*Russian*), Agrimonia (*I.*).

This is an indigenous perennial plant, common about the borders of fields and hedges, flowering in June and July.

The stem rises two feet in height; is erect, leafy, angular, hairy, and of a reddish hue. The leaves are alternate, interruptedly pinnate, with three pair of opposite leaflets, and a terminal one somewhat lyre-shaped and hairy; the leaflets are obovate, pointed, deeply serrated; the terminal one is petiolate; the intermediate ones are much smaller, entire or trifid. Stipules two, deeply cleft. The flowers, which are small, golden yellow, and peduncled, are in long, terminal, erect, hairy spikes, with trifid bractes. The calyx is persistent, inclosed within an involucre at the base of the germen, furrowed, and divided into five ovate, pointed segments, surrounded with rigid, hooked awns: the corolla consists of five ovate, spreading petals inserted into a glandular substance at the base of the germen. The filaments are from five to twelve; the anthers two-lobed. The germen is inferior, supporting two styles with blunt stigmas. The seeds two, one often abortive, contained in a capsule formed of the calyx.

For medical use, this herb should be cut when it is in full flower.

<sup>1</sup> Εὐπατωρία, Dioscoridis.



*Qualities*.—Agrimony, when fresh, has an agreeable aromatic odour, depending on a volatile, essential oil, which is lost when the herb is dried<sup>1</sup>: the taste is bitterish and subastrigent. The infusion of it in water reddens the more delicate vegetable blues, and strikes a black colour with sulphate of iron. Potass and its carbonate first change it to a yellow, then to an orange-colour, and, lastly, throw down a white precipitate.

*Medical properties and uses*. — Agrimony was formerly regarded as a remedy of much importance, as a tonic and deobstruent: but it is now very seldom or never prescribed. The dose, in powder, is from ℥j. to ʒj. given two or three times a-day.

ÆRUGO. See *Cuprum*.

ALLIUM. *Spec. Plant. Willd.* ii. 63.

Cl. 6. Ord. 1. Hexandria Monogynia. Nat. ord. Spathaceæ, Linn. Asphodeli, Juss.

G. 626. Corolla six-parted, spreading. Spathe many-flowered. Umbel heaped together. Capsule superior.

\* Stem-leaves plane. Umbel bearing a capsule.

Species 2. *Allium porrum*. The Leek.

\*\* Stem-leaves plane. Umbel bulbiferous.

Species 14. *Allium sativum*. Garlic. *Med. Bot.* 2d edit. t. 256.

\*\*\*\* Leaves radical. Stem naked.

Species 43. *Allium Cepa*. The Onion.

#### 1. ALLIUM PORRUM.<sup>2</sup>

*Officinal*. PORRI RADIX, *Lond.* Leek root (bulb).

*Syn.* Poireau (*F.*), Spanische lauch (*G.*), Porro (*I.*).

The leek is a biennial plant, a native of Switzerland, flowering in June.

The bulb consists of concentric circles, seated on a radical plate, from which spring fibrous roots. The leaves are broad and gramineous: the stem is a naked scape, bearing a congested capital, or rather spherical umbel of flowers; with rough-keeled petals, shorter than the stamens, which are alternately trifid. It is better known as an alimentary or pot herb than as an article of *Materia Medica*.

*Qualities*.—All the parts of the plant have a pungent offensive odour, and an acrimonious taste: properties depending on an essential volatile oil, which is much dissipated by boiling, and can be consequently separated by distillation.

*Medical properties and uses*.—The leek is stimulant and diuretic. The expressed juice has been given with advantage in ascites and other dropsies. The dose is from f ʒ ss. to f ʒ ij. mixed with mucilage or syrup.

<sup>1</sup> The oil, which is yellow, can be obtained by distillation with water. — *Lewis*.

<sup>2</sup> Πράσον, Theophrasti et Dioscoridis.



2. ALLIUM SATIVUM.<sup>1</sup>

*Officinal.* ALII RADIX<sup>2</sup>, *Lond.* ALLII SATIVI RADIX, *Edin.* ALLIUM; RADIX, *Dub.* Garlic root (bulb).

*Syn.* Ail (*F.*), Knoblauch (*G.*), Aglio (*I.*), Ajo sativo (*S.*), Knoflook (*Belg.*), Kvidlögen (*Dan.*), Hvitloken (*Swed.*), Czosnek (*Polish*), Tschesnok (*Russian*), Seer (*Pers.*), Soom (*Arab.*), Bawang (*Jav.*), Lasúna (*San.*), Lehsen (*H.*), Vullay poondoo (*Tam.*), Bavangpootie (*Malay*).

Garlic is a perennial bulbiferous plant, found wild in Sicily, and cultivated in most parts of Europe, for culinary and medicinal use. It flowers in July.

The bulbs of this species of allium are numerous; three or more being inclosed in one covering, forming a nucleus, round which others are disposed, and the whole enveloped in a common membrane, from the base of which proceed long white fibrous roots. The stem rises two feet in height, surrounded with many long, flat, linear, grass-like leaves, proceeding chiefly from the young bulbs; and is terminated by a mixed cluster of flowers and bulbs, inclosed in a spathe, which opens at one side and withers. The flowers are small, consisting of six oblong white petals, with tapering filaments, shorter than the corolla, and supporting erect anthers: the germen is superior, short, angular, bearing a simple style with a sharp stigma; and becoming a short, broad, three-celled capsule, containing roundish seeds.

Garlic is dug up for use in the month of August, then cleaned and dried in the sun, and preserved in bunches in a dry place. In this state the exterior membrane is of a dirty white colour, and of a withered aspect; but the bulbs, which are called *cloves*, are white, succulent, and juicy. On drying, they lose nine parts in fifteen of their weight.

*Qualities.*—All the parts of the plant, but particularly the bulbs, have a pungent offensive odour, and an acrimonious biting taste. These properties depend on an essential oil, which can be obtained separate by distillation with water; of a thick and ropy consistence, a yellow colour, heavier than water, and possessing, in an eminent degree, the sensible qualities of the garlic. It blisters the skin when applied to it, and strikes a black colour when triturated with oxide of iron. Simple coction with water renders garlic mild and inert. The acrid principle is

<sup>1</sup> Σκόροδον, Theophrasti et Dioscoridis.

<sup>2</sup> The root, strictly speaking, is not the part of the plant intended to be ordered. This error has arisen from the bulbs of plants having been generally placed by botanists among the roots, under the title “radices bulbosæ:” they, however, have no affinity to roots, but the closest to leaf buds. The roots of bulbiferous plants are fibrous, and issue from a radical plate at the base of the bulb. The clearest definition of the bulb is given by Gärtner, in his work *De Fructibus*, &c. Introduction, p. 111. “Bulbus est germen compositum, subglobosum, subaphyllum, ex carina brevissima, et squamis succulentis, crassis compaginatum, quod tandem sponte a matre sua solvitur.” The Colleges, therefore, should have used the word *bulbus* instead of *radix*.



obtained also by expression : and it is in a less degree extracted by water, by alcohol<sup>1</sup>, and by acetic acid. The odour is so penetrating, that when garlic is applied to the soles of the feet it is perceived in the breath, the urine, and the perspiration. From 1406 parts of fresh garlic, Cadet<sup>2</sup> obtained 520 of mucilage, 37 of albumen, 48 of fibrous matter, and 801 of water by estimate. Bouillon la Grange<sup>3</sup> found sulphur also, with vegetable albumen, and sugar.

*Medical properties and uses.*—Garlic is stimulant, diaphoretic, expectorant, diuretic, and anthelmintic, when exhibited internally; and rubefacient when externally applied.

It has been successfully given in intermittents, and in fevers of the typhoid type. If the body be kept warm during its use, it acts powerfully by diaphoresis. It has long been esteemed a valuable remedy in pituitous asthma, chronic catarrh, flatulent colic, calculus, and dropsies; and as a preventive of worms. Externally, it is applied bruised to the soles of the feet, in the coma of typhus; and in confluent small-pox when the determination to the head is considerable. A poultice made of it is a good resolvent of indolent tumors. A clove of it, wrapped in cotton or gauze, or a few drops of the juice introduced into the external ear, is said to be extremely efficacious in atonic deafness; and applied to the pubis as a poultice in retention of urine, owing to a want of action in the bladder, it sometimes is effectual in stimulating that viscus to discharge its contents. The juice is also applied, united with oil, to herpetic eruptions.

Garlic may be exhibited in substance, the whole clove or pieces of it being dipped in oil and swallowed; or it may be formed into pills. The expressed juice also is given mixed with sugar: or the bulb may be infused in milk, which was Rosenstein's mode of administering it to children afflicted with worms. It is frequently united with calomel in the form of pill or bolus, in hydropic cases. An ointment is formed by mixing the juice with oil. The bruised bulb has also been used as a suppurative.

The dose, in substance, is from ʒss. to ʒij.; or from one to six cloves, swallowed whole, twice or thrice a day; and in pills, united with soap or calomel, from grs. xv. to ʒj. Of the juice, f ʒss. is given for a dose in any proper vehicle.<sup>4</sup>

An overdose, or the too liberal use of it as a condiment, is apt to occasion head-ache, flatulence, thirst, fever, inflammation, and discharges of blood from the hæmorrhoidal vessels.

Officinal Preparation. *Syrupus Allii*, D.

<sup>1</sup> With alcohol, a reddish yellow tincture is obtained, which leaves, when evaporated, a very acrid brown extract, that attracts moisture from the air.

<sup>2</sup> *Ann. de Chimie*, lix. p. 106.

<sup>3</sup> *Journ. de Pharm.* No. viii. p. 357.

<sup>4</sup> Its use as a condiment is of great antiquity. Thus, when the Israelites remembered the flesh pots of Egypt, among other deprivations, which the malecontents enumerated, were "the leeks, and the onions, and the garlick," Numbers, c. 11. v. 5.



3. ALLIUM CEPA.<sup>1</sup>*Officinal.* CEPA ; RADIX, *Dub.* The Onion.

*Syn.* Ognon (*F.*), Swiebel (*G.*), Uijen (*Dutch*), Cipolla (*I.*), Cebolla (*S.*), Cebola (*Port.*), Lögen (*Dan.*), Rödlek (*Swed.*), Cebula (*Pol.*), Luck (*Russian*), Vungayum (*Tam.*), Loong (*Cing.*), Pecaz (*Pers.*), Bavangmera (*Malay*), Palandoo (*San.*), Pecáj (*H.*), Bassul (*A.*), Kanda (*Sans.*), Brangbang (*Jav.*).

The onion is a perennial, bulbiferous plant, cultivated all over Europe for culinary purposes : flowering in June. It is so well known, as scarcely to require a particular description. The bulb is simple, formed of concentric circles, with a radical plate at the base and fibrous roots. The stem is a naked, swelling scape, with fistular, pointed, spreading leaves, sheathing at the base. The flowers are produced in a capital or head, inclosed in a deciduous spathe.

*Qualities.*—The odour and taste of the onion do not materially differ from those of garlic, but are much weaker. A little acrid volatile essential oil, combined with sulphur<sup>2</sup>, is obtained by distillation; and the recent juice contains sugar, mucus, phosphoric acid, phosphate of lime, and citrate of lime.<sup>3</sup>

*Medical properties and uses.*—The onion, “considered as an article of food, when eaten liberally, is said to produce flatulencies, occasion thirst, headache, and turbulent dreams.” As a medicine, it is stimulant, diuretic, and expectorant, and may be used in the same cases as garlic. On account of the free phosphoric acid it contains, the juice is supposed to be useful in calculous cases, as it dissolves phosphate of lime out of the body. Onions are, however, scarcely ever employed, except externally, as suppurative cataplasms; for which purpose they are generally roasted, split, and applied to tumours.

ALOE. *Spec. Plant. Willd.* ii. 184.

*Cl.* 6. *Ord.* 1. Hexandria Monogynia. *Nat. ord.* Coronariæ, *Linn.* Asphodeli, *Juss.*

*G.* 659. *Corolla* erect, mouth spreading, bottom nectariferous.

*Filaments* inserted into the receptacle.

*Species* 2. Aloe *spicata*. Spiked Aloe.

——— 3. Aloe *vulgaris*. Common Aloe. *Sibthorp. Flor. Græc.*

1. ALOE SPICATA.<sup>4</sup>

*Officinal.* ALOES SPICATÆ EXTRACTUM, *Lond.* ALOES EXTRACTUM, *Edin. Dub.* Extract of Aloes.

*Syn.* Suc d'Aloes (*F.*), Glausinde Aloe (*G.*), Azevre Succotrina (*Portug.*), Aloe (*I.*), Aloe (*S.*), De doorblädige Aloe (*Belg.*), Aloe (*Russian*), Elwa (*H.*), Sehber (*Pers.*), Musebber (*A.*), Corriabolum (*Tam.*).

The spiked aloe is undoubtedly the species which yields the best extract brought from the Cape of Good Hope; and it is

<sup>1</sup> *Κρόμμον*, Dioscoridis. The specific appellation is derived from *caput*, a head, on account of the form of its bulb.

<sup>2</sup> It is this sulphureted oil which blackens silver plate in which onions are placed, and which occasions the disagreeable odour of this bulb in putrefying.

<sup>3</sup> Fourcroy and Vauquelin, *Ann. de Chim.* lxx. 161.

<sup>4</sup> Ἀλόη, Dioscoridis?



also supposed to yield the extract brought from the island of Zocotora; which was formerly the only place of export for the best aloes, thence named Socotrine aloes. It grows abundantly in the interior of the Cape, particularly at Zwellendam, near Mossel Bay. The stem is round, about four inches in diameter, leafy at the top, and rising three or four feet in height. The leaves are spreading, about two feet long, subverticillate, broad at the base, gradually drawn to a point, channelled, acute, with remote teeth. The flowers spread horizontally, in very close spikes. Under each flower is a single ovate, acute, broad, membranaceous bracte, white, with three green streaks, and a little shorter than the corolla. This is bell-shaped, and six-petalled: the three inner segments are white, marked with three green lines, not connected together, ovate, blunt, and broader than the three outer; which are connected with them at the base, and resemble them, but are narrower and less concave. The flower contains a large portion of a purple honey juice. The seeds, which are numerous, with a membranaceous appendage attached to each, are contained in a superior capsule.

At the island of Zocotora the leaves are cut off close to the stem, then cut in pieces, and the juice expressed: this is allowed to remain at rest for forty-eight hours, during which time a feculent matter is deposited: after which the supernatant liquor is poured off into flat dishes and evaporated in the sun. At Zwellendam, in the month of July, the leaves are pulled, then cut into pieces, the juice expressed, and inspissated by means of heat.

The real Socotrine aloes, which are now scarce in the market, and, as has been already stated, are supposed to come from the island of Socotora, or Zocotora<sup>1</sup>, near the straits of Babel-mandel, are brought to this country by way of Smyrna and Malta, in chests and casks. Those from the Cape are brought in similar packages. The greater part of what are now sold as Socotrine aloes are brought from Bombay, and are the real Hepatic aloes. The Bombay aloes are rather duller and browner, but in other respects have nearly the same characters as the Socotrine, which are sometimes mixed with the Indian. They are imported in casks containing from two to eight hundred weight, and sometimes in skins.<sup>2</sup>

*Qualities.*—The real *Socotrine* extract has a peculiar aromatic odour, not unlike that of the russet apple decaying; and a very permanent intensely bitter taste. It is in pieces of a deep reddish-brown colour, glossy as if varnished; breaking with a

<sup>1</sup> Zocotora is an island subject to the Prince of Hadecamaut, an Arabian province contiguous to Yemen.

<sup>2</sup> Mr. Barrow states, that the quantity of Cape aloes sent to London, from 1799 to 1802 inclusive, was 341,927 lbs. Vide *Travels in Africa*.



smooth conchoidal fracture. The thin edges and small fragments are reddish, or golden yellow, and semitransparent. It softens in the hand, and is adhesive; yet is sufficiently pulverulent; and the powder has a bright golden yellow hue.

The *Cape* aloes have a stronger and more disagreeable odour than the Socotrine and the Hepatic; the taste is nearly the same. The outside of the pieces is more friable, has more of a yellow cast, and is less glossy; but the inside is apt to continue soft and pliable. The colour of the powder is a beautiful greenish yellow, resembling gamboge, but less bright. They are imported in chests and casks.

## 2. ALOE VULGARIS.<sup>1</sup>

*Officinal.* ALOE HEPATICA; EXTRACTUM, *Edin.* GUMMI RESINA, *Dub.* Extract of the Common Aloe, or Barbadoes Aloes.

The British Pharmacopœias formerly considered the plant which yields the Barbadoes aloes as a variety of the *Aloe perfoliata*; but Sir E. Smith, the learned editor of Sibthorp's *Flora Græca*, has marked the above-named plant, which is a distinct species, as the one that Sloane describes, in his History of Jamaica, as producing the Barbadoes extract. The volume of Sibthorp's work, in which the plant will be fully described, and proved to be the true *Ἀλόη* of Dioscorides, is yet unpublished, and therefore we cannot avail ourselves of the description.

The month of March is the period for cutting the aloes in the island of Barbadoes. The leaves are cut off close to the stem, and disposed in tubs, in a such a manner that the juice runs out. After a sufficient quantity of it is collected, it is exposed to heat in copper boilers; and, as it becomes more inspissated by a constant and regular fire, it is ladled from one boiler to another, and fresh juice added, until that in the last, which is called the *teache*, acquires the consistence of honey; when it is poured into calabashes, and hardens by age. It is brought home in these calabashes, or large gourd-shells, each of which contains from sixty to seventy pounds' weight. They are often passed off as Hepatic aloes.

There is still another kind of aloes, named Foetid or Caballine, but on account of its foetid odour it is not used in medicine. A Mocha aloe resembling the Cape, but less purgative, is also brought to this country.

*Qualities.*—The odour of the Barbadoes aloes is stronger and less pleasant than that of the Socotrine, and has some resemblance to the odour from the human axilla. The taste is nauseous and intensely bitter. The pieces are also of a duller brown colour, less glossy, not so smooth in the fracture, but easily splinter. The edges are not so sharp and transparent;

<sup>1</sup> Ἀλόη, Dioscoridis, l. 3. c. 25.



but rather blunt, and of a dull yellowish hue. It softens in the hand, and is adhesive. The colour of the powder is a dull olive yellow.

All the kinds of aloes, when analysed, yield a small portion of vegetable mucus, resin, and a peculiar extractive matter. Braconnot found aloes to consist chiefly of a peculiar bitter matter, which he has termed the resinous bitter principle.<sup>1</sup> The odour, taste, and medical virtues of the drug reside chiefly in the extractive; and the superiority of the Socotrine, the Cape, and the Bombay aloes is correctly supposed to arise from their containing a larger proportion of it, and consequently less resin, than the Barbadoes. Boiling water dissolves nearly the whole of any of the kinds; but as the solution cools, the resinous part is deposited; and by boiling aloes in water the extractive is altered, rendered insoluble in water, and approaches in its properties to the nature of resin. The alkalies and their carbonates greatly assist its solution. All the kinds dissolve, also, in proof spirit. When the Socotrine aloes is distilled, a volatile oil is obtained, which is not procured from the Barbadoes.

*Medical properties and uses.*—Although all these kinds of aloes differ in their sensible qualities, yet they agree in their medical properties. They are warm, stimulating cathartics, of slow solution, and thence they are supposed to act chiefly on the colon, particularly its sigmoid flexure, and the rectum. By the extension of their stimulus to the uterine vessels they produce, also, emmenagogue effects. This operation is slow and moderate, but tolerably certain. From the stimulant property of aloes, they are useful in cases where the intestines are in a sluggish, relaxed, and insensible state, attended with viscosity of the abdominal secretions: as in the habitual costiveness of the sedentary and hypochondriacal; or that arising from a paucity of bile, in jaundice, chlorosis, and scrophula: and by their powerful effects on the rectum they have been found very serviceable in expelling ascarides. Their use is contraindicated in hæmorrhoidal cases, the symptoms of which they are apt to aggravate; but it has been well remarked that their effect upon the rectum is more owing to the manner of prescribing the remedy, than any specific action which it exerts on that portion of the intestines, which is but little affected when the aloes are prescribed in a soluble form. They are also improper in very irritable and plethoric constitutions in phthisis pulmonalis, and during the flow of the menses. Aloes, and aloetic compounds, have been likewise regarded as improper in pregnancy; but Dr. Denman has justly remarked, that “they are in common use among the lower class of peo-

---

<sup>1</sup> *Ann. de Chimie*, lv. 152.



ple, because they are cheap, and conveniently given in the form of pills<sup>1</sup> ;” and no bad effects are observed to follow.

Aloes may be given in substance, in doses from grs. v. to grs. xx.; larger doses not operating more effectually. Whether in the simple state, or when compounded with soap, bitters, and other substances, the form of pill is to be preferred on account of the nauseous taste of the medicine.<sup>2</sup>

Officinal preparations. *Pulvis Aloes comp.* L. *Pilulæ Aloes*, E. D. *Pil. Aloes comp.* L. *Pil. Aloes cum Myrrha*, L. E. D. *Pilul. Aloes et Assafætidæ*, E. *Pil. Aloes c. Colocynthide*, E. *Pil. Rhei comp.* E. *Pil. Scammonii comp. cum Aloe*, D. *Extractum Aloes*, D. *Extractum Colocynthidis comp.* D. *Tinctura Aloes*, L. E. D. *Tinct. Aloes comp.* E. D. *Tinct. Aloes ætherea*, E. *Tinct. Benzoini comp.* E. D. *Tinct. Rhei et Aloes*, E. *Vinum Aloes*, L. E. D.

ALTHÆA.<sup>3</sup> *Spec. Plant. Willd.* iii. 770.

Cl. 16. Ord. 8. Monadelphia Polyandria. Nat. ord. Columniferae, Linn. Malvaceæ, Juss.

G. 1289, Cal. double; the exterior 6 or 9-cleft. Caps. numerous, 1-seeded.

Sp. 1. *A. officinalis*. Common Marsh Mallow. *Med. Bot.* 2d edit. 552. t. 198. *Eng. Bot.* t. 147. *Smith's Flora Britan.* 3. 739.

Officinal. ALTHÆÆ FOLIA, ET RADIX, Lond. RADIX, Edin.

The leaves and root of Marshmallow.

Syn. Guimauve (F.), Eibisch (G.), Gemeene heemst (Belg.), Ibisk (Danish), Alte (Swed.), Slaz wielkilesny (Polish), Proswurujak (Russian), Altea (I.), Malvarisco (S.).

The marshmallow is an indigenous plant, which grows, as its name imports, in marshy places, particularly salt-marshes, and on the banks of rivers, throughout Europe. It flowers in June and July, and ripens its seeds in September. The root is perennial and fusiform. The stems are annual, herbaceous, upright, rising from two to five feet in height, round, naked and purplish below, but leafy and branching above. The leaves are alternate and petiolate, longer than they are broad, slightly five-lobed, and unequally serrated: both surfaces are downy, and give a soft velvety feeling when rubbed between the fingers. From the axillæ of the leaves the flowers spring in short thick panicles. Both the calyxes are persistent: the exterior has 7, 9, 10, or 12 very narrow unequal divisions; the interior is more regularly, but less deeply, cleft into five broader and sharper segments. The petals are five, cordate, coalescing

<sup>1</sup> *Introduction to Midwifery*, vol. i. 287.

<sup>2</sup> Dr. Paris, (see *Pharmacologia*), has enumerated the following empirical preparations as owing their efficacy chiefly to the aloes they contain. *Anderson's pills*, consisting of aloes, jalap, and oil of anniseed; *Hooper's pills*, formed of Pil. Aloes c Myrrha, sulphate of iron and Canella bark; *Dixon's antibilious pills*, a compound of aloes, scammony, rhubarb, and tartar emetic; *Speediman's pills*, of aloes, myrrh, rhubarb, the extract and the essential oil of camomile; and *Lady Webster's dinner pills*, for which the following is the formula, extracted from the old Paris-codex. R. aloes optimæ, ʒvj. mastiches, et rosarum rubrarum, ā a ʒij. syrupi de Absinthio q. s. ut fiat massa, in pilulas 120 dividenda.

<sup>3</sup> Ἀλθαία, Dioscoridis. Named from Althea, the mother of Meleagar.



at their bases, of a pale blush colour. The stamens are many, united at their bases into a tube, and support reniform anthers. The germen is orbicular, bearing a cylindrical style, divided into many stigmas, which rise above the anthers. The capsules, generally about twenty in number, are of a rounded kidney shape, united laterally in a circle, so as to form a flattened wheel-shaped seed-vessel; and each contains a solitary reniform, flattened, smooth, brown seed. The roots, which are the parts medicinally used, are dug up in autumn.

*Qualities*.—Marshmallow root is inodorous, mucilaginous when chewed, externally tough and of a yellowish colour, internally white and fibrous; and contains a very considerable portion of mucus<sup>1</sup>, which is yielded to water by coction.

*Medical use*.—The preparations of this plant, which derive their virtues from its mucus, are useful demulcents in visceral inflammations and calculous complaints. The roots well boiled, and bruised, are sometimes used as an emollient suppurative cataplasm; and a decoction of the leaves forms a useful fomentation in external abrasions; and in cutaneous eruptions accompanied with a sharp ichorous discharge.

Officinal preparations. *Decoctum Althææ officinalis*, E. *Syrupus Althææ*, L. E.

ALUMEN. *Lond. Edin. Dub.* Alum.

*Syn.* Alum (*F.*), Alaun (*G.*), Aluin (*Dutch*), Alun (*Dan. Swed.*), Halun (*Polish*), Kevaszu (*Russ.*), Allume (*I.*), Alumbre (*S.*), Pedrathune (*Portug.*), Sp'hatica (*San.*), P'hitcari (*H.*), Sheb (*Arab.*), Paddicarum (*Tam.*).

This salt is a ternary compound of alumina, potash, and sulphuric acid. It is found native in some places, either effloresced on bituminous schistus, as at Göttwig in Austria; or united with the soil in volcanic regions, as at the Solfatara near Naples, where the only processes requisite for its extraction are lixiviation and evaporation.<sup>2</sup> But the greater quantity of the alum of commerce is prepared by a peculiar management of schistose pyritic clays, usually denominated alum ores. At La Tolfa, near Civita Vecchia, where the best Roman alum is made, the ore is *alum stone* or sulphureted clay, found in large stratified masses among compact iron-shot argillaceous limestone<sup>3</sup>; but at other places, both on the continent and in Great Britain, it is manufactured from *pyritaceous clay*, which is in black, hard, brittle masses; *volcanic aluminous ores*, a white saline earth; *shale alum slate*, which occurs amorphous, or in concentric balls. At Hurlett, near Paisley, the largest alum

<sup>1</sup> Mucus differs from mucilage of gum arabic, in not being precipitated by silicated potash, nor affected by red or oxysulphate of iron. Bostock. *Nicholson's Journ.* xviii. 31.

<sup>2</sup> These processes are performed in pans sunk in the ground, the heat of which is sufficient to carry on the evaporation.

<sup>3</sup> An alum is found near Moscow which contains much sulphate of iron. *Mem. de la Soc. Imp. de Moscow*, t. i. p. 22.



mine in this country, the schistus lies ten inches thick above coal. But the alum is in every instance the result of the decomposition of what is termed its ore; and a subsequent synthetical union of its components.

To prepare the alum, the ore is properly exposed to the atmosphere, when the sulphur of the sulphuret of iron, present in it, is oxygenized by the air and converted into sulphuric acid, which, combining with the alumina, produces an aluminous efflorescence. In general, however, the ore is first calcined with a low heat, so as to destroy the bituminous matter of the slate, and partly convert the sulphur into sulphuric acid: the oxygenizement is then completed by exposing the roasted ore to air and moisture, by which means a supersulphate of alumina is formed, which is extracted by lixiviation. To the solution concentrated, until it acquire the spec. grav. 1.35, is added the impure subcarbonate of potass of commerce, or supersulphate of potass, or putrid urine, or muriate of potass, after which it is run into coolers to be crystallized. After standing four days the mother waters are drained off, and the crystals being washed, are redissolved in boiling water to saturation, and then *roched*, that is, run into casks. On taking asunder the casks at the end of sixteen days, the alum exteriorly is found in a solid cake, but interiorly crystallized in large octahedral pyramids, inserted into one another.

The ancients were probably unacquainted with alum, for the *στυπτηρία* of the Greeks, and the *alumen* of the Romans, were merely vitriolic earths: and the first regular alum-works appear to have been established by the Asiatics, in the middle ages, particularly at Roccha, in Syria, whence the name *Roch alum*; and from them Europe was supplied till the fifteenth century. After this period, works were begun at Tolfa and Volterra in Italy, at Oberkaufungen and several other places in Germany, and at Almacoran in Spain. In England, in the reign of Elizabeth, Sir Thomas Chaloner established the first alum work at Gisborough, in Yorkshire. The largest alum works at present in this country, are those on the estate of Lord Glasgow at Hurlett, and those of Lords Dundas and Mulgrave, at Whitby, in Yorkshire.

The best alum is the Roman, which is in irregular, octahedral, crystalline masses, powdery on the surface. The English is in large, irregular, semitransparent, colourless masses, having a glassy fracture, not efflorescent, and difficult to pulverize; and that from the Levant, or Roch alum, is in small morsels, about the size of an almond, rather friable, and of a pale rose colour. The form of the regular crystal of alum is an octahedron. Its specific gravity is 1.7109. According to Vauquelin, its constituents are, acid 30.52, alumina 10.50, potass 10.40, and water 48.58, in 100 parts: but by a more



recent analysis, Berzelius makes them sulphate of alumina 36.85, sulphate of potass 18.15, and water 45: it also generally contains ammonia; and none of the alum of commerce, except the Roman, is free from a minute portion of iron.

*Qualities.* — Alum is inodorous, and has a sweetish, acidulous astringent taste. Its specific gravity is about 1.71. It reddens litmus and tincture of cabbage<sup>1</sup>; is in a small degree efflorescent; and soluble in six parts of water at 60°, and in three-fourths of its weight of boiling water. When exposed to a gentle heat it undergoes the watery fusion, and in a stronger heat swells, loses 44 per cent. of its weight, which is water, and becomes an opaque, white, friable spongy mass. It is decomposed by the alkalies and alkaline earths, which attract the greater part of its acid, and precipitate the alumina united with a small portion of acid and potass. According to the analysis of Mr. R. Phillips, alum is composed of sulphuric acid 34.94 parts, alumina 11.18, potassa 10.33. and water 43.55 in 100 parts. According to Dr. Thomson, its atomic composition is 3 atoms of alumina, 1 atom of sulphate of potass, and 25 atoms of water: or in numbers 60.875 grains contain 21.75 of alumina, 11 sulphate of potass, and 28.125 of water.<sup>2</sup> Gallic acid also precipitates its earth: hence the alkalies and their carbonates, muriate of ammonia, magnesia, lime, carbonate of magnesia, chalk, tartrate of potass, and infusions of galls and of cinchona, are incompatible in prescriptions with solutions of alum; as are also superacetate of lead, and the salts of mercury.

*Medical properties and uses.* — Alum is a powerful astringent. It is used both as an internal and external remedy for restraining violent hæmorrhages; and also in cases of obstinate diarrhœa, diabetes, and fluor albus; but we agree with Dr. Cullen, that it is not to be depended upon in the two latter diseases. It has been given as an auxiliary to Cinchona in intermittents, and in confluent small-pox when the pustules are bloody; and Dr. Percival regarded it as a prophylactic in colica pictonum, and a cure for slighter cases.<sup>3</sup> It is used locally in gargles, in cases of cynanche, relaxation of the uvula, and aphthæ; and as the basis of injections, in cases of gleet and leucorrhœa, and of eye-waters in chronic ophthalmia.

The dose in hæmorrhages is from grs. v. to ℥j., repeated every hour or two till the bleeding abate: in other cases, smaller doses are more advisable, larger being apt to nauseate

---

<sup>1</sup> Much of the English alum which we have examined strikes a green with syrup of violets, and most other vegetable colours, with the exception of the two above mentioned.

<sup>2</sup> *First Principles of Chemistry*, vol. i. p. 310.

<sup>3</sup> *Observations on Lead*, &c.



the stomach, and occasion violent constipation. The addition, however, of an aromatic, prevents it, to a certain degree, from exciting nausea. It is sometimes administered dissolved in the serum of milk, in the form of whey, (*serum lactis aluminosum*), which is prepared by boiling ʒij of powdered alum in a pint of milk, and straining. A small piece of alum, briskly agitated with the white of an egg, forms a coagulum, which, applied between two pieces of gauze or thin rag, proves very serviceable in ecchymosis, and some species of ophthalmia. The dose of alum in substance, is from gr. v. to ʒj.; of the whey fʒij. or fʒiij.

Official preparations. *Alumen exsiccatum*, L. E. D. *Liquor Aluminis comp.* L. *Pulvis Aluminis comp.* E.

AMMONIÆ MURIAS, *Lond.* MURIAS AMMONIÆ, *Edin.* SAL AMMONIACUM, *Dub.* Muriate of Ammonia.

*Syn.* Sel Ammoniac (*F.*), Salmiak (*G.*), Salmiakken (*Dan. Swed.*), Sale Ammoniaco (*I.*), Sal Armoniaco (*S.*), Nosader (*H.*), Navacharum (*Tam.*), Urmiena (*Arab.*), Nowshader (*Pers.*), Vayvagarra Loonoo (*Cyng.*), Nuosadur (*Sans.*).

This salt, which is a compound of muriatic acid and ammonia, is found as a product of volcanoes;<sup>1</sup> but the greater part of that which is employed in medicine and the arts is artificially prepared.

Muriate of ammonia was originally manufactured in Egypt, by sublimation from the soot of fuel, formed of the dung of phytivorous animals kneaded with straw into clods, and dried in the sun. From this source all the European states were formerly supplied: but since the manufacture of it in Europe the importation of Egyptian sal ammoniac has been discontinued. The process differs in different places, and is generally kept secret; but the following is a sketch of the actual practice at a large establishment, which was carried on some years ago near London, and is probably the mode usually adopted in this country.<sup>2</sup>

Bones, chopped into small pieces, and boiled in order to extract the marrow and fat, were distilled from an iron cylindrical still into a leaden receiver, cooled by a refrigeratory, which was its cover, and contained about four inches in depth of water. Six parts of impure alkaline liquor and five of foetid oil were thus procured; the oil was skimmed off, and the alkali mixed with pulverized gypsum. By double decomposition sulphate of ammonia and carbonate of lime were thus formed; the liquor which contained the former was then mixed with common salt (*muriate of soda*); and thus, by a se-

<sup>1</sup> The eruption of Etna in 1811 afforded as much Sal ammoniac as supplied all the manufactories and apothecaries' shops in Sicily. *Annales de Mines*, tom. v. p. 135. It is exhaled also from the solfatara of Pozzuola; from one of the great apertures of which it has been extracted for several years.

<sup>2</sup> *Aikin's Dictionary of Chemistry*, art. Sal ammoniac.



cond decomposition, the sulphuric acid of the sulphate of ammonia uniting with the soda of the muriate of soda, and the muriatic acid with the ammonia; muriate of ammonia and sulphate of soda were formed in the liquor. This solution was clarified by subsidence and decantation; and by a skilfully managed evaporation in leaden boilers, the two salts were separated as they crystallized. The water of crystallization was then driven off from the muriate of ammonia, by exposing it to heat in a kind of oven; and the spongy, friable, ash-coloured mass, into which it changed, was put, while hot, into globular bottles, or glazed earthen jars furnished with a moveable perforated cover, in which the muriate was sublimed by exposing them to a heat of  $320^{\circ}$  in iron pots filled with sand. The cakes of salt produced, after being placed “for a day or two in a damp atmosphere,” to soften their surface, and facilitate “the removal of any superficial impurities,” were packed in casks for sale. As soot of coal affords, by maceration in water, a quantity of sulphate of ammonia, it is used in the Scotch manufactories instead of bones.

The cakes of muriate of ammonia are hemispherical, about an inch thick; elastic; and when broken are towards the convex surface white, striated, and opaque; but towards the concave have a more crystallized appearance, and are nearly semi-transparent. This salt is also, sometimes, crystallized in conical masses, that are deliquescent, owing to the presence of muriate of lime, which renders it unfit for medicinal purposes.

The greater part of the sal ammoniac in the London market is made in the north of England; but an inferior sort is imported in chests from the East Indies.

*Qualities.* — This salt is inodorous; has a salt, bitterish, acrid and cool taste; very slightly attracts moisture from the air; and has a specific gravity of 1.450. It is rather ductile, and therefore not very easily pulverized. It requires 3.25 times its weight of water at  $60^{\circ}$ , and its own weight at  $212^{\circ}$ , to dissolve it; and during its solution a great reduction of temperature takes place. It is also soluble in  $4\frac{1}{2}$  parts of alcohol. At a high temperature it sublimes without melting, and is unchanged. When dissolved in boiling water, it forms, as the solution cools, in tetrahedral, or in flaky plumose crystals. Its components are 31.5 of ammonia, and 68.5 of acid, independent of the water of crystallization. It combines with oxymuriate of mercury, and increases its solubility in water. The sulphuric and nitric acids unite with its alkali, and set free the muriatic acid. Potass and its carbonate, carbonate of soda, barytes, strontites, lime, chalk, magnesia, carbonate of magnesia, combine with its acid, and set free the ammonia, which is rendered sensible by its odour. Superacetate of lead, when added to a solution of it, throws



down a precipitate of muriate of lead; it is decomposed also by nitrate of silver, the base of which forms an insoluble compound with muriatic acid: hence these salts are incompatible in prescriptions with muriate of ammonia: but it may be combined in solution with the sulphate of copper, or of zinc.

*Medical properties and uses.* — This salt was formerly considered a powerful aperient and attenuant of viscid humors, acting as a diaphoretic, diuretic, purgative, and emetic, according to the mode of exhibition, or the extent of the dose; but it is now scarcely ever ordered as an internal medicine. Externally, it is advantageously employed, when mixed with its weight of nitre, and dissolved in eight parts of water, on account of the cold produced during its solution, to abate the pain and heat of inflammation and to allay violent headache. It is useful, also, in cases of mania, plethoric apoplexy, and injuries of the head, and to assist in the reduction of hernial tumors; but when it is employed for this purpose the solution should be made immediately before applying it to the affected part. It is also a useful application in dropsy of the thyroid gland.<sup>1</sup> Owing to its stimulant qualities, it forms an excellent discutient, when dissolved in the proportion of ℥j of the salt, in f℥ix of water, with f℥j of alcohol, in indolent tumors, gangrene, scabies, and chilblains; in which cases it is better not to be too recently dissolved: and as a gargle, it is occasionally useful in cynanche. A plaster formed with ʒss of the muriate, ℥j of soap, and ʒij of acetate of lead, is highly recommended by Dr. Paris<sup>2</sup> as a rubefacient in pulmonary affections. Its efficacy depends on the extrication of ammonia by the decomposition of the muriate, on which account it should be renewed every twenty-four hours.

*Officinal preparations.* *Ammonia subcarbonas*, L. E. D. *Liquor Ammonia*, L. *Aqua Ammonia*, E. D. *Hydrarg. præcip. alb.*, L., *Alcohol ammoniatum*, E. D. *Ferrum ammoniatum*, L. E. D. *Aqua Cupri ammoniati*, D. *Sulphuretum Ammonia*, D. *Murias Ammonia et Ferri*, D.

AMMONIACUM. Vide *Heracleum gummiferum*.

AMYGDALUS. *Species Plant. Willd.* ii. 982.

Cl. 12. Ord. 1. Icosandria Monogynia. Nat. ord. Pomaceæ, Linn. Rosaceæ, Juss.

G. 981. Cal. 5-cleft, inferior, Pet. 5. *Drupe* with a nut perforated. *Species* 2. A. communis.<sup>3</sup> The Common Almond-tree. *Med. Bot.* 2d edit. t. 183.

*Varieties.* β. *Amygdalus sativa*. Sweet Almond-tree.

———— γ. *Amygdalus amara*. Bitter Almond-tree.

<sup>1</sup> Burns on the Anat. of the Neck, p. 191.

<sup>2</sup> Pharmacologia.

<sup>3</sup> Πάδος, Theophrasti.



*Officinal.* AMYGDALÆ AMARÆ.—DULCES, *Lond.* AMYGDALUS COMMUNIS, NUCLEI, *Edin.* AMYGDALÆ DULCES, *Dub.* Bitter and Sweet Almonds.

*Syn.* Amandes douces et ameres (*F.*), Bittere und Süsse Mandeln (*G.*), Amandel (*Dutch*), Mandel (*Dan.* and *Swed.*) Migdalowe (*Polish*), Mandorli dolce ed amare (*I.*), Almendra (*S.*), Amendo (*Portug.*), Bādāmie Parsie (*Hind.* and *Pers.*), Lowz (*A.*), Parsie Vadomcotta (*Tam.*), Lonzan (*Mala.*).

The almond-tree is a native of Syria and Barbary; but it is now naturalized in the south of Europe, and even in England<sup>1</sup>; where, however, the fruit seldom ripens. The flowers display themselves in March and April before the leaves are expanded. It rises to the height of twenty feet, and divides into many spreading branches, which are covered with a dark grey bark. The leaves stand upon short foot-stalks, are about three inches long, and three-fourths of an inch broad, elliptical, pointed at both ends, minutely serrated, with the lower serratures glandular, and of a bright green colour. The flowers, which are supported on very short peduncles, are of a pale rose or blush colour, varying to white: the calyx is tubular, with the lip divided into five blunt segments; the petals are five, oval, and convex; the filaments about thirty, inserted into the calyx, tapering, spreading, of unequal lengths, and furnished with orange-coloured, simple anthers: the germen is downy, with a simple style, supporting a round stigma. The fruit is of the peach kind, but flatter, with a tough coriaceous covering instead of the rich pulp of the peach, which opens spontaneously at the longitudinal furrow when ripe. The kernel or almond, which is inclosed in a tender, thick, brittle, spongy shell, is oblong, flattish, rounded at one end, and pointed at the other, and composed of two white cotyledons, enveloped with a thin, pale brown, veined, bitter skin, covered with an acrid meal.

These two varieties of the *Amygdalus communis* are not distinguished from each other, but by the taste of the kernel of their fruit. The *Valentia* almond is a sweet, large, flat almond, pointed at one extremity, and compressed in the middle as if with the thumb. The *Italian* are not so sweet, smaller, and less depressed in the middle. The *Jordan* almonds, which come from Malaga, and are the best sweet almonds brought to England, are said to be the produce of a distinct species of *Amygdalus*. They are longer, flatter, less pointed at one end, and less round at the other, and have a paler cuticle than those we have described.

Sweet almonds are imported in mats, casks, and cases: the bitter, which come chiefly from Mogadore, arrive in boxes.

---

<sup>1</sup> It was cultivated in England by Lobel before 1570.



When the almond is not well preserved, it is preyed on by an insect that eats out the internal part; or, if this does not happen, the oil it contains is apt to become rancid.

*Qualities.* — The cuticle of both kinds of almonds has an unpleasant bitterish austere taste; but it is easily detached by putting the almonds into boiling water. When thus decorticated, they are said to be blanched.

The blanched *sweet-almond* is inodorous; has a sweet, pleasant, bland taste, and consists chiefly of fifty-four parts of fixed oil, three of mucus, six of saccharine matter, and twenty-four of albumen. When eaten as food, it is not very digestible, probably owing to the albumen; and requires to be well masticated. The *bitter-almond* is also inodorous when entire, but when triturated with water it has the odour of the peach blossom; and the taste is the pleasant bitter of the peach kernel. It contains less fixed oil and more albumen than the sweet almond; and a portion of Prussic, or Hydrocyanic acid, upon which its narcotic power is supposed to depend. This variety of the almond is said to operate as a poison on dogs and some other animals, but not generally on the human species. The distilled water, however, of the bitter almond exerts an action not less deleterious than that of laurel water on the human frame. It produces vertigo, head-ache, or rather a sense of weight at the summit of the head, tinnitus aurium, dizziness of sight, and vomiting, when taken to the extent of thirty drops only; and a drachm of it has killed a stout dog.<sup>1</sup> When a large dose is taken, a paralytic state of the extremities supervenes, the pupil remains unalterably dilated, and the excitability of every organ of sense is diminished; indeed, death almost instantly follows. In order to counteract its poisonous effects, when that can be done, we must have recourse to diffusibles, as brandy and ammonia; or three or four spoonsful of oil of turpentine may be given at intervals of half an hour. The essential oil of bitter almonds which contains the Prussic acid, is prepared from the cake remaining after the expression of the fixed oil, by submitting it to distillation with water. One hundred weight of the cake generally yields from two ounces to two and a half ounces of the essential oil. From the experiments of Mr. Hennel it appears, that the Hydrocyanic acid may be separated from this oil by digesting it with red oxide of mercury, which is converted into a cyanuret. The fixed oil, which both varieties of

---

<sup>1</sup> Much information on this subject may be obtained from the works of Fodere, Langrish, Heberdeen, Watson, and a recent Treatise on Hydrocyanic Acid, by Dr. Granville. See also the *Appendix* to this work.



the almond yield by expression in large quantity, is insipid and inodorous when heat has not been employed.

*Medical properties and uses.* — Sweet almonds are used more as food than as medicine; but they afford little nourishment. Heartburn is said to be relieved by eating six or eight of them decorticated. When triturated with water, milky mixtures or emulsions are formed, which shall be afterwards noticed; and they are also used in pharmacy for assisting, by trituration, the combination of some substances, such as camphor and the resins, with water. Bitter almonds are scarcely ever used medicinally, although Bergius<sup>1</sup> mentions a case of intermittent having been cured by them, when the Peruvian bark had failed; and reflecting on the effects which have been found to result from the use of the Hydrocyanic acid, I have employed the bitter almond in pulmonary and dyspeptic affections, hooping cough, and asthmatic complaints, with the best success. As a local application, I have found the emulsion extremely beneficial when used as a lotion in acnè rosacea, and in impetigo. Owing to a peculiar idiosyncrasy of some habits, the smallest quantity of the bitter almond taken into the stomach produces urticaria, and other unpleasant effects.

Official preparations. *Oleum Amygdalæ*, L. E. D. *Emulsio Amygdalæ comp.* E. D. *Emulsio Acaciæ arabicæ*, E. D. *Emulsio Camphoræ*, E. *Confectio Amygdalarum*, L.

AMY'LUM. Vide *Triticum hybernum*.

AMYRIS. *Spec. Plant. Willd.* ii. 333.

*Cl.* 8. *Ord.* 1. Octandria Monogynia. *Nat. ord.* Terebintaceæ, *Juss.* G. 755. *Cal.* four-toothed. *Pet.* four oblong. *Stig.* four-cornered. *Berry* drupaceous.

*Species* 2. *Amyris elemifera*. Elemi tree.

— 6. *Amyris gileadensis*. Balsam of Gilead tree. *Med. Bot.* 2d edit. 603. t. 214. *Bruce's Abyssinia*, vol. v. p. 16. t. 2, 3.

1. AMYRIS ELEMIFERA.

*Official.* ELEMI, *Lond.* ELEMI; RESINA, *Dub.* Elemi.

*Syn.* Eleme (*F. G. I.*), Goma de Limon (*S.*).

The Elemi tree, of the botanical characters of which we know very little, is a native of Carolina and the Brazils. It does not, according to Catesby, rise to a great height; and the trunk is small, and covered with a grey bark. The leaves are opposite, on footstalks; ternate, and sometimes pinnate; with stiff, pointed leaflets, of a bright green colour, shining, and downy underneath. The flowers are in terminal corymbs; small, white, with the petals inflex at the tips. The fruit is the size and figure of an olive.

The resin is obtained by making incisions in the bark in dry weather, and is left to dry in the sun as it exudes. It

<sup>1</sup> *Mat. Med.* art. *Amygdalus*.



used to be brought from Turkey in long roundish cakes, wrapped in flag-leaves; but the elemi now brought, comes in mats and chests, each containing from four to six pounds' weight.

*Qualities*. — True elemi has a fragrant aromatic odour, not unlike that of fennel seeds, but stronger. The taste is very slightly bitter, and warm. The cakes are of a pale yellow colour, semitransparent, brittle on the outside, soft and tenacious within, and very fusible. Spec. grav. 1.0182. When distilled with water, it affords  $\frac{1}{10}$ th of a thin pale-coloured essential oil, on which its fragrance and softness depend; and the residue is a brittle, inodorous resin. Alcohol dissolves the greater part of elemi; but a white, flaky, inodorous matter remains, which is almost entirely soluble in water; hence we may consider the constituents of elemi to be gum, and an intimate combination of resin and essential oil. But no true elemi is now to be found in the shops.

*Medical properties and uses*. — This resin is stimulant, but is very rarely used as an internal remedy, being chiefly employed for forming the mild digestive ointment which bears its name.

Officinal preparation. *Unguentum Elemi compositum*, L. D.

## 2. AMYRIS GILEADENSIS.<sup>1</sup>

*Officinal*. —, RESINI LIQUIDA, *Edin.* Balsam of Gilead.

*Syn.* Balsamier de la Mecque (*F.*), Opobalsamo (*I.*), Balsamo (*S.*), Akooy-eelarsemoonroome (*Arab.*), Roghen Bulsan (*Pers.*).

This species of amyris is a native of Abyssinia, growing, according to Bruce, among the myrrh-trees behind Azab, all along the coast to the Straits of Babelmandel.<sup>2</sup> It appears, however, to have been transplanted into Judea 1730 years before Christ: and as it was from Gilead in Judea that the merchants brought its resinous product, in early times, to Egypt, it thence derived its appellation *Balasan*, or *Balsam of Gilead*.

This tree rises above fourteen feet in height; has a flat top and stunted aspect, with many spreading crooked branches going off nearly at right angles; the wood is light, open, and covered with a smooth bluish white bark. The leaves are thinly scattered, small, composed of one or two pair of opposite leaflets, with an odd one; these are obovate, entire, veined, and of a bright green colour. The flowers are white, appearing upon the young shoots, three upon one stalk; but two generally drop, and one only produces fruit. The calyx

<sup>1</sup> Βαλσάμου δένδρον, Theophrasti et Dioscoridis.

<sup>2</sup> Bruce's *Abyss.* App. p. 16. The whole of Mr. Bruce's account of this tree is highly interesting, and we regard his authority undoubted.



is permanent, divided into four expanded sharp teeth: the petals are four, oblong, concave, spreading; the filaments eight, erect, supporting oblong anthers: the germen is superior, ovate, with a thick style, the length of the filaments crowned with a quadrangular stigma. The fruit opens with four valves, and contains a smooth nut.

The ancients held the balsam obtained from this tree in great esteem, but it does not appear that they were well acquainted with the tree itself. To obtain the balsam, the bark is “cut by an axe, when the juice is in its strongest circulation, in July, August, and the beginning of September. It is then received into a small earthen bottle, and every day’s produce gathered and poured into a larger, which is kept closely corked.” The first that flows, called opobalsamum<sup>1</sup>, “is of a light yellow colour, apparently turbid.” It afterwards becomes clear, fixed, and heavier; and the colour, by degrees, deepens to a golden yellow. The opobalsamum of the ancients was the green liquor found in the kernel of the fruit: the carpobalsamum, the next in esteem, was made by the expression of the ripe fruit; and xylobalsamum, or the worst kind, by the expression or decoction of the small twigs. The real balsam rarely finds its way into this country, dried Canada balsam being generally substituted for it; but it wants the peculiar odour of the true balsam.

*Qualities.*—The odour is at first strongly pungent; but the pungency is lost by exposure to the air, and by age; and the balsam gradually acquires the consistence of turpentine. The colour is yellowish exteriorly, and paler in the inside; the taste is acrid, rough, and pungent. When pure, it dissolves easily in water.

*Medical properties and uses.*—This balsam was esteemed in the earlier ages as a medicine possessed of almost universal virtues; and at the present day the Arabs use it “in all complaints of the stomach and bowels,” reckoning it a powerful antiseptic, and preventive of the plague. Its chief use, however, is as a cosmetic by the Turkish ladies. It is never brought genuine to this country, and we know not why the Edinburgh College retain it in the list of *Materia Medica*.

ANCHUSA.<sup>2</sup> *Spec. Plant. Willd.* i. 756.

*Cl.* 5. *Ord.* 1. Pentandria Monogynia. *Nat. ord.* Asperifoliæ, *Linn.* Boragineæ, *Juss.*

*G.* 277. *Corolla* funnel-shaped: the throat closed with arches. *Seed* engraved at the base.

<sup>1</sup> Willdenow has a distinct species under the name *Amyris opobalsamum*; but, in a note, says—“sunt forte non distinctæ species, sed varietates ab ætate vel solo ortæ.” *Spec. Plant.* vol. ii. p. 334.

<sup>2</sup> Ἀγχονσα ab ἀγχω strangulo, suffoco; the ancients believing that this species of plants choked and destroyed serpents. Vide *Bod. in Theophrast.* p. 835.



*Species* 7. *Anchusa tinctoria*. Dyer's Alkanet. Lobel. *Icon*. 578.  
*Officinal*. —, RADIX, *Edin*. ANCHUSA; RADIX, *Dub*. Alkanet root.

*Syn*. Orcanette (*F.*), Alcanna (*I.*)

This species of anchusa is perennial, a native of the south of Europe, where it is cultivated in great abundance, particularly near Montpellier. It is found in our gardens as an ornamental plant; but its roots do not acquire in Britain the beautiful colour for which the foreign are prized. It flowers from June until October. The root is long, round, fibrous, white within, and covered with a purplish red bark. The stem rises eighteen inches in height; is round, rough, hairy, and branched; with long, alternate, sessile, lanceolate, obtuse, hairy leaves. The flowers are of a reddish purple colour, and terminate the branches in close clusters. The calyx is persistent, divided into five oblong erect segments; the corolla funnel-shaped, consisting of a cylindrical tube the length of the calyx, and a five-tooth expansion, closed by five scaly leaflets. The filaments are shorter than the corolla, bearing simple anthers; the germens four, with filiform styles the length of the filaments, each crowned with an obtuse notched stigma.

Alkanet root is brought to this country chiefly from France. It is in twisted pieces, with a withered, dusky red bark, easily separated. The smaller roots are the best, as they have proportionally more bark than the larger.

*Qualities*. — It has a very faint odour, and a bitterish astringent taste when fresh; but the dried root is inodorous and insipid. It imparts a fine, deep-red colour to alcohol, ethers, oils, fats, and wax; but to water, even when hot, it yields only a brown colour. Sulphate of iron strikes a black with the watery infusion; and sulphate of zinc throws down a copious dark-coloured precipitate.

*Medical properties and uses*. — Alkanet root was formerly prescribed as an astringent in several diseases; but it is properly rejected from modern practice, and is used as a colouring matter only for oils, ointments, and plasters.

ANETHUM.<sup>1</sup> *Spec. Plant. Willd.* i. 1469.

*Cl.* 5. *Ord.* 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

*G.* 560. *Fruit* nearly ovate, compressed, striated. *Pet.* involuted entire.

*Species* 1. *A. graveolens*. Common Dill. *Med. Bot.* 2d ed. 125. t. 48.

*Species* 3. *A. fœniculum*. Sweet Fennel. *Med. Bot.* 2d ed. 127. t. 49.

*Smith Flor. Brit.* 329.

#### 1. ANETHUM GRAVEOLENS.

*Officinal*. ANETHI SEMINA. *Lond. Edin.* Dill seed.

*Syn*. Fenouil puant, ou persil odorant (*F.*), Dill (*G.*), Dille (*Belg.*), Dil (*Dan.*), Dill (*Swed.*), Kopr (*Polish*), Endro (*Portug.*), Eneldo de olor pesado

<sup>1</sup> *Ανηθον*, Dioscoridis.



(*S.*), Appio palustre, sellano, Sellero (*I.*), Sadācoopei (*Tam.*), Sowa, (*Hind.*), Moongsi (*Javanese*), Buzralshibbet (*Arab.*).

This plant is an annual, a native of Spain and Portugal, growing generally in corn-fields, and flowering in June and July. It is cultivated in this country.<sup>1</sup> The root is fusiform and long, striking deep into the ground; and sending up several erect, striated, jointed stems, about two feet in height, and branched. The leaves are glaucous and odorous, upon sheathing foot-stalks; doubly pinnated, with the pinnæ linear and pointed. The flowers are in large, flat, terminal umbels, without either universal or partial involucre; the corolla consists of five ovate, obtuse, concave, yellow petals, with the apex inflected: the filaments yellow, and longer than the corolla: with an inferior germen, covered by the nectary, and supporting two short styles, terminated by obtuse stigmas.

The seeds of dill, which are the parts of the plant medicinally used, are scarcely the length of a caraway seed, but broader and flatter. They are oval, concave on one side, convex and striated on the other; of a brown colour, and surrounded with a dull pale-yellow or straw-coloured membranous expansion. When good they are heavy, of a bright colour, and an aromatic odour.

*Qualities.*—The dried seeds have an aromatic, sweetish odour, not very agreeable, nor yet unpleasant; the taste is moderately warm and pungent. These qualities depend on an essential oil, which is extracted by distillation with water, (according to Mr. Brande, in the proportion of 2 lb. to 1 cwt.<sup>2</sup>) and imparted to alcohol by digestion. The bruised seeds yield their flavour to boiling water by infusion.

*Medical properties and uses.*—Dill seeds are carminative and stomachic. They are scarcely ever employed except in hic-cough and the flatulent colic of infants. The dose of the powdered seed is from grs. xv. to ʒj.

Officinal preparations. *Aqua Anethi*, L.

## 2. ANETHUM FENICULUM.<sup>3</sup>

*Officinal.* FENICULI SEMINA, *Lond. Edin.* FENICULUM DULCE; SEMINA, *Dub.* The seed and root of Sweet Fennel.

*Syn.* Fenouil ou Anis douce (*F.*), Fenchelsamen (*G.*), Venkel (*Dutch*), Fennikel (*Dan.*), Fänkol (*Swed.*), Kopo Wlosky (*Pol.*), Eneldo hinojo (*S.*), Finocchio (*I.*), Hinojo (*S.*), Funcho (*Port.*), Razecanuj (*Arab.*), Badeeyan (*Pers.*), Perumsiragam (*Tam.*), Mayuri (*Hind.*), Adas (*Javanese*).

Fennel is a biennial plant, originally found in the south of Europe only, but now growing abundantly on our chalky soils and cliffs, and flowering in July and August. The root is fusiform, elevating a stem about four feet in height, erect, branching, leafy, striated, and smooth. The leaves are alternate, tripinnate, composed of long, smooth, depending, linear leaf-

<sup>1</sup> It was first cultivated by Gerarde in 1597.

<sup>2</sup> *Manual of Pharmacy.*

<sup>3</sup> *Μάραθρον*, Hippocratis et Dioscoridis.



lets, of a very deep green colour. The flowers are in large, terminal, many-rayed, flat umbels: the petals five, ovate, emarginate, with their points turned inward; and of a yellow colour: the filaments shorter than the petals, spreading, also yellow, and bearing double anthers. The germen is similar to that of *dill*: the seed ovate, very little compressed, of a brownish olive colour when ripe, three-ribbed, and encircled with a membranous margin.

There are three varieties of fennel; the root of the first of which, the *common fennel*, and the seed of the second, the *sweet fennel*, are officinal. The roots found in the shops are the produce of our own country, and are taken up in the spring; but the seeds are generally imported from Italy.

*Qualities*.—The roots are covered with a brown bark, are woody and white within, have scarcely any odour, and only a slightly sweetish taste, with very little aromatic warmth; but the *seeds* have a fragrant odour; and a sweet, warm, aromatic taste. These qualities depend on an essential oil, which is dissipated by decoction in water, and separated by distillation: they are completely imparted to alcohol, but only imperfectly to boiling water, by infusion. The seeds contain also a fixed inodorous, insipid oil.

*Medical properties and uses*.—Fennel was formerly esteemed as a remedy; and supposed to be resolvent, diuretic, carminative, and stomachic; but even as a carminative it is not superior to anise-seed and caraway; and it is therefore now seldom employed. The dose of the bruised seed may be from ℥j to ʒj.

Officinal preparations. *Aqua Fœniculi*, L. D. *Oleum Seminum Fœniculi dulcis*, D.

ANGELICA. *Spec. Plant. Willd.* i. 1428.

Cl. 5. Ord. 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

G. 343. *Fruit* roundish, angular, solid, with reflected styles. *Corollas* equal. *Petals* bent inward.

*Species* I. *Angelica archangelica*, Garden Angelica. *Med. Bot.* 2d ed. 86. t. 35. *Smith Flor. Brit.* 1. 311.

*Officinal*. ———, RADIX, *Edin.* The root of Angelica.

*Syn.* Angelique (*F.*), Angelikawurzel (*G.*), Tamme Engelwortel (*Belg.*), Angelik (*Dan.*), Angelikerot (*Swed.*), Dziegiel ogrodni (*Polish*), Djajilnik (*Russian*), *Angelica domestica* (*I.*), Anjelica (*S.*).

This species of angelica is a native of the more northern parts of Europe; but although it has been found growing wild in England, as at Broadmoor near Birmingham, and some other parts, yet it is uncertain whether it be indigenous. It is however abundantly cultivated for medicinal and other purposes; flowering in June and August.<sup>1</sup> The root is biennial, thick, fleshy, and resinous; the stem erect, hollow, round,

<sup>1</sup> It was first described by Joann. Jacob. de Manlius, a writer of the 15th century; and was cultivated in England before 1568.



smooth, furrowed, of a purplish hue, rising upwards of five feet in height, and sending off many branches, which terminate in globular many-rayed umbels, composed of dense, hemispherical umbellets. The leaves are numerous, petiolated, large, pinnated; with the leaflets ovate, pointed, cleft, and acutely serrated, smooth, somewhat decurrent, and the terminal ones three-lobed: the petioles membranous at the base, nerved, greatly dilated, and bellying. The involucre is deciduous; the involucre short, consisting of five linear lanceolate leaves. The calyx is five-cleft, minute: the corolla small, of a greenish-white colour; petals five, with the points turned inward: the stamens spreading, longer than the petals; and the germen inferior, supporting two reflected styles with obtuse stigmas. The seeds are large, elliptical, flat on one side, convex on the other, emarginate at both ends, and acutely three ribbed.

The roots of angelica, when wounded in the spring, yield an odorous yellow juice, which being slowly desiccated, proves an elegant gum-resin, very rich in the qualities of the plant. For medicinal purposes, the roots should be dug up in the autumn of the first year; in which case they are more easily preserved; but when gathered in the spring, they become mouldy, and are preyed on by insects. They should be thoroughly dried, and kept in a well-aired, dry place: and in order to secure their preservation, Lewis suggests “the dipping them in boiling spirit, or exposing them to steam, after they are dried.” The leaves and seeds do not retain their virtues when kept. The stems are cut, when tender, in May, and made into an agreeable sweatmeat by the confectioners.<sup>1</sup>

*Qualities*.—The odour of every part of the recent plant is fragrant and aromatic; the taste sweetish at first, then aromatic, warm, and slightly bitter. The dried root is corrugated, and of a greyish-brown colour externally; breaks short with a starchy fracture, and presents a firm interior, whitish, with many resinous brown and yellow points. It has the same odour and taste as the recent plant; and yields these qualities to alcohol, and in some degree to boiling water.

*Medical properties and uses*.—The leaves and seeds when recent, and the root both in the fresh and dried state, are tonic and carminative; but although the most elegant aromatic of northern growth, yet they are scarcely ever prescribed in modern practice. The dose in substance is from ʒss to ʒj, which may be given three or four times a-day.

ANISI SEMINA. Vide *Pimpinella Anisum*.

ANTHEMIS. *Spec. Plant. Willd.* iii. 2174.

---

<sup>1</sup> The Icelanders eat the stems and roots of angelica raw with butter. Vide *Sir Geo. Mackenzie's Travels in Iceland*, 4to. p. 255.



Cl. 19. Ord. 2. Syngenesia Superflua. Nat. ord. Compositæ Discoideæ, Linn. Corymbiferæ, Juss.

G. 1517. Receptacle chaffy. Seed down none, or a membranaceous margin. Calyx hemispherical, nearly equal. Florets of the ray more than five.

\* With a colourless or white ray.

Species 15. *A. nobilis*. Common Chamomile. Med. Bot. 2d ed. 47.

t. 19. Smith Flor. Brit. 904.

— 25. *A. Pyrethrum*. Pellitory of Spain. Med. Bot. 2d ed. 50. t. 20.

#### 1. ANTHEMIS NOBILIS.<sup>1</sup>

Officinal. ANTHEMIDIS FLORES, Lond. Edin. CHAMÆMELUM; FLORES, Dub. Chamomile Flowers.

Syn. Camomille Romaine (*F.*), Roemische hamiller (*G.*), Romische Kamille (*Dutch*), Romerske Kamulblomster (*Dan.*), Romerske Kamillblommer (*Swed.*), Camomilla Romana (*I.*), Manzanilla de Botera (*S.*), Marcella Romana (*Portug.*), Chāmaindoopoo (*Tam.*), Baboeneh (*Pers.*), Ehdaklmerza (*Arab.*).

This species of anthemis is an indigenous perennial plant, growing in dry pastures, and flowering in August and September. The greater part of the chamomile, however, which is medicinally used, is cultivated by the growers of physical plants.<sup>2</sup> The roots are woody, fibrous, and spreading: the stems trailing, about a span in length, foliaceous and downy: the leaves verticillately bipinnate, the pinnæ distant, and the leaflets small, threadlike, sharp, generally cleft into three segments; odorous, and of a pale green colour. The flowers are on solitary, terminal, unifloral, naked, striated, hairy peduncles. The calyx is common to all the florets, hairy, with broad membranaceous edges: the disc is yellow and convex; the florets of the radius white, spreading, long, and somewhat elliptical, three-toothed, and turned down; and the seed obscurely crowned.

Both the single, and the double-flowered varieties are cultivated; but as the sensible qualities of the flower reside chiefly in the disc florets, the single kind is preferred: and, as these qualities are also stronger before the tubular florets are blown, the flowers are then picked, and carefully dried for use. Those which are large and whitish are to be preferred.

*Qualities*.—The whole of the plant is gratefully odorous. The smell of the flowers is strong and fragrant; their taste bitter and aromatic, with a slight degree of warmth; and both the odour and the taste are extracted by water and alcohol. By distillation with water, they yield a small quantity of brownish yellow essential oil,<sup>3</sup> on which the odour and the antispasmodic powers of the plant seem to depend. Hot water takes up nearly one-fourth of the weight of the dry flowers, and when

<sup>1</sup> *Ανθεμης*, Dioscoridis. *Ανθεμην*, Theophrastis.

<sup>2</sup> Much of what is brought to the London market is grown about Mitcham in Surry. The soil best adapted for it is a dry sandy loam. A wet summer weakens the flavour of the flowers. — *Stevenson's Survey of Surry*, 379.

<sup>3</sup> The quantity obtained is about 3jss. from 1 cwt. of the flowers. *Brande's Manual*.



the infusion is evaporated, a bitter extractive matter and a small portion of resin remain. The active principles, therefore, of chamomile flowers appear to be bitter extractive, resin, and essential oil.

*Medical properties and uses.* — Chamomile flowers are tonic, carminative, and slightly anodyne: yet when a strong infusion of them is taken in a tepid state, it proves powerfully emetic. When given in substance, united with opium and astringents, if the bowels be easily affected, they have been successfully used for the cure of intermittents: and the infusion, in combination with ginger, or other aromatics, and the alkalies, is an excellent stomachic in dyspepsia, chlorosis, gout, flatulent colic and chronic debility of the intestinal canal.<sup>1</sup> The tepid strong infusion is a ready emetic, and is often employed to promote the operation of other emetics. By coction in water the essential oil is dissipated; chamomile flowers, therefore, ought never to be ordered in decoctions. Externally they are used as fomentations in colic, intestinal inflammation, and to phagedenic ulcers: and their infusion is also found to be an useful addition to emollient anodyne glysters in flatulent colic, and in irritations of the rectum producing tenesmus. The dose of the powdered flowers is from ʒss to ʒj, twice or thrice a day.

*Offi. preparations.* *Decoctum Anthemidis nobilis*, E. D. *Infusum Anthemidis*, L. *Extractum Anthemidis*, L. E. *Oleum Anthemidis*, L.

## 2. ANTHEMIS PYRETHRUM.<sup>2</sup>

*Officinal.* PYRETHRI RADIX, *Lond. Edin.* PYRETHRUM; RADIX *Dub.* Pellitory root.

*Syn.* Pyrèthre (*F.*), Bertram Wurtzel; Zahn wurtzel (*G. Dutch, Dan., Swed.*), Zebne Ziele (*Pol.*), Piretro (*I.*), Anthemis pelitri (*S.*), Akākrācārum (*Tam.*), Akurkurha (*A.*), Akkaraputta (*Cyng.*).

This is a perennial plant, a native of the Levant, Barbary, and the south of Europe. It is sometimes cultivated in Britain<sup>3</sup>; flowering from June to July. The root is long, tapering, about the thickness of a finger, with a brownish cuticle, sending off several lateral fibres; and throwing up many trailing stems, more commonly simple and unifloral than branching. The leaves are doubly pinnate, with narrow linear segments of a pale green colour. The flowers are large, with the florets of the radius white on the upper, and purple on the under side, and those of the disc yellow. In form they resemble the flowers of *Anthemis nobilis*.

Pellitory root is brought into this country from the Levant, and the coast of Barbary, packed in bales. It is frequently mixed with other roots, from which, however, it is easily distinguished.

<sup>1</sup> Selle's celebrated Pulvis Ecephracticus consisted of equal parts of chamomile flowers, rhubarb, carbonate of potass, magnesia, sulphur, and oleo-saccharum of fennel.—*Selli lib. de Cur. Stom. Morb.* vol. i. p. 131. (quoted in *Med. Rep.* vol. i.)

<sup>2</sup> Πυρεθρον, Dioscoridis.

<sup>3</sup> It was cultivated in England by Lobel in 1570.



*Qualities.*—The dried root, as we receive it, is inodorous. When chewed it appears at first insipid, but after a few seconds, excites a glowing heat, and a pricking or thrilling sensation on the tongue and lips, which remains for ten or twelve minutes. The pieces break with a short resinous fracture; the transverse section presenting a thick brown bark studded with black shining points, and a pale-yellow, radiated inside. The pungency appears to depend on a fixed oil, which is deposited in vesicles in the bark. M. Gautier describes it as solid, having a reddish colour and strong odour.<sup>1</sup> It is completely extracted by alcohol and sulphuric ether.

*Medical properties and uses.*—Pellitory root possesses powerful stimulant properties. Its chief use is as a sialagogue, to stimulate the excretories of the salivary glands, and excite an increased flow of saliva; by which inflammations and congestions of the neighbouring parts are relieved. Hence it has been found useful when chewed in some kinds of head-ache, apoplexy, chronic ophthalmia, rheumatic affections of the face, and tooth-ache; and by its direct stimulus in paralysis of the tongue and muscles of the throat. A gargle made by boiling ℥ss of the bruised root in Oj. of water until the fluid is reduced one-half, and to the strained liquor adding f℥ii. of Liq. Ammoniae has been long employed in St. Bartholomew's Hospital, in cases of relaxation of the uvula and soft palate.

ANTIMO'NIUM. Στιμμι. *Stibium*. Antimony.

*Syn.* Antimoine (*F.*), Spiessglanzmetall (*G.*), Antimonie (*I.*), Antimónio (*S.*).

The term antimony was formerly given to an ore, in which antimony was combined with sulphur; but it is now solely appropriated to express the pure metal. It is found in various parts of the world in different states of combination.

*A. Metallic.* (at Stalberg, Sweden, and Allemont in France.)

g. i. combined with silver and iron. Sp. 1. *Native antimony*.

ii. ——— with sulphur. - - - - - 1. *Grey sulphuret of antimony*.

Var. *a.* compact.

*b.* foliated.

*c.* striated.

*d.* plumose.

2. *Nickeliferous sulphuret*.

*B. Oxidized* - - - - - { 1. *White antimony*.

iii. combined with oxide of iron. - { 2. *Antimonial ochre*.

{ 3. *Red antimony*.

The grey sulphuret is the ore in which it is most abundantly procured, and from which the pure metal is generally obtained. Pure antimony is of a white colour, with a bluish shade, brilliant, and very slowly tarnished in air of a low

<sup>1</sup> *Ann. de Chim. et Phys.* viii. p. 101.



temperature. The texture is foliated; moderately hard; brittle and pulverulent. It is fusible at  $809^{\circ}$ : in a higher temperature it volatilizes in close vessels; but, if exposed to the air, is very rapidly oxidized. It decomposes water when ignited, and is oxidized by, and combines with, the sulphuric, nitric, and muriatic acids; but the other acids unite with its oxides only. It readily combines with sulphur and phosphorus. Its specific gravity, according to Brisson, is 6.702. The pure metal when rubbed between the fingers communicates to them a peculiar taste and smell, but it exerts no action on the body, nor is it used for officinal purposes.

*Officinal.* ANTIMONII SULPHURETUM, *Lond.* SULPHURETUM ANTIMONII, *Edin. Dub.* Sulphuret of Antimony.

*Syn.* L'Antimoine sulfure (*F.*), Spiessglanz (*G.*), Spiesglas (*Belg.*), Spidseglas (*Dan.*), Rå Spetsg'ans (*Swed.*), Szpiglas (*Polish.*), Unjienakulloo (*Tam.*), Sulfuro d'Antimonio (*I.*), Antimonio (*Portug.*), Kohul (*Arab.*), Surmeh (*H. Pers. Duk.*), Sauvira (*San.*).

Sulphuret of antimony is commonly sold in loaves, under the name of *crude antimony*; and is the *grey ore*, separated from the stony matter and other gross impurities with which it is naturally combined. It is the *striated variety*, the most common of all the antimonial ores, found both in masses and crystallized in Hungary, Saxony, France, Tuscany, Spain, Cornwall in England, and Dumfriesshire in Scotland; generally "in micaceous schistus and clay porphyry, mixed with pyrites and oxides of iron." In its natural state, its colour is light lead grey; its internal lustre splendid; its fracture radiated, affording splintery fragments. It is soft; not very brittle; but easily frangible.

Sulphuret of antimony is fitted for the market by the following process. The ore is separated from the greater part of the stony gangue by hand, and then placed in the bed of a reverberatory furnace, covered with charcoal powder. As it is brought to a low red heat, the sulphuret of antimony is fused, while the earthy parts float on the surface, and are taken off with a rake or ladle: and the fluid portion, when cast into the form of loaves or large cakes, is fit for sale, and forms the *crude antimony* of commerce.<sup>1</sup> Sometimes the ore is broken into small pieces, then washed and put into a pot perforated with holes, which is let into the mouth of another pot, so that the fluid sulphuret flows into the undermost, while the infusible matter remains in the uppermost. These loaves are dark-grey externally, but internally they have a striated structure, and considerable brilliancy. Their goodness depends on their compactness and weight, the largeness and distinctness of the striæ, and the volatility of

<sup>1</sup> *Journal des Mines, Aikin's Dictionary of Chemistry.*



the sulphuret. When carelessly prepared, they contain lead, sometimes arsenic, and occasionally manganese and iron. When they contain much lead, the structure is more foliated, and less distinctly striated, while, at the same time, the volatility is so much diminished that a portion, which is the lead, remains fixed; arsenic is discovered by the garlic odour emitted when the sulphuret is thrown on live coals; manganese and iron, by their not being volatilized when it is exposed to a red heat, and iron in particular by the brown colour produced by deflagrating them with Nitre. The specific gravity of the sulphuret is about 4·368; and its constituents are antimony 73·8, sulphur 26·2, in 100 parts; or, according to the atomic doctrine, 1 atom of antimony = 45, and 1 atom of sulphur = 16, the weight of an atom of the sulphuret being = 61.<sup>1</sup> The greater part of the sulphuret used in this country is imported from Germany and Holland. It should never be purchased in the form of powder.

*Qualities*.—Sulphuret of antimony is inodorous, insipid, of a leaden grey or steel colour, staining the fingers; has a rough spicular fracture, and is insoluble in water and alcohol. Its brilliancy is dulled by long exposure to the air; in a red heat it melts, and is partly dissipated along with its sulphur in the form of a white smoke; and what remains in the crucible is a grey, ash-coloured oxide. It is slightly acted upon by the vegetable acids<sup>2</sup>; and decomposes the sulphuric and nitric acids when assisted with heat; the metallic part of the sulphuret is oxydized, and sulphureous acid and nitrous gases are disengaged, the muriatic, even in the cold, decomposes it, and sulphuretted hydrogen is extricated. The fixed alkalies combine with it, and form compounds used in medicine.<sup>3</sup>

*Medical properties and uses*.—Sulphuret of antimony is not an active medicine when taken into the stomach, unless it meets with acid in that viscus and the bowels, when it acts with extreme violence; before it is prescribed, therefore, the bowels should be opened. It was not employed internally until the middle of the fifteenth century; and now, owing to the uncertainty of its operation, its occasional violent action, and

<sup>1</sup> Dr. Thomson's proportions, are antimony 100, sulphur 35·572; nearly a mean of all the other analyses that have been published: Vauquelin stated them to be antimony 100, sulphur 33·333; Wenzel, antimony 100, sulphur 29·870; Proust, antimony 100, sulphur 33·333; Dr. J. Davy, antimony 100, sulphur 34·960; Berzelius, antimony 100, sulphur 37·000; Bergman, antimony 100, sulphur 35·035. *Thomson's Chemistry*, 5th edit. i. 556.

<sup>2</sup> Wine was formerly put into cups made of sulphuret of antimony, and owing to the acid acting upon sulphuret, it acquired an emetic quality.

<sup>3</sup> The sulphuret was used by the Greek ladies, and is still employed by the Turkish ladies for staining the eyelashes black, which softens the appearance of the eye. It was a custom among the Jews also; for, although in our translation of the Bible, Jezebel is described as having, "painted her face," (2 *Kings*, ix. 30.) yet the expression in the Hebrew means, literally, "put her eyes in painting."



the difficulty of obtaining it perfectly free from other noxious metals, as copper, lead, and arsenic, it is almost entirely discarded from modern practice. It has been given in gouty and rheumatic affections, in scrofula, and other glandular obstructions, and in chronic cutaneous eruptions.

It produces perspiration; and in a few instances in which it was given in large doses, Dr. Cullen found that some nausea, and even vomiting were excited. It is freely used in veterinary practice, and is given to horses, mixed with their food, to produce a smooth coat. Its chief use is for the preparation of the other antimonial remedies.<sup>1</sup>

The dose of the sulphuret may be from ten grains to one drachm, or more if the stomach can bear it.

*Officinal preparations.* The table drawn up by Dr. Black has generally been given as presenting the best view of the officinal preparations of which antimony is the basis; but as many of those mentioned in it have been long since disused, and the nomenclature of all is changed, we have altered it so as to present, on the same plan, a distinct view of the preparations now found in the British Pharmacopœias.

Medicines are prepared from SULPHURET of ANTIMONY.

I. By trituration in the metallic state united with sulphur.

1. *Sulphuretum Antimonii præparatum*, E. D.

II. By the action of heat; (*oxidized*.)

2. *Antimonii vitrum*, L.

*Oxidum Antimonii cum sulphure vitrificatum*, E.

3. *Oxidum Antimonii vitrificatum cum cera*, E.

III. By the action of heat with phosphate of lime; (*oxidized*.)

4. *Oxidum Antimonii cum phosphate Calcis*, E.

*Pulvis antimonialis*, L. D.

IV. By the action of alkalies; (*oxidized*.)

5. *Antimonii Sulphuretum præcipitatum*, L. E.

*Sulphur antimoniatum fuscum*, D.

V. By the action of acids; (*oxidized*.)

6. *Oxydum Antimonii nitro-muriaticum*, D.

*Murias Antimonii*, E.

7. *Antimonium tartarizatum*, L.

*Tartras Antimonii*, olim *Tartarus emeticus*, E.

*Tartarum antimoniatum*, sive *emeticum*, D.

8. *Vinum Antimonii tartarizati*, L.

*Vinum Tartratis Antimonii*, E.

ANTIMONII VITRUM, Lond. *Glass of antimony*.

This is a vitrified protoxide of antimony, combined with a small portion of the undecomposed sulphuret, from which it is prepared. The sulphuret, reduced to a coarse powder, is

---

It is an ingredient in *Spilsbury's Drops*, which, according to Dr. Paris, consist of *Corrosive Sublimate* ℥ij., *prepared Sulphuret of Antimony* ℥j., *Gentian root* and *Orange-peel*, of each, ℥ij., *Shavings of red Saunders* ℥j., made into a tincture, with a pint of proof Spirit digested and strained.



exposed to the action of a gentle heat, which is gradually augmented so as to drive off a large proportion of the sulphur, which rises in vapours of a white colour and sulphureous odour. The heat is then much augmented, until the powder, when brought to a red heat, exhales no more vapours. During the roasting, the powder is continually stirred to prevent it running into lumps; and, when this part of the process is finished, it is melted with an intense heat. As soon as it has assumed the appearance of fused glass, it is poured on a heated brass plate.

It may be more readily prepared, in a small way, by de-flagrating sulphuret of antimony with twice its weight of nitrate of potass, and melting the protoxide, thus obtained, in a crucible, with less than one-eighth of its weight of sulphur. In this process, the protoxide of antimony, in melting, partially loses its oxygen; and this deoxygenized portion, uniting with the sulphur and being converted into sulphuret, combines, at the moment of its formation, with the remaining undecomposed protoxide, and forms the glass of antimony.

Glass of Antimony, thus prepared, is sold in small flat fragments, glazed on one side and dull on the other, heavy, and having a metallic character.

*Qualities.*—Glass of Antimony is inodorous, insipid, semi-transparent, and by transmitted light of a fine hyacinthine colour. Its specific gravity is 4·95.<sup>1</sup> It is fusible in the flame of a candle, and soluble in the mineral acids. On dissolving it in muriatic acid, sulphuretted hydrogen gas is given out. According to Proust, it consists of eight parts of protoxide of antimony, and one part of sulphuret of antimony. Thenard has shown that 100 parts of this oxide contain 16 parts of oxygen. The glass always contains silicious earth; generally, according to Vauquelin, in the proportion of from 9 to 10 parts in 100, an ingredient of the product which is probably derived from the crucible, during the vitrification of the protoxide. In a late analysis, M. Soubeiran has stated the following to be its constituents—protoxide of antimony 91·5, silica 4·5, peroxide of iron 3·2, sulphuret of antimony 1·9.

*Medical properties and uses.*—Glass of Antimony is so acrid, harsh, and uncertain in its operation, that it is never medicinally employed; and is retained in the Pharmacopœia merely for the purpose of preparing some other antimonial preparations.

Officinal preparations. *Antimonium Tartarizatum*, L. *Oxidum Antimonii vitrificatum cum cera*, E.

<sup>1</sup> The sp. grav. of glass of lead is 6·95, which offers an easy mode of detecting it when mixed with glass of antimony.



All the preparations of antimony contained in the table, have one general mode of action, and possess, therefore, the same medicinal virtues. Their general operation is evacuant, either by the stomach, the bowels, or the skin; but their determination to these particular parts depends more on the dose, and the constitution and state of the patient, than on the nature of the preparation. In small doses they produce nausea, and diaphoresis; in larger doses, vomiting and purging.

Antimonials, prior to the time of Basil Valentine, were used only in veterinary medicine: but ever since they were introduced by that learned Benedictine<sup>1</sup> into the *Materia Medica*, they have been very generally employed for the cure of febrile and inflammatory diseases, when the excitement is great. In the latter stage of fever, however, when much debility prevails, their use is contraindicated. Some have imagined that the preparation which produces the least sensible evacuation, the antimonial powder of the London College, or its prototype James's powder, is to be preferred in typhus, and the tartarized antimony in synochus; believing that the benefit in the first disease is greater when no sensible evacuation is produced: but as this implies some inexplicable specific action of that preparation, we are not inclined to admit the distinction.

ARBUTUS. *Spec. Plant. Willd.* ii. 616.

*Cl.* 10. *Ord.* 1. Decandria Monogynnia. *Nat. ord.* Bicornes, *Linn.* Ericæ, *Juss.*

*G.* 871. *Cal.* five-parted. *Corolla* ovate, the mouth pellucid at the base. *Berry* five-celled.

*Species* 7. *A. Uva Ursi*. Trailing Arbutus, or Bearberry. *Med. Bot.* 2d edit. 287. t. 100. *Smith's Flora Britan.* i. 403.

*Officinal.* UVÆ URSI FOLIA, *Lond. Dub.* ARBUTI UVÆ URSI FOLIA, *Edin.* Leaves of *Uva Ursi*, Bearberry, or Trailing Arbutus.

*Syn.* Bousserole; Raisin d'ours (*F.*), Baerentraube; Sandberren (*G.*), Beerendruif (*Dutch*), Mulbär-riis (*Dan.*), Mjölönris (*Swed.*), Borowekowe (*Pol.*), *Uva Orsina* (*I.*), Madronna *Uva de Orso*; Guaynha (*S.*), *Uva de Urso* (*Port.*), Kleh (*Chipewyan*), Attoonagaweeat, (*Esquimaux*).

This shrub is a native of the north of Europe and is found growing wild on the heathy mountains of Scotland, flowering in June. It is a low shrub, with the branches nearly trailing; woody, and the bark smooth. The leaves are not unlike those

---

<sup>1</sup> BASIL VALENTINE was a Benedictine monk at Erford in Germany. He was born in the year 1394, and was the first person who applied chemistry, which prior to his time was considered merely as the art of making gold, to the purposes of medicine. He was the discoverer of the virtues of antimonial preparations as medicines; and has celebrated them in his "*Currus Triumphalis Antimonii*," a work written in high Dutch; but of which there is an elegant Latin translation by Kirkringius. To Basil Valentine we are also indebted for the discovery of *Ammonia*, and of *Ether*. He recommended a *fixed alkali*, made from the shoots of the vine, cut in the beginning of March, for the cure of the gout and gravel. He was the chief of the medical alchymists.



of the myrtle, thick, evergreen, alternate, obovate, or wedge-shaped, on short petioles; with a net-work of veins on the under surface, which is pale green, whilst the upper is of a very deep green colour, and glossy. The flowers are in small clusters, each supported on a red pedicel. The calyx is small and obtusely five-toothed; the corolla tubular, oval, flesh-coloured, or whitish with a red lip, divided at the margin into five minute, obtuse, reflex segments: containing ten short, downy filaments crowned with erect reddish anthers; and an oval germen, bearing a style longer than the anthers, with a simple stigma. The fruit is a small, round, smooth, glossy, red berry, with a depressed umbilicus, five-celled, of an austere taste, and containing five angular seeds.

The plant should be procured in autumn; and “the green leaves alone selected and picked from the twigs, and dried by a moderate exposure to heat.”<sup>1</sup> The leaves are sometimes adulterated with those of *Vaccinium Vitis Idæa*, which however are easily detected by wanting the reticulated surface of the *Uva Ursi* leaves, and by their infusion not yielding either tannin on the addition of a solution of isinglass, nor displaying the presence of gallic acid on the addition of sulphate of iron.

*Qualities.*—The fresh leaves are inodorous, and have a slightly bitter, astringent taste, leaving a sweet sensation in the mouth. When properly dried and powdered, they acquire an odour similar to that of Hyson tea; but the taste remains the same, the degree of bitterness only being increased. The colour of the powder is a light brown, with a shade of greenish yellow. Both water and alcohol extract its virtues, and the watery infusion strikes a deep black colour with sulphate of iron. According to the analysis of Melandri and Moretti, the leaves yield tannin, mucus, bitter extractive, gallic acid, some resin, lime, and an oxygenizable extract.

*Medical properties and uses.*—*Uva Ursi* possesses astringent properties<sup>2</sup>, on which account it was employed by the ancients in several diseases; but it was not till after the middle of the last century that the attention of modern practitioners was directed to it, as a remedy for calculous complaints, and ulcerations of the urinary organs, by De Haen. His observations were confirmed by Cullen; who, however, referred the good effects it produced to its action on the stomach. It has also been employed in menorrhagia, cystirrhœa, diabetes and other fluxes; and Dr. Bourne has lately recommended it in phthisis pulmonalis. He combined it with Cinchona and Opium, but

<sup>1</sup> *Cases of Pulmonary Consumption, &c.* by Robert Bourne, M.D. 8vo. Lond. 1806.

<sup>2</sup> It is used in Russia for tanning leather.



the cases he published were scarcely sufficiently decisive to confirm its use in this complaint. The dose of the powdered leaves is from ʒj. to ʒj. to be taken two or three times a-day.

ARCTIUM.<sup>1</sup> *Spec. Plant. Willd.* iii. 1630.

*Cl.* 19. *Ord.* 1. Syngenesia Æqualis. *Nat. ord.* Compositæ Capitatae, *Linn.* Cinarocephalæ, *Juss.*

*G.* 1429. *Receptacle* chaffy. *Calyx* globular; the scales at the apex with inverted hooks. *Seed-down* bristly chaffy.

*Species* 1. *Arctium Lappa*. Common Burdock. *Med. Bot.* 2d edit. 32. t. 13. *Eng. Bot.* 1228. *Smith's Flora Britan.* ii. 844.

*Officinal.* ARCTII LAPPÆ SEMINA ET RADIX, *Edin.* BARDANA; RADIX, *Dub.* The root of Burdock.

*Syn.* Bardane (*F.*), Bardana (*I.*), Bardána (*S.*)

This is an indigenous biennial plant, common on the sides of roads and in waste places; flowering in July and August. It is so well known as scarcely to require a description. The root is spindle-shaped, simple, externally of a brown colour, and internally white; the stem succulent, rising three or four feet in height, with spreading branches; and very large, undulated, cordate leaves, of a dark green colour above, and whitish underneath, supported on long footstalks. The flowers are in terminal panicles; the calyx is common, globular, composed of imbricated scales, with hooked extremities, by which they adhere to clothes, and the fur of animals; the corolla is compound, with purple uniform florets, tubular, five-cleft, and all fertile. The receptacle is punctured; the seed downs are rough and prickly, and the seeds quadrangular.

*Qualities.* — The roots of Burdock are inodorous, the taste sweetish, with a slight degree of bitterness and astringency. The seeds, which are sometimes used, are aromatic, bitterish, and subacid.

*Medical properties and uses.* — The seeds and roots of this plant possess some diuretic powers, and are said to determine also to the surface, without exciting nausea, or increasing irritation. They have been employed, and, as far as report can be credited, with advantage, in scurvy, arthritic affections, lues venerea, phthisis, and nephritic complaints. We have no experience of their efficacy; but are ready to believe that the remedy is at least safe. A decoction, made by boiling two ounces of the fresh root in three pints of water to two, should be taken, in divided doses, in twenty-four hours.

ARGENTUM, *Lond. Edin.* ARGENTUM; IN LAMINAS EXTENSUM, *Dub.* Silver. Silver Leaf.

*Syn.* Argent (*F.*), Silber (*G.*), Silwer (*Swed.*), Argento (*I.*), Pláta (*S.*), Villie (*Tam.*), Rupáh (*H.*), Fezzeh (*A.*), Nokra (*Pers.*), Perak (*Malay*), Peddie (*Cyng.*), Rajata (*Sans.*), Im-root (*Esquimaux*).

<sup>1</sup> Ἀρκτίον, Dioscoridis.



Silver exists native, and mineralized, in different parts of the globe, at Barnaoule in western Siberia; but not in any very great abundance. It is found,

A. In its metallic state;

- a. pure, crystallized.
- b. alloyed with gold.
- c. ——— with antimony.
- d. ——— with iron and arsenic.
- e. ——— bismuth.

- Sp. 1. *Native silver*.  
 2. *Auriferous silver ore*.  
 3. *Antimonial silver*.  
 4. *Arsenical silver*.  
 5. *Bismuthic silver*.

B. Sulphurets;

- f. combined with sulphur.
- g. ——— with lead, antimony, }  
     and iron.

1. *Sulphuret of silver*.  
 2. *White silver ore*.

C. Oxidized;

- h. combined with antimonial sulphuret of silver. }

- { 1. *Red silver ore*. Subsp.  
 { a. dark red. b. light red.

D. Salts;

- i. combined with muriatic acid.

- { 1. *Horn silver, common*  
 { and earthy.

- k. ——— with carbonate of }  
     antimony.

2. *Carbonate of silver*.

Besides these ores, there are many metallic ores which contain silver in sufficient quantity to render the extraction of it profitable. In its native state, it is in small lumps, or crystallized in cubes, hexahedrons, octahedrons, or dodecahedrons; and occasionally assumes the form of leaves, threads, or twigs.<sup>1</sup> Its colour is white, its lustre metallic, and fracture hackly. Its specific gravity is from 10 to 10·338. It is not perfectly pure, but contains from ·03 to ·05 of gold, or arsenic or antimony. But silver is obtained in its pure metallic state generally either by fusion or by amalgamation. By the first process the ore is roasted to expel the sulphur, antimony, arsenic, or other volatile principles; the residuum is then fused with lead, and exposed in a cupel, (a vessel made of bone or of wood ashes,) to a strong heat in the hearth of a refining furnace: when the lead and the foreign metals being thus oxidized, are in part absorbed by the porous cupel and in part volatilized and driven off by the current of air from the bellows or the blast pipe. An experienced eye knows when the silver is sufficiently pure; but in general it requires a second cupellation at a higher temperature to purify it completely from the lead with which it is combined. By the second process, the ore is first roasted, then ground to a fine powder, washed, and formed into an amalgam with mercury, by being mixed in small barrels made to revolve very rapidly on their axes by means of machinery. The silver is then separated from the mercury by distillation.

<sup>1</sup> It is found in this form in the famous mine of *Potosi*, and is called *dendrites*.



*Qualities.* — Pure silver is a brilliant-white, insipid, inodorous, sonorous metal, with a very rich lustre, which it loses when long exposed to the air, owing to sulphureted hydrogen being almost always present in the atmosphere. It is in hardness between iron and gold, of considerable malleability, the finest silver leaf being only one-third thicker than gold leaf. It is of inferior ductility to gold, platina, and iron. Its specific gravity is 10·47. Silver is fusible at 28° Wedgwood; volatilized by a stronger heat: but difficult of oxidizement by the action of heat and air. It is oxidized by several of the acids, and combines with them; but none of the compounds, except that produced with the nitric acid, are used in medicine.

*Medical properties and uses.* — Metallic silver has no action on the human body; but when combined with nitric acid, it forms a very powerful remedy. Many of the instruments used by the surgeon require to be made of silver.

Officinal preparation. *Argenti Nitras*, L. E. D.

ARISTOLOCHIA.<sup>1</sup> *Spec. Plant. Willd.* iv. 151.

Cl. 20. Ord. 4. Gynandria Hexandria. *Nat. ord.* Sarmenaceæ, *Linn.* Aristolochiæ, *Juss.*

G. 1609. *Cor.* of one petal, strap-shaped, ventricose at the base.

*Cap.* six-celled, inferior, containing many seeds. *Stem* twining, frutescent.

*Species* 27. *A. serpentaria*. Virginia Snakeroot, or Birthwort.

*Med. Bot.* 2d edit. 152. t. 59. *Veg. Mat. Med. of the United States*, pl. 25.

*Officinal.* SERPENTARIÆ RADIX, *Lond.* — RADIX, *Edin.* SERPENTARIA VIRGINIANA; RADIX, *Dub.* Serpentaria Root.

*Syn.* Serpentaire (*F.*), Virginische Schlangenzwurzel (*G.*), Slangenwortel (*Dutch*), Slangrod (*Danish*), Ormrot (*Swed.*), Serpentaria de Virginia (*I. S. Port.*).

This plant is a native of North America, from Pennsylvania to Florida; flowering in May and June, and ripening its seeds in September. The root is perennial, consisting of bundles of fibres, of a yellow-ochre colour, which changes to brown on drying, attached to a contorted horizontal trunk; from which several stems rise about ten inches in height, slender crooked, and jointed; supporting on long footstalks at each knot, thin, cordate, entire, pointed, trinerved leaves, of a yellowish green colour. The flowers proceed from the joints near the root, and stand upon long sheathed articulated peduncles, which bend down and almost bury the flowers beneath the decayed leaves near the roots. There is no calyx; the corolla is of a brownish purple colour, globular at the base, contracted and bent in the middle, and terminating in a triangular lip. The anthers are sessile, attached to the under-side

---

<sup>1</sup> The *Αριστολοκία* Dioscoridis gives name to the genus; but is not the officinal plant, which was introduced only since the settlement of Europeans in America.



of the stigma, which is roundish, divided into six parts, and almost sessile, rising from an oblong, angular, hairy, inferior germen. The seeds are flat, and contained in a six-celled hexagonal capsule.

Dried serpentaria root is imported into this country in bales, each containing from two to five hundred weight. It is frequently mixed with the roots of *Collinsonia præcox*.

*Qualities.* — The dried root has an aromatic odour, not unlike that of valerian; and a sharp, warm, bitter, pungent taste, resembling in some degree that of camphor. Water extracts all the sensible qualities of the root, affording a yellowish brown infusion, which is not altered by sulphate of iron or zinc, nitrate of silver, oxymuriate of mercury, tartarized antimony, the mineral acids, and the alkalies, nor is it precipitated by gelatine or tannin. The superacetate of lead throws down a flocculent precipitate, which is not soluble in acetic acid, shewing the presence of mucus. With alcohol, it affords a bright greenish tincture, which is rendered turbid by the addition of water. The active principles of *Serpentaria*, therefore, appear to reside in a bitter resin, and an essential oil.

*Medical properties and uses.* — *Serpentaria* root is a stimulating diaphoretic and tonic. It is beneficially employed in typhoid and putrid fevers, whether idiopathic, or accompanying the exanthemata, to excite diaphoresis, and support the powers of the system; and is found frequently to increase the efficacy of *Cinchona* in removing protracted intermittents. It is also an excellent remedy in dyspepsia, particularly when the skin is dry and parched; and is sometimes used as a gargle in putrid sore throat. On account of its stimulant properties, it is contraindicated in the inflammatory diathesis: and, previous to its exhibition, the bowels should be well evacuated.

It may be given in substance, or in infusion made by macerating ℥j. of the bruised root in f ℥xij. of boiling water, in a covered vessel for two hours, and straining. Decoction is a bad form of preparation of *Serpentaria*, as the boiling dissipates the essential oil, on which the virtues of the remedy chiefly depend. The dose of the powdered root is grs. x. or grs. xx., increased to ʒss.; that of the infusion f℥jss. to f℥ij. given every fourth hour.

Officinal preparations. *Tinctura Serpentariæ*, L. E. D. *Tinctura Cinchonæ composita*, L. D. *Electuarium opiatum*. E.

ARMORACIÆ RADIX. Vide *Cochlearia Armoracia*.

ARNICA. *Spec. Plant. Willd.* iii. 2106.

Cl. 19. Ord. 2. Syngenesia Superflua. Nat. ord. Compositæ Discoideæ, Linn. Corymbiferae, Juss.



G. 1491. *Recep.* naked. *Seed-down* simple. *Cal.* with equal leaflets. *Corol.* of the ray have more frequently five filaments without anthers.

*Species* 1. *A. Montana*, Mountain Arnica. *Med. Bot.* 2d edit. 41. t. 17. *Flor. Dan.* t. 728. *Nuttall. Gen. Aus.* ii. p. 164.

*Officinal.* ARNICÆ MONTANÆ FLORES ET RADIX, *Edin.* ARNICA; FLORES, RADIX, *Dub.* The flowers and root of Arnica.

*Syn.* Arnique (*F.*), Arnika, Woheverleih, Falkraut (*G.*), Valkruid (*Belg.*), Wolverley (*Dan.*), Häst sibler, St. Hansblomster (*Swed.*), Arnica (*I.*).

This species of Arnica is a native of the northern parts of the continents of Europe, and America, and of Siberia; flowering in July. It is also found on the Pyrennees, and is cultivated in our gardens.<sup>1</sup> The root is perennial, brown, woody, præmorse, with bundles of long fibres attached to it: the stem, which rises about a foot in height, is obscurely angular, striated, rough, hairy, and terminated by two or three upright peduncles, each bearing one flower. The radical leaves are ovate, entire, ciliated, and more obtuse than those of the stem, which are in opposite pairs, and lance-shaped. The flowers are of a deep copper-yellow colour, tinged with brown; the calyx is a dirty green, composed of fifteen or sixteen lancet-shaped hairy scales, with purple points: the ray consists of about fourteen ligulate flowers, twice as long as the calyx, striated, three-toothed, and hairy at the base; the disc of tubular florets, with a five-lobed margin. The seeds are oblong, striated, hairy, and crowned with a russet-coloured down.

The herbaceous part of the dried herb, which is used equally with the flowers and root, seems as if covered with a hoary powder.

*Qualities.* — The dried plant has a pleasant, weak, aromatic odour, and excites sneezing. The taste of the *leaves* and *flowers* is slightly aromatic, bitter, and pungent; that of the root, bitter and acrid. The leaves and flowers, macerated in boiling water, yield an olive-brown infusion, which has an odour not unlike that of senna, and a bitter, hot taste. It reddens tincture of litmus; but does not precipitate glue, nor alter solutions of tartarized antimony and of oxymuriate of mercury. With sulphate of iron and of zinc, it strikes a deep green colour, and gives dark precipitates. Superacetate of lead coagulates it. The mineral acids render it muddy, and of a dirty white colour, occasioning brown precipitates; but the alkalies only deepen its proper colour. Both alcohol and sulphuric ether take up from the flowers and leaves a resinous matter, which can be separated from the alcohol by water, and

---

<sup>1</sup> It was introduced by Mr. P. Miller in 1759.



from the ether by evaporation. Hence we may conclude, that Arnica contains a peculiar acid,<sup>1</sup> resin, a nauseous bitter matter, tannin, and mucus; and that sulphates of iron and of zinc, superacetate of lead, and the mineral acids, are incompatible in prescriptions with infusions of its leaves and flowers.

*Medical properties and uses.* — The *leaves* and *flowers* of Arnica are narcotic, stimulant, and diaphoretic; and in large doses, emetic and cathartic: the *root* is tonic and aromatic. The former have been used with advantage in paralytic affections, amaurosis, gout, rheumatism, and chlorosis. They have been extolled also in convulsive diseases, diarrhœa, and dysentery; but in the latter, their stimulant properties prove often hurtful. In paralysis, their good effects are generally preceded by a pricking sensation in the affected part; but in general they do not produce any sensible operation, unless when exhibited in too large doses: in which case they produce great anxiety, pain, vomiting, and the other deleterious effects of powerful narcotics. The *root* has been much extolled in Germany, as a succedaneum for Cinchona in intermittents, putrid fevers, and gangrene; particularly by Dr. Collin of Pazman; but in the hands of British practitioners it has not deserved the high encomiums which he has bestowed on it in these cases. It is regarded by the French practitioners as an excellent tonic in paralysis.<sup>2</sup>

Externally the powdered leaves may be used as an errhine.<sup>3</sup>

Arnica may be exhibited in substance; or in an infusion, made by macerating ʒjss. of the leaves and flowers, or ʒij. of the bruised root, in fʒxij. of boiling water, and straining through linen. The infusion soon ferments. A dose of the powder is from grs. v. to grs. x.; that of the infusion, fʒjss. twice or thrice a day.

#### ARSENICUM<sup>4</sup>, Arsenic.

*Syn.* Arsenic (*F.*), Arsenick (*G.*), Arsenico (*I.*), Arsénico (*S.*).

This metal is found in most parts of the world, accompanying other metals, and occasionally uncombined, forming distinct and peculiar veins. The following are the states in which arsenic is found:—

<sup>1</sup> Bouillon la Grange thinks it is the Gallic acid.

<sup>2</sup> Vide *Nouv. Elemens de Therapeutique*, par I. L. Alibert, 2d ed. vol. i. p. 141.

<sup>3</sup> The Savoyards, and the inhabitants of the Vosgeses, both snuff and smoke the leaves; and thence the plant is known on the Continent by the name of *Tabac de Savoyards et de Vosges*.

<sup>4</sup> From *αρσενικον*, Dioscoridis, which, however, is not the metal, but realgar, one of the species of the sulphuret; *σανδαραχη* of the other Greeks.



*A.* In its metallic state:

- i. Alloyed with iron, or  
silver, or gold. *Sp.* 1. *Native arsenic*.

*B.* United with sulphur and iron.

- ii. sulphurets. *2. Arsenical pyrites.*  
*3. Orpiment.* Var. *a.* Realgar.  
*b.* Yellow orpiment.

*C.* United with oxygen:

- iii. oxide. *4. Native oxide.*

*D.* Acidified; and

- iv. Combined with lime. *5. Arseniate of lime. Pharmacolite.*

- v. ————— copper. *6. ————— copper.*  
Var. *a.* Foliated,  
*b.* Lenticular,  
*c.* Oliven ore.

- vi. ————— iron. *7. ————— iron. Cube ore.*

- vii. Combined with lead. *8. Arseniate of lead.*

- viii. ————— cobalt. *9. ————— cobalt. Red cobalt ore.*

- Var. *a.* Cobalt crust.  
*b.* ————— bloom.

As metallic arsenic is not used in the arts, it is not extracted from its ores, but is prepared for the purposes of experiment or of curiosity from the white oxide, which is commonly procured in roasting the arseniate of cobalt. It is necessary, however, to be acquainted with the appearances and properties of metallic arsenic, as one mode of ascertaining whether the white oxide has been used as a poison, in cases of suspected death, is to reduce the oxide.

Its colour is bluish gray, something like that of steel, with much brilliancy. It is quickly tarnished by exposure to the air, becomes black, and falls into powder. It is extremely brittle, and pulverulent. Its specific gravity is 5.763. It volatilizes at a heat of 356° Fahrenheit, in dense white fumes, which have the odour of garlic, although the solid metal is inodorous. In its metallic state, Arsenic exerts no action on the animal system; but when oxydized it is a virulent poison.

*Officinal.* ARSENICUM ALBUM, *Lond.* OXIDUM ARSENICI, *Edin.*

ARSENICUM (*Oxydum album*), *Dub.* White Arsenic.

*Syn.* Arsenic oxydé natif (*F.*), Naturlicoer Arsenickhalk (*G.*), Arsenik (*Dan. Swed. Polish*), Muschjak (*Russian*), Arsenico iixneo (*I.*), Arsenico Turabulhalik (*Arab.*), Samuel-k'har (*H.*), Sanc'hya (*San.*), Vallay Paskanum (*Tam.*), Sumulfar (*Pers.*), Wrangon (*Malay.*).

The greater part of the white oxide of Arsenic of commerce is obtained in Bohemia and Saxony, in roasting the cobalt ores, in making zaffre, and sometimes by sublimation from arsenical pyrites. The roasting is performed in furnaces with long flues, in which the impure oxide is condensed, and this is purified by sublimation in the following method. Large square boxes of cast iron, furnished with conical heads, which



are closely luted to them with clay, are disposed in a brick area, heated by the flues of two furnaces placed a little beneath them. When these boxes are red-hot, the impure arsenic, by fifteen pounds at a time, is put into them, where it melts, and soon sublimes in the conical head. Successive additions are thus submitted to the action of heat, till about 150 pounds have been used to each vessel; and then the apparatus is allowed to cool. The conical head is now separated from the box, and carried with its contents into another place, where the workmen break off with hammers the sublimed oxide, separating the impurities for a second operation.<sup>1</sup>

The oxide, or *arsenious acid*, thus obtained is a dense, semi-transparent, solid cake; which becomes opaque, of a snowy whiteness, and pulverulent, when exposed to the air. It is met with in both these forms in the shops; and often is sold in powder, in which state it is sometimes adulterated with white sand, chalk, and gypsum; but the fraud is easily detected by heating a small portion of the suspected powder; by which the oxide is entirely dissipated, and the impurities are left behind.

The greater quantity of the oxide of arsenic used in this country is brought from Germany, in casks, each containing from two to five hundred weight.

*Qualities.* — White oxide of arsenic is inodorous; has an acrid taste, leaving on the tongue a sweetish impression; and is highly corrosive.<sup>2</sup> When pure, if it has not been freely exposed to the action of the air, it is in semi-transparent, colourless, shining masses, which break with a conchoidal fracture. It is soluble in 400 parts of water at 60°, and in 13 parts of boiling water; and the latter solution, on cooling, retains three parts of the white oxide for every 100 of water, and deposits the remainder in tetrahedrous crystals.<sup>3</sup> Both solutions redden infusion of litmus, and combine with the alkalies. It is soluble also in solution of pure potass, in alcohol and in oils. When heated in the open air, this oxide is volatilized in a temperature of about 383° Fahr., and the vapour has no odour: but if it be heated in contact with any substance, which has a strong affinity for oxygen, the vapours have an alliaceous colour, owing to the partial reduction of the oxide.<sup>4</sup> The

---

<sup>1</sup> *Journal de Physique*, tom. i. p. 44.

<sup>2</sup> The majority of Chemists now regard it as an acid.

<sup>3</sup> Klaproth, *Schweigger's Journal*, vol. vi. p. 932; and *London Med. Repository*, vol. ii. p. 250.

<sup>4</sup> The smelters of copper in Cornwall and Wales, although much exposed to the vapours of arsenic, yet suffer very little from them; but they are sometimes attacked with cancer in the scrotum. Dr. Paris (*Pharmacologia*) remarks, that they rely upon oil being an antidote; and are consequently supplied with it by their employers.



specific gravity of the oxide in its ordinary state is 3·706, that of the glass 3·699. According to the average of the experiments by different chemists, 100 parts of the oxide consist of 74·6 of arsenic, and 25·4 of oxygen.<sup>1</sup> On the simple watery solution of the oxide, no change is produced by a solution of sulphate of iron, of oxymuriate of mercury, tartarized antimony, the mineral acids, or the alkalies: but nitrate of silver throws down a yellowish precipitate, which gradually passes to a brown colour; and a white precipitate is produced by superacetate of lead. Lime water also precipitates it white; sulphurets of the alkalies, pale yellow; and sulphuretted hydrogen gas, golden yellow.

*Medical properties and uses.* — Although White Oxide of Arsenic is the most virulent of the mineral poisons, yet when properly administered, it is a medicine of great efficacy; and is employed internally as a tonic, and externally as an escharotic. It had been long used as an internal empirical remedy in cancer, and some cutaneous affections, both in Europe and the East Indies; and for the cure of intermittents in Hungary; and in Lincolnshire under the name of “the ague drop:” but its effects were not clearly understood, nor the proper mode of administering it known, till Dr. Fowler of Stafford published his Observations on its use in the cure of remitting fevers and periodic head-aches.<sup>2</sup> Since that time the authority of many respectable practitioners has been brought forward in confirmation of its efficacy in these diseases; and in lepra, chronic rheumatism, intermittent hemicrania or *megrim*, scirrhus, and some local painful affections “of the ends of the bones, cartilages, or ligaments, or of all three together.” It has also been used in dropsy, hydrophobia, syphilis, visceral and glandular obstructions, and in many other diseases, in which, however, its efficacy is by no means established.<sup>3</sup> In the East Indies the native physicians employ arsenic (*sanc’-hya*) made into pills with six parts of black pepper, for the cure of confirmed lues (*Persian fire*), and a species of elephantiasis (*Judham*).<sup>4</sup> It is also used in cases of the bite of the hooded snake, *cobra del capello*.

The internal use of white oxide of arsenic is contraindicated

<sup>1</sup> Proust, Davy, and Dr. Thomson’s proportions are, arsenic 100 × 34·930 oxygen. *Annals of Phil.* iv. p. 176.

<sup>2</sup> It is a curious fact that previous to the introduction of copper-works in Cornwall, agues were very frequent; but since that period the disease is extremely rare. “I have heard it,” says Dr. Paris (*Pharmacologia*), “remarked by the men in the works, that the smoke kills all fevers.” Is this owing to the arsenical fumes?

<sup>3</sup> For a list of these diseases see a paper by Mr. Hill of Chester. *Edinburgh Med. Journal*, v. 19. 312. and vi. 55.

<sup>4</sup> Asiatick Researches, 8vo. 5th edit. vol. ii. p. 158.



in all cases attended with strong arterial action ; and where there are any pulmonary symptoms : and should a cough even intervene during its use, it should be instantly discontinued. When it is exhibited in proper cases, and with necessary precaution, the effects it produces must be carefully observed : “ the feeling of swelling and stiffness of the palpebræ and face, heat, soreness and itching of the tarsi, or tenderness of the mouth,<sup>1</sup>” are indications that the dose of the remedy has been carried to its full extent, and should then be diminished. If erythema or salivation appear, the use of it must be suspended : and it should be altogether abandoned if pain of the stomach, or nausea, vomiting, head-ache, vertigo, or cough be induced.

The white oxide is exhibited either in substance or in solution. The best mode of giving it in substance is in the form of pills, formed by rubbing one grain of the oxide with ten grains of sugar, and then beating the mixture with a sufficient quantity of crumb of bread, so as to form ten moderately-sized pills ; one of which is a dose. The solution, however, is more manageable. The most common form of it is that of the London College ; (vide *Liquor arsenicalis*), but the simple solution in distilled water, in the proportion of four grains to a pint, is also given according to M. Le Febvre’s method. A table spoonful of the solution, mixed with a little syrup of poppies and half a pint of milk, is directed to be taken in a morning fasting, and the frequency of the dose increased until six spoonful be daily taken.

As an external application the oxide of arsenic has been long employed in cases of cancer : and has certainly done more to improve the ulceration, and give it a disposition to contract and heal, than any other external application. It has been sprinkled, in the form of powder, upon the sores ; but the most violent pain follows this mode of applying it ; and in some instances, probably from its absorption, the general system has been dangerously affected. The more usual mode of using it is in the form of a lotion, composed of eight grains of the oxide, and the same quantity of subcarbonate of potass, dissolved in four fluid ounces of water ; or as an ointment, formed by rubbing together one drachm of the oxide and twelve drachms of spermaceti ointment. These applications produce little pain and irritation, cause the diseased parts to slough off, and amend the fetid discharge ; but, although, to a certain extent they produce the most beneficial effects, yet, the instances in which a cure has been effected are very rare.

---

<sup>1</sup> Dr. Kellie, *Edin. Med. Journ.*



The white oxide of arsenic is not unfrequently the cause of death; from accidents occurring to those artists who use it in their manipulations; as glass-makers, dyers, and workers in gold: or from ignorance of the proper dose of its preparations when medicinally used; or from the employment of it as a poison. The symptoms which occur are those of inflammation of the stomach, incessant vomiting, purging, and pain of the stomach<sup>1</sup>; constriction of the throat, an increased flow of saliva and great heat of the mouth; rapid sinking of the pulse, which is at first full, hard and frequent, cold sweats, convulsions, and death: but if the quantity be not sufficient to produce speedy dissolution, the first-mentioned symptoms are succeeded by paralysis, hectic, and other symptoms of extreme debility.<sup>2</sup> When death takes place, symptoms of putridity are said soon to present themselves; but this is not always the case, although the body is often marked with livid stripes, and covered with enchymoses; and on dissection the stomach often, although not always, appears either abraded, or completely eroded in several parts; with appearances of inflammation extending through the whole abdominal viscera: but from M. Brodie's experiments there is great reason for supposing that death is less the consequence of inflammation than of a peculiar and immediate influence of the poison upon the nervous system. Particles of the arsenic are occasionally found adhering to the abraded parts of the villous coat of the stomach.

Various methods of counteracting the poison of arsenic have been recommended. Whatever antidote is adopted, the stomach should, in all cases, be immediately evacuated: and the best mode of doing this is by administering large draughts of tepid mucilaginous fluids. In order to render the Arsenic inert, solutions of the alkaline sulphurets, and vinegar, have been advised; but the experiments of Renault have demonstrated how little reliance is to be placed on these articles. Hahneman orders one pound of soap to be dissolved in four pounds of water, and a cupful taken, tepid, every three or four minutes; and as this is the antidote most readily procured, if lime-water or chalk and water cannot be at hand, it should always be the first employed. Lime-water proves useful by coating the particles of the arsenic with an arseniate of lime, which is insoluble, and consequently inert.<sup>3</sup> Dr. Yelloly,

---

<sup>1</sup> In a case detailed by Dr. Yelloly, no pain of the stomach, convulsions, nor delirium occurred, although it terminated fatally. *Edin. Med. and Surg. Journ.* v. 389.

<sup>2</sup> On this subject our readers will find much practical information in the *London Medical Repository*, vol. v. p. 97.

<sup>3</sup> See our experiments on this subject, in the *London Med. Repos.* vol. viii. 157.



reasoning on the probability that the inflammation induced is often the cause of death, even after the stomach is freed from the whole of the poison, suggests the propriety of early blood-letting in these cases.<sup>1</sup> Opium, camphor, and ether may be employed to quiet the nervous irritability, and ammonia to stimulate the heart.

As medical men are often called upon in courts of law to establish the fact of White Oxide of Arsenic having been used as a poison, it is necessary to know the best tests by which it may be recognised. If on searching in the stomach, or among its vomited contents, any considerable quantity of the suspected poison be discovered, a little of it must be mixed with three times its weight of black flux, composed of one part of finely-powdered charcoal, and two parts of dry carbonate of potass; or, to a grain of the poison add half a grain of charcoal, and a grain of dried carbonate of potass. These must be put into a thin glass tube, about eight inches in length and 1-4th inch in diameter, hermetically closed at one end. The open extremity must then be slightly plugged with a piece of paper (taking care to clean the upper portion of the tube by means of a feather), and the tube kept for a quarter of an hour in a well-burnt coal fire; or the flame of a spirit-lamp: when, if the powder introduced into the tube contain Arsenic, metallic Arsenic will sublime and be found lining the inside of the tube with a brilliant crust. That it was Arsenic may be further proved by volatilizing a small portion of the reduced metal on a red-hot iron, and observing whether it present the garlic odour peculiar to the vapour of metallic Arsenic. The white oxide may also be detected in the following manner: Mix some of the suspected matter with the black flux; place the mixture between two pieces of polished copper; and after binding them tightly together with iron wire, place them in an ordinary fire: if Oxide of Arsenic be present, a white stain will be left on the surface of the copper, which is an alloy of metallic Arsenic and the copper.

When the poison is found in very small quantity only, let it be dissolved in two drachms of hot rain or distilled water, with three grains of subcarbonate of potass, or, what is to be preferred, the subcarbonate of ammonia; then add to this a warm solution of five grains of sulphate of copper, which will produce a lively grass-green precipitate if Arsenic be present, which, if heated in a test-tube with a little wax will sublime in the form of metallic Arsenic. When no powder is discovered in the stomach, its contents and the vomited matter

---

<sup>1</sup> *Edinburgh Med. and Surg. Journal*, v. 392.



must be washed with hot water, and filtered, carbonate of potass added to the filtered fluid, and then a warm solution of the sulphate of copper, as above described. Mr. Phillips, however, has shewn that if the sulphate contain any peroxide of iron, it will afford a green precipitate, although no Arsenic be present. To prevent this fallacy, he directs the potass to be added to the sulphate of copper first, and to a part of this the suspected solution, when the blue precipitate will become green, if arsenious acid be present. A still more delicate test than any of those already mentioned has been proposed by Mr. Hume<sup>1</sup>: one part of the suspected poison, and three parts of subcarbonate of potass, are to be dissolved in a sufficient quantity of rain or distilled water at 212°; and the surface of this solution slightly touched with a piece of nitrate of silver. If Oxide of Arsenic be present, a sulphur-yellow coloured precipitate will be seen falling rapidly from the point where the nitrate is applied. In our experiments we have found that the sixtieth part of a grain of the oxide is clearly discovered in two ounces of water by this test. This precipitate should afford, when heated with wax, indications of metallic Arsenic, as in the former case. All these experiments should be performed in the day-time, and the precipitated fluid examined by reflected, not transmitted light.<sup>2</sup> — Objections have been raised against this test, because the presence of the alkaline phosphates in the suspected fluid would produce precipitates of a similar colour with nitrate of silver; and if muriate of soda, or of any other alkali, were present, the test could not be employed, on account of the copious precipitates which these produce with the nitrate. The first objection is obviated by making the trial on paper, as recommended by Dr. Paris: Drop a little of the suspected fluid on writing-paper, and draw several times over it a stick of lunar caustic; which, if arsenic be present, will leave a streak of colour, that becomes a very bright queen's yellow, if brushed with some liquid ammonia, which continues unchanged; but if no arsenic be present, and only alkaline phosphates, the streak will be uniform, and in a few minutes fade into a sad green, and gradually become black. Dr. Marcet has shown us how to obviate the difficulty with regard to the muriates, by adding to the suspected fluid dilute nitric acid, and then to apply the nitrate of silver to its surface until no more precipitation is produced; by which means the whole of the muriatic acid is removed: and as the arseniate of silver re-

---

<sup>1</sup> *Philosophical Mag.* May 1809.

<sup>2</sup> Bostock, *Edin. Med. and Surg. Journ.* v. 170.



mains in solution, it is rendered evident by a yellow precipitate being instantly formed on the addition of ammonia. — But the great difficulty is to detect the presence of the poison in the stomach. Besides examining the contents of this viscus by filtration and dilution, if no Arsenic can be detected, Orfila advises the viscus to be cut in pieces and then separately examined<sup>1</sup>, by boiling them in water and testing the decoction. In this case, the best test is a current of sulphuretted hydrogen gas passed through the filtered decoction, which will diffuse a fine yellow colour in the fluid if any Arsenic be present. If no colour appear, which may occur, although Arsenic be in the decoction, owing to the presence of phosphate of soda, the addition of very dilute pure nitric acid will produce it.<sup>2</sup>

When any of the vehicle in which the poison has been exhibited can be procured, more satisfactory results will be obtained from the examination of it, than from that of the contents of the stomach. If a white powder be found in it, the most satisfactory proof is that of reducing this to the metallic form, as already described; but, if the whole of the arsenic be dissolved, it must then be tested by different reagents. One of the simplest methods which I have tried is the following. Into the suspected solution stir a moderate quantity of charcoal powder; allow it to settle; then pour off the clear supernatant fluid, or filter the mixture; and when the powder which remains on the filter is dry, sprinkle some of it on a red hot poker: if the solution contain arsenic, the odour of garlic will be rendered sensible. This effect becomes more obvious, if a few grains of dried subcarbonate of potass be added to the dried charcoal powder.<sup>3</sup> The results from no

---

<sup>1</sup> *Traité des Poisons*, &c. par M. P. Orfila, vol. ii. p. 169.

<sup>2</sup> Vide *An Essay on Chemical Analysis*, &c. By J. G. Children. Lond. 8vo. 1817. p. 393.

<sup>3</sup> To ascertain the delicacy of this test, the following experiments were made.—  
Ex. 1. Half a drachm of white oxide of arsenic being boiled in two ounces of water, and the fluid filtered when cold, it was found to retain twenty-eight grains of the white oxide in solution. Exp. 2. One drachm of this solution being mixed with two ounces of water in a cylindrical glass vessel, so as to form a solution which contained about one part of the oxide for 592 parts of water, a scruple of finely-powdered charcoal was added, and the mixture being well agitated with a glass rod, and allowed to settle, was filtered. The powder, when dry, on being thrown upon a red hot shovel, emitted a *very faint* odour of garlic. Exp. 3. The same as the former, except that two drachms of the solution were employed, making the proportion of the white oxide to the water in the diluted solution, as 1 to 286, the garlic odour was *very perceptible*. Exp. 4. Four drachms of the arsenical solution being employed, making the proportion of the white oxide to that of the water in the diluted solution as 1 to about 143, the garlic odour was *extremely strong*. From these experiments it is evident that this test will detect arsenic in any solution strong enough to act as a poison.



single test should, however, be relied upon : and as a knowledge of the appearances produced by the four principal reagents usually employed for the detection of arsenic, must greatly facilitate such an examination, we have constructed the following table from actual experiments ; comparing the results obtained from solutions of arsenic with those from solutions of corrosive sublimate, tartarised antimony, and muriate of barytes, which are the only substances likely to be mistaken for arsenic. It is necessary to remark, that the broth employed was made with beef, and contained a moderate proportion of carrots, turnips, and onions, and that the coffee and the tea contained milk and sugar in the usual proportions employed in these beverages.



COMPARATIVE TABLE of the Precipitates obtained from Solutions of White Arsenic, of Oxymuriate of Mercury, of Tartarized Antimony, and of Muriate of Barytes, with different Tests.

1st TEST.—WATER SATURATED WITH SULPHURETTED HYDROGEN GAS.				
Solvents.	Precipitates from Solutions of WHITE ARSENIC.	Precipitates from Solutions of CORROSIVE SUBLIMATE.	Precipitates from Solutions of TARTAR EMETIC.	Precipitates from Solutions of MURIATE of BARYTES.
Water - - - -	Bright golden yellow, which was deepened by the addition of a few drops of strong acetic acid. <sup>1</sup>	Yellow at the instant of its formation, but soon becoming blackish. On shaking the tube it changes to a dirty white.	Orange, curdy, partly suspended, partly thrown down. Ultimately bright orange.	Heavy, and of a dirty, dark brown colour.
Broth - - - -	Scarcely any at first, but on adding a few drops of strong acetic acid, a pale yellow.	Whitish yellow at first, quickly changing to mixed clots of yellow, black and white.	Pale orange at first, soon changing to a deeper bright orange.	Dirty pale brown, heavy.
Milk - - - - -	Little change, but on the addition of a drop of strong acetic acid, a straw-coloured precipitate.	Light ochre, requiring for its formation a large quantity of the test.	Golden yellow, with a shade of orange.	Dirty nankeen, with a shade of brown.
Tea - - - - -	At first, very pale yellow; after some time, a pale greenish yellow. The precipitate was curdy.	Brownish white and yellow, mixed.	Deep orange, curdy, slowly formed: the supernatant fluid yellow.	Dirty light brown, deepening as it fell.
Madeira Wine	Turbid, pale yellow, the colour of the wine destroyed.	Muddy, gradually displaying small floating black flocculi.	Pale orange, long suspended	The muriate mixed with white wine is milky.—Not tested.
Port Wine -	Turbid, paler yellow: the precipitate in both is very slowly formed.	Nearly as in the white wines, like clouds through the purple of the wine.	Dark, dirty brown.	Pale brown, heavy.
Coffee - - - -	A deep golden yellow.	Brownish black.	Deep orange brown.	Not tried.
Gruel - - - - -	Pale yellow suspended.	Light brown, slowly formed.	Pale orange.	Not tried.

<sup>1</sup> This precipitate, dried upon a filter, and heated with some caustic potash in a slender test tube, is decomposed in a few seconds, forming a sulphuret of potash, whilst the arsenic is volatilized in its metallic form, and adheres to the sides of the tube. (*Orfila.*)



## II. HYDRO-SUPHURET OF POTASH.

Solvents.	Precipitates from Solutions of WHITE ARSENIC.	Precipitates from Solutions of CORROSIVE SUBLIMATE.	Precipitates from Solutions of TARTAR EMETIC.	Precipitates from Solutions of MURIATE of BARYTES.
Water - - -	White, with the faintest tint of sulphur yellow, when a large quantity of the test was used. <sup>1</sup>	Black, mottled with yellow.	Bright orange.	Deep olive green.
Broth - - -	Pale, but bright, sulphur yellow.	Clotted, heavy, black, mottled with gray. Black, clotted.	Dull orange, heavy. Orange.	Lt. brown, partially suspended Brown, greenish when the mixture was shaken.
Milk - - -	Bright golden yellow.	Brownish black.	Reddish orange, flocculent.	Not tried.
Tea - - -	A beautiful yellow.	Nearly black.	Deep brownish orange.	Not tried.
Coffee - - -	A deep golden yellow. <sup>2</sup>	Dirty white, or slate colour.	Beautiful bright orange.	Vide 1st Table.
Madeira Wine	Sulphur yellow.	Slate colour, with violet supernatant liquor. <sup>3</sup>	Dark brown, with a tinge of orange.	Violet, heavy.
Port Wine -	Fawn colour.	Black dense clots. <sup>4</sup>	Orange-clotted.	Dusky yellowish green.
Gruel - - -	Bright queen's yellow.			

## III. SOLUTION OF SULPHATE OF COPPER, with the addition of Ammonia in excess.

Water - - -	Beautiful grass green. It completely disappeared on the addition of a few drops of strong acetic acid. <sup>5</sup>	White thick, and heavy.	Pale whitish blue, very little thrown down.	Copious whitish blue.
Broth - - -	Beautiful pale green, suspended. <sup>6</sup>	White, curdy, partly suspended, partly thrown down.	Pale whitish blue, with a tint of green. Whitish blue.	Opaque, glaucous.
Milk - - -	Pale greyish green.	Blueish white, curdy.	Muddy, pale blueish green.	Curdy white, with a tinge of blue.
Tea - - -	Obscure olive, but scarcely a precipitate.	Dirty yellowish white, curdy.	Dirty blueish green. inous blue.	Greyish, heavy, supernatant fluid, yellowish green.
Coffee - - -	Dark grass green. <sup>7</sup>	Dirty white.		Not tried.
Madeira Wine	Greenish, with a slight tinge of green.	Heavy clotted white, with a tint of green.		Vide 1st Table.
Port Wine -	Clotted, heavy, dark greenish grey.	Heavy, clotted, blueish grey.	Heavy, dirty slate blue, Pale blueish green.	Dirty violaceous grey. Pale blueish green.
Gruel - - -	Beautiful grass green.	Pale blueish white.		



Solvents.	Precipitates from Solutions of WHITE ARSENIC.	Precipitates from Solutions of CORROSIVE SUBLIMATE.	Precipitates from Solutions of TARTAR EMETIC.	Precipitates from Solutions of MURIATE of BARYTES.
Water - - -	Copious bright sulphur yellow, falling in flocculi from the point of contact. <sup>8</sup>	Dull yellowish white, clotted, changing to dirty white.	Pale brown.	White, heavy; soon blackening.
Broth - - -	White, (owing to the muriate of soda) but yellow when treated with nitric acid.	White, copious.	Brownish, mixed with much muriate of silver.	White, dense, curdy.
Milk - - -	White, with a tint of yellow.	Dirty white.	Very pale, scarcely visible brown.	Not tried.
Tea - - -	Yellowish white, which soon blackens.	Dirty white.	Dirty brown.	Not tried.
Coffee - - -	Yellow, remaining unchanged.	White, changing to black.	Not tried.	Not tried.
Madeira wine	Pale sulphur yellow.	Dirty white, changing to black.	White.	Vide 1st Table.
Port Wine	White, becoming brown on exposure to the light.	Ibid.	Dirty white.	Heavy, dirty white.
Gruel - - -	Yellowish.	Dense, dirty white clots.	Not tried.	Dense, clotted, white.

<sup>1</sup> The hydrosulphuret, added to a solution of the phosphates, throws down a greenish yellow precipitate, the supernatant fluid being yellow and turbid.

<sup>2</sup> Lime water, also, added to coffee containing arsenic, throws down a yellow precipitate; although it precipitates the watery solution of arsenic white. (*Orfila.*)

<sup>3</sup> Corrosive sublimate cannot be exhibited in port wine with an intention to commit murder, (except by a self-murderer,) as it changes the colour of the wine to pale violet.

<sup>4</sup> All the precipitates by the sulphuret, when dried, and heated in a tube with iron filings, afford metallic mercury, which is volatilized, and shews itself in globules upon the sides of the tube.

<sup>5</sup> This test is capable of detecting arsenic in a solution containing  $\frac{1}{100000}$  of its weight. (*Orfila.*)

<sup>6</sup> It has been suggested, that onions boiled in broth, or eaten so as to impregnate with their peculiar qualities the contents of the stomach, might produce the same effects on sulphate of copper, as if arsenic were present; but although a green colour is produced, yet no precipitate falls as when arsenic is present.

<sup>7</sup> Dr. Porter, of South Carolina, says that sulp. of copper with ammonia produced the same coloured precipitate in coffee, which contains no arsenic? *American Journ. of Science*, vol. iii. p. 354.

<sup>8</sup> A similar precipitate is formed by nitrate of silver, in a solution of any of the phosphates; and with chromate of potass; but the fact of the precipitate being occasioned by arsenic, is easily ascertained by testing a fresh portion of the solution with lime water. If it contain arsenic, a copious white precipitate will be thrown down; if a phosphate only, there is scarcely any change, or at the most a translucent flocculent precipitate, which remains long suspended. A new method of employing this test was suggested by Dr. Paris. It is to put upon a piece of clean white paper a broad streak of the suspected fluid; and then run lightly over it a stick of lunar caustic. This is an excellent test, when modified as I have elsewhere (*London Med. Repository*, vol. viii. p. 178.) suggested, by brushing the streak lightly over with liquid ammonia, immediately after the application of the caustic. If arsenic be present, a bright queen's yellow is instantly produced, which remains permanent for nearly an hour: but, when the lunar caustic produces a bright yellow before the ammonia is applied, we may suspect the presence of some phosphate rather than arsenic.



ARTEMISIA. *Spec. Plant Willd.* iii. 1815.

Cl. 19. Ord. 2. Syngenesia Superflua. *Nat. ord.* Compositæ Nucamentaceæ, *Linn.* Corymbiferæ, *Juss.*

G. 1473. *Receptacle* subvillous or almost naked. *Seed-down* none. *Cal.* imbricate, with roundish converging scales. *Cor.* without rays.

\* *Shrubby.*

*Species* 8. *A. Abrotanum*, Southernwood. *Med. Bot.* 2d edit. 52. t. 21.

\*\*\* *Herbaceous, with the stem somewhat branching, the flowers in panicles, the leaves compound.*

*Species* 26. *A. santonica*, Tartarian Southernwood. *Med. Bot.* 2d edit. 61. t. 23.

——— 42. *A. maritima*. Sea Wormwood. *Med. Bot.* 2d edit. 60. edit. 24. *Smith Flora Brit.* 864.

——— 63. *A. Absinthium*. Common Wormwood. *Med. Bot.* 2d edit. 54. t. 22. *Smith's Flora Brit.* 864.

#### 1. ARTEMISIA ABROTANUM.<sup>1</sup>

*Officinal.* ABROTANUM; FOLIA, *Dub.* Southernwood leaves.

*Syn.* Citronelle Auronne (*F.*), Eberaute, Stabwurz (*G.*), Averoon (*Dutch*), Abröd (*Dan.*), Abrodd (*Swed.*), Bozedrzwkko (*Polish*), Abrotano (*I.*), Limbri-guera (*S.*).

This a perennial undershrub, a native of the south of Europe, Siberia, China, and Cochinchina. In England, where it is abundantly cultivated, it resists the winter, but very rarely flowers. It rises about three feet in height, with a shrubby branching stem covered with smooth brown bark. The leaves are alternate on long footstalks, irregularly doubly pinnate; the pinnæ linear, concave on the upper surface, convex below, tomentose, and of a pale green colour. The flowers are compound, of a greenish yellow colour, and produced on one-flowered peduncles in axillary spikes at the extremities of the branches. The seeds are naked and solitary.

*Qualities.* — Southernwood has a strong fragrant odour; and a warm, bitter nauseous taste. Both water and alcohol extract these qualities; but the alcohol more perfectly than the water, the infusion having scarcely any bitterness. The tincture is of a beautiful green colour, the infusion of a pale olive. The latter strikes the black with sulphate of iron and precipitates acetate of lead. A small quantity of essential oil is procured by distillation; on which, and a bitter resinous matter, the qualities of the plant appear to depend.

*Medical properties and uses.* — Southernwood is said to pos-

<sup>1</sup> Ἀβροτόνον, Dioscoridis, ab ἀβροτον inhumanum; vel ἀβρωτον cibo inutile. Vide *Alston's Mat. Med.* ii. 65.

<sup>2</sup> Dr. Burdach of Wiebal, near Serase, and Professor C. G. Huffeland have used the root of *Artemisia vulgaris*, Mugwort, taken up in October, and administered in powder in doses, of from gr. l. to gr. lxx., with great success in epilepsy. *Journ. der Pract. Heilkunde.* 1824.



sess tonic, diaphoretic, anthelmintic, and deobstruent properties. It was formerly much used in debilities of the stomach, chlorosis, and jaundice. Externally it has been employed as a discutient and anodyne fomentation for inflammations, pains, tumours and gangrenous ulcers. But it is very rarely used in modern practice. The dose may be from ℥j. to ʒj. of the leaves in substance; or, of an infusion made with ʒvj of the leaves and f℥x of water, a cupful, taken twice or thrice a-day.

## 2. ARTEMISIA SANTONICA.

*Officinal.* ARTEMISIÆ SANTONICÆ CACUMINA, *Edin.* SANTONICUM<sup>1</sup>; CACUMINA, *Dub.* The tops of Tartarian Southernwood.

*Syn.* Sementine (*F.*), Tartarisches Beyfus (*G.*), Santonico (*I.*).

This species of artemisia is a native of Tartary and Persia; but it is cultivated in our gardens, flowering in September. The root is perennial; and the plant has the habits of indigenous field southernwood, but is erect. The stem is paniced, rising two feet in height, and rather hoary. The lower leaves are pinnate, much cut, linear, and hoary. The branches are wand-like, with alternate racemes, recurved, and having flowers all looking the same way. The flowers are solitary, and cylindrical. In the fruiting plant all the stems are erect, and lose their hoariness. The leaves on the branches are very small, linear, and undivided. The receptacle is naked.<sup>2</sup>

The *qualities* and *medical properties* of this plant are nearly the same as those of the former species of artemisia; and it may be used for the same purposes. The worm seeds (*semina Santonici*) of the former pharmacopœias, which were supposed to be the production of this plant, are now properly rejected, as their place can be well supplied with anthelmintics of more certainty.

## 3. ARTEMISIA MARITIMA.<sup>3</sup>

*Officinal.* ABSINTHIUM MARITIMUM; CACUMINA, *Dub.* The tops of Sea Wormwood.

*Syn.* Absinthe Maritime (*F.*), Seebeifuss (*G.*), Zeealsem (*Dutch*), Vild Cypress (*Danish*), Svenski marum (*Swed.*), Assenzio Maritimo (*I.*).

This is an indigenous, perennial plant, growing near the sea shores, and in salt marshes, flowering in August. The root is fibrous, and somewhat woody. The stems rise two or three feet in height. They are erect, leafy, furrowed, and hoary. The inferior leaves are pinnate, with three-cleft pinnae; the upper ones variously divided; the highest simple, and the whole entire in the margin, downy on both sides. The flowers are ovate, of a brownish-yellow colour, in nodding racemes. The calyx is woolly on the outside, with a scaly margin; the receptacle is naked, and the ray florets are very few.

<sup>1</sup> Σαντονιον, Dioscoridis.    <sup>2</sup> Willdenow, iii. 1827.    <sup>3</sup> Σεριφιον, Dioscoridis.



*Qualities*. — The odour is slightly fragrant, and the taste bitter and weakly aromatic. Like the first described species, its activity seems to depend on a bitter resin and essential oil.

*Medical properties and uses*. — These are in every respect the same, in a diminished degree, as those of the next species. It is scarcely ever used.

#### 4. ARTEMISIA ABSINTHIUM.<sup>1</sup>

*Officinal*. ABSINTHIUM, *Lond*. ARTEMISIÆ-ABSINTHII FOLIA, SUMMITATES, *Edin*. ABSINTHIUM VULGARE; FOLIA, CACUMINA. The leaves and flowering tops of Wormwood.

*Syn*. Absinthe commun (*F.*), Wormuth (*G.*), Alsem (*Dutch*), Malurt (*Danish*), Malert (*Swed.*), Piolun (*Polish*), Polin (*Russian*), Assenzio (*I.*), Artemisio axenjo (*S.*), Losna (*Portug.*).

Common wormwood is an indigenous perennial plant, growing in dry waste places, and flowering in August. The greater part, however, of that which is used for medicinal purposes is cultivated in the physical gardens.<sup>2</sup> The root is somewhat woody, and branched. The stems rise nearly erect to the height of two or three feet; are branching, angled, and furrowed, with the summits panniced. The lower leaves are bipinnate; the upper pinnatifid or digitated; with oblong obtuse very entire segments. The racemes are erect, and the flowers pedicellated, nodding, hemispherical, and of a brownish-yellow colour. The florets of the disk are numerous, but those of the ray few: and the receptacle is covered with white silky hairs, shorter than the calyx.

*Qualities*. — The odour of common wormwood is strong, and although fragrant, yet to many persons it is very disagreeable: the taste is intensely bitter, slightly pungent, and nauseous. These qualities are given out both to water and alcohol: and a dark-green essential oil, on which the odour depends, is obtained by distillation with water. The watery infusion of the plant has a pale olive colour: sulphate of iron, and of zinc, slowly deepen it to a black; and superacetate of lead throws down a yellowish-green flocculent precipitate: but it is not affected by Tartarized Antimony. The active parts of the plant seem to be extractive, an essential oil which is not in the least bitter, and a small portion of resin. Kuns-muller<sup>3</sup> found in the residue of 12 ounces of the plant after infusion, besides other things, 59 grains of carbonate of lime.

*Medical properties and uses*. — Common wormwood is tonic, and anthelmintic; and, when externally applied, is said to be

<sup>1</sup> Ἀψινθιον, Dioscoridis.

<sup>2</sup> A good deal is cultivated at Mitcham in Surry, chiefly for the seed, which is sold to the rectifiers of British spirits at about 30s. per cwt. — *Stevenson's Survey*, p. 378.

<sup>3</sup> *Ann. de Chim.* vi. p. 35.



discutient and antiseptic. It has been used with advantage in intermittents, gout, scurvy, and dropsy; and although modern practitioners will scarcely rely on its efficacy in these complaints, yet it is undoubtedly of some value as a stomachic in dyspepsia and hypochondriacal affections. When it is desirable to free the remedy from its narcotic property, it should be given in decoction, as the boiling dissipates the essential oil on which this depends. The dose in substance is ℥j. to ℥ij. and of the infusion, made by macerating ʒvj. of the plant in fʒxij. of boiling water, fʒj. to fʒxij., which may be given three or four times a-day.<sup>1</sup>

Official preparations. *Extractum Absinthii*, D.

ARUM. *Spec. Plant. Willd.* iv. 477.

Cl. 21. Ord. 7. Monœcia Polyandria. Nat. ord. Piperitæ, Linn. Aroideæ, Juss.

G. 1705. *Spathe* one-leafed, cowled. *Spadix* naked above, female below, staminate in the middle.

\* *Stemless with compound leaves.*

Sp. 17. *A. maculatum*.<sup>2</sup> Arum, or Cuckow-pint. *Med. Bot.* 2d edit.

728. t. 249. *Eng. Bot.* 1298. *Smith Flora Britan.* iii. 1024.

Official. Arum; RADIX RECENS. *Dub.* The recent root of Arrum.

*Syn.* Gouet (F.), Aronswurzel (G.), Arro manchado (S.), Iurro (Port.), Gw-lakt kalfsvort (Dutch), Tydsk ingefaer (Dan.), Dansk Ingefær (Swed.), Aronowa broda (Polish), Aro (I.).

This is a perennial indigenous plant, growing under hedges and on the sides of banks in many parts of Britain<sup>3</sup>; flowering in May, and ripening its berries in August. The root is tuberous, about the size of the first joint of the thumb, with many radical fibres issuing from every side. The leaves, which seldom exceed four in number, are radical, supported on grooved sheathing petioles about nine inches long: they are triangular and barbed, five inches in length, and two in breadth at the base, smooth, glistening, of a deep green colour above, and often sparsely spotted with dark brown or black blotches. The flower stem is a simple, erect scape, from eight to twelve inches high, and obscurely channelled. The spathe is erect, bellied, pointed above, of a pale green colour, sometimes spotted, withering, and covering the fruit till it is nearly ripe, when it drops. The spadix is enclosed in the sheath, is club shaped, obtuse, of a purple colour above, and whitish below; collared towards the middle with many sterile filaments; a little lower with numerous sessile anthers; and at the base with many roundish germens, crowned with sessile, simple stigmas. The berries are succulent, of a bright scarlet colour when fully ripe, and contain two or more hard seeds.

<sup>1</sup> Purl is an infusion of wormwood in ale.

<sup>2</sup> Ἀρον μέγα, Hippocratis.

<sup>3</sup> There is great abundance of it in the lanes near Ewell and Dorking in Surry.



For medical use the roots of arum should be dug up in autumn, after the leaves are completely decayed. They may be preserved fresh for nearly a year if buried in sand in a cool cellar.

*Qualities*.—The arum root is white, and inodorous. When chewed the taste is at first sweetish and soft, but it soon excites a burning, pricking sensation on the tongue and in the mouth, which continues many hours, and is attended with great thirst. Butter, milk, and oily fluids allay these unpleasant sensations. The sliced root applied to the skin, reddens, and excoriates or vesicates it. The acrimonious matter, however, can be washed off from the bruised root by water; is completely dissipated by drying; and abstracted by a mixture of water and alcohol by distillation, although the fluid receive no sensible impregnation; so that it may be regarded as a vegetable principle *sui generis*. The recent expressed juice reddens vegetable blues; and has been found to contain malate of lime.<sup>1</sup> The dried root is chiefly fecula, perfectly inert, and saponaceous; and is used in France as a cosmetic, under the name of Cypress powder.

*Medical properties and uses*.—Arum root in its recent state is stimulant, diaphoretic, and expectorant. It has been employed in cachetic, chlorotic, and rheumatic cases; and in humoral asthma. Bergius says, he found it a never-failing remedy for cephalæa sympathica, which resisted all the other means he employed. But the difficulty of procuring Arum root always in a state to be depended on, prevents it from becoming a remedy of general utility.

The dose of Arum, in substance, may be from grs. x. to ℥j, three or four times a day, combined with any thing which can sheath its acrimony, as mucilage, milk, or thick barley-water; or triturated with gum and water, so as to form an emulsion.

ASARUM. *Spec. Plant. Willd.* ii. 858.

*Cl.* 11. *Ord.* 1. Dodecandria Monogynia. *Nat. ord.* Sarmentaceæ, *Linn.* Aristolochiæ, *Juss.*

*G.* 925. *Calyx* three or four-cleft, placed on the germen. *Corolla* none. *Capsule* coriaceous, crowned.

*Spec.* 1. *A. Europæum*.<sup>2</sup> Asarabacca. *Med. Bot.* 2d edit. t. 66.

*Eng. Bot.* t. 1083. *Smith. Flora Brit.* 509.

*Officinal.* ASARI FOLIA. *Lond. Edin.* ASARUM; FOLIA. *Dub.* Asarabacca leaves.

*Syn.* Asaret; Cabaret (*F.*), Hazelwurtzel (*G.*), Hazelwort (*Dutch*), Hasselurt (*Dan.*), Kopytnik (*Polish*), Asaro, la bacchera (*I.*), Asaro de Europa (*S.*), Haselört (*Swed.*), Asāroon (*Arab.*), Tuckir (*Hind.*), Mootriennjavia (*Tam.*).

<sup>1</sup> *Ann. de Chimie.* xxxv. 153.

<sup>2</sup> Ἀσάρον, Dioscoridis. The Arabic word *Asāroon* signifies astringency.



This is a perennial plant, the geographical limits of which extend from  $60^{\circ}$  to  $37^{\circ}$  N. latitude; and is consequently a native of several parts of England, particularly Lancashire and Westmoreland: growing in woods and shady places; and flowering in May. The root is creeping, fleshy, and fibrous. The stem short, round, simple, pubescent, generally bearing two leaves only, and one flower. The leaves are opposite, on footstalks three inches long, of a kidney shape, entire, somewhat hairy, and of a deep shining green colour. The flower is on a short terminal peduncle of an herbaceous colour on the outside, and dusky purple within; and is in some degree hid under the leaves: the calyx is bell-shaped, and three-cleft, with the points of the segments, which are erect, turned inwards: there is no corolla: the filaments are produced beyond the anthers into a hook or little horn; and the style is a cylindrical column crowned with a six-parted reddish stigma. The seeds are few, contained in a six-celled, inferior, coriaceous capsule, egg-shaped, and crowned with a persistent calyx.

As a great deal of the acrimony of *Asarabacca* is lost with keeping, the leaves should be used in as recent a state as possible; and dried without the application of much heat.<sup>1</sup>

*Qualities.* — The recent leaves are nearly inodorous; their taste slightly aromatic, bitter, acrid, and nauseous. The decoction is inert, but the watery infusion, which has the colour of brandy, possesses the sensible qualities of the leaves. Sulphate of iron changes the colour to a deep olive, throwing down a greyish precipitate. The recent root when distilled yields a volatile oil which smells like camphor; but this is not obtained from the dried root. The recent root possesses, also, emetic properties.

*Medical properties and uses.*—The leaves of *Asarabacca* are emetic, cathartic, and diuretic; but in modern practice they are never used except as an errhine; and, perhaps, as Dr. Cullen has remarked, they form the most useful species of this genus of local stimulants. A proper dose snuffed up the nose for a few successive evenings at bed-time, occasions a copious discharge from the nostrils, which continues to flow for several days. They have been found particularly beneficial in cephalæa, obstinate tooth-aches, chronic ophthalmia, and lethargic affections. The dose of the powdered leaves is grs. iij. to grs. v., which should be repeated every night untill the full effect is produced, avoiding exposure to cold during its use.

---

<sup>1</sup> The roots, which are not ordered in the British pharmacopœias, contain the same acrid principle as *Arum*; and are violently emetic and cathartic. Their odour, which is not unlike that of *Valerian*, is said to prove fatal to moles. *St. Hilaire, Expos. des Fam. Nat.* vi. 174.



Officinal preparation. *Pulvis Asari compositus.* E. D.

ASSAFŒTIDÆ GUMMI RESINA. Vide *Ferula Asafœtida.*

ASPIDIUM. *Flora Britannica, Smith, 1118.*

Cl. 24. Ord. 1. Cryptogamia Filices. Nat. ord. Filices Linn.

G. 429. (Smith.) Fructification in roundish points, scattered, not marginal. Involucre umbilicated, open almost on every side.

\*\* Frond. nearly bipinnate.

Species 4. *A. Filix mas.*<sup>1</sup> Male Fern root. *Med. Bot. 1st edit. t. 49.*

(*Polypodium Filix mas.*) *Eng. Bot. 1458.*

Officinal. FILICIS RADIX. Lond. ASPIDII FILICIS MARIS RADIX.

Edin. FILIX MAS; RADIX. Dub. Root of the Male Fern.

Syn. Fougère (F.), Johanniskwurtzel (G.), Felie Maschia (I.), Polypodio heleo masculino (S.).

This is a common indigenous, perennial plant, growing in woods and shady places, and flowering in June and July. The root consists of many matted fibres, forming a turfy or cespitose head, of a blackish colour and scaly. The leaves or fronds grow in circular tufts, from a foot to four feet in height, with the stipe and mid-rib chaffy. They are of a bright green colour, lanceolate and pinnate. The pinnæ are at first alternate, increasing in size from the base towards the middle, then decreasing towards the summit of the leaf; each being deeply subdivided into linear obtuse parallel lobes, crenate on the edges. The fructification is like small dots on the back of each lobe, placed in two rows near the base, and distant from the edges; composed of a kidney-shaped shield or involucre, and a pale brown capsule, with a saffron-coloured elastic ring.

*Qualities.* — The dried root is nearly inodorous; the taste at first sweetish, then slightly bitter, subastringent; and mucilaginous when chewed. By a late analysis of M. Mirca it was found to contain a fatty substance of a nauseous odour and disagreeable taste, heavier than water; — gallic and acetic acids, uncrystallizable sugar, tannin, starch, and a variety of gelatine soluble in water and alcohol. He found that it also contained subcarbonate, sulphate, and hydrochlorate of potass, carbonate et phosphate of lime, alumine, silex and oxide of iron.<sup>2</sup> The internal part of the root, which yields a reddish powder, is the portion that is medicinally used.

*Medical properties and uses.* — This root is astringent, and has been celebrated both by the ancients and the moderns as a powerful anthelmintic. It appears to have been used as such by Theophrastus, Dioscorides, and Galen; but although recommended by Hoffman, yet it was neglected by the moderns, until the publication of Madame Noufer's specific for the tape worm, by the French government, again brought it into notice.

<sup>1</sup> Ὠλυπτερίς, Dioscoridis.

<sup>2</sup> *Ann. de Chim.* xxvi. 219



According to her plan of administering it, from one to three drachms of the powdered root are to be taken in a large cupful of water, in the morning while the patient was in bed: and, two hours afterwards, a strong cathartic of calomel and gamboge, proportioned to the age and strength of the patient, is to be given; and, if necessary, the further operation promoted by a dose of purging salts; nothing but broth being taken till the worm come away. If this, however, did not happen on the same day, the process was ordered to be repeated on the next day.

Notwithstanding the celebrity of this remedy, there is every reason for ascribing more efficacy to the cathartic than to the fern root; and it may now be rejected altogether from the materia medica, oil of turpentine being a more certain remedy for expelling tænia.

ASTRAGALUS. *Spec. Plant. Willd.* iii. 1256.

Cl. 17. Ord. 4. Diadelphia Decandria. *Nat. ord.* Papilionaceæ or Leguminosæ *Linn.*

G. 1379. *Legume* generally two-celled, gibbous.

*Species.* — *A. verus*. True Astragalus. *Olivier, Voy. dans l'Empire Ottoman*, v. 342. pl. 44.

*Officinal* TRAGACANTHA. *Lond.* ASTRAGALI TRAGACANTHÆ GUMMI. *Edin.* Tragacanth.

*Syn.* Gommi Adraganti (*F.*), Traganth (*G.*), Gom Drangant (*Dutch*), Draganti (*I.*), Gomo Tragacantho (*S.*), Alcatira (*Port.*), Sumégh ulkassael (*Arab.*), Xuttivah (*H.*), Vaeberno cottuy pisin (*Tam.*).

This shrub is a native of the north of Persia, where it is called Kurn, flowering in July and August. It rises two or three feet only in height, on a stem about an inch in thickness; with many branches closely crowded together, and covered with imbricated scales and spines, formed from the petioles of the former year. The leaves, which scarcely exceed half an inch in length, are composed of six, seven, or eight pairs of opposite, villous, stiff, pointed leaflets; and the mid-rib is terminated with a sharp yellowish point. The flowers are small, yellow, and proceed from the axilæ of the leaves, with cottony bractes. The calyx is five-toothed, and shorter than the corolla, which is papilionaceous.<sup>1</sup>

The gum exudes in summer, more or less copiously according to the heat of the weather, in tortuous filaments, which are allowed to dry on the plant before being collected. A large portion of the Tragacanth collected in Persia is sent to

---

<sup>1</sup> Before Oliver discovered that the species of *Astragalus* above described yields the Tragacanth of commerce, this gum was supposed to be yielded by the *A. Tragacantha* of Linnæus, on the authority of Tournefort; or the *A. gummiifera*, on that of M. de la Billardiére. Perhaps all these species yield it, although the *A. verus* be that from which it is more generally procured. The Kuttirah gum from India has been found not to answer the purposes of the ordinary Tragacanth.



India, Bagdad, Bussorah, and Russia. But what we receive is sent to Aleppo, whence it is exported, packed in cases.

*Qualities.* — Good Gum Tragacanth is inodorous; impressing a very slightly bitter taste as it dissolves in the mouth. It has a whitish colour, is semitransparent, and in very thin, wrinkled, vermiform, pieces: brittle but not easily pulverized, except in frosty weather, or in a warmed mortar. It swells and softens in water, but does not form a homogeneous fluid-mucilage, unless triturated after digestion with a large portion of water: but when the water is acidulated with any of the mineral acids, a small portion of it is dissolved. It is also insoluble in alcohol and ether. Dr. John has given the name of *Cerasin* to this species of Gum, from its being exuded pure from the Cherry-tree, *Prunus cerasus*. Its mucilage differs from that of acacia gum in being precipitated by the superacetate of lead and oxymuriate of tin; and not by silicated potass<sup>1</sup>, or the oxysulphate of iron.

*Medical properties and uses.* — Gum Tragacanth is demulcent; and may answer the purposes of the Acacia gum; being even better adapted for allaying tickling cough, and sheathing the fauces in catarrhal affections, owing to its greater viscosity. It is chiefly, however, employed for pharmaceutical purposes. The dose is grs. x. to 3 j. or more.

*Officinal preparations.*—*Mucilago Astragali Tragacanthæ*. E. D. *Pulvis Tragacanthæ comp.* L.

ATROPA. *Spec. Plant. Willd.* i. 1016.

Cl. 5. Ord. 1. Pentandria Monogynia. *Nat. ord.* Luridæ Linn. Solonaceæ Juss.

G. 381. Cor. bell-shaped. Stam. distant. Berry globular, 2 celled. *Species* 2. A. *Belladonna*. Deadly Nightshade, or Dwale. *Med.*

*Bot.* 2d ed. 230. t. 82. *Eng. Bot.* 592. *Smith. Flor. Brit.* 253.

*Officinal.* BELLADONNÆ FOLIA, *Lond. Dub.* ATROPÆ BELLADONNÆ FOLIA. *Edin.* Deadly Nightshade leaves.

*Syn.* Balladone (*F.*), Tollkraut (*G.*), Doodkruid (*Dutch*), Wargbar (*Dan. Swed.*), Belladonna (*I. S. Portug.*), Psinki (*Pol.*), Sug-ungger (*H.*), Robuturbuc (*Pers.*), Inubus sahel (*Arab.*).

*Belladonna* is an indigenous perennial, found in many parts of Great Britain, particularly in shady places where the soil is calcareous, flowering in June, and ripening its berries in September. The root is thick, fleshy, and creeping; sending up several erect, purple-coloured, herbaceous, annual stems, about three feet in height, branching, leafy, round, and somewhat fleshy. The leaves are lateral, in pairs of unequal size, decurrent, on short petioles, egg-shaped, pointed, entire; of a dusky green colour above, and paler below; soft and fatty to the touch. The flowers are supported on one-flowered, soli-

<sup>1</sup> Bostock, *Nicholson's Journ.* lviii. 30.



tary, axillary peduncles; large, drooping, and having a faint narcotic odour: the calyx is green, persistent, and deeply divided into five ovate segments; the corolla bell-shaped, of a lurid hue externally, and within dusky or brownish violet, with a yellow variegated base, inclosing five filaments shorter than the corolla, nodding, and bearing large anthers; with a pyramidal germ, supporting a long simple style and two-lobed stigma. The ripe berry is large, seated within the calyx, roundish, with a longitudinal furrow on each side, shining, smooth, and of a deep purple colour; containing many seeds, and a sweetish, violet-coloured juice.

*Qualities.* — The leaves of Belladonna are inodorous; the taste is slightly nauseous, sweetish, and subacid. They do not lose their active properties by drying. Vauquelin found that they contain a substance resembling animal albumen, salts with a base of potash, and a bitter principle, on which their narcotic quality depends; which has since been ascertained by M. Brandes to be an alkali, which he has named *Atropia*<sup>1</sup>. Every part of the plant is poisonous; and children and the ignorant have often suffered from eating the berries, the beautiful appearance and sweet taste of which render them very alluring. The symptoms which they induce are those of intoxication, accompanied with fits of laughter and violent gestures<sup>2</sup>; great thirst, difficulty of deglutition, nausea, dilatation of the pupil, with the eyelids drawn down; redness and tumefaction of the face, stupor or delirium, a low and feeble pulse, paralysis of the intestines, convulsions and death. Dissections show that the stomach and intestines have been inflamed; and after death the body swells, blood flows from the nose, mouth, and ears; and the most rapid decomposition

---

<sup>1</sup> To obtain it boil the dried leaves in distilled water, press the decoction out and filter, after the albumen has been thrown down by a little sulphuric acid; then add potash as long as a precipitate is produced: wash this precipitate in pure water, redissolve it in muriatic acid, and again precipitate by ammonia. This is *Atropia*. It is in white acicular crystals, insipid, little soluble in cold water, or even alcohol, but very soluble in boiling alcohol, from which, however, it is deposited on cooling. — *Ann. of Phil.* vol. i. p. 2, 3. new series. *Schweigger's Journ.* vol. xxviii. p. 1. M. Runge, however, asserts that *Atropia* is destroyed by alkalies, and consequently they cannot be used to obtain it. He recommends hydrate of magnesia, procured by decomposing the sulphate by potash. This hydrate just precipitated is added to the aqueous infusion of Belladonna; and the whole evaporated to dryness. The residue is then pulverized, and treated with alcohol. The crystalline mass obtained from the spontaneous evaporation of the alcoholic solution filtered, is *Atropia*. *Ann. de Chim.* xxvii. p. 32:

<sup>2</sup> Buchanan, the Scottish historian, states that the victory of Macbeth over the Danes was obtained chiefly by mixing this plant in a donation of wine and ale, which was sent by the Scots to Sweno during a truce. He describes very accurately the botanical characters of the plant, and adds “vis fructui, radici, ac maxime semini somnifera, et quæ in amentiam si largius sumantur, agat.” — *Rerum Scot. Hist.* lib. vii. sect. 6.



ensues. The best mode of averting these fatal effects is by exhibiting emetics of sulphate of zinc, or of copper, and assisting their operation by irritating the fauces; then evacuating the bowels by active purgatives and glysters; and following these by large doses of vinegar and other vegetable acids. The recovery is always slow. If M. Runge's experiments be correct, lime water should be the antidote for poisoning by Belladonna.

*Medical properties and uses.* — The deleterious effects we have enumerated demonstrate that Belladonna is a very powerful narcotic. It is besides diaphoretic, and diuretic. When injudiciously or incautiously given, or when it is taken for a considerable length of time, even in small doses, it is apt to induce a dryness and stricture of the pharynx and adjoining parts of the œsophagus, sickness, vertigo, and dimness of sight; symptoms sufficiently indicative of the necessity of suspending its use for some time, and giving it in smaller doses when it is resumed. The internal administration of Belladonna appears to have been suggested by the advantages resulting from its external application. Cullen, De Haen, Junker, and others, found it very serviceable in scirrhus and cancerous affections; and it has also been given with advantage in obstinate intermittents, chronic rheumatism, gout, paralysis, amaurosis, and pertussis, in which we can speak of its efficacy from our own experience. Hufeland asserts that it has the power of allaying convulsions arising from scrophulous irritation; and its beneficial effects in neuralgia facialis have been well ascertained.<sup>1</sup> Its narcotic powers are certainly great; but they have not been found sufficiently constant and permanent to insure its general use. Externally, used either as a fomentation, or the dried leaves powdered and sprinkled over the parts, it is of singular efficacy in diminishing the pain of cancerous and ill-conditioned sores: and as the infusion, when dropped into the eye, produces a great dilatation of the pupil, it was proposed by professor Reimar, for dilating the pupil previous to the extraction of the cataract, and the extract is now commonly used in this country, for the same purpose.<sup>2</sup> M. Runge<sup>3</sup> ascertained that this power is destroyed by alkaline solutions. "Its operation appears to be limited to the radiated fibres of the iris."<sup>4</sup> By continued use it loses its effect: but regains it after the application has been, for a short time, suspended. Professor Koreff has stated that Belladonna given during the prevalence of

<sup>1</sup> *Observations on the use of Belladonna, &c.* by John Bailey, 8vo. 1817.

<sup>2</sup> *Med. and Phys. Journal*, No. xxxii.

<sup>3</sup> *Ann. de Chim.* xxvii. 32.

<sup>4</sup> *Adams's Practical Observations on Ectropium, &c.* 8vo. p. 44.



Scarlatina, has the power of protecting the individual who takes it from the infection.

Belladonna may be given in substance, beginning with one grain of the dry leaves powdered, and gradually increasing the dose to 12 or 14 grains; or of an infusion, made with one scruple of the dried leaves in ten fluid ounces of boiling water, two ounces may be given daily, and the dose cautiously increased. From what has been stated regarding the effect of alkalies on Belladonna, these bodies are incompatible in formula with it.

Officinal preparations.—*Extract. Belladonnæ*, L. *Succus spissatus Atropæ Belladonnæ*, E.

AURANTII BACCÆ. Vide *Citrus urantium*.

AVENA. *Spec. Plant. Willd.* i. 443.

Cl. 3. Ord. 2. Triandria Digynia, *Nat. ord.* Gramina, *Linn.*

G. 142. *Calyx* two-valved, many-flowered; with a twisted awn on the back.

*Species* 13. *A. sativa*.<sup>1</sup> Common Oat.

Off. AVENÆ SEMINA. *Lond. Edin.* The seeds of the Oat decorticated, called Grits.

*Syn.* Gruau d'avoine (*F.*), Habergriize (*G.*), Gewoone haver (*Dutch*), Havre (*Dan.*), Hafra (*Swed.*), Avena (*I.*), Avena (*S.*), Avea (*Portug.*).

The oat was found by Anson growing wild upon the island of Juan Fernandez, on the coast of Chili; but the place whence it was first brought to Europe has never been satisfactorily ascertained. The root is annual and fibrous, pushing up a culm or straw, which rises above two feet in height. The inflorescence is a loose panicle, with the subdivisions on long pendulous peduncles. The glumes of the calyx are two, marked with lines, pointed, unequal, and larger than the flower. There are usually two flowers and seeds in each calyx; they are alternate, conical; the smaller one is awnless: the larger puts forth a strong, two-coloured, bent awn from the middle of the back of the glumule: both seeds are fertile.

There are many varieties of this species of grain cultivated in the north of Europe.<sup>2</sup> In this country that which is called *the potatoe oat* is considered the best; its pickle is short and plump, with a thin, clean, bright, pale straw-coloured cuticle.

Oats, when freed from their cuticle only, are named grits; in which state, and ground into meal, they are dietetically and medicinally used. In both states they yield their fecula to water by coction; and form a nutritious amylaceous gruel. The nutrient qualities of oats are well known. In many

<sup>1</sup> Βρώμος, Dioscoridis.

<sup>2</sup> In Scotland, some parts of England, a part of Siberia and in the northern parts of Norway, and Sweden, oats form the chief part of the vegetable nutriment of the inhabitants.



places the meal forms the chief support of the poor: and for infants who are unfortunately deprived of their natural and proper nourishment, the breast milk, no better substitute can be adopted than thin grit gruel mixed with good cow's milk. The gruel should not be kept longer than forty-eight hours, as it becomes acescent after that period.<sup>1</sup>

*Qualities.* — Oats are inodorous; and taste very slightly, but not unpleasantly, bitter. They have not been chemically examined; but the greater part of their substance appears to consist of fecula or starch.

*Medical properties and uses.* — Gruels, or decoctions of grits or of oatmeal, are excellent demulcents, and therefore very frequently prescribed in inflammatory diseases, diarrhoea, cholera, dysentery, calculus, and febrile affections. They may be sweetened, acidified, or used plain. They are also used locally in glysters; and the meal boiled with water into a thick paste forms an excellent, suppurative poultice.

**BALSAMUM PERUVIANUM.** Vide *Myroxylon Peruvianum*.

**BALSAMUM TOLUTANUM.** Vide *Toluifera Balsamum*.

**BARYTA.** Barytes.

*Syn.* Baryte (*F.*), Baryterde, Schwerede (*G.*), Barite (*I.*).

This mineral substance does not exist, as far as we know, in an uncombined state; and its native combinations hitherto discovered are very few. It is found

A. Combined with carbonic acid:

Sp. 1. *Carbonate of barytes, or Witherite.*<sup>2</sup>

B. ——— with sulphuric acid:

2. *Sulphate of barytes, or Heavy spar.*

Barytes is obtained by decomposing these fossils. It is not a simple substance, but a compound of a peculiar metallic base named *barium* by Sir H. Davy<sup>3</sup>, and oxygen, in unknown proportions.

#### 1. CARBONATE OF BARYTES.

*Officinal.* **CARBONAS BARYTÆ, Edin.** Carbonate of Barytes.

*Syn.* Carbonate de Baryte (*F.*), Kohlonsaure Baryterde (*G.*), Ossicarbonato di Barite (*I.*)

This fossil is found native in Sweden, Scotland, and Cumberland, but, in greatest abundance at Anglesark, in Lanca-

<sup>1</sup> The following is the simplest mode of making gruel: Put three ounces of grits which have been washed, into four pints of water, and boil slowly until the water be reduced one half; then strain through a sieve to separate the undissolved part of the grits from the gruel.

<sup>2</sup> So named by Werner, after Dr. Withering, who discovered it native on Alston moor in Cumberland, in 1783.

<sup>3</sup> Mr. Murray proposes *barytum* instead of barium. *Sys. of Chem.* ii. 205.



shire. It usually occurs massive in veins, which traverse the independent coal formation; and sometimes, although rarely, it is found crystallized. The crystals are small, and their primitive form is yet undecided.

*Qualities*.—Carbonate of barytes is inodorous and insipid, but is nevertheless poisonous. Its colour is white, or yellowish grey: it is translucent, with a shining, somewhat resinous lustre; and breaks in one direction with a fracture intermediate between radiated and foliated, and in another uneven: the fragments wedge-shaped. The lustre of the principal fracture is glimmering; that of the cross glimmering and resinous. Its specific gravity is 4.331. When heated it becomes opaque; and is fused into a white enamel by the blow-pipe; and, when pulverized, its powder phosphoresces when thrown on burning coals. It is nearly insoluble, requiring for its solution 4304 times its weight of cold water; and 2304 of boiling water. It dissolves with effervescence in diluted nitric acid, although the strong acid exerts no action on it. It consists of 9.75 of barytes and 2.75 of carbonic acid in 100 parts.

*Use*.—It is only used for preparing the muriate. It may, however, be exhibited as a poison; and in this case the antidote is diluted sulphuric acid, as the sulphate which is thus formed is an inert salt.

Officinal preparation. *Murias Barytæ*, E.

## 2. SULPHATE OF BARYTES.

*Officinal*. SULPHAS BARYTÆ, *Edin*. Sulphate of Barytes.

*Syn*. Sulfate de Baryte (*F.*), Schwefelsaures Baryterde (*G.*), Tungspat (*Swed.*), Ossisolfato di Barite (*I.*).

This combination of barytes, which was formerly named *Ponderous spar*, is found native at Freyberg in Germany, and in many parts of the world. It is found almost always in veins; but sometimes it occurs in beds. It is occasionally obtained in powder, frequently in amorphous masses, and often in crystals.

*Qualities*.—Sulphate of barytes is inodorous and insipid. Its colour is white, with shades of yellow, red, blue, or brown. It occurs transparent, semitransparent, or only translucent; and is hard, brittle, and heavy, its specific gravity being from 4.484 to 4.500. The varieties of form of its crystals are numerous; but the primitive form is a rectangular prism, the basis of which are rhombs with angles of  $101^{\circ} 30'$  and  $78^{\circ} 30'$ .<sup>1</sup> The most common varieties of its crystals are the octahedron with oblique summits, the six or four-sided prism, the hexangular table with bevelled edges, sometimes

<sup>1</sup> Häüy, *Thomson's Chemistry*, iv. 369.



they are needle form.<sup>1</sup> It breaks with a straight foliated fracture; the fragments are nearly rhomboidal, and having a shining, pearly, almost vitreous lustre. It is fused by the blow-pipe, and converted into the sulphuret; and is soluble in sulphuric acid only, from which it is precipitated by water. The results of experiments to ascertain its constituents have been very discordant: according to Berzelius, they are 9.573 of barytes, and 5 of acid.

*Use.*—This barytic salt is introduced into the list of materia medica, merely as a substitute for preparing the muriate of barytes, when the carbonate cannot be procured.

BELLADONNÆ FOLIA. Vide *Atropa Belladonna*.

BENZOINUM. Vide *Styrax Benzoin*.

BISTORTÆ RADIX. Vide *Polygonum Bistorta*.

BISMUTHUM. Bismuth.

*Syn.* Etain gris, bismuth (*F.*), Wismuth (*G.*).

This metal is not very widely diffused. Its ores are found chiefly in Saxony; and, less abundantly, in Sweden, France, and Cornwall. It is usually accompanied by cobalt. It most commonly occurs in the state of metal, and, therefore, its ores are not much diversified. The following are the states in which Bismuth is found:—

A. In its metallic state:

i. Alloys.

*Sp. 1. Native Bismuth.*

B. United with other metals and sulphur:

ii. Sulphurets.

{ 2. Common Sulphuret.  
3. Needle Ore.  
4. Cupreous Sulphuret.

C. United with oxygen:

iii. Oxyde.

5. Bismuth ochre.

The ancient German miners regarded it as incomplete silver, or silver beginning to form, and termed it *tectum argenti*: and so late as the close of the 17th century it was considered a species of lead.

Bismuth was for some time considered as an alloy by chemists; but this opinion was gradually discovered to be erroneous; and it now ranks as a distinct metal.<sup>2</sup>

BISMUTHUM. *Lond.* Bismuth.

This metal appears as if formed of broad shining plates adhering to one another, of a reddish white colour; and is both insipid and inodorous. Its specific gravity is 9.822. It is not very brittle; and is rather softer than copper, but is not very malleable, and breaks when struck smartly by a ham-

<sup>1</sup> Thomson's *Chemistry*, 5th edit. iii. p. 413.

<sup>2</sup> The Greeks and Arabians were not aware of the existence of Bismuth; but it was very early distinguished by the Germans. It is mentioned as a peculiar metal by Agricola, in his treatise intitled *Bermannus*, written in 1529.



mer: it soon loses its lustre when exposed to the air, but remains unaltered under water. It cannot be drawn out into wire. It melts at  $476^{\circ}$  of Fahr., in which state its surface is oxydized in pellicles like lead; it evaporates in a higher temperature, and may be distilled over in close vessels. It is inflammable in a strong red heat, burning with a faint blue flame and emitting a yellow smoke, which condenses into an oxyde, insipid, unvolatilizable, insoluble in water. The oxyde when strongly heated melts and becomes darker coloured; and it may be sublimed at a heat below that which is required to fuse it. It is easily reduced when heated in conjunction with combustible bodies; the affinity between bismuth and oxygen being weak. According to Lagerhjelm this oxyde consists of 100 parts of Bismuth and 11.275 of oxygen: or, according to Mr. Phillips, of Bismuth 90 and oxygen 10 in 100 parts.<sup>1</sup>

Bismuth inflames in chlorine gas, and forms a chloride. It also combines readily with Iodine when heated, and forms an iodide of an orange-yellow colour, and soluble in pure potass. Bismuth does not combine with azote, hydrogen, carbon, boron, silicon, nor phosphorus. Bismuth in fusion readily combines with sulphur, and forms a sulphuret of a bluish grey colour, which is not unlike sulphuret of antimony. It crystallizes in tetrahedral crystals, which cross one another: and is very brittle and fusible. It is readily alloyed with other metals, forming with several of them compound metals of great fusibility.<sup>2</sup> Its constituents, according to Dr. John Davy, are 100 of Bismuth and 22.34 of sulphur, which nearly agrees with the analysis of Lagerhjelm, who makes them to be 100 of Bismuth and 22.52 of sulphur.<sup>3</sup>

Bismuth, in its metallic state, has no action on the animal economy. It is used merely for preparing the *subnitrate*; a salt which has long been employed, with great advantage, in cardialgia and similar affections of the stomach, but was not admitted into the Pharmacopœia, until the present edition of that work appeared.

Officinal preparation. — *Bismuthi subnitratis*, L.

## BITUMEN.

*Syn.* Bitume (*F.*), Erdharze (*G.*), Bitume (*I.*), Bitumen (*S.*).

In the limited signification of this term, it is meant to imply those mineral inflammable bodies, which resemble, in a certain degree, oily and resinous substances. They have been divided

<sup>1</sup> *Translation of the London Pharmacopœia*, 1824. p. 91.

<sup>2</sup> An alloy composed of 8 parts of Bismuth, 5 of lead, and 3 of tin, will melt in water below the temperature of  $212^{\circ}$ ; and is rather useful for taking casts and similar purposes.

<sup>3</sup> *Thomson's Chemistry*, 5th edit. vol. i. p. 459.



into two classes: the first containing bitumens, or, properly speaking, *bituminous oils*, which possess nearly the same properties as the essential oils; the second, *Bitumens*, strictly so called, which possess properties peculiar to themselves<sup>1</sup>; and a third class may be formed of those substances in which bitumen predominates with other components.

*A. Bituminous oils:*

- |               |  |   |
|---------------|--|---|
| <i>fluid.</i> | Sp. 1. Petroleum.                              | Var. <i>a. Naphtha.</i><br><i>b. Petroleum.</i> |
| <i>solid.</i> | { 2. Maltha, or Sea wax.<br>3. Mineral tallow. |   |

*B. Proper bitumens:*

- |                   |                            |
|-------------------|----------------------------|
| <i>solid.</i>     | 1. Asphaltum.              |
| <i>semifluid.</i> | 2. Mineral tar, or tallow. |
| <i>solid.</i>     | 3. Mineral Caoutchouc.     |

*C. Bituminous compounds:*

- |                                 |   |
|---------------------------------|---|
| 1. <i>with resinous matter.</i> | Sp. 1. Resin-asphaltum.   |
| 2. <i>with charcoal.</i>        | 2. Pit-coal. Var. <i>a. Brown coal.</i><br><i>b. Black coal.</i> <i>c. Glance coal.</i> |

According to Hatchett the elements of bitumens are carbon, hydrogen, sometimes azote, and probably oxygen.<sup>2</sup>

*Officinal.* PETROLEUM, *Lond.* BITUMEN PETROLEUM, *Edin.* PETROLEUM BARBADENSE, *Dub.* Petroleum. Barbadoes Tar.

*Syn.* Petrole (*F.*), Steinöhl (*G.*), Bergolja (*Swed.*) Petrolio (*I.*), Neft (*Arab.*), Mitti tel (*H.*), Munty lum (*Tam.*), Minnia tanna (*Malay*), Ippoo (*Sumatran*), Kesosonoabra (*Japanese*).

The officinal names imposed by the three British colleges are not to be regarded as synonymous of the same species of bitumen. The first species of the bituminous oil is properly named by the London college, the second variety of that species being the real Petroleum of the shops; but the Dublin college has incorrectly given the second species of the proper bitumens as the synonyme of Bitumen Petroleum.

Petroleum is found in many parts of the world, in various states of purity. When free from foreign ingredients, and before it has been long exposed to the action of the air, it is named *naphtha*; of which the purest kind that is brought to Europe comes from Monte Ciaro, near Piacenza in Italy. "This hill consists of horizontal beds of argillite, in which pits are sunk till the water comes in; after which the naphtha oozes out of the sides and floats on the surface of the water, whence it is skimmed off every week."<sup>3</sup> It usually flows from rocks of the coal formation. There is also a fountain of very pure Naphtha at the base of the mountains of Buck-

<sup>1</sup> Thomson's *Chemistry*, 5th edit. vol. ii. p. 384.

<sup>2</sup> *Linn. Trans.* iv. 129.

<sup>3</sup> *Mem. Sci.* 1756, p. 57, quoted by Aikin, *Diet. of Chem.* art. *Bitumen*.



tiavi, betwixt the city of Bruster and the valley of Ramhov-mouz. The less pure petroleum of the shops is procured from Monte Festino, near Modena. In the Birman empire there are 520 wells in one district, which yield annually more than 400,000 hogsheads of petroleum.

*Qualities.* — Naphtha is of a pale yellowish colour, thin, fluid, light, transparent, odoriferous, unctuous to the touch, volatile, and very inflammable. By long exposure to the air, it passes into the second variety *petroleum*; which is thicker than naphtha, unctuous to the feel, semi-transparent, of a reddish or blackish brown colour, and when long exposed to the air changes into true *bitumen*. It has a strong, penetrating, not disagreeable odour, and a bitter, pungent, acrid taste; and is not quite so inflammable as naphtha. Its sp. gr. varies from 0·730 to 0·873.<sup>1</sup> It is insoluble in water; but is soluble in alcohol, ether, the volatile and the fixed oils. When distilled with water, it comes over nearly as clear and fluid as naphtha. Both these varieties of bituminous oil combine with fat, resins, essential oil, and camphor; with alkalies they form soapy compounds; and sulphuric and nitric acids, change them into solid resins.

*Medical properties and uses.* — Petroleum is a stimulating antispasmodic and sudorific; and as such has been given in asthma and coughs unattended with inflammation; and in Germany has been extolled as a specific for tape worm; but it is chiefly used externally as a stimulant in diseases of the hip-joint, in rheumatic and other chronic pains, in chilblains, por-rigo, and to paralytic limbs, applied by friction.<sup>2</sup> It is, however, scarcely ever employed; and on this account is not often to be procured in the shops. The dose of petroleum may be from ℥x. to fʒss., given in any convenient vehicle.

**BOLETUS.** *Spec. Plant. Willd.*

*Cl.* 24. *Ord.* 13. Cryptogamia Fungi, *Nat. ord.* Fungi, *Linn. Juss.*  
*G.* — *Fungus* horizontal, porous beneath.

\* *Parasitic, stemless.*

*Spec.* 3. *B. Ignarius.* Agaric of the oak. *Sowerb. Fung. t.* 34.

*Officinal.* BOLETUS IGNARIUS. AGARICUS, *Edin.* Agaric.

*Syn.* Agaric de chêne (*F.*), Feuerschwamm (*G.*), Tontelgezwam (*Dutch*), Tyndersvamp (*Dan.*), Fröske (*Swed.*), Gubka (*Polish*), Esca o fungo preparata (*I.*), Agarico (*S.*), Boleto da isca (*Portug.*), Garikoon (*Arab. and Tam.*).

This species of fungus is found in Britain, growing upon decayed trunks of the ash and the oak. The pileus, or hat, is scaly and convex, but depressed in the centre. When young

<sup>1</sup> *Kirwan.*

<sup>2</sup> In the West Indies the Barbadoes tar is used as an internal remedy. Of the Burma Petroleum, Dr. Fleming remarks, "in chronic rheumatism I found much greater benefit from it than from the more costly Cajeput oil." — *Asiatic Researches.*



it is of a light brown colour above, and soft like velvet; white underneath, and covered with a slimy matter; but when mature it changes to dark brown, approaching to black. It is from six to ten inches in diameter; and although generally stemless, yet it is sometimes supported on a footstalk an inch in length.<sup>1</sup>

The boletus which grows upon the oak is said to be the most valuable. It should be gathered in August or September, and be kept in a dry room. "The way of preparing it is to take off with a knife the white and hard part, till you find a substance so soft as to yield under the finger like shammy leather."<sup>2</sup> This must be divided into different pieces, and these beaten with a hammer till they become so soft as to be torn with the finger.

*Qualities.*—Prepared agaric is inodorous, and has a slightly astringent taste. According to Bouillon la Grange, by whom it has been chemically examined, it contains resin, extractive, something similar to animal gelatin, and different salts.

*Medical properties and uses.*—Agaric has been much celebrated as a styptic, when externally applied to bleeding arteries and veins. It was introduced by Brossard, a French surgeon, in 1750, and was for some years used both on the continent and in this country: but if it really possess styptic powers greater than those of lint or sponge, which does not appear to be the case, the improved practice of surgery renders all such applications less necessary than they were when the use of agaric was introduced.

BONPLANDIA. *Plantæ Equinoctiales.* Tom. ii. p. 5.

Cl. 5. Ord. 1. Pentandria Monogynia. *Nat. ord.* Quassiæ, *Juss.*

*Gen. Char.* Calyx Monophyllous, campanulate, five-toothed.

*Corolla* five petals cohering near the base, funnel-shaped.

*Spect.* 1. *B. trifoliata.* Three-leaved Bonplandia. *Humboldt*, l. c. tab. 97. *Mem. de l'Institut.* 184. Part I. p. 82. pl. 10.

*Officinal.* CUSPARIÆ CORTEX, *Lond.* BONPLANDIÆ TRIFOLIATÆ CORTEX, *Edin.* ANGUSTURA; CORTEX, *Dub.* Cusparia bark, Bonplandia bark, or Angustura bark.

*Syn.* Angusture (*F.*), Angusturarinde (*G.*), Angustura (*I.*).

Although the London College has designated the tree which yields this bark, by the name of *Cusparia febrifuga*, as originally given to it by Humboldt, yet we have preferred the name imposed by Willdenow, and since adopted by Humboldt.

The *Bonplandia trifoliata* is a native of South America, growing abundantly in the woods, five or six leagues from the eastern bank of the Carony, at the foot of the hills that surround the missions of Capassui, Upata, and Alta Græcia.

<sup>1</sup> Withering, *Bot. Arrange.* ii. 767.

<sup>2</sup> *Phil. Trans.* xlix. Part. I. 29.



It grows also west of Cumana, in the gulf of Santa Fe; and, as Humboldt remarks, may become an article of export from New Andalusia. It is an elegant evergreen, rising to the height of from sixty to eighty feet, having a cylindrical *trunk*, covered with a grey-coloured bark, and branching towards the summit. The branches are alternate, the upper ones spreading nearly horizontally. The *leaves*, which are ranged alternately on the branches, are about two feet long, independent of the petiole, and composed of three oblong ovate leaflets, pointed at each extremity, and attached at their bases to a single channelled petiole, from ten to twelve inches in length. The *leaflets* are glandular, and, when fresh, exhale an agreeable aromatic odour. The *inflorescence* is a terminal raceme, composed of alternate peduncles, bearing from three to six flowers each: the calyx is inferior, persistent, five-toothed, and tomentose; the corolla is funnel-shaped, and composed of five petals, so united below as to appear as one tube, with a five-cleft spreading tube. The *nectary* consists of fine glandular bodies. The *stamens*, which are shorter than the petals, have white filaments supporting oblong yellow anthers; the *pistil* is formed of five, oval, hairy ovaries, from the centre of which a single style rises, supporting five, fleshy, green stigmata. The *fruit* consists of five, oval, bivalve capsules, each enclosing a single seed. The first parcels of Cusparia bark were imported from Dominica in 1778, and the tree yielding it was supposed to be a native of Africa<sup>1</sup>; but importations from Cadiz and the Havannah, and the travels of Humboldt and Bonpland, have led to the knowledge of the real place of its growth. It is brought to this country packed in casks; but the original package, Mr. Brande informs us, is curiously formed of the large leaves of a species of palm, surrounded by a kind of network made of sticks. It is in pieces of different lengths, some nearly flat, and others in partial quills of all sizes intermixed.

*Qualities.*—The odour of this bark is not strong, but peculiar; the taste bitter, slightly aromatic, and permanent, leaving a sense of heat and pungency in the throat. The pieces are covered with a whitish wrinkled thin epidermis; the inner surface is smooth, of a brownish yellow colour, and the intermediate substance mottled fawn colour, and of a compact texture. It breaks with a close, short, resinous fracture; is easily pulverized, and affords a powder which, when triturated with lime or calcined magnesia, gives a smell of ammonia.—The ac-

---

<sup>1</sup> See Brande's *Experiments and Observations on the Angustura Bark*, a name which it received only because it came from Nueva Guayana, or Angustura.



tive matter is taken up by cold and hot water in infusion, and is not injured even by coction, but the addition of alcohol precipitates part of the extractive. The alcoholic tincture reddens litmus paper, and becomes milky on the addition of water. The watery infusion precipitates the infusion of galls, and of yellow cinchona, but not gelatine.<sup>1</sup> I found that it precipitates sulphate of iron, tartarized antimony, sulphate of copper, acetate and superacetate of lead, oxymuriate of mercury, and pure potass yellow; which confirm Vauquelin's analysis. Nitrate of silver also precipitates it yellow, but assumes a violet colour after some time. Ammonia deepens the colour, but is not precipitated. Sulphuric acid gives the infusion a brown colour, and gradually a lemon-yellow precipitate is deposited; whilst nitric acid deepens the colour to a blood red, and after some time affords a lemon-yellow precipitate. The muriatic acid does not affect it. Sulphuric ether takes up one part from ten of the powder, and when evaporated on water, leaves a greenish yellow, very acrid resin, and renders the water milky: the addition of nitro-muriatic acid changes this milky appearance to red, slowly producing a lemon-yellow coloured precipitate, and giving the resin on the side of the glass a brownish pink colour. By distillation with water, the bark yields a small portion of a white essential oil. These experiments ascertain the substances which are incompatible in prescriptions with infusion, or tincture of cusparia bark; and show that it contains *resin*, a peculiar variety of *extractive*, *carbonate of ammonia*, and *essential oil*.<sup>2</sup>

*Medical properties and uses.*—Cusparia bark is stimulant and tonic. It was introduced in the West Indies with very high pretensions; and, although it is not superior or even equal to Cinchona bark in intermittent fevers, yet it is a remedy possessed of very considerable powers. It does not oppress the stomach, but gives to it a degree of warmth, expels flatus, keeps the bowels open, and increases the appetite for food. It is particularly efficacious in bilious diarrhœa and dysentery, after due evacuations; and also proves useful in dyspepsia,

<sup>1</sup> Vauquelin. *Annals de Chimie*, lix. 150.

<sup>2</sup> A species of bark, in some respects resembling the Cusparia, has lately been introduced upon the Continent, possessing the most deleterious quality. Planche has examined it, and named it *Angustura ferruginea*. It is readily distinguished from the true bark, by its greater thickness and weight, and the epidermis being of a brownish, olive hue, and warty. It impresses also the most nauseous and permanent bitter when chewed. By agitating the powder in very dilute muriatic acid, it assumes a beautiful green, owing to the iron contained in the cuticle of this bark. The narcotic, deleterious matter has been ascertained to be an alkali, which has been named *Brucine*. For particulars regarding its poisonous properties, vide *Orfila's Traité des Poisons*, tom. ii. p. 351., and *The Lond. Med. Repository*; and for the characters of *Brucine*, vide *Ann. de Chim. et Phys.* xii. 115.



hysteria, leucorrhœa, and most of the diseases in which the use of a general tonic is indicated. Mr. Brande published several cases which came under his own observation, and some from the communications of others, in which its usefulness, as a remedy for intermittents, appears to be confirmed: but this is disputed, particularly by Alibert, who gave it a fair trial in the hospital of St. Louis. My own experience does not enable me to give an opinion on the subject. Its employment is contra-indicated in directly inflammatory complaints, in hectic fever, and colliquative diarrhœa.

It may be exhibited in substance, in watery infusion, in tincture, and in the form of watery extract. The powdered bark is given in doses of from grs. v. to grs. xx., beyond which it is apt to induce nausea. It may be combined with rhubarb, neutral salts, magnesia, and testaceous medicines, or with powdered cinnamon, which covers its nauseous taste better than any other adjunct. Of the aqueous extract, grs. x. is a full dose. In large doses all the forms are apt to excite nausea.

Officinal preparations. *Infusum Cuspariæ*, L. *Tinctura Bonplandiæ trifoliatæ*, E. *Tinctura Angusturæ*, D.

**BOSWELLIA.** *Roxburgh.*

*Cl.* 10. *Ord.* 1. Decandria Monogynia. *Linn.*

*Gen. Char.* *Cal.* beneath, five toothed. *Cor.* five petals. *Nect.* a crenulated fleshy cup, surrounding the lower part of the germ, with stamens inserted on its outside. *Capsule* three-sided, three valved, three-celled. *Seeds* solitary, membranous, winged.

*Species.* — *B. Serrata.* *Asiatic Researches*, 8vo. vol. ix. p. 377.

*Officinal.* OLIBANUM<sup>1</sup>, *Lond.* JUNIPERI LYCIÆ GUMMI-RESINA, *Edin.* OLIBANUM; GUMMI-RESINA, *Dub.* Olibanum.

*Syn.* Encens (*F.*), Weirauch (*G.*), Olibano (*I.*), Paringhi Sambrani (*Tam.*), Looban (*Malay.*), Koondir Zuckir (*Hind.*), Cundur (*Arab.*), Labúniyá (*Syr.*).

Olibanum was supposed, on the authority of Linnæus, to be the production of the *Juniperus Lycia*<sup>2</sup>: but this opinion appears to be erroneous; for, as Mr. Colebrooke has observed, “this species of juniper is a native of the south of France;” and the French botanists deny that it yields the resinous gum in question.<sup>3</sup> On this account, therefore, and influenced by other proofs brought forward by Mr. Colebrooke, we have been induced to regard olibanum, at least that brought from India, as the production of the *Boswellia serrata* of Roxburgh, although it is still referred to the *juniperus Lysia* in the British pharmacopœias.<sup>4</sup> Lamark supposes that the Arabian oliba-

<sup>1</sup> Quasi Oleum Libani. Colebrooke, *Asiatic Researches*, 8vo. vol. ix. p. 382.

<sup>2</sup> Λιβανός, Dioscoridis.

<sup>3</sup> *Asiatic Researches*, vol. ix. p. 377.

<sup>4</sup> Although so much used in the early ages as incense in sacrifices, and, latterly, in the ceremonies of the Greek and Roman churches, yet, both ancient and modern writers have differed regarding the plant yielding it.



num is the production of the *Amyris Gileadensis*; but his reasons are not very conclusive.

The *Boswellia serrata* is a native of the mountains of India, and is vulgarly known under the name of *Sálaï*. It is a large tree, with the foliage crowded at the extremities of the branches. The leaves are impari-pinnate, consisting of ten pairs of leaflets, sessile, each an inch, or an inch and a half in length, obliquely ovate, oblong, obtuse, serrate, and villous, supported by round downy petioles. The flowers, which are produced in axillary racemes, shorter than the leaves, are numerous, small, and of a pale pink colour, accompanied with minute bracteas. The calyx is monophyllous, five-toothed, and downy: the corolla composed of five oblong, spreading, exteriorly downy petals; and the nectary a fleshy, crenulate, coloured cup, adhering to the calyx. The stamens are ten, alternately shorter, supporting oblong anthers: the pistillum consisting of an ovate germen, a cylindrical style, and trilobate stigma. The capsule is smooth, three-sided, trilocular, three-celled, and three-valved: each cell containing one perfect seed only, which is broad, cordate, and winged.

Olibanum is imported in chests and casks from the Levant, and is also sold at the East India Company's sales; but the Indian olibanum is not much esteemed.

*Qualities.*—Olibanum is a translucent, whitish-yellow, brittle substance, generally covered with whitish powder, produced by the friction of the pieces against each other. Its odour, when burnt, is fragrant; its taste is acrid, bitterish, and somewhat aromatic. When heated, although it melts with difficulty, yet it burns brilliantly, and leaves a whitish ash, composed of phosphate, carbonate, and sulphate of lime, with muriate and carbonate of potass. When distilled alone it affords a volatile oil; but, when in conjunction with water or alcohol, no oil comes over. Alcohol dissolves three-fourths of it, forming a transparent solution; and when triturated with water a milky solution is produced, from which the resinous matter is deposited after some time, and three-eighths only remain dissolved. Ether takes up rather more than one half, and when evaporated on water leaves a very pure transparent resin; while the part undissolved by it becomes white and opaque, and is almost entirely soluble in water, forming a milky solution. Hence, olibanum appears to consist of resin, gum, and a volatile oil. This opinion has been lately confirmed by Braconnot, who has analysed olibanum, and found in 100 parts of it, 8 of fragrant volatile oil, 56 of resin, 30 of gum, and 5.2 of a matter resembling gum, but insoluble in water and alcohol. The oil resembled the oil of lemons in odour and colour.



*Medical properties and uses.* — Olibanum is stimulant and diaphoretic. It was formerly much used in gleet and affections of the chest; and, externally, as a vulnerary; but it is now employed only as a perfume in sick rooms.

**BUBON.**<sup>1</sup> *Spec. Plant. Willd.* i. 1439.

*Cl.* 5. *Ord.* 2. Pentandria Digynia. *Nat. Ord.* Umbellatæ.

*G.* 546. *Fruit* ovate, striated, villose.

*Species* 2. *B. Galbanum*.<sup>2</sup> Lovage-leaved Bubon. *Med. Bot.* 2d. edit. 98. t. 40.

*Officinal.* GALBANI GUMMI RESINA, *Lond. Dub.* BUBONIS GALBANI GUMMI RESINA, *Edin.* Galbanum Gum-resin.

*Syn.* Galbanum (*F.* and *Dutch*), Mutterharz (*G.*), Galbano (*I.*), Galbâne (*S.*), Bārzud (*A.*), Bireejā (*H.*), Beeznd (*Pers.*).

This species of Bubon is a perennial plant, a native of Africa about the Cape of Good Hope, and of Syria, flowering in June and July. It rises eight or ten feet in height, with the stem ligneous at the base, and there furnished with a purplish bark; but the upper part is jointed, branching, leafy, and covered with a glaucous exudation, which can be easily wiped off. The lower leaves are nearly tripinnate, on vaginant footstalks; the uppermost are almost simple, tri-lobed, thickish, irregularly serrated, and of a greyish colour. The flowers are in terminal umbels, the principal umbel being large, and plano-convex. The involucre is composed of twelve narrow, lancet-shaped, membranous, whitish leaflets, bent downwards; but the involucre of only six leaflets. The flowers are all fertile; the petals yellow, with inflected tips: the stamens are longer than the petals, supporting yellow anthers; and the styles two, short and tapering. Each flower is succeeded by two oblong-channelled seeds, having a thin membranous border.

When the stem of the growing plant is broken, or wounded by a knife, a cream-coloured juice flows out; and in this manner the gum-resin is procured, by making an incision, or cutting the stem across a few inches above the root; when it soon concretes, and is fit to be gathered. A small quantity exudes spontaneously from the joints of the stem. The gum-resin is brought to this country from the Levant, in cases or chests, containing from one to three hundred weight each. The best is in ductile masses, composed of distinct whitish tears agglutinated together by a pale brown or yellowish substance. It is generally much mixed with stalks, seeds, and other impurities. The separate tears are considered to be the best part of the mass. When the colour is dark brown or blackish, it must be rejected as bad.

<sup>1</sup> Βουβωνιον, Dioscoridis.

<sup>2</sup> Χαλβάνη, Hippocratis.



*Qualities.* — Galbanum has a strong peculiar odour, slightly resembling that of turpentine; and a bitterish, warm, acrid taste. Its specific gravity is 1.212.<sup>1</sup> When triturated with water, about one-fourth of its weight is dissolved, forming a milky solution; but after standing for a little time, four parts are again deposited, and what remains undissolved by the trituration is, exclusive of the impurities, almost completely soluble in alcohol. Wine and vinegar act on it nearly in the same manner as water. Alcohol takes up one-fifth of its weight; and a yellow tincture is produced, which has the sensible qualities of the galbanum, and becomes milky on the addition of water; but there is no precipitate. Proof spirit acts slowly on it, and dissolves the whole, the impurities excepted. Sulphuric æther dissolves a considerable portion of galbanum, forming a bright golden-coloured tincture, which, when evaporated alone, or floating on the surface of water, leaves a yellow, tenacious resin, that retains in perfection the sensible qualities of the galbanum. The part insoluble in æther is nearly wholly soluble in water. Oxymuriatic acid added to the solutions of galbanum, throws down an insoluble matter which appears to be oxidized extractive. By distillation the gum-resin “yields half its weight of volatile oil, which has at first a blue colour.”<sup>2</sup> From our experiments, galbanum appears to consist of resin, volatile oil, gum, and extractive.

*Medical properties and uses.* — Galbanum is antispasmodic, expectorant, and deobstruent; and may be placed between ammonia and assafoetida. It has been found useful in hysteria, particularly when attendant on difficult menstruation; and in chlorosis. Externally it is applied as a resolvent and a stimulating suppurative to indolent tumors.

The dose is from grs. x. to ℥j. in pills; or triturated with water and gum arabic so as to form an emulsion.

Officinal preparations. *Pilulæ Galbani comp.* L. *Pilulæ Assafoetidæ comp.* E. *Pilulæ Myrrhæ comp.* D. *Tinctura Galbani*, D. *Emplastrum Galbani*, D. *Emplast. Galbani compositum*, L. *Emplastrum Assafoetidæ*, E. *Emplastrum Gummosum*, E.

CAJUPUTI OLEUM. Vide *Melaleuca Cajuputi*.

CALAMI RADIX. Vide *Acorus Calamus*.

CALAMINA. Calamine. See *Zincum*.

CALUMBÆ RADIX. Vide *Menispermum*.

CALX. *Edin.* Lime.

*Syn.* Chaux (*F.*), Kalk (*G.*), Calce (*I.*), Calviva (*S.*), Chunāmboo (*Tam.*), Capoor (*Malay*), Noorch (*Pers.*), Chunna (*H.*), Ahúck (*A.*), Kakote-tung-ō-ă (*Esquimaux*).

<sup>1</sup> Brisson.

<sup>2</sup> Thomson's *Chemistry*, 4th edit. v. 142.



This earth is very rarely found in an uncombined state<sup>1</sup>; but very abundantly in combination with other substances. It forms a part of the bodies of animals and of vegetables; exists in the water of most rivers, and of the ocean; and is a principal constituent of many fossils, soils, and mountains. The following species only of the fossils in which it is found in combination with carbonic acid require to be noticed.

It is nearly pure in :

Sp. 1. *Chalk*.

2. *Limestone*. Subsp. 1. *Compact Limestone*.

Var. *a*. Common. *b*. Roe-stone.

3. *Foliated limestone*. Var. *a*. Granular foliated, or statuary marble. *b*. Calcareous spar.

4. *Fibrous limestone*. Var. *a*. Common fibrous, or satin spar. *b*. Calcsinter, or Stalactite.

5. *Pea stone*.

By exposing any of these carbonates to a strong heat the carbonic acid is driven off, and lime, or quicklime, as it is commonly called, is obtained; the properties of which shall be noticed under the title *Calx*, among the preparations. It is not, however, perfectly pure, but contains generally portions of silex, argil, or magnesia. To obtain pure lime, let white marble be dissolved in dilute muriatic acid, leaving an excess of marble undissolved. A solution of pure ammonia being added to the solution of marble will indicate by a precipitate the presence of argil and magnesia, which are to be separated by filtration; and the lime itself precipitated in the form of carbonate by a solution of pure sub-carbonate of potass. This precipitate, after it is washed with water and dried, and exposed to a very violent heat in a platinum crucible, is pure lime.<sup>2</sup> It is a compound of a peculiar metallic base, named *calcium* by Sir H. Davy, and oxygen, in unknown proportions.<sup>3</sup> We have here to notice the calcareous fossils only which are medicinally used.

#### 1. CHALK. *Friable Carbonate of Lime*.

*Officinal*. CRETA, *Lond. Dub.* CARBONAS CALCIS. *a*. MOLLIOR, *Creta alba*. *Edin.* Chalk.

*Syn.* Craie (*F.*), Kreide (*G.*), Wite kryt (*Dutch*), Kride (*Danish*), Kalabastrace (*Swed.*), Mjel (*Russian*), Gasso (*I.*), Greda (*S. and Portug.*), Khurree muttee (*H.*), Kilo (*A.*), Sirnic Chunamboo (*Tam.*), Gal sifid (*Pers.*), Capoor-engrees (*Malay*).

This mineral is found in the north of France, Poland, some of the Danish islands, and in great abundance in the south of England, within a range which commences at Flamborough Head, in Yorkshire, and is continued, with irregular interrup-

<sup>1</sup> Monnet affirms that it exists in the mountains of Upper Auvergne, mixed, however, with a little oxide of iron. *Monnet's Mineralogy*, 515.

<sup>2</sup> Chenivix, *Memoirs of the Irish Academy*, 1802.

<sup>3</sup> *Phil. Trans.* 1808.



tions, through the midland counties, to Surrey, Sussex, Hampshire, and into Dorsetshire. It occurs massive in beds; and contains numerous relics of land and marine animals.

*Qualities.* — Chalk is inodorous and insipid; but adheres slightly to the tongue. Its colour is either white, or yellowish, or grayish white. It feels meagre and rough: is not very hard, but is pulverulent; breaks with an earthy fracture; stains the fingers, and marks. Its specific gravity is from 2.3 to 2.6. It effervesces with acids; and generally contains a small portion of argil. The average proportion of lime is 53 per cent.

*Medical properties and uses.* — Chalk is antacid; but it must undergo levigation and washing, before it can be internally administered. In powder it is externally advantageously employed as an absorbent in burns and excoriations.

Officinal preparations. *Creta preparata*, L. E. D.

## 2. LIMESTONE. *Hard carbonate of Lime.*

*Officinal.* LAPIS CALCAREUS, *Lond.* CARBONAS CALCIS. *b.* DURIOR, *Marmor album*, *Edin.* Limestone. White Marble.

*Syn.* Pierre à chaux; chaux carbonatée (*F.*), Kalkstein (*G.*), Marino; ossicarbonato di calce (*I.*), Chunambo kullo (Tam.), Kakote-tung-ō-ā (*Esquimaux*).

Although all the varieties of limestone may be regarded as officinal, yet the two varieties particularly designated are *var. a.* of the first subspecies, *common compact limestone*, and *a.* of the second subspecies, *granular foliated limestone*, or *white Carrara marble*. The first is found abundantly in Britain, in extensive strata connected with floetz and coal formations; the second is brought from Carrara and Paros, and belongs exclusively to the primitive and transitive mountains.

*Qualities.* — *Common limestone* is inodorous and insipid: of a grey colour, sometimes variegated with veins, stripes, and clouds of yellow, flesh red, and greenish grey. It is hard and brittle; the fracture splintery; the fragments sharp-edged, and scarcely translucent. Its specific gravity is from 2.6 to 2.7. *White marble* has a granular texture, white colour, and foliated fracture. Its specific gravity is from 2.7 to 2.84. Both varieties dissolve in acids with effervescence; and contain about 65 per cent. of lime.

*Use.* — Limestone is chiefly used for obtaining pure lime.

CAMBOGIA. Vide *Stalagmatis Cambogioides*.

CAMPORA. Vide *Laurus Camphora*.

CANCER. *Syst. Nat. Gmelin*, 2963.

*Ord.* 7. Insecta, Aptera.

*G.* 270. *Feet* eight, (sometimes six or ten,) two of them with claws.

*Palpi* six, nearly equal. *Eyes* two, distant, moveable, in many of the species standing on elongated peduncles. *Mandible* horny, thick. *Lip* triple. *Tail* jointed and unarmed.

*Sp.* 27. *C. pagurus*. Black-clawed Crab. *Brit. Zoology*, iv. 4. t. 3.

*Sp.* 63. *C. ostacus*. The Crawfish. *Brit. Zoology*, iv. 9. t. 15. f. 27.



## 1. CANCER PAGURUS.

*Officinal.* CHELÆ CANCRORUM, *Edin.* CANCER; CALCULI OCULI DICTI<sup>1</sup>; CHELÆ, *Dub.* Crab's Claws.

*Syn.* Patte d'Ecrevisse (*F.*), Klaua an Krabbe (*G.*), Forbici di Granchi (*I.*), Pierna de Cangrejos (*S.*), Kreeftschaaren (*Dutch*), Kräftkler (*Swed.*).

The black-clawed crab frequents the rocky coasts of the North Sea, and the British isles; and is considered delicious food. The thorax is obtusely scalloped; the body smooth; and the front five-toothed. The hind feet are subulate; but the fore are furnished with large claws tipped with black. It annually casts its shell, between Christmas and Easter.

Mr. Hatchett found that the crustaceous covering of crabs and lobsters consists of carbonate of lime, phosphate of lime, and a cartilaginous matter, possessing the properties of coagulated albumen. The first of these constituents predominates; and it is on it that the medical properties of the claws depend. They are now deservedly rejected by every judicious practitioner, chalk answering, much better, every purpose for which they can be prescribed.

## 2. CANCER ASTACUS.

*Officinal.* LAPILLI CANCRORUM, *Edin.* Crab Stones.

*Syn.* Yeux d'Ecrevisse (*F.*), Krabsangen Krebssteine (*G.*), Kraftsteene (*Dutch*), Kraftstenar (*Swed.*), Ojos de Cangrejos (*S.*).

The crawfish frequents rivers, forming its holes in their clayey banks. It is small, and in some degree, resembles the lobster in shape. The snout is projecting, and serrated on the sides; the thorax is smooth; as is also the back, which has two small spines on each side. The large claws are beset with small tubercles: the two first pairs of legs are clawed; the two next subulated; and the tail has five joints, with rounded fins.

The concretions, called eyes, are found in the stomach, one on each side, before the fish casts its shell in July, at which time the inner coat of the stomach also is renewed. They are said to be destined for assisting in the formation of the new shell. At Astracan, where the greatest number of these concretions are procured, the crawfish are bruised with mallets, and allowed to putrefy in heaps; after which their remains are washed, and the stones picked out.

*Qualities.* — Crabstones are whitish, or reddish, hard and stony, of very different sizes, weighing from one grain to twelve grains each; round and convex on one side, and a little concave on the other: the texture is laminated; inodorous and insipid. Their constituents are the same as those of the crab's claws. They effervesce in acids; but instead of dissolving altogether they become soft, transparent, and retain their original

<sup>1</sup> This is an error of the Dublin College, as these calculi are never procured, at least for medical use, from the crab, but always from the crawfish.



form; by which means the real stones are easily distinguished from counterfeited imitations.

*Medical properties and uses.*—These concretions are absorbent and slightly antacid: and, when prepared by trituration and levigation, are employed in dyspepsia and other diseases attended with acidity of the *primæ viæ*; but as chalk answers better in these cases, they may well be dispensed with.

The dose is ʒj or ʒij, suspended in a proper fluid.

Official preparations. *Cancrorum Lapilli præparati*, E.

CANELLA. *Spec. Plant. Willd.* ii. 857.

Cl. 11. Ord. 1. Dodecandria Monogynia. *Nat. ord.* Oleraceæ, *Linn.* Meliaceæ, *Juss.*

G. 942. *Cal.* Three-lobed. *Pet.* five. *Anthers* 16, adhering to a pitcher-shaped nectary. *Berry* one-celled, with two or four seeds. *Species* 1. *C. alba*.<sup>1</sup> White or Laurel-leaved Canella. *Med. Bot.* 2d edit. 694. t. 237. *Trans. Linn. Soc.* vol. i. 96. t. 8.

*Officinal.* CANELLÆ CORTEX, *Lond.* CANELLÆ ALBÆ, CORTEX, *Edin.* CANELLA ALBA, *Dub.* Canella Bark.

*Syn.* Cannelle blanche (*F.*), Weisser Zimmet (*G.*), Hwit Kanel (*Swed.*), Cannella bianca (*I.*), Canella blanca (*S.*).

This tree is a native of the West India islands, growing in the inland woods. It rises very straight, from ten to fifty feet in height. The branches are erect, not spreading, and only at the top of the tree; furnished with petiolated leaves, irregularly alternate, oblong, obtuse, entire, nerveless, of a dark green colour, thick and shining like those of the laurel, and emitting a similar odour. The flowers, which exhale a powerful aromatic perfume, are small and of a violet colour, seldom opening, and grow in clusters upon divided footstalks at the summits of the branches. The calyx is of one piece, small, persistent, and deeply tripartite; the petals are five times as long as the calyx, oblong, sessile, concave, erect, two a little narrower than the others; the nectary is pitcher-shaped, antheriferous, and deciduous. The anthers are twenty-one in number, distinct, fixed longitudinally to the outside of the nectary, and discharge a yellow pollen. The germen is superior, ovate; the style cylindrical, with two rough, convex, blunt stigmas. The fruit is an oblong, one-celled, glossy black berry.

The inner bark of the branches is freed from the cuticle, and dried in the shade. It is brought to this country packed in casks and cases, in long pieces, some rolled in quills, and

---

<sup>1</sup> This plant has been often confounded with the *Wintera aromatica*, an error authorized in some degree by Linnæus, who combined the two genera of *Winterana* and *Canella* under the name of *Laurus Winterana*; but afterwards made this a distinct genus under the title *Winterania*, a name by which it was known till Professor Murray corrected the error and made a distinct genus of *Canella*. Vide *Syst. Veg.* 14th edit. 443. Sir Hans Sloane stated the error of confounding this bark with the *Cortex Winteranus*, in his description of the tree in the *Phil. Trans.* xvii. 465.



others flat; the quilled sort is considerably thicker than cinnamon, and the flat nearly one-fourth of an inch in thickness.

*Qualities*. — The quilled pieces of *Canella* are of a whitish yellow colour on both sides, and break with a starchy fracture; the flat pieces, which appear to be the bark of the largest branches or of the stem, are yellow on the outside, and pale brown within. The odour of both kinds, when fresh broken, is aromatic, something like a mixture of cloves and cinnamon: and the taste slightly bitter, extremely warm, and pungent. Although boiling water takes up nearly one-fourth of the weight of the bark, yet the infusion possesses but little of its warmth and pungency; the bitter chiefly predominating. Alcohol extracts all its qualities in perfection: the tincture is bright yellow, and becomes milky on the addition of water. The infusion is not altered by infusion of galls, sulphate of iron, zinc, muriate of mercury, nor tartarized antimony; but nitrate of silver and acetate of lead render it milky, and throw down precipitates. By distillation with water, *Canella alba* affords a thick, heavy, yellow, very pungent, gratefully odorous essential oil; on which and a little bitter resinous matter its virtues seem to depend.

*Medical properties and uses*. — This bark is stimulant, and slightly tonic. It is a useful adjunct to bitters in some cases of dyspepsia and atonic gout; but it is employed chiefly on account of its flavour, and to correct the griping quality of the resinous cathartics. It is said to prove useful in scurvy.<sup>1</sup>

The dose of the powdered bark is from grs. x. to ʒss.

Officinal preparations. *Tinctura Gentianæ composita*, E. *Vinum Aloes*, L. E. D. *Pulvis Aloes cum canella*, D.

**CANTHARIS.** *Latreille. Gen. Crustaceorum et Insectorum.*

*Cl. 5. Ord. 1. Insecta Coleoptera, Linn. Eleuterata, Fabric.*

*G. 215. Feelers* filiform. *Palpi* four, unequal; the posterior ones clubbed. *Thorax* nearly round. *Head* inflected, gibbous. *Elytra* soft, flexible.

*Species 1. Cantharis vesicatoria.* Blistering Fly. *Latreille, tom. 2. p. 220.*

*Officinal. CANTHARIS, Lond. CANTHARIS VESICATORIA, Edin.*

*CANTHARIS, Dub.* Blistering or Spanish Fly. *Cantharides.*

*Syn.* *Cantharides* (F.), *Spanische Fliegenoder Kanthariden* (G.), *Spaansche Vliegen* (Dutch), *Spanka* (Swed.), *Machy Hisapanskie* (Pol.), *Cantarelle* (I.), *Cantharidas* (S.).

This insect is found on the privet, ash, elder, lilac, white

<sup>1</sup> This bark and the fruit of the capsicum were formerly common ingredients in the food and drink of the Caraihs, the antient natives of the Antilles; and at present enter the meagre pot of the negroes. — *Linn. Trans.* l. c.



poplar, and the tartarian honeysuckle, in Spain, Italy, France, and to a certain extent over the greater part of Europe. They are two-thirds of an inch in length, and one-fourth of an inch in breadth, oblong, and of a green, gold-shining colour; with soft elytra or wing sheaths, marked with three longitudinal raised stripes, and covering brown membranous wings. The body is terminated by two small, callous, sharp spines, and on the head are two black, jointed feelers. When alive they have a foetid odour.<sup>1</sup> They are gathered by smoking with brimstone the trees on which they are found, and catching them on a cloth spread underneath. They are sometimes simply shaken from the trees, and then killed by the steams of boiling vinegar, and dried either by the sun or in a stove.

Blistering flies are imported from Sicily, but chiefly from Astracan, packed in casks and small chests. The best are of a lively, fresh colour, a small size, and not mouldy, nor mixed with the *Melolontha vitis*, an insect resembling them in some degree, but possessing no vesicating property. It may be distinguished by its form, which is altogether more square than that of the *Cantharis*, and by its black feet.<sup>2</sup> If the blistering flies have been properly dried, and are kept in a well-stopped glass bottle, they will remain unchanged in appearance, and retain their acrimony for a great length of time<sup>3</sup>; but sometimes, in spite of every precaution, they are attacked by a small worm, which, however, feeds on the inactive part only of the fly, reducing it to a powder, that still possesses the active quality of the entire insect. They soon putrefy when kept in a damp place, and therefore should be occasionally spread out to the air.

*Qualities.* — Blistering flies have a heavy disagreeable odour, and an acrid taste. Lewis found that their active constituents are soluble both in water and in alcohol, and that the residuum is inert. Thouvenel, Beaupoil, and Robiquet, have analysed the insect; but their inquiries lead to no very certain conclusions.

Thouvenel treated the entire flies with *water*, *alcohol*, and *ether*, separately, submitting them to the press; and obtained

<sup>1</sup> It is asserted that a person who sits under a tree on which many of these insects are, particularly at the time of copulation, experiences ardor urinæ, pain of the bladder, and sometimes ophthalmia.

<sup>2</sup> Fabricius thus describes the *Melolontha*; "*Maxilla brevis cornea; apice multidentata. Antennæ lamellatæ. Melolontha vitis. Viridis, thoracis lateribus flavis, pedes nigri.*" Vide *Ræmer Gen. Insect.* t. 1. fig. 11.

<sup>3</sup> Van Swieten kept them upwards of thirty years in a glass vessel, not particularly well corked, and they still produced vesication.



the following results: 1st, Three-eighths of reddish yellow, very bitter extractive, affording by distillation an acid liquor: 2d, One-tenth of concrete, waxy, green oil, having the odour of the flies, and yielding by distillation a very sharp acid and a thick oil: 3d, One-fiftieth of concrete yellow oil, apparently the colouring matter of the insect; and, 4th, One-half of solid parenchymatous matter. He imagines that the blistering principle resides in the green waxy oil, and that the strangury produced by blisters is the effect of the acid obtained from this oil by distillation.<sup>1</sup>

Beaupoil found that an aqueous infusion of the flies, when exposed to the air, lets fall a yellow precipitate, exhales an ammoniacal odour, and reddens tincture of turnsole: the addition of ether or alcohol divides it into two parts; viz. a black gluey matter, insoluble in alcohol, and a yellowish-brown, very soluble matter.<sup>2</sup> The black matter blistered the skin without affecting the urinary organs; the yellow matter did not blister when applied alone, but blistered quickly when united with wax; and a green matter, which he also obtained, acted under similar circumstances, but less actively.

Robiquet asserts, that the flies, when recently collected, yield some uric acid. By treating them with water, alcohol, and ether, he obtained a peculiar matter, in the form of small, crystalline, micaceous plates, insoluble in water and in cold alcohol, but soluble in boiling alcohol, in ether, and in oils; on the presence of which the vesicatory property of the flies depend, and which, in combination with oil, might supersede their use. Dr. Thomson<sup>3</sup> has named it *Cantharidin*.<sup>4</sup>

*Medical properties and uses.* — Blistering flies, internally exhibited, are powerfully stimulant and diuretic; and, externally applied, rubefacient and epispastic. Notwithstanding their acrimony, they appear to have been given as an internal remedy by Hippocrates, who prescribed them chiefly in cases of dropsy and amenorrhœa.<sup>5</sup> They have a considerable effect

<sup>1</sup> *Annales de Chimie*, xlvii. 280.

<sup>2</sup> From one ounce of cantharides he obtained, Of black matter, 2 gros. 2 grs.; yellow matter, 1—2; green matter, 1—8; parenchyma, 4—36; phosphate of lime, 12 grains; carbonate of lime, 2 grains; sulphate and muriate of lime, 4 grains; oxide of iron, 2 grains; and an acid, the quantity of which was not ascertained. *Annales de Chimie*, xlviii. 35.

<sup>3</sup> Vide *System of Chemistry*, 5th edit. iv. p. 436., and *Ann. de Chim.* lxxvi. p. 308.

<sup>4</sup> Dioscorides and Galen imagined that the active principle of the fly was contained in its body, and that the head, wings, and feet, contained its antidote.

<sup>5</sup> Dr. Groenvelt was prosecuted for using them internally, and published his tract, "*de tuto Cantharidum uso interno*," as his vindication; but, although it proved to his prosecutors the safety of his practice, yet (says Quincy, *Pharm.* p. 152.) it ruined the unhappy doctor.



on the urinary organs, even when externally applied; and unless their internal exhibition be conducted with great caution, they act with so much violence on the kidneys, bladder, and small intestines, as to produce bloody urine, purulent stools, insupportable pains of the abdomen, vomiting, and other symptoms of intestinal inflammation; convulsions, delirium, syncope, and death. They have, however, been successfully employed in dropsy, obstinate gleet,<sup>1</sup> leucorrhœa, and incontinence of urine arising from paralysis of the sphincter vesicæ. The free use of diluents, as milk, almond emulsion, and mucilaginous solutions, is absolutely necessary during their employment to moderate their action. The tincture is the most proper form for internal use; or, if given in substance, the dose should not exceed one grain of the powdered flies, formed into a pill with opium or extract of henbane. They require to be used for some time, in order to prove beneficial.

Blistering flies, when applied to the skin, act as a local stimulant, first reddening and inflaming the part, and then producing from the exhalents a copious discharge of serum under the cuticle. These effects they produce more certainly completely than any vegetable acrid, and therefore they are generally employed to raise blisters.

It is uncertain whether blisters were used by the ancients; but modern practitioners daily and successfully employ them. Although their first operation is local, yet, under certain circumstances, the stimulus is sufficient to rouse the whole nervous energy, and excite the general system, so as to render their application useful in diseases of diminished excitement: on which account, in deep-seated local affections, when the inflammatory diathesis is considerable, the force of the circulation must be diminished by bleeding, purging, or other evacuates, before blisters can be advantageously applied. The diseases of debility in which they are useful, are low nervous fever, when accompanied with delirium, pale urine, frequent sighing, great anxiety, deafness, a fixed stare and glistening eyes. In palsy, and gutta serena, they are applied to the forehead over the supra-orbital nerve. They are found efficacious also in spasmodic and convulsive affections, from the irritation they produce, overcoming the morbid irritation which induced the spasm. Blisters, by their local action, relieve internal inflammatory diseases, by altering the balance

---

<sup>1</sup> Probably gleets were included in the term gonorrhœa by the old writers, who frequently mention cantharides as a remedy for gonorrhœa. Thus Boccone (*Museo di Fisica*, 1699) says, they were much used by the Sicilians in gonorrhœa.



of the circulation ; partly also, by diverting the attention from the prior seat of pain, contrary to the opinions of the ancient physicians, who attributed much of their efficacy to the serous effusion which they induce. Hence, their utility in ophthalmia, applied behind the ears, on the temples, or the forehead ; in phrenitis, over the head ; in cynanche tonsillaris, and in small-pox, when the swelling of the fauces affects respiration, upon or near the neck ; and in phthisis, catarrh, hepatitis, pneumonia, gastritis, and other intestinal inflammations, immediately over the seat of pain. In acute rheumatism, particularly that variety of it named *sciatica*, they have been found very useful. On the same principle, caries in the bones and joints, or a disposition to it, is often cured by the repeated application of blisters. “ Under their application the enlargements obviously subside ; the crepitation between the bones, the consequence of the abrasion of the cartilages, ceases to be felt when the blister begins to operate, the use of the joint is effectually recovered, and ankylosis prevented.”<sup>1</sup> A succession of blisters, also, to the vicinity of an inflamed organ, is more beneficial than a protracted discharge from one ; and a second blister often relieves after the first has failed. Blisters are contra-indicated in diseases of great debility, where there is a tendency to mortification ; as in the low stages of petechial fevers, cynanche maligna, confluent small-pox, and malignant measles ; and in dropsy, in which they are apt to occasion a very painful, dangerous erysipelas, and gangrene. Peculiar idiosyncrasies forbid their use in some persons, as they irritate, heat, produce thirst, pain, tremors, and sometimes convulsions. In those of irritable temperaments, their application is often attended with strangury and bloody urine ; and this effect is much increased if the blister-plaster be applied over a newly-shaved part, or if it be allowed to remain too long on after the blister has risen. To prevent strangury from the application of blisters, camphor has been erroneously regarded as a specific. It is more effectually prevented and relieved by copious dilution with milk, and mucilaginous fluids, and by fomentations of warm milk and water to the blistered part after the removal of the plaster : and much inconvenience of this nature may be prevented by interposing between the vesicatory and the skin, a piece of gauze, wetted with vinegar, and applied smooth and close over the plaster.<sup>2</sup>

The dose of the flies, is from gr. j. to grs. iii.

---

<sup>1</sup> Ford on Diseases of the Hip-joint, 55.

<sup>2</sup> In America, *Lytta cinerea* and *vittata* have been used instead of *Cantharides*. They raise a blister as speedily, and are said not to occasion strangury.



Officinal preparations. *Tinctura Cantharidis*, L. *Emplastrum Cantharidis*, L. *Emplastrum Cantharidis vesicatoriæ*, E. *Ceratum Lyttæ*, L. *Unguentum infusi Cantharidis vesicatoriæ*, E. *Unguentum Cantharidis*, D. *Unguentum Pulveris Cantharidis vesicatoriæ*, E.

CAPSICUM. *Spec. Plant. Willd.* i. 1050.

Cl. 5. Ord. 1. Pentandria Monogynia. Nat. ord. Luridæ, Linn. Solaneæ, Juss.

G. 334. Corolla wheel-shaped. Berry without juice.

Sp. 1. *C. annuum*. Annual Capsicum.<sup>1</sup> *Med. Bot.* 2d ed. 226. t. 80.

Officinal. CAPSICI BACCÆ, Lond. CAPSICI ANNUI FRUCTUS, Edin.

Dub. Berries of the Capsicum, or Cayenne Pepper.

Syn. Poivre d'Inde (F.), Spanisdier oderturkircher pfeffer (G.), Iaarlykse Spaansche peper (Dutch), Baelg-peber (Dan.), Spansk peppar (Swed.), Pieprzyea (Pol.), Pepperone (I.), Pimienta de Indias (S.), Pimentao da India (Port.), L'ul Mirch (H.), Brähn Maricha (San.), Mollaghai (Tam.), Filfil surkh (Pers.), Felfel (Arab.), Lornbak (Jav.), Chabai (Malay).

This is an annual plant, a native of both the Indies. It flowers in June or July. The stem is herbaceous, roundish, smooth, crooked, branching, and rising two or three feet in height. The leaves are ovate, smooth, entire, placed on long footstalks in an irregular order. The flowers are peduncled, axillary, solitary and white: the calyx is persistent, tubular, and divided at the edge into five short segments; the corolla wheel-shaped, five-cleft, the segments pointed and plaited: the filaments are short, tapering with oblong anthers; and the germen is ovate, supporting a slender style, which is longer than the filaments, and terminated by a blunt stigma. The fruit is a long, conical, pendulous pod-like berry of a shining, orange scarlet, or sometimes yellow colour, two-celled, and containing a dry spongy pulp with several flat, kidney-shaped seeds.

Many varieties of this species of *Capsicum* enter into the composition of Cayenne pepper; but, certainly, the best which is brought home from the West Indies, ready prepared, is made from the *Capsicum baccatum* (Bird pepper). Cayenne pepper is often mixed with muriate of soda; and, sometimes, with, a less innocent substance, the red oxide of lead. This fraud may be discovered by boiling some of the suspected pepper in vinegar, and, after filtering the decoction, adding to it a solution of sulphate of soda. If the pepper contain oxide of lead, a white precipitate will be produced, which, after being dried, and exposed to heat, mixed with a little charcoal, will afford a globule of lead: or the sulphuret of ammonia may be added to the acetic solution, which will then throw down a dark precipitate if lead be present.

---

<sup>1</sup> Sprengel, in his *History of Botany*, under the head "*Plinius*," says "*Capsicum annuum sine dubio est ea Piperitis, quam et Siliquastrum vocat* — (20-17.)"



*Qualities.*—Capsicum berries have an aromatic odour, which is somewhat impaired by drying; and an aromatic, extremely pungent, acrimonious taste, setting the mouth, as it were, on fire, and the impression remaining long on the palate. These sensible qualities are imparted to water, alcohol, and ether. Half a drachm of the powder infused in f℥jss. of boiling water lost grs. xij. The infusion was precipitated by infusion of galls, and alcohol dissolved the precipitate. It was also precipitated by nitrate of silver, oxymuriate of mercury, acetate of lead, the sulphates of iron, zinc, and copper, the alkaline subcarbonates, and alum: but was not altered by the mineral acids, the solution of potass, nor silicized potass. The ethereal tincture, when evaporated on the surface of water, left an orange coloured resin, in which the pungency of the capsicum was concentrated. These experiments point out the substances which are incompatible in formula with infusions of capsicum; and lead to the conclusion that it contains chiefly cinchonia, resin, vegetable mucus, and an acrid principle, perhaps alkaline, in which the acrimony resides.<sup>1</sup> When sold in powder, capsicum is sometimes adulterated with red lead, which, however, may be readily detected by digesting it in diluted nitric acid, and adding to the solution sulphate of soda, which will produce a white precipitate if red lead have been present in the pepper.

*Medical properties and uses.*—The fruit of the capsicum, or Cayenne pepper, is a powerful stimulant, unaccompanied with any narcotic property. It has been successfully given in atonic gout; in dyspepsia, when accompanied with much flatulence; in tympanitis, and paralysis. In dropsies, and other cachectic complaints, when chalybeates are indicated, a small portion of powdered capsicum is recommended as an excellent addition by Dr. Wright; and Bergius says, that he used it with success in obstinate intermittents.<sup>2</sup> I have had sufficient experience of its efficacy as an adjunct to cinchona in intermittents. It has also been found beneficial in lethargic affections<sup>3</sup>: but the diseases in which capsicum has been found most useful are cynanche maligna, and scarlatina maligna, in which it is given both internally, and used as a gargle. Its sensible effects are heat in the stomach, and a general glow over the body, without much affecting the pulse; and, as a gargle, it cleans without impeding the healing of the ulcers of the fauces. Cataplasms

---

<sup>1</sup> Journ. de Physique, 1820, p. 173.

<sup>2</sup> Mat. Med. e Regno Veg. i. 144.

<sup>3</sup> Dr. Paris says that *Rymer's Cardiac Tincture* is an infusion of capsicum, camphor, cardamom seeds, rhubarb, aloes, and castor, in proof spirit, with a small quantity of sulphuric acid. *Pharmacologia*.



of capsicum operate as powerful rubefacients without blistering the skin, and are used in the West Indies to relieve the coma and delirium which, almost constantly, attend tropical fevers. The diluted juice of the fruit is said to be a sovereign remedy in ophthalmia from relaxation.

Capsicum may be given in the form of pills, in doses from gr. vj. to grs. x.; or fʒss. to fʒij. of a tincture made with ʒiv. of *capsicum* and fʒviij. of *alcohol*. The gargle usually employed is made by kneading into a paste ʒj. of *Cayenne pepper* and ʒj. of *common salt*; then adding fʒvj. of boiling water; and to the solution, strained when cold, fʒiv. of vinegar. But a simple addition of fʒij. of the tincture to fʒvj. of water, or of infusion of roses, answers equally well.

Officinal preparations. *Tinctura Capsici*, L. D.

**CARBO LIGNI.** *Lond. Edin. Dub.* Charcoal.

*Syn.* Charbon de bois purifié (*F.*), Reine Kohle (*G.*), Koole (*Dutch*), Kol (*Swed.*), Carbone di legna, (*I.*), Carbon de lena (*S.*), Adapoo Currie (*Tam.*), Fuhm Chobic (*Arab.*), Zegal Chobic (*Pers.*), Koyla (*H.*), Arang (*Malay*).

Charcoal is prepared for the common purposes of fuel, by piling up billets of wood into conical heaps, which are covered with earth and sods, and then burned, with as little exposure to the action of the air as possible: but for the preparation of the finer kinds of charcoal, fit for medicinal use, the following process is employed. The wood to be charred is put, in the form of chips or of sawdust, into a large, cast-iron cylinder, fixed in masonry over a grate. This cylinder terminates at one end in a curved pipe, and the other end is furnished with a door, which is accurately closed after the wood is introduced, when a fire is lighted in the grate: the water, empyreumatic acid, and volatile parts of the wood, are driven off through the curved tube by the heat, which is increased until the contents of the cylinder become red-hot. The fire is then withdrawn, the cylinder is allowed to cool; and a black, shining, pure charcoal is thus obtained.<sup>1</sup> Ivory and bone shavings, treated in the same manner, make the preparation termed *Ivory black*. For internal use, however, it is perhaps necessary to have wood charcoal still purer; and to effect this the process of M. Lowitz is to be preferred. The charcoal is to be reduced to fine powder, and put into a crucible (so as to fill it), on which a pierced cover must be luted. This vessel is then to be heated red-hot, and kept so, as long as a blue flame appear to issue from the hole in the cover; and when this stops, it is to be taken from the fire, cooled in a dry place, and the charcoal, instantly, put into well-stopped bottles for use.<sup>2</sup>

<sup>1</sup> This process was invented by Bishop Watson, for the use of the gunpowder manufacturers, who require a very pure charcoal. — *Aikin's Chem. Dict. art. Carbon.*

<sup>2</sup> *Crell's Chemical Journal*, ii. 270.



In whatever manner charcoal is prepared, the purest contains, generally, about one-fiftieth of its weight of earths, salts, or metallic matters; its other constituents are, according to Doberiener<sup>1</sup>, 68.4 of carbon, with 1.5 of hydrogen, and a minute portion of oxygen.

*Qualities.*—Pure charcoal is inodorous and insipid; black, shining, and brittle. When newly prepared, it absorbs air and moisture from the atmosphere, so as to increase its weight from 10 to 18 per cent. It is insoluble in water and every other fluid<sup>2</sup>; and is easily pulverized. When excluded from air, it is not affected by the highest degree of heat. It corrects the foetid odour of putrifying animal and vegetable substances; and destroys the odour, taste, and colour of others, particularly of mucilages and oil, and matters in which extractive abounds. Thus common vinegar becomes colourless when it is boiled in charcoal powder; water, which has become foetid at sea, is purified by filtering it through charcoal; that intended for long voyages may be preserved perfectly pure by thoroughly charring the insides of the casks; and the empyreumatic odour and tastes of oils, as well as their adventitious colour, and the colour of the varieties of ardent spirits, of litmus, indigo, and other colouring matters, when they are dissolved or suspended in water, are destroyed by running them through newly-prepared charcoal powder. It also deoxidizes most of the acids.

*Medical properties and uses.*—Charcoal is evidently an anti-septic; and, as such, has been given internally to correct the putrid eructations of some kinds of dyspepsia. But, in order that it may produce this effect, it should either be newly prepared, or such as has been preserved in very well-stopped bottles. It is probable that it operates both by correcting the foetor, and absorbing the gas generated in the stomach, as well as checking the decomposition of the undigested aliment. Dr. Caliagno, an Italian physician, proposed to employ it instead of cinchona in intermittents<sup>3</sup>; but this suggestion has not been supported by British practitioners. It has been applied, advantageously, mixed up in powder with boiled bread, or linseed meal and water, as a poultice, to foul ulcers and gangrenous sores; and it is, undoubtedly, in combination with powdered catechu, kino or rhatany root, the best toothpowder known. I have discovered that it is also an excellent test for arsenic. (See the article *Arsenic*.)

---

<sup>1</sup> *Schweigger's Journal*, xvi. p. 92.

<sup>2</sup> There is, nevertheless, a quack preparation for cleaning the teeth, sold under the name of "*Concentrated solution of charcoal*."

<sup>3</sup> *Vide London Med. Repos.* vol. iii. p. 7.



The dose of charcoal may be from gr. x. to ʒj., combined with rhubarb.

CARDAMINE. *Spec. Plant. Willd.* iii. 481.

Cl. 15. Ord. 2. Tetradynamia Siliquosa. *Nat. ord.* Siliquosæ.

G. 1237. Pods opening elastically, with revolute valves. Stigma entire. Calyx somewhat gaping.

\*\*\* With pinnate leaves.

Sp. 19. *C. pratensis*.<sup>1</sup> Cuckow Flower. *Med. Bot.* 2d edit. 396. t. 133. *Smith's Flora Britan.* ii. 699.

Officinal. CARDAMINES FLORES, *Lond. Edin.* CARDAMINE; FLOS, *Dub.* The flowers and leaves of Cuckow Flower.

Syn. Cresson de Pres (*F.*), Wasserkresse (*G.*), Kockkesbloem (*Dutch*).

Cuckow flower is a perennial, indigenous, herbaceous plant, which grows in moist meadows, and flowers in April and May. The root is tuberous, and somewhat toothed. The stem rises about nine inches in height, is erect, smooth, stiffish, somewhat angular, and a little branched at the top: the leaves are dark-green, pinnated; the radical ones petiolate and spreading; those of the stem almost sessile; the leaflets are four pair or more, opposite, with a terminal one; on the lower leaves they are roundish, and irregularly dentated; but become more entire, linear, pointed, and concave, the nearer they are to the top of the stem. The flowers terminate the stem in a corymb, and stand upon smooth naked peduncles. The calyx is yellowish green, composed of four concave, oblong, nearly obtuse, deciduous scales, alternately larger, and protuberant at the base: the corolla is cruciform, the petals large, of a very pale purple colour, or white, ovate, veined, and slightly emarginate; with a yellowish green base. The filaments are six, four long, standing above the corolla, and two short, almost hid, supporting small oblong, yellow anthers; and invested at the base with four nectareous glands. The germen is the length of the stamens, slender and round, with a sessile stigma: it becomes a bivalved compressed pod, an inch in length, which, when the seeds are ripe, opens elastically, and rolls back in a spiral form. The seeds are many and round.

*Qualities.*—Every part of the plant is inodorous; but the flowers and leaves are slightly bitter and pungent, having in an inferior degree the taste of water-cresses. The leaves are often added to spring salads.

*Medical properties and uses.*—Cardamine flowers are said to be diuretic, and antispasmodic. They were supposed to be useful in all affections of the head by the ancients<sup>2</sup>: but their efficacy in spasmodic diseases was first mentioned by Dale<sup>3</sup>;

<sup>1</sup> Σισυμβριον ἱερὸν Dioscoridis.

<sup>2</sup> Cuckoo flowers form a part of the wreath “of idle weeds” with which Shakspeare has decorated the head of Lear, in his madness.

<sup>3</sup> *Pharmacol.* 204.



on the authority of a MS. of Doctor Tancred Robinson; and they were afterwards, in the year 1767, strongly recommended by Sir George Baker<sup>1</sup>, who had successfully used them in the cure of chorea, spasmodic asthma, and some other convulsive affections. Dr. Odier of Geneva<sup>2</sup> mentions a case of *incubus* which was cured by their use, although it had resisted several other antispasmodic medicines. They sometimes produce diaphoresis, but have otherwise little sensible operation. They are seldom used. The leaves have been regarded as possessing antiscorbutic qualities, but they have very little efficacy. The dose of the dried flowers powdered is from one drachm to three drachms, given twice or thrice a day.

CARDAMOMI SEMINA. Vide *Matonia Cardamomum*.

CARICÆ FRUCTUS. Vide *Ficus Carica*.

CARUM. *Spec. Plant. Willd.* i. 1470.

Cl. 5. Ord. 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

G. 561. Fruit ovate-oblong, striated. Involucre one-leafed. Petals keeled, inflex emarginated.

Sp. 1. *C. Carvi*.<sup>3</sup> Common Carraway. *Med. Bot.* 2d ed. 102.

t. 41. *Eng. Bot. Smith's Flora Britan.* 330.

Officinal. CARUI SEMINA, *Lond.* *Edin.* CARUON; SEMINA, *Dub.* Carraway seeds.

Syn. *Carvi* (*F.*), Kummelsamen (*G.*), Veldkemyn (*Dutch*), Kommer (*Dan.*), Brödkummin (*Swed.*), Karny (*Pol.*), Tmin (*Russian*), *Carvi* (*I.*), Alcaravea (*S.*), Alcaravia (*Portug.*), Dshintan (*Javanese*).

Carraway is an indigenous, biennial, umbelliferous plant, growing wild in meadows and pastures: but cultivated in several parts, particularly in Essex, for the sake of its seed. The flowers expand in May and June, and the seeds ripen in August. The root is fusiform; the stem smooth, channelled, branching; and seldom exceeding three feet in height; with smooth, doubly pinnate, incised leaves, the pinnulæ or segments of which are narrow, linear, pointed, and of a deep green colour. The flowers are in numerous terminal erect umbels, generally of ten rays, furnished with an involucre consisting of narrow leaflets, solitary, or two to three together, often altogether deficient; and without any partial involucre. The petals of the flowers are five, nearly equal, keel inward, obtuse, inflected, white, or of a pale blush colour; the filaments slender, rather longer than the petals, and bearing small roundish anthers; and the germen inferior, supporting very short capillary styles with simple stigmas. The seeds are two, oblong, bent, about one-fourth of an inch in length, of a brown colour, with five moderately elevated, longitudinal, straw-coloured ridges, the interstices being also obscurely furrowed. Carraway plants

<sup>1</sup> *Med. Trans.* i. 442.

<sup>2</sup> *Manuel de Médecine Pratique, &c.* Lect. xvi.

<sup>3</sup> *Καστροπος ορχυς* Dioscoridis. Careum, non Carum, Latine dici deberet. Conf. *Plin.* l. xix, sect. 49. *Gærtner*,



do not perfect their seeds until the second year. They are cut down in July, and the seed thrashed out on a cloth. The seeds are used by the London confectioners and bakers, as well as for medicinal purposes.

*Qualities*.—Carraway seeds have a pleasant aromatic odour, and a sweetish, warm, pungent taste; depending on an essential oil which is almost completely extracted by rectified spirit, and in an inferior degree by water. By distillation with water the whole is elevated, and an insipid extract remains.

*Medical properties and uses*.—These seeds are carminative and stomachic. They are used in flatulent colic and hysteria; and to give warmth to purgatives and other active remedies.

The dose in substance is from grs. x. to ʒij.

Officinal preparations. *Oleum Carui*, L. D. *Aqua Carui*, L. *Spiritus Carui*, L. E. D.

CARYOPHYLLA. Vide *Eugenia caryophyllata*.

CASCARILLÆ CORTEX. Vide *Croton Cascarilla*.

CASSIA. *Spec. Plant. Willd.* ii. 513.

Cl. 10. Ord. 1. Decandria Monogynia. Nat. ord. Lomentaceæ, Linn. Leguminosæ, Juss.

G. 813. Cal. five-leaved. Petals five. Anthers three superior, barren; the three lower ones beaked. Lomentum.

\* *Sennas*.

Sp. 18. C. *Fistula*. Purging Cassia. *Med. Bot.* 2d ed. 445. t. 160. (Conna) *Hort. Malabar*, Part i. p. 37. fig. 22.

Sp. 24. C. *Senna*. *Senna*. *Med. Bot.* 2d ed. 442. t. 159.

#### 1. CASSIA FISTULA.<sup>1</sup>

Officinal. CASSIÆ PULPA, Lond. CASSIÆ FISTULÆ FRUCTUS, Edin. CASSIA FISTULARIS; FRUCTUS PULPA, Dub. Cassia pulp.

Syn. Casse (F.), Rohnkassie (G.), Pypkassie (Dutch), Cassievër (Dan. Swed.), Polpa di Cassia (I.), Fistularis (S.), Ameltás (H.), Suvernaca (San.), Konnekai (Tam.), Khyar Sheber (Arab.), Khyar Chirber (Pers.), Drangu (Jav.), Mentus (Malay).

This tree is a native of both the East and West Indies and of Egypt. It rises to the height of forty or fifty feet, with a large trunk, covered with a soft cineritious bark, and is much branched at the top. The leaves are composed of six pairs of ovate, pointed, undulated pinnæ, of a pale green colour, with many transverse nerves, and peduncled: the stipules are scarcely apparent. The flowers, which appear in June, are of a golden colour, placed upon long pendent terminal spikes.<sup>2</sup> The leaves of the calyx are crenated, blunt, and greenish; the petals unequal, spreading, and waved. The three undermost filaments are long and incurved; the others exhibit large anthers, three of which are rostrated, or like the open beak of a bird, at the

<sup>1</sup> Γλυκοκάλαμον Myrepsici, ultimi fere Græcorum medicorum. Chaiarxambar of the Egyptians. Prosper Alpinus, de Plantis Ægypti, cap. ii.

<sup>2</sup> Alpinus says, "Sunt etiam hi valde odorati, præsertimque oriente sole." Ibid. l. c.



extremity. The fruit is a long, woody, dark-brown pod, about the thickness of the human thumb, and nearly two feet in length, cylindrical, with two longitudinal furrows on one side, and one on the other; and divided into numerous transverse cells, each containing one smooth, oval, yellowish, shining seed, with red lines dividing it longitudinally, imbedded in a soft black pulp.<sup>1</sup>

The pods are said to undergo a kind of fermentation, to prepare them for keeping. Those which are brought to this country come principally from the West Indies, packed in casks and cases; but a superior kind is brought from the East Indies; and is easily distinguished by its smaller, smooth pod, and by the greater blackness of its pulp. The heaviest pods, and those in which the seeds do not rattle on being shaken, are the best, and contain the greatest quantity of pulp, which is the part used.

*Qualities.* — The pulp has a slight, rather sickly odour, and a sweet mucilaginous taste. It is viscid; almost entirely soluble in water, and partially so in alcohol and sulphuric ether. The watery infusion, which shows a tendency to gelatinize, has, when filtered, a deep brown colour, and yields a precipitate with alcohol, and the solution of the superacetate of lead. The alcoholic and ethereal tinctures are not effected by the addition of water; although, when they are evaporated, a thin pellicle of resin remains. No alteration is produced on the alcoholic and watery infusions by infusion of galls, nitrate of silver, sulphate of iron, nor the nitric nor sulphuric acids; but oxymuriatic acid throws down a yellow-coloured precipitate which is insoluble in ether. Hence there is reason for concluding, with Vauquelin, that this pulp contains sugar, gelatin, gluten, mucus, a small portion of resin, extractive, and some colouring matter.

*Medical properties and uses.* — Cassia pulp is gently laxative; but, although it is adapted for children and very delicate women, yet, it is apt to induce nausea, flatulence, and griping, when taken in doses sufficient for stronger habits. To assist its operation, and prevent the griping, it is usually conjoined with some neutral salt and an aromatic; but it is now rarely prescribed in any case. The dose is  $\mathfrak{z}$ ijj to  $\mathfrak{z}$ j or more.

Officinal preparations. *Pulpa Cassiæ fistularis expressa*, E. *Confectio Cassiæ*, L. E. D. *Confectio Sennæ*, L. E. D.

## 2. CASSIA SENNA.

*Officinal.* SENNÆ FOLIA, *Lond.* FOLIA CASSIÆ SENNÆ, *Edin.* SENNA; FOLIA, *Dub.* Senna Leaves.

<sup>1</sup> *Gærtner de Fruct.* i. 313. t. 147.



*Syn.* Séné (*F.*), Sennabläter (*G.*), Sennet (*Swed.*), Liscie Senesowe (*Pol.*), Senne (*Dutch*), Senet (*Dan.*), Senna (*I. & Port.*), Sen (*S.*), Sená (*Arab.*), Sená Mecci (*H.*), Nilaverei (*Tam.*).

This species of *Cassia*, which yields the Senna of commerce, is an annual plant, a native of Upper Egypt. The best grows in the valleys of Nubia<sup>1</sup>, where it is named *Abyreyga*; flowering in July and August. It rises with an erect branching stem about two feet in height. At the base of the leaves, which are pinnate, and placed in alternate order, are two narrow pointed stipules: the leaflets, of which each leaf has five or six pair, are sessile, oblique at the base, oval, pointed, scarcely an inch in length, rather more than one-fourth of an inch broad, and of a yellowish green colour. The flowers are yellow, in loose axillary spikes: the calyx is deciduous, consisting of five narrow, obtuse, concave leaflets; the petals are roundish, concave, entire; the three lower ones larger than the two upper: the undermost filaments also are longer than the others, and furnished with large rostrated curved anthers; and the germen is long and flat, with a short incurved style terminated by an obtuse stigma. The fruit is an ovate, reniform, membranous, leafy, compressed legume, torose and marked with capillary, transverse, parallel striæ; bivalve, with six or nine cells, divided by very thin transverse partitions, and each containing one oblong heart-shaped seed.<sup>2</sup>

The best Senna, named in Nubia *guebelly*, or *sena mekke*, grows wild, and yields two crops of leaves, the abundance of which depends on the periodical rains. The first crop is collected after the first rains about the middle of September; the second in the following March, at which time the fruit is at its full maturity. The plants are cut when the flowers begin to fall, and exposed on the rocks to dry in the sun. — The leaves are then picked, packed up in bales, and sent to Bou-lac, the great entrepôt of senna<sup>3</sup>, where they are mixed with two other species of cassia; one the *C. senna* of Forskal with obtuse leaves; the other probably the *C. angustifolia* of Willdenow, the leaves of which are longer, narrower, and sharper pointed than those of the proper Senna, and come from Mocha; but the leaf with which Senna is chiefly adulterated is that of the *Cynanchum Oleafolium*, known in Egypt by the name of Argel or Arguel. The proportions, according to Dr. Cal-loden, are five hundred parts lance-leaved senna, two hundred


<sup>1</sup> C. Nectoux. Vide *Phil. Mag.* xv. 55.    <sup>2</sup> *Gærtner de Fruct.* ii. 312. t. 146.

<sup>3</sup> Burckhardt says that the Bedouin Arabs, who are the chief collectors of senna, sell it to the merchants of Esne at about one pound sterling per camel load (from 400 to 500 weight.) *Travels in Nubia*, 4to, p. 31.



of obovate senna, and two hundred of argel. The two first are equally good, but the last is truly an adulteration. It can be readily distinguished by attending to the following rules :

1. The leaf of argel (*a*) is an inch or fourteen lines long, while that of Senna (*c*) never exceeds nine lines. 2. The leaf of argel has a straight side; and the lateral nerves are not seen on the under disk, while those of Senna are conspicuous. 3. The leaf of argel is regular at its base (*b*), the two sides terminating at the same point on the petiole: but the Senna leaflet is oblique, one of the sides being larger, and produced lower on the petiole than the other. (*d*)<sup>1</sup> There is also reason for thinking that it is further adulterated with the leaves of *Colutea arborescens*, bladder senna, and of box: but these are easily distinguished from Senna leaves. The Senna, after being thus mixed, is re-packed in bales at Alexandria, whence it is exported to Europe.<sup>2</sup>



*Qualities.* — The odour of Senna leaves is faint, rather disagreeable, and sickly; the taste slightly bitter, aromatic, sweetish, and nauseous. Boiling water extracts about one-third of the weight of the leaves employed, but it requires a pint of boiling water to extract all the active matter from  $\bar{3}$ j. of Senna leaves. The infusion has a deep reddish brown colour, with the odour and taste of the leaves. This infusion, when exposed to the atmosphere, deposits a lemon-yellow coloured, insoluble matter; and a precipitate is produced by the strong mineral acids, oxalic acid, the carbonates of the alkalies, and several other substances. (See *Infusum Sennæ* among the preparations.) Alcohol and sulphuric ether, digested on the powdered leaves, acquire a deep olive-green colour. When the ethereal tincture is poured on the surface of pure water, a dark olive pellicle remains after the evaporation of the ether, which is almost insipid, and has all the properties of resin; and a golden yellow colour is communicated to the water.<sup>3</sup> The alcoholic tincture is rendered only slightly milky by the addition of water, and scarcely any precipitate

<sup>1</sup> *Hist. Nat. et Med. des Casses, &c.* Par L. T. Fred. Calloden, de Geneve, M.D. 4to. Planches. Montpellier, 1816.

<sup>2</sup> Nectoux says, the palthier, or senna-manager of Alexandria, acknowledged that the product of the two crops varies from 700 quintals to 1100 or more, one-third of which is *arguel*, the obtuse pointed cassia, and the sale is 1400 or 1500 quintals (more probably from 15 to 1600). *Phil. Mag.* l. c. Burckhardt says, that for many years the senna trade has been exclusively in one hand, being farmed by Mohammed Aly; and that "M. Rosetti has paid for the monopoly of senna 150 purses per annum, or about 3500*l.*" *Travels in Nubia*, 4to. 1819, p. 53.

<sup>3</sup> This colour may be produced by some extractive being taken up by the ether, closely united to the resin.



is produced; but a copious one is thrown down by oxymuriatic acid. The active principle of Senna, according to the experiments of M. M. Lassaigne and Fenuelle, is a saline substance, which they have named *cathartine* (cathartia). It is uncrystallizable, of a reddish-yellow colour, and has a bitter, nauseous taste. It is soluble in alcohol and in water, but insoluble in ether.<sup>1</sup> According to Bouillon Lagrange, the residue of the watery infusion, evaporated to dryness and burnt, yields potass, sulphate of potass, carbonate of lime, magnesia and silica.

*Medical properties and uses.*— Senna is purgative, generally operating under four hours after it is taken; and is well adapted for all cases in which the bowels require to be certainly, yet moderately, evacuated. In many habits it is apt to occasion griping, and therefore requires the addition of some aromatic, as carraway or cardamom seeds, or ginger; and its operation to be assisted by drinking plentifully of weak broths or gruel. The griping seems to be occasioned by the resinous matter, as the infusion made with cold water does not gripe, although it purges. Senna may be given in substance powdered; but the more usual form is that of infusion. Decoction is a bad form, as the activity of the medicine is much impaired by the boiling: owing, according to Gren, to the total dissipation of the nauseous and volatile principles; but, in our opinion, it is owing to the oxidizement of the extractive, which also accounts for the severe gripings induced by the decoction. The dose of the powder of the leaves is from ℥j. to ʒj.; but it is seldom given alone.

*Officinal Preparations.* *Confectio Sennæ*, L. E. D. *Extractum Cassiæ Sennæ*, E. *Infusum Sennæ*, L. D. *Infusum Tamarindicum Senna*, E. D. *Pulvis Sennæ compositus*, L. *Tinctura Sennæ*, L. E. D. *Tinctura Sennæ composita*, E. *Syrupus Sennæ*, D.

CASTOR. *Syst. Nat. Gmelin*. 124.

*Cl.* 1. *Ord.* 4. Mammalia, Glires.

*G.* 23. *Fore-teeth* in the upper jaw truncated, hollowed with a transverse angle; in the lower transverse at the point. *Grinders* in both jaws four. *Tail* long, depressed, scaly. *Clavicles* perfect.

*Species* 1. *C. Fiber*. The Castor Beaver. *Jonst. Quadr.* p. 147. t. 68.

<sup>1</sup> *Annales de Chim. et Phys.* xvi. 20. Cathartine is obtained by precipitating a filtered decoction of senna by acetate of lead: diffusing this in water, and passing through it sulphuretted hydrogen gas. The fluid is next evaporated to dryness, the extract digested in alcohol, and the alcoholic solution evaporated to dryness. This result is to be digested in alcohol acidulated with sulphuric acid, the effect of which is to throw down an insoluble sulphate of potass, which must be separated by the filter. Any excess of sulphuric acid in the fluid must be precipitated by acetate of lead, after which sulphuretted hydrogen is to be passed through the fluid; and this, being filtered and evaporated to dryness, is the cathartine.



*Officinal.* CASTOREUM<sup>1</sup>, *Lond. Edin.* CASTOREUM ROSSICUM et CANADENSE, *Dub.* Castor; Russian and Canadian.

*Syn.* Castoreum (*F.*), Kastoreunt (*G.*), Bevergeil (*Dutch*), Baevergel (*Dan.*), Bäfvergall (*Swed.*), Stroy Bobrowy (*Polish*), Castoro (*I.*), Castoreo (*S.*), Ash batchegan (*A.*), Goond beyduster (*Pers.*).

The beaver is an amphibious quadruped, found in the northern parts of Europe, Asia, and America, inhabiting the wooded banks of uninhabited rivers and lakes; in which situations it is gregarious, and constructs its habitation with greater skill than any other animal except man. The body is thick, under three feet in length, and covered with short iron-brown and chesnut-coloured hair: the feet are five-toed and webbed; the eyes small, and the ears short and hairy. The tail is about half the length of the body, flat, horizontal, scaly, with that part only of it which is next to the body covered with hairs. Between the anus and the external genitals are four follicles, of an oblong shape, smaller above and larger below; the two upper are filled with a fatty substance, whilst the two larger contain each about two ounces of an oily, viscid, strong-smelling substance, inclosed in membranous cells, which is the officinal castor.

When the beaver is taken, the follicles are cut off entire, and dried either by exposure to the sun, or in smoke. The castor is at first nearly fluid, but gradually becomes solid and viscid, occasionally perfectly dry and pulverulent. The best comes from Russia; but of late years it has been very scarce, and all that is now found in the shops is the produce of Canada. The cods of the Russian castor are large, dry, roundish, heavy, and solid, appearing, when cut, of a reddish liver colour; those of the Canadian are smaller, hard, oblong, thin, and corrugated on the outside. The Canadian castor, when treated with ammonia, affords an orange-coloured product; the Russian a white. In each beaver there is a large and small bag, and the castor in the larger bag is always the best. The goodness of the castor is determined by its sensible qualities; that which is quite black is insipid, inodorous, oily, and unfit for use. Castor is said to be sometimes counterfeited by a mixture of some gummy and resinous substances, with a little real castor, artificially interspersed with membranes, and stuffed into the scrotum of the goat.<sup>2</sup> The fraud is easily detected by comparing the smell and taste with those of real castor, and by the deficiency

<sup>1</sup> *Kapos*, Dioscoridis. The ancients erroneously believed that the castor follicles were the testicles of the beaver.

<sup>2</sup> *Duncan's New Edinburgh Dispensatory*, 5th edit. 220.



of the sebaceous follicles, which are always attached to the real cods.

*Qualities.* — The odour of castor is strong, heavy, and aromatic; the taste bitter, sub-acrid, and nauseous. It feels slightly unctuous, and is of a red-brown colour. Its odorous principle is dissipated by coction with water; but when it is simply infused in boiling water, its sensible qualities are in a small degree imparted to the infusion, which has a yellow colour, and shows the presence of an alkali, by changing to green the vegetable blues. Alcohol and sulphuric ether dissolve the resinous part of the castor, which remains after the evaporation of the menstrua, and retains all the odour and taste of the drug. According to the analysis of Bouillon Lagrange, castor contains the carbonates of potass, lime, and ammonia, iron, resin, a mucilaginous extractive matter, and a volatile oil. Canadian castor contains benzoic acid, both free and combined.<sup>1</sup> Bonn, who examined Russian castor, found in it a volatile oil, adipocere, resin, and lime.<sup>2</sup>

*Medical properties and uses.* — Castor is antispasmodic, and emmenagogue. It was formerly given in combination with myrrh as a specific in quartans<sup>3</sup>; and is still prescribed, with seeming advantage, in low nervous fevers, hysteria, epilepsy, and spasmodic affections: and from the idea of its action being particularly determined to the uterine system, it is supposed to prove useful in amenorrhœa and chlorosis. It may be exhibited either in powder, or in the form of tincture; but owing to the scarcity and the high price of good castor, it is seldom ordered; and the materia medica certainly contains many better antispasmodics.

The dose of powdered castor is grs. x. to ℥j., given as a bolus.

Officinal preparation. *Tinctura Castorei*, L. E. D.

CENTAURIA. *Spec. Plant. Willd.* iii. 2277.

Cl. 19. Ord. 3. Syngenesia Frustranea. Nat. ord. Compositæ Capitatæ, Linn. Cinarocephalæ, Juss.

G. 1548. *Receptacle* bristly. *Seed-down* simple. *Corolla* of the ray funnel-shaped, longer, irregular.

\* \* \* \* \* *Calcitrapæ*; with the spines of the calyx compound.

*Species* 89. *C. Benedicta*.<sup>1</sup> Blessed Thistle. *Med. Bot.* 2d edit. 34. t. 14.

*Officinal.* CENTAURIÆ BENEDICTÆ HERBA, *Edin.* CARDUUS BE-

<sup>1</sup> Laugier. *Ann. de Mus. d'Hist. Nat.* t. ix. p. 323.

<sup>2</sup> Bonn. *Anatomie Castoris atque Chemica Castorei Analysis*. 1806.

<sup>3</sup> Sennertus de Febribus, lib. ii. cap. 20.

<sup>4</sup> Ἀκοννα, Theophrasti.



NEDICTUS, FOLIA, *Dub.* The herbaceous part, or the leaves of Blessed Thistle.

*Syn.* Chardon benit (*F.*), Kardo benediktenkract (*G.*), Kardebenedict (*Dutch*), Korbenedikt (*Dan.*), Kardebenedikt (*Swed.*), O set Wloski (*Polish*), Cardo santo (*I.*), Cardo bendito (*S.*).

This is an annual plant, a native of Spain and the Grecian islands, flowering in June and September; and cultivated in the gardens of this country, where it thrives as well as in its native soil.<sup>1</sup> The root is whitish, cylindrical, and branched; the stem erect, roundish, channelled, and rough, about two feet in height, and branched towards the top. The lower leaves are peduncled, but the upper sessile, and in some degree decurrent; the whole are long, elliptical, rough, runcinate, and barbed with sharp points; of a bright green colour above, whitish underneath, and reticulated. The flowers are surrounded by an involucre of ten leaves, the five exterior of which are larger: the calyx is oval and woolly, and each scale terminated by pinnate, spinous points: the florets are of a yellow colour, those of the ray small, trifid, and sterile. The seeds have a paleaceous receptacle, are brown, pyriform, a little curved, deeply striated, and crowned with a double pappus, the outer one calyculate, the inner spinous.<sup>2</sup>

This plant is in greatest perfection when in flower, at which time it should be cut, quickly dried, and preserved in a dry airy place.

*Qualities.* — The odour is weak, yet unpleasant; the taste intensely bitter, but not very permanent. Its virtues are extracted both by water and alcohol. The watery infusion has a pale greenish-yellow colour, which is changed to a deep olive by sulphate of iron, and an orange-brown by the pure alkalies, although the carbonates do not affect it. Nitrate of silver and superacetate of lead occasion copious precipitates, and are therefore incompatible with this infusion.

*Medical properties and uses.* — *Carduus benedictus* is either emetic, diaphoretic, or tonic, according to the form and strength of the preparation in which it is administered. The decoction and strong infusion provoke vomiting; the less strong warm infusion determines powerfully to the surface, occasioning a copious flow of sweat; and the light infusion, made with six drachms of the leaves, and one pint of cold water, is an elegant and efficacious bitter in loss of appetite, and the dyspepsia which is occasioned by irregularities. It was formerly supposed to possess such extraordinary medicinal powers as to deserve the appellation *benedicta*; but it is sel-

<sup>1</sup> It was described as being cultivated in England by Gerarde in 1597.

<sup>2</sup> *Gærtner de Fruct.* ii. 385. t. 162.



dom used in modern practice. The dose of the powdered herb is from grs. xv. to ʒj; that of the infusion fʒij, given every three hours.

CENTAURII CACUMINA. Vide *Chironia Centaurium*.

CEPHAELIS. *Spec. Plant. Willd.* i. 977.

*Cl.* 5. *Ord.* 1. Pentandria Monogynia. *Nat. ord.* Aggregatæ, *Linn.* G. 357. Flowers in an involucred head. Corolla tubular. Stigma two-parted. Berry two-seeded. Receptacle chaffy.

*Species nova.* Cephaëlis vel Callicocca *Ipecacuanha*.<sup>1</sup> *Ipecacuan*, *Linn. Soc. Trans.* vi. p. 137. t. 2.

*Officinal.* IPECACUANHÆ RADIX, *Lond. Edin. Dub.* The root of *Ipecacuan*.

*Syn.* *Ipecacuanne* (*F.*), *Brechwerzel* (*G.*), *Kräkrot* (*Swed.*), *Ipecacuana* (*I.*), *Ipecacuanha* ó *Bejuquillo* (*S.*).

This plant is a perennial, found growing in shadowy moist situations in the forests of the provinces of Pernambuco, Bahia, Rio Janeiro, Paulensia, Mariannia, and other provinces of the Brasils; flowering in December, January, February, and March; and ripening its berries in May. The root is simple, or somewhat branched, and furnished here and there with short radicles; it is roundish, three or four inches in length, and two or three lines in thickness; bent in different directions, externally brown, and annulated with prominent, unequal, roughish rings. The stem is procumbent at the base, rising from five to nine inches in height, round, the thickness of a hen's quill; smooth, brown, leafless, and knotted in the lower part, but leafy towards the apex: after the first year it throws out runners, from which, about six inches apart, new erect stems arise. The inferior leaves are caducous, so that not more than eight generally remain at the summit of each stem when it flowers: they are almost sessile, opposite, spreading, ovate, pointed at both ends, three or four inches long, and less than two broad; of a deep green colour on the upper surface, and of a whitish-green, downy, and veined on the under. At the base of each pair of leaves are sessile, fimbriated, short, withering stipules, embracing the stem. The flowers are aggregated in a solitary head, on a round, downy footstalk, terminating the stem, and encompassed by a four-leaved involucre. The florets are sessile, from 15 to 24 in number, interspersed with little bractes: the calyx is very small, five-toothed, superior, and persistent; the corolla monopetalous, the expansion shorter than the tube, and divided into five ovate, acute, recurved segments: the filaments are short, capillary, inserted into the upper part of the tube, and

---

<sup>1</sup> As Willdenow, following Swartz, has united the genus *Callicocca* with that of *Cephaëlis*, we have referred the *Ipecacuanha* to this genus.



bearing long erect anthers : the germen inferior, supporting a filiform style, with two obtuse stigmas the length of the anthers ; becoming a soft one-celled berry, of a reddish-purple colour changing to black, and containing two oval seeds.

According to Decandolle, the term *Ipecacuanha* in South America implies generally *vomiting root* ; and, therefore, it is applied to the roots of very different species of plants. The plant, however, which we have described from Professor Brotero's description published in the sixth volume of the Linnean Transactions, and the *Psycotria emetica*, which Mutis says yields the Peruvian gray Ipecacuan, are the plants that yield the varieties of the root brought to this country.<sup>1</sup> We have found very little of the white Ipecacuan in any of the specimens of the Ipecacuan of the shops which we have examined. Both the gray and the brown varieties of the root are brought to this country packed in bales from Rio Janeiro. Both are in short, wrinkled, variously bent and contorted pieces which break with a resinous fracture. The gray is about the thickness of a small quill, full of knots and deep circular fissures, that nearly reach down to a white woody vascular cord that runs through the heart of each piece ; the external part is compact, brittle, and looks smooth ; the brown is smaller, more wrinkled, of a blackish brown colour on the outside, and white within : the white is woody, and has no wrinkles.

In choosing *Ipecacuanha*, the larger roots, which are compact, and break with a resinous fracture, having a whitish gray somewhat semitransparent appearance in the inside of the cortical part, with a pale straw-coloured medullary fibre, are to be preferred.

It is impossible to ascertain at what period the effects of this root were first known in America, where the Indians used it as an emetic before their connexion with Europeans : but although Piso described its uses fully in his Natural History of Brazil so early as 1618, and brought the root to Europe, yet it was scarcely used by Europeans before the year 1700. It was carried to France by a French physician of the name of Le Gras in 1672 ; but it did not attract general notice until it was, a third time, introduced by a French merchant of the name of Grenier, who brought 150lbs. of it

---

<sup>1</sup> The title of Ipecacuan is generally given to the roots of the following plants, besides those mentioned above, in South America : *Viola parviflora*, *V. Ipecacuanha*, *V. Calceolaria*, *Cynanchum Ipecacuanha*, *C. tomentosum*, and *Asclepias currassavica* ; and sometimes to *Euphorbia Ipecacuanha*, *Dorstenia Brasiliensis*, and *D. arifolia*. In St. Domingo several species of *Ruellia*, which provoke vomiting, are named false Ipecacuan. *Nouveau Dictionnaire d'Histoire Naturelle*, art. *Ipecacuanha*.



from Spain in 1686, with which trials were made at the Hotel Dieu. Helvetius first made known its use in dysentery, and was rewarded by Louis XIV. with 1000*l.* sterling for the discovery.

*Qualities*. — The entire root is inodorous, but the powder has a faint disagreeable odour. The taste is bitter, subacid, and extremely nauseous. Water at 212° takes up rather more than eight parts in twenty of ipecacuan, but decoction destroys the emetic power of the root: alcohol takes up four parts, and proof spirit six and a half; and the alcoholic is more emetic than the aqueous solution. Various analyses of Ipecacuanha have been made in order to detect its emetic principle, but the most satisfactory is that of MM. Majendie and Pelletier.<sup>1</sup> After digesting the powdered root in double its weight of ether, in order to separate any fatty matter, the remainder was treated with four times its weight highly rectified alcohol, until it ceased to become coloured even when aided by heat. These tinctures, after being allowed to cool, and to deposit some flakes of wax which were separated by filtration, were then evaporated to dryness, and the residue re-dissolved in water: acetate of lead being added to the watery infusion, a precipitate formed, which, when edulcorated and diffused through water, was exposed to a current of sulphuretted hydrogen gas to separate the lead; after which the liquid being filtered and evaporated to dryness, afforded a substance of a peculiar nature, which they termed *Emetine*, and on which it was experimentally demonstrated that the emetic properties of the root depend. *Emetine*<sup>2</sup>, when pure, is of a reddish brown colour, solid, and pulverulent, nearly inodorous, and has a slightly bitter, acrid, but not nauseous taste. When exposed to a heat stronger than that of boiling water, it is decomposed, furnishing water, carbonic acid, some oil, and acetate acid, charcoal being left. It is little soluble in water, and does not deliquesce in a moist atmosphere. It is soluble in alcohol, but not in ether. To detail the action of other chemical agents on this body is here unnecessary; the results are sufficient to characterize it as a substance *sui generis*. Besides emetine, ipecacuanha has been found, by the experiments of the above chemists, to contain oil, wax, gum, starch, and lignin.

The medicinal value of ipecacuanha depends, undoubtedly, on the quantity of emetine it contains; and this varies in the three varieties of the root found in the shops. MM. Majen-

<sup>1</sup> Vide *Ann. de Chim. et de Phys.* iv. 172. and *Lon. Med. Repository*, viii. p. 252.

<sup>2</sup> The name is derived from *εμεω*, vomo.



die and Pelletier obtained 16 parts of it in 100 of the cortical part of brown Ipecacuanha, the root of the *Psycotria emetica* of Mutis<sup>1</sup>, 14 in 100 of the gray Ipecacuanha, the root of the *Callicocca Ipecacuanha*<sup>2</sup>, and 5 only in 100 of the white Ipecacuanha, the root of the *Viola emetica*.<sup>3</sup> The woody pith even of the brown-variety contains very little emetine, and hence it should be separated in reducing the root to the form of powder.

Experiments made with emetine on animals, prove that it is emetic and purgative, in doses of half a grain, and exerts a specific action on the lungs and mucous membrane of the intestinal canal, and has also narcotic properties: that it may be employed instead of Ipecacuanha in every case in which this medicine is useful, the dose being more easily regulated, and the effects more certain. When taken in an over-dose, its action can be instantly paralysed by decoction of galls. These experiments are at variance with those of Dr. Irvine, which led him to conclude that the watery solution of Ipecacuan is more emetic than the alcoholic, the reverse being found to be the case.

The powder of Ipecacuanha is apt to become inert by keeping; and therefore it should be preserved in small phials, well corked, and not exposed to the light. Long-continued boiling also renders it inert.

*Medical properties and uses.* — Ipecacuanha, when administered in large doses, is emetic; in smaller ones diaphoretic and expectorant; and in still smaller doses, it acts as a stomachic, stimulating and giving energy to the digestive organs. As an emetic, it is mild, safe, and certain in its operation; but it is a mistake, that when given in larger doses than are necessary, it does not operate more violently, but only in a shorter space of time. It does not act so quickly as many other emetic substances; but it evacuates, completely, the contents of the stomach, and does not so much weaken it as antimonial emetics. It is given at the commencement of continued fevers, the progress of which is sometimes cut short by its operation; and it is also, frequently, found to stop the paroxysm of an intermittent, when given immediately before the accession of the cold stage. At the commencement of inflammation of the

<sup>1</sup> The components procured from 100 parts of brown Ipecacuanha were as follows: of fatty and oily matter 2; emetic matter (emetine) 16; wax 6; gum 10; starch 42, and ligneous matter 20; the remaining four parts being regarded as loss.

<sup>2</sup> 100 parts of the gray variety yielded, of fatty matter 2; emetine 14; gum 16; starch 18; woody matter 48; with merely a trace of wax, and two of loss.

<sup>3</sup> From 100 parts of the white Ipecacuanha, were obtained of emetine 5; gum 35; vegeto-animal matter 1; and woody matter 57; besides three of loss.



pharynx, larynx, and trachea, when the inflammation does not run very high; in cynanche tonsillaris; purulent ophthalmia, abscess, and every case in which it is necessary to evacuate the stomach, or to increase the energy of the absorbent system by full vomiting, ipecacuan has been found useful. As an emetic, however, it is contra-indicated when there is any reason for suspecting inflammation of the encephalon, passive hæmorrhagy, or hernia; and in the advanced stage of typhus fevers, when the pulse is feeble, and the strength much diminished; but in these instances all emetics are hurtful. In doses sufficient to excite nausea without producing vomiting, Ipecacuan is given with excellent effects in dysentery<sup>1</sup>, and obstinate diarrhœa: in which cases its efficacy seems to arise in a great degree from the nausea, which is kept up by the repetition of the small doses, diminishing the arterial excitement, and determining to the surface; and partly also, as Cullen supposed, from its producing a steady determination of the peristaltic motion of the intestines downwards.<sup>2</sup> Perhaps also to these first-mentioned effects of the nausea, may be attributed much of the benefit which results from the use of Ipecacuan in spasmodic asthma, dyspnœa, pertussis, and epilepsy. In the first of these diseases its emetic power is taken advantage of to relieve the paroxysm, after which it is given in repeated small doses to prevent its return.<sup>3</sup> In nauseating doses also, owing to the nausea lessening the force of the circulation, it has been employed with the best success in uterine and pulmonary hæmorrhages. As a sudorific, it is used in acute rheumatism, arthritic affections, dropsy, and other diseases in which sweating is necessary. It is generally given, in these cases, in combination with opium and neutral salts, according to the mode introduced by *Dover*; (see *Pulvis Ipecacuanhæ compositus*.) But we have found it in combination with opium alone in a larger proportion, more efficacious, particularly in rheumatism. Its expectorant powers have been found exceedingly useful in catarrhal affections, pneumonia after bleeding, and in the early stage of phthisis, in which its diaphoretic effect is also beneficial.

The emetic operation of Ipecacuan is quickened by combining it with tartarized antimony: and, on the contrary, it is counteracted by opium, by vegetable infusions containing tannin, and by vegetable acids. Opium, however, is rendered less narcotic when combined with Ipecacuan, although its power of allaying pain is not diminished, while the sudorific

<sup>1</sup> *Piso, Helvetius, Cleghorn, Pringle.*

<sup>2</sup> *Materia Med.* ii. 477.

<sup>3</sup> *Akenside.*



effect of the Ipecacuan is much augmented by the combination. We do not, however, agree in opinion with those who think, that it is to be relied upon as an antidote against the deleterious effects of opium; its emetic effect being too slow, and checked by the opium. The infusion of nutgalls is the only certain and powerful antidote for an over-dose of Ipecacuan, instantly rendering it inert. Idiosyncrasy occasions some persons to be affected with the most distressing sensation of suffocation, by the effluvia of this root.

Ipecacuan is exhibited in substance, and in aqueous and vinous infusions: and, on the continent, a syrup of it is used for children.<sup>1</sup> The dose of the powder to produce full vomiting, is from grs. xv. to ʒss.; and of the aqueous infusion, which is made by macerating for an hour ʒij of the powdered root in fʒvj. of boiling water, and filtering, fʒj. or fʒjss. may be given every half hour till it excite vomiting. The emetic effect is continued, and rendered easier to the patient, by drinking, in the intervals of vomiting, large draughts of tepid water. For producing the other effects of Ipecacuan, it is given in doses of one, two, or three grains, generally in the form of pills, and repeated every four or five hours: but although its sudorific effect, when begun, is aided and kept up by the use of warm fluids, yet these must not be drank soon after the dose has been taken.

Officinal preparations. *Pulvis Ipecacuanhæ compositus*, L. D. *Vinum Ipecacuanhæ*, L. E. D.

#### CERA. Wax.

*Syn.* Cire (*F.*), Wachs (*G.*), Wasch (*Dutch*), Vox (*Dan.*), Wax (*Swed.*), Work (*Pol.*), Cera (*I.*), Cera (*S.*), Shuma (*Arab.*), Mom (*H. Pers.*), Medhúch-hista (*Sans.*), Mellughoo (*Tam.*), Miellie (*Cyng.*).

Bees, as the experiments of Huber have proved<sup>2</sup>, produce the wax of which the delicate partitions of the cells of their combs are constructed, from honey, sugar, and the sweet secreted juice found in the nectaries of plants; but they do not collect it ready formed from the anthers of flowers, as has been generally supposed. The organ by which it is secreted has not yet been discovered; but it is deposited in what have been termed wax pockets, situated under the four, intermediate, ventral segments. These wax pockets are trapeziform, whitish, and of a membranaceous texture; and on them the laminæ of wax are found. It is, nevertheless, also produced as a secretion by many plants forming the silvery powder or

<sup>1</sup> The following is the mode of preparing the syrup. Take oz. vj. of ipecacuan, in fine powder, and pour over it lbs. vj. of cold water, and after twenty four hours decant it off; then add lbs. vj. more of water; and again lbs. vj. more, a third time, proceeding always as at first. Mix the decanted liquors, and filter; and then with a moderate heat dissolve in them lbs. xij. of refined sugar. One ounce is equivalent to twelve grains of the powder. *Annales de Chimie*, xlvi. 55.

<sup>2</sup> *Nicholson's Journal*, ii. 182.



bloom, which often covers their leaves and fruit, and is found in great abundance combined with resin, covering the trunk of the wax-palm (*Ceroxylon Andicola*) of South America<sup>1</sup>, and very pure, encrusting the seeds of the *Myrica cerifera*, or wax-tree of Louisiana and other parts of North America.<sup>2</sup> Hence wax, in the extended meaning of the term, may be regarded both as an animal and a vegetable product. But it is the former species only of it, or bees' wax, which is officinal, and demands our present consideration. It is admitted into the list of materia medica under two forms: 1st, As it is procured originally from the combs, combined with colouring matter, or unbleached; and, 2d, Deprived of colour, and purified or bleached.

#### 1. UNBLEACHED WAX.

*Officinal.* CERA FLAVA, *Lond. Edin. Dub.* Yellow Wax.

*Syn.* Cire jaune (*F.*), Wachs (*G.*), Gult wax (*Swed.*), Zotty Wosk (*Pol.*), Cera gialla (*I.*), Cera qualda (*S.*), Munjie Mellughoo (*Tam.*).

Yellow wax is prepared immediately from the honeycomb.<sup>3</sup> The honey is obtained by dripping and pressing the comb, which is then soaked for some days in clear water to extract all the remaining honey, and afterwards melted in a clean vessel with boiling water, and pressed through cloth bags. It is then re-melted and cast into round cakes, in which form it is brought to market.<sup>4</sup>

*Qualities.*—Good and recent yellow wax has a slight odour of honey, is insipid, and of a bright, pale-yellow hue. It is brittle, yet soft, somewhat unctuous to the touch, but without adhering to the fingers, or to the teeth when it is chewed; acquires tenacity when heated; melts at 142°, and burns entirely away. Its specific gravity varies from 0.9600 to 0.9650. (For the other properties of wax, see *Cera alba*.)

Wax in this form is often adulterated with earth, peas' meal, or resin and tallow. *Earth*, or *peas' meal*, may be suspected when the cake is very brittle, and the colour inclines more to gray than bright pale yellow: they may be separated by re-melting and straining the wax. The presence of *resin* may be suspected when the fracture appears smooth and shining, instead of being granulated; and it may be detected by put-

<sup>1</sup> This palm is found in the Quindin mountains only, rising 180 feet in height, and having leaves twenty feet long. The waxy secretion covers the trunk to the thickness of about two inches, and consists of two-thirds of resin and one of wax. *Humboldt, Plantæ Æquinoctiales*, &c. fasc. i.

<sup>2</sup> The *pe la*, ornatural white wax of the Chinese, is an animal wax, produced by a species of *Coccus*; and the *white lac* of India appears also to be a variety of wax.

<sup>3</sup> There are bees in India who prepare a black wax. *Jacquin, Elem. Chim.* p. 34.

<sup>4</sup> Large quantities of wax are imported from the Baltic, the Levant, and the Barbary coast.



ting small pieces of the wax in cold alcohol, which will readily dissolve the resinous part, without acting on the real wax. *Tallow* is discovered by the greater softness and unctuousity of the cake, and its disagreeable suffocating smell when melted.

*Medical properties and uses.* — Yellow wax is scarcely ever ordered for internal use, although its colouring matter does not affect its medical properties. It is chiefly employed in the composition of external applications.

Officinal preparations. *Cera flava purificata*, D. *Emplastrum Ceræ*, L. E. *Cerata*, L. E.

## 2. BLEACHED WAX.

*Officinal.* CERA ALBA, *Lond. Edin. Dub.* White Wax.

*Syn.* Cire blanche (*F.*), Hwitt wax (*Swed.*), Biaty Wosk (*Pol.*), Cera bianca (*I.*), Cera blanca (*S.*), Vulay Mellughoo (*Tam.*), Suffiad mooru (*Duk.*).

When yellow wax is exposed, with an extended surface, to the action of light and air, and sprinkled with water, the yellow colour and peculiar odour are lost, and it becomes white. This process is thus performed: The yellow wax is melted with a very little water in a copper vessel, and then run off, through a plug-hole in the bottom, into another vessel, which is covered with a cloth to retain the heat until the water and the impurities settle. The clarified melted wax is next suffered to flow into a vessel, the bottom of which is full of small holes, through which it runs in small streams upon a cylinder kept constantly revolving over, and partly dipping in cold water, into which the wax falls, drawn out into thin shreds or ribbands, and is instantly cooled. These are spread upon cloths stretched on frames exposed to the light and air, and occasionally watered and turned; so that after some days the colour nearly disappears. After being thus half-bleached, the wax remains heaped up in a solid mass for a month, when the whole process is again repeated. It is, lastly, generally melted and cast into thin discs about five inches in diameter, in which form it is found in the shops.

White wax is sometimes adulterated with white oxide of lead, in order to increase its weight, and with white tallow. The former is detected by melting the wax in water, when the oxide falls to the bottom of the vessel; and white wax is known to contain tallow when it is of a dull opaque white, and wants the translucency which distinguishes pure wax.

*Qualities.* — Pure white wax is perfectly insipid, inodorous, and somewhat translucent. It is harder, less unctuous to the touch, heavier, and less fusible than yellow wax; its specific gravity being from 0.8203 to 0.9662, and its melting point 155°. It melts into a colourless transparent fluid, which concretes again as it cools, resuming its former appearance. Wax is perfectly insoluble in water, and nearly so in cold alcohol, although this fluid takes up about one-twentieth of its weight



at a boiling temperature; which, however, is again deposited as the fluids cool. Ether acts in the same manner as alcohol. Wax dissolves in the fixed oils, forming the base of cerates and ointments; and unites in some degree, when boiled, with alkalis, forming soaps. The acids at an ordinary temperature scarcely affect it; but when the sulphuric acid is boiled on it, a thick blackish mass is produced. The products of its decomposition by heat, in close vessels, show that, like the fixed oils, it is a triple compound of carbon, 81.784; hydrogen, 12.672; and oxygen, 5.544, in 100 parts.<sup>1</sup> Dr. John affirms, that 100 parts of wax digested in boiling alcohol is divided into two distinct substances; eighty parts consisting of a body soluble in hot alcohol and oils, and deposited by cooling, and thirteen of a substance completely insoluble in alcohol, the first of which he has named *cerin*, the second *myricum*.<sup>2</sup>

*Medical properties and uses.*—Wax is regarded as a demulcent, and is sometimes exhibited in obstinate cases of diarrhœa and dysentery, with the view of sheathing the bowels; but its place may be better supplied by simple mucilages and gelatinous solutions. It is generally exhibited diffused in mucilaginous fluids by means of soap, in the proportion of one-third part of the wax, with which it is first melted, and then rubbed in a mortar with the fluid, which is gradually added: but Poerner's method, which is, first to melt the wax with olive oil, and then to mix the oily compound while hot with the mucilaginous fluid, by triturating with the yolk of an egg, is a preferable one. The dose is a cupful of the emulsion, containing about ʒj. of wax, given every four or five hours.

Officinal preparations. *Cerata varia*, L. E. D.

CERVUS. *Syst. Nat. Gmelin*, 175.

Cl. 1. Ord. 5. Mammalia Pecora.

G. 29. *Horns* solid; when tender covered with a velvety coat, and growing at the apex; shed annually; forked. *Fore-teeth*, eight in the lower jaw. *Tearing-teeth* none (sometimes solitary in the upper jaw).

*Species* 1. C. *Elaphus*.<sup>3</sup> The Stag, or Hart. *Johnst. Quadr.* 82. t. 32. 35.

*Officinal.* CORNUA, *Lond.* CERVİ ELAPHİ CORNU, *Edin.* CORNU CERVINUM, *Dub.* Hart's horns.

*Syn.* Corne de Cerf (*F.*), Hirschhorn (*G.*), Hertshoorn (*Dutch*), Hjorthorn (*Swed.*), Corno di Cervo (*I.*), Cuerno de Ciervo (*S.*), Corne de Veado (*Port.*).

The stag, of which there are three known varieties, is a native of almost every part of Europe, and of the northern parts of America and Asia. In Britain its numbers have been much

<sup>1</sup> Thenard. *Recherches Phys. Chim.* ii. p. 316.

<sup>2</sup> *Tableau Chim. du Règne Animal.* p. 209.

<sup>3</sup> Ελαφος. *Aristot. Hist. Animal.* ii. c. 7. 18.



reduced by the progress of civilization; but it is still found wild in the highlands of Scotland, the moors bordering on Devonshire and Cornwall, and on the Kerry mountains in Ireland. It is a very beautiful animal, about three feet and a half in height, of a rust-brown colour on the upper part of the body, and whitish below. The horns are annually shed, about the end of February and March; but are soon reproduced in a soft, tender state, full of blood vessels, and covered with a velvety skin, which is lost as they increase in size; and at length, about the month of July, they become hard, compact, and bony. They have no horns till they are above a year old, and these do not branch till the third year; after which the branches increase in number every year, so that the age of the animal may in some degree be determined by them.<sup>1</sup>

These horns differ from those of most other animals, and approach nearer to the nature of bone, containing only less of the phosphate of lime in their composition, and yielding a much larger proportion of gelatine. It is for the sake of the gelatine that their shavings are medicinally used. These are often adulterated with shavings of mutton bones, which, however, are easily detected by their greater degree of brittleness.

*Qualities.* — Hartshorn shavings when good are inodorous and insipid, pliant, of an ivory-yellow colour; and contain 27 parts of gelatine in 100 parts.<sup>2</sup> Four ounces of the shavings boiled in two pints of water until one pint be dissipated, and the remainder strained, afford, when the decoction cools, a clear, transparent, colourless, insipid, inodorous jelly, which is a compound of gelatine and water.

*Medical properties and uses.* — The gelatine yielded by stags' horns is considered as a demulcent; but its nutrient properties are more useful than its medicinal virtues. It forms, when united with orange-juice, sugar, and a little wine, a good article of diet for the sick and convalescent; and when mixed with an equal portion of cows' milk, it is very useful in the irritations of infants arising from acidities in the primæ viæ.

*Officinal preparations.* *Cornu ustum*, L. D. *Liquor vol. Cornu cervini*, D. *Oleum Cornu cervini rectif.* D.

**CEREVISIÆ FERMENTUM**, *Lond. Edin.* Yeast.

*Syn.* Leveure (*F.*), Güscht (*G.*), Fermento di cervogia (*I.*), Espuma de cerbeza (*S.*).

This substance is the scum or frothy matter which collects on the surface of beer while fermenting. It soon undergoes the putrefactive fermentation, but may be preserved by drying it to the consistence of a slightly cohesive paste; in which state

<sup>1</sup> The castrated fawn never gets horns at any period of its life.

<sup>2</sup> *Annales de Chimie*, xxxiv. 71.



it is sold in Paris. It has been chemically examined by Westrumb, who obtained from it a variety of ingredients<sup>1</sup>; but its essential constituent, or the fermenting principle, is supposed to be gluten, or something very analogous to that vegetable principle. Its medical properties may, perhaps, be attributed to its containing the bitter of the hop, some ready-formed alcohol, and carbonic acid.

*Qualities.* — Yeast has a vinous, sour odour; a bitter taste; and reddens the vegetable blues. When it is filtered, a matter remains on the filter which possesses properties similar to those of vegetable gluten; and by this separation the yeast loses the property of exciting fermentation, but recovers it again when the gluten is added. The addition of yeast to any vegetable substance containing saccharine matter excites fermentation in it, and carbonic acid gas is evolved.

*Medical properties and uses.* — Yeast is tonic and antiseptic. Some years ago it was given with seeming advantage in typhoid fevers attended with symptoms of putridity; but the facts brought forward in support of its efficacy require further confirmation.<sup>2</sup> As an external application, however, to foul and sphacelating ulcers, when united with farinaceous matters in the form of cataplasm or poultice, it is productive of the best effects. It corrects the fœtor of the discharge, assists sloughing, and promotes the formation of a benign and healthy pus.

The dose of yeast is a table-spoonful or two (about f ʒ ss), repeated every second or third hour. It is generally combined with porter, or wine, and sugar.

Officinal preparations. *Cataplasma Fermenti*, L.

CETACEUM. Vide *Physeter macrocephalus*.

CHIRONEA. *Spec. Plant. Willd.* i. 1065.

Cl. 5. Ord. 1. Pentandria Monogynia. Nat. ord. Rosaceæ, Linn. Gentianæ, Juss.

G. 349. Cor. wheel-shaped. Pistil declined. Stam. seated on the tube of the corolla. Anthers spiral at the end. Pericarp two-celled.

Species 9. *C. Centaurium*.<sup>3</sup> Common Centaury. *Med. Bot.* 2d edit. 275. t. 96. *Smith. Flor. Brit.* 257. *Eng. Bot. t.* 417.

Officinal. CENTAURII CACUMINA, Lond. CHIRONEÆ CENTAURII SUMMITATES, Edin. CENTAURIUM MINUS; CACUMINA FLORENTIA, Dub. The flowering tops of Common Centaury.

Syn. Petite Centaurée (F.), Tansendgüldenkraut (G.), Centaúra (I.), Gentiana Centaura (S.).

<sup>1</sup> From 15.142 parts of yeast, he obtained the following substances: potass 13, carbonic acid 15, acetic acid 10, malic acid 45, lime 69, alcohol 240, extractive 120, mucilage 240, saccharine matter 315, gluten 480, and water 13.595 parts; besides some traces of phosphoric acid and of silica. *Crell's Annals*, 1796, and *Thomson's Chemistry*, 4th edit. v. 406.

<sup>2</sup> It was suggested as a remedy in these complaints by the Rev. Mr. Cartwright.

<sup>3</sup> Named, according to Pliny, *κενταυριον*, from Chiron the centaur, l. 25. c. 6.



This is an indigenous annual plant, growing in dry gravelly pastures, and flowering in July and August. The root is small, woody, and branching: the stalk, which rises about a foot in height, is erect, smooth, and quadrangular; divided above into a dichotomous panicle, and branched at the base. The leaves are opposite, sessile, elliptical, three-nerved, and smooth: the flowers spring from the angles of the division of the stem, sessile, and erect, expanding only in the sunshine: the calyx is half the length of the tube of the corolla, five-cleft, with the segments subulate and erect; the corolla is pink, or rose-coloured, divided at the limb into five elliptical equal segments, spreading, and somewhat concave. The filaments are bent down, and furnished with oblong, yellow, three-times twisted anthers; the germen is oblong, bearing a simple style, with a clubbed stigma.

*Qualities*.—Common Centaury is almost inodorous; but the petals, leaves, and stalk have an intensely bitter taste. Both water and alcohol in sufficient quantity extract the whole of its active principles, leaving the insoluble part perfectly insipid. It appears to contain a bitter resin and mucus.

*Medical properties and uses*.—Common Centaury is tonic and antiseptic. Before the discovery of Cinchona, it was much used for the cure of fevers, and was one of the ingredients of the celebrated Portland-powder.<sup>1</sup> It is a useful bitter and tonic; and may well supply the place of some of the more expensive remedies of this description in dyspeptic complaints. The dose of the powder is from ʒss. to ʒj.; and of an infusion, made by macerating ʒvj. of the dried tops in Oss. of boiling water, and straining, from fʒjss. to fʒij. may be given three or four times a-day.

CINCHONA.<sup>2</sup> *Spec. Plant. Willd.* i. 957.

Cl. 5. Ord. 1. Pentandria Monogynia. *Nat. ord.* Contortæ, *Linn.*  
*Rubiaceæ, Juss.*

G. 346. *Corolla* funnel-shaped. *Capsule* inferior, two-celled, bipartite, with a parallel partition. *Seed* winged.

\* *Corollas* downy, with the *stamens* included.

<sup>1</sup> It is amusing to observe the fate of the various specifics for the gout, which have each held for a time its sway over the public opinion. The following were the ingredients of the Portland powder: equal quantities by weight of the roots of birthwort (*Aristolochia rotunda*), and of gentian (*Gentiana lutea*); the tops and leaves of germander (*Chamædrys*), ground pine (*Chamæpytis*), and lesser centaury (*Chironea Centaurium*), powdered and mixed together. Regarding its effects, Heberden says, “Dum fama ejus vigeret, in tot ægris, qui eo usi sunt, podagra vel mitior facta est, vel rarius repetit, ut vix possit dubitari hos effectus isti medicamento esse tribuendos. Quod autem ulla mala ex illo orta sint, præter fastidium, quod modus nimius attulit, mihi quidem nec certum unquam visum est, nec verisimile.” *Comment. de Morborum Hist.* 50.

<sup>2</sup> Supposed to be named after the Countess del Chinchon, wife of a viceroy of Peru, who introduced it into Europe, on her return to Spain in 1640.



*Species 1. C. Lancifolia*, Mutis. *Papel Periodici de Santa Fé*, p. 465. *Rhode. Monog. Cinchonæ Generis Tentamen*, p. 513. *Zea, Annales de Historia Natural*, tom. ii. p. 207.<sup>1</sup> *Flora Peruv.* tom. ii. p. 50. t. 191. *Humboldt, Plantæ Æquinoctiales*, p. 33. t. 10. *Lambert's Description of the Genus Cinchona*, plate 1. *ibid. Illustration of the Genus*, &c. p. 2.

*Species 2. C. Oblongifolia*, Mutis. *Per. de Santa Fé. Zea*, l. c. ii. 211. *C. Magnifolia*, *C. lutescens. Flor. Peruv.* ii. 53. t. 196. *Quinologia*, art. vi. 71.

*Species 3. C. Cordifolia*, Mutis. *Per. de Santa Fé. Zea*, l. c. ii. 214. *C. purpurea. Flor. Peruv.* tom. ii. 52. t. 193. *C. ovata. Ruiz. Quinologia. C. micrantha. Flor. Peruv.* 52. t. 194. *Lambert*, p. 21. plate ii.<sup>2</sup> *Illustration*, &c. p. 3.

This important genus, of which twenty-four species have been described, is not yet altogether freed from the ambiguity which has so long involved it; and although much has been effected by the industry of the Spanish botanists, whom their government sent out to make enquiries concerning it, yet many species remain undescribed<sup>3</sup>, from which it is very probable the bark-gatherers collect some part of the large cargoes which are annually sent to Europe. The three kinds medicinally used have been distinguished and named as above by Mutis, a celebrated botanist, who resided in the neighbourhood of Santa Fé de Bogota, as director of the exportation of bark<sup>4</sup>; and his observations have been fully detailed by his pupil Zea; whilst the travels of Humboldt and Bonpland have afforded them an opportunity of ascertaining accurately, and describing the species first delineated by Condamine in 1738, in the *Mém. de l'Academie*<sup>5</sup>, and named by Linnæus *officinalis*, un-

<sup>1</sup> Zea adds the following synonymes: *Quinquina*, Condam. A. A. Paris, 1738. *C. officinalis*, Linn. Syst. Veg. ed. 10. p. 929. Spec. Plant. p. 244. Gen. Plant. ed. 7. p. 91. *C. officinalis*, Vahl. Act. Soc. Nat. Haum. 1. fasc. p. 17. t. 1. *C. nitida*, Flora Peruv. et Chil. ii. p. 50. t. 191.; and *Ruiz*, Quinologia, art. ii. 56. *C. lanceolata*, Flora Peruv. 51. *C. glabra*, Ruiz. Quinol. art. iv. 64. *C. rosea*, Flora Peruv. 54. *C. fusca*, Ruiz. Quinol. art. viii. 77.

<sup>2</sup> Besides these synonymes, Zea adds, *C. officinalis*, Linn. Suppl. p. 144. s. v. edit. Persoon, p. 222. *C. pubescens*, Act. Haum. 1. fasc. p. 17. t. 2.

<sup>3</sup> In a large collection of dried specimens of the genus *Cinchona* in my possession, which were collected in 1805, both near Loxa and Santa Fé, I find many species which are not mentioned in the works of any of the Spanish botanists; nor even by Mr. Lambert, to whom I gave specimens of many of the species.

<sup>4</sup> Mutis is a native of Cadiz, and went to Santa Fé in 1760, as physician to the viceroy Don Pedro Misia de la Cerda. He discovered the *Cinchona*, in the forests between Guaduas and Santa Fé, in 1772: although the credit of this discovery was attempted to be wrested from him by Don Sebastian Jose Lopez Ruiz; who, however, from his own documents, transmitted by his brother to Baron Humboldt, to prove the priority of his discovery, appears to have known the *Cinchona* about Honda, only since 1774.

<sup>5</sup> Condamine made the first, and the only attempt that has been made to bring young *Cinchona* trees alive to Europe. He nursed them for eight months, du-



der which term, however, no less than two very distinct species were confounded by that distinguished naturalist. Under this trivial name, *officinalis*, also, the British pharmacopœias placed, as varieties, the three kinds of barks known in the shops; and this error is still retained by the Dublin college: but, in the last editions of their Pharmacopœias, the London and the Edinburgh colleges have adopted the arrangement and names of the three officinal species according to Mutis.

Prior to the year 1772, all the *Cinchona* bark brought to Europe was shipped at the ports of the Pacific; but since Don Jose Celestino Mutis discovered the *Cinchona* about Santa Fé, much of it has come by the way of Carthagena de Indias to Cadiz.<sup>1</sup> Before describing the officinal species, it is proper to state, that although they are named from the form of their leaves, yet, as Humboldt justly remarks, “no tree varies more in the shape of its leaves than the *Cinchona* ;” and, in examining dried specimens, he who has not seen them in their native forests, “will be led to discover different species by leaves which are of one and the same branch :” a remark which I am enabled to confirm, by the extensive collection of dried specimens of the genus in my possession.

1. CINCHONA LANCIFOLIA, *Mutis*.<sup>2</sup>

*Officinal*. CINCHONÆ LANCIFOLIÆ CORTEX, *Lond. Edin.* CORTEX PERUVIANUS, *Dub.* Lance-leaved *Cinchona*. The *Pale Bark* of the shops.<sup>3</sup>

*Syn.* Quinquina (*F.*), Chinarinde (*G.*), Kina (*Dutch*), Kinabark (*Dan. & Swed.*), China (*I.*), Quina Naranjada (*S.*), Quina (*Portug.*).

This tree is found chiefly in the neighbourhood of the village Ayavaca, at heights from 6250 to 8300 feet, where the mean temperature varies between 59 and 62 degrees, on a bottom of micaceous schist in the woods of Caxanuma,

ring a passage of 1200 leagues, but they were washed out of the boat into the sea and lost, near Cape Orange, north of Para. *Lambert's Illust. of the Genus Cinchona*, 4to. 1821. p. 24.

<sup>1</sup> Humboldt informs us, that the quantity of *Cinchona* bark annually exported from America is 12,000 or 14,000 quintals. The kingdom of Santa Fé furnishes 2000 of these, which are sent from Carthagena; 110 are furnished by Loxa; and the provinces of Huamanga, Cuença, and Jean de Bracamoros, with the thick forests of Guacabamba and Ayavaca, furnish the rest, which is shipped from Lima, Guayaquil, Payta, and other ports on the South Sea. *Plantæ Æquinoc.* p. 34. Perhaps more bark is used in Rome than in any other part of Europe. According to M. Valentin, (*Voy. Med. en Italie*, 1820), 4000 lbs. are consumed annually in the Hospital, *Lo Spirito Santo*; and about 10,200 lbs. in the whole city.

<sup>2</sup> *Synonimes.* *C. nitida*, Ruiz. *C. Condamine*, Humb. *C. angustifolia*, Ruiz. *C. hirsuta*, Lopez.

<sup>3</sup> I have to acknowledge my error, in regarding this as the yellow bark of the shops, in the first and second editions of the *Dispensatory*. The mistake arose from my imperfect knowledge of the Spanish language at that time; and the ambiguity of the term *Amarilla*, which may be translated either *pale* or *yellow*.



Uritusinga, Villonaco, and Monge.<sup>1</sup> It is a lofty, handsome tree, always in leaf; from thirty to forty feet in height, and standing generally single; whereas most of the other species are found in groups. It exudes, wherever it is wounded, a yellow astringent juice. The trunk is about eighteen feet in height, and fifteen inches in diameter, erect, with a cracked blackish-brown or ash-coloured bark: the branches are round, in opposite pairs, erect, brachiated; with the younger ones obscurely quadrangular at the nodes. The leaves are of a lively green, shining, ovate-lanceolate, about four inches long, with a little pit in the axillas of the nerves on the under surface, which is filled with an astringent, aqueous fluid, and having the orifice shut with hairs: they stand on purplish footstalks one-sixth of their length, flat above, and convex below; but the form of the leaf varies according to the altitude at which it grows; particularly before the tree comes into flower. The stipules are two, supra-axillary, acute, silky, contiguous, and caducous. The flowers, which appear in May, June, and July, are odorous, of a whitish rose colour, and furnished with little bractes; appear in terminal, brachiated, leafy, trichotomous panicles, supported on round peduncles and pedicels, which are powdered and silky. The calyx is of a globular bell-shape, five-toothed, powdered and silky like the peduncles, with the teeth ovate, acute, very short, contiguous, and violaceous. The corolla is somewhat salver-shaped, longer than the calyx, with the tube obscurely pentagonous, silky, more frequently of a rose colour; the *limb* wheel-shaped, with linear lanceolate segments, much shorter than the *tube*, white and woolly above. The anthers are twice the length of the free portion of the filaments, and the free parts are two-thirds shorter than the adherent.<sup>2</sup> The germen is globular, changing to an ovate, woody, longitudinally ten-striated capsule, crowned with the calycinal teeth, two-celled, many-seeded, oppositely twice-furrowed, and opening from the base to the apex with two valves.

This tree affords the original *Cinchona* of Peru, which is now very rare, 110 quintals only being cut, instead of 4000, which was the quantity cut in 1779, and reserved for the use of the Spanish government.<sup>3</sup> Zea says it is the *lancifolia* of

<sup>1</sup> As the *Condaminea* of Humboldt, notwithstanding the reasonings of this author to prove the contrary, is evidently a variety, if not the same species meant by Mutis, we have availed ourselves of Mutis's accurate description.

<sup>2</sup> Humboldt.

<sup>3</sup> Estan raro, que apenas corresponde á uno par mil de las otras especies juntas. *Ann. de Hist. Nat.* tom. ii. p. 210.



Mutis, under which we have placed it; and there is also a great affinity between it and the *scrobiculata* of Humboldt, according to that celebrated traveller. The bark of the *lancifolia* is the pale bark of the shops, the *Quina Naranjada*, *Cascarilla fina de Uritusinga*, and *Quino fino* of the Spaniards. It is known in commerce by the name of *Calisaya*<sup>1</sup>; and is preferred in South America to all the other Cinchonas. Two other varieties of it, probably produced by distinct species, are also known in commerce by the names of *lagatijada* (lizardlike), and *negrilla* (blackish), from the colour of their epidermis. It has always been known in this country by the vague name of Peruvian or officinal bark. The branches are decorticated in the dry season, from September to November, which is the period when all the kinds are barked, and the bark is carefully dried in the sun. The trees frequently die after the operation.

The bark arrives in Europe packed in chests made of slips of wood roughly fastened together, and covered with skins; each of which contains about 200lbs. weight, well packed, but generally mixed with a quantity of dust and other heterogeneous matter. It consists of pieces eight or ten inches in length, some of them scarcely one-tenth of an inch in thickness; singly and doubly quilled, or rolled inward, the quills, generally, being in size from a swan's quill to an inch and a half<sup>2</sup>; and others of a coarser texture, thicker and nearly flat. It has a chopped, greyish or cineritious epidermis, often covered with flat, sometimes stringy lichens<sup>3</sup>; and is internally of a pallid fawn or cinnamon hue. This colour is brightened when the bark is moistened, approximating to pale orange. Both the quilled and the flat varieties are evidently the bark of the same tree; the quilled sort being that of the smaller branches, and the flat that of the larger and of the trunk. But the chests probably contain similar barks obtained from different species.

*Qualities.* — Good bark of this description has scarcely any odour when in substance; but during decoction the odour is sensible, and agreeably aromatic. The taste is bitter, but not

<sup>1</sup> The name *Calisaya* is that of a province producing this bark, in the most southern part of Peru, in the Intendencia de la Paz; but the term is also used as a generic name by which the Peruvian Indians distinguish the superior barks. There are three varieties of *Calisaya* known in South American commerce, 1. *Calisaya arrollenda*, rolled *Calisaya*; 2. *Calisaya de plancha*, flat *Calisaya*; 3. *Calisaya de Santa Fé*, which is a thick bark. MSS. of Dr. Devoti.

<sup>2</sup> The great desire of our bark merchants to procure quilled bark has induced the bark-gatherers often to produce this effect by heat, which always diminishes the virtue of the bark. MSS. of Don Felix Devoti of Lima, in my possession.

<sup>3</sup> On this account the inhabitants of Peru name it *Quinacana*, hairy Quina.



unpleasant, slightly acidulous and austere, resembling in some degree that of a dried rose. It is light, and breaks with a close fracture, with the internal fibres somewhat drawn out. The powder of the quilled kind is paler than the bark, being of an uniform pale cinnamon hue; but the flat kind yields a deeper-coloured and browner powder. The best specimen of this bark which could be procured by me, and subjected to experiment, gave the following results: Water at  $212^{\circ}$  extracted all its active principles; affording an infusion, which, when filtered, was of a pale yellow or straw colour, and had the odour and taste of the bark. The infusion reddened litmus paper; was instantly and copiously precipitated by solution of galls; and, in a smaller degree, and more slowly, in yellowish flocculent flakes, by solution of isinglass. A solution of tartar emetic was rendered turbid, and slowly precipitated by it; but this effect was quickly and copiously produced on a solution of acetate of lead. Sulphate of iron changed its colour to bright olive-green, but was scarcely precipitated. Decoction afforded a more saturated solution, with a colour resembling the cold infusion of the yellow bark; and a yellowish precipitate was deposited. The powder macerated in sulphuric ether afforded a golden-yellow tincture, which reddened litmus paper, and left a pellicle of bitter resin when evaporated on the surface of water, to which it gave the colour of the tincture. This coloured water had the flavour of the watery infusion, but differed from it, in not precipitating the solution of galls and that of tartar emetic; and in throwing down a copious precipitate from the solution of sulphate of iron. With alcohol the powder afforded a tincture of a deep orange hue, which precipitated sulphate of iron, tartarized antimony, and tannin; became turbid when added to water, and let fall a light reddish precipitate. From the effects of these re-agents on the aqueous infusion of this bark, it appears to be the same as the 3rd and 15th species examined by Vauquelin; which he names *superior grey Cinchona*, and *common Cinchona of Peru*.<sup>1</sup> But these experiments throw no light on the nature of the active principle of this species of cinchona; and it was not until the subject was investigated by MM. Pelletier and Caventou, in 1818, that that principle was ascertained to be an alkaline substance, which has been named *Cinchonia*, combined with a peculiar acid, the Kinic. It is obtained by boiling a pound of bruised *Cinchona lancifolia* in a gallon of water, previously mixed with three fluid drachms of sulphuric acid. On completing this decoction, the residue should be again boiled with half the quantity of water and acid, and so on until all the

---

<sup>1</sup> *Annales de Chimie*, lix, 116.



soluble matter be extracted. To these decoctions mixed together, quick lime is to be added to saturate fully the acid, and the precipitate, which is a mixture of sulphate of lime and cinchonia, being dried, is boiled for some minutes in strong alcohol, which is decanted off whilst it is hot. The residue is again to be boiled in fresh alcohol, and these boilings repeated until the alcohol ceases to act on it. By evaporating these alcoholic solutions in a retort, impure crystals of cinchonia are deposited. These are purified by dissolving them in boiling alcohol and recrystallization. In this state the crystals are acicular and white, have an intensely bitter taste, and are scarcely soluble in water, alcohol, or ether, at common temperatures: when heated to  $450^{\circ}$  in a glass tube, cinchonia fuses like a resin, and on increasing the heat is decomposed, evolving ammonia, and depositing carbon: from which it appears to be a compound of carbon, hydrogen, and nitrogen.<sup>1</sup>

Cinchonia is also found, in small quantities, in the two other officinal species of bark; in Cusparia bark, and some other vegetable productions.

Mutis and Zea regard this species of *Cinchona* as directly febrifuge; as chiefly applicable in intermittent fevers of long standing; and also assert, that it never fails to cut short an ague when administered at its accession.<sup>2</sup>

## 2. CINCHONA OBLONGIFOLIA. *Mutis*.<sup>3</sup>

*Officinal.* CINCHONA OBLONGIFOLIAE CORTEX, *Lond. Edin.* CORTEX PERUVIANUS, *Dub.* Oblong-leaved Cinchona Bark. Red Bark.  
*Syn.* Quinquina rouge (*F.*), Rothe Chinarinde (*G.*), China (*I.*), Quina Roxa (*S.*), Quina Vermetha (*Portug.*).

The tree yielding this bark is found on the Andes, growing in the woods on the banks of the mountain-streams in great abundance, at Riobamba, Chinchao, Cuchero, and Chacahnassi; flowering in June and July. It rises to a very considerable height on a single, erect, round stem, which is covered with smooth, brownish, ash-coloured bark. The older branches are round, smooth, and of a rusty colour; the younger are obtusely four-cornered, leafy, and of a diluted reddish colour. The leaves are opposite, large, the full-sized ones being one or two feet in length, of an oblong oval shape, and supported on short, semi-round, purple petioles. They are entire, pale, and shining on the upper surface, on the under marked with veins that turn to a purplish colour; and at the base of each are numerous bundles of white bristles: the stipules are supra-axillary, interfoliaceous, opposite, contiguous, united at the base, and of an

<sup>1</sup> See Mr. Brande's experiments. — *Quarterly Journal of Science*, vol. xvi. p. 281, 282.

<sup>2</sup> *Annal. de Historia Natural.* ii. 609.

<sup>3</sup> *Synonymes.* *C. Magnifolia*, *Flor. Peruv.* Lambert denies its affinity with *Magnifolia*. See *Illustration*, &c. p. 12.; but from the specimens in my possession I cannot admit this opinion.



obovate figure. The flowers appear in large, erect, much compounded, terminal panicles, somewhat branched, on long, brachiated, many-flowered peduncles; the calyx is small, five-toothed, and of a purple colour; the corolla white and odorous, with the limb spreading, and hairy within: the filaments are very short, inserted into the tube of the corolla, supporting oblong anthers, bifid at the base, and are situated below the middle of the tube of the corolla. The capsules are large, oblong, obscurely striated, slightly curved, and crowned with the calyx.<sup>1</sup>

This tree is named in the vernacular Spanish, *Palo de requeson*, and *Cascarilla de flor de Azahar*, from the flowers resembling in odour those of the orange. Its bark is the *Quina roxa* and *colorada* of commerce. The bark is brought to this country in chests, which contain from 100 to 150 lbs. each. It consists of various sized pieces, covered with a thin and rough, entire, reddish-brown epidermis. The greater number of the pieces are flat, but some are partially quilled, as if taken from half the circumference of the branches to which they belonged. Under the epidermis there is an intermediate layer, which is dark coloured, compact, brittle, and seemingly resinous; and within it the internal part is woody, fibrous, and of a rust-red colour. The fracture, examined by a lens, consists of close, longitudinal, parallel, needle-form fibrillæ of a pale red colour, with a deep-red, agglomerated powder in the interstices. The powder is of a deeper colour than the internal part of the bark.

*Qualities.* — Red Cinchona bark has a weak peculiar odour; and its taste is much less bitter, but more austere and nauseous, than the barks of the other species. The aqueous infusion has a pale ruby colour, a slight degree of bitterness, and a decided astringency. It lets fall a sediment of a brighter hue than that of the dry powder. It reddens litmus paper<sup>2</sup>, is slowly precipitated by the solution of galls, the supernatant liquor being perfectly colourless; and a very light, flocculent, ruby-coloured precipitate is produced by the solution of isinglass. It is not altered by tartarized antimony, nor by the acetate of lead; and the sulphate of iron makes it assume a dirty, yellow-olive colour only, little being precipitated. The ethereal tincture is of the same colour and exhibits the same appearances as that of the former species, when treated in a similar manner. The alcoholic is of a very deep brownish red colour; when diluted with water a red flocculent matter falls down; and it precipitates the solutions of sulphate of iron, and of tartarized antimony, the former of a black colour, and the

<sup>1</sup> *Flora Peruv.* ii. 53. t. 196.

<sup>2</sup> Fourcroy found in it a portion of citric acid, some muriate of ammonia, and muriate of lime. See *Thomson's Chem.* v. 216.



latter red. It comes nearest to the second species examined by Vauquelin, which he calls *Santa Fé Cinchona*; and differs from his *Cinchona magnifolia* in reddening litmus paper, and precipitating tannin. Its virtues depend on two alkaline substances; one the *Cinchonia* already described, the other *Quinia*, about to be described in the account of the next species; and both are combined with kinic acid.

This bark was introduced by Don Sebastian Josef Lopez Ruiz, in 1778; and is considered by Zea and Mutis as the least directly febrifuge of the three kinds which are described in this work. It possesses powerful astringent and antiseptic properties; consequently its use is contra-indicated in inflammatory and bilious affections: but the Spanish physicians regard it as highly beneficial as an external application in suppurating and sphacelating ulcers. An extract prepared from it is much used in Spain, in putrid fevers.

### 3. CINCHONA CORDIFOLIA. *Mutis*.<sup>1</sup>

*Officinal*. CINCHONÆ CORDIFOLIÆ CORTEX, *Lond. Edin.* CORTEX PERUVIANUS, *Dub.* Heart-leaved Cinchona. The yellow bark of the shops.

*Syn.* Quinquina jaune (*F.*), Gelbe Chinarinde, Rieberrinde (*G.*), Gul Kina-bark (*Swed.*), China (*I.*), Quina amarilla (*S.*).

The tree which affords this bark is found on the mountains of Loxa, in the kingdom of Quito, and those of Santa Fé, growing along their skirts, and on the plains, under the 4th degree of north latitude, on heights betwixt 900 and 1440 toises; flowering from May to September. It is a spreading tree, rising on a single, erect, round stem of no great thickness; and covered with a smooth bark, externally of a brownish grey colour. The younger branches are quadrangular, smooth, leafy, sulcated, and tomentose: the leaves, which are about nine inches in length, are opposite, petiolate, spreading, of an oblong oval, cordate or egg-shape, entire, shining on the upper surface, ribbed and pubescent on the under: with the petioles flat on one side, and roundish on the other, about a thumb's breadth in length, and of a purple colour; but the leaves of this species vary even more than those of *lancifolia*. The flowers appear in large, terminal, leafy panicles, supported on long, compressed, tetragonous peduncles. The calyx is five-toothed, downy, and of a dull purple colour; the corolla internally tomentose; the tube of a diluted-red colour; the limb shaggy, white above and purplish below; and the segments spreading, with reflected tips. The filaments are short, supporting linear anthers, bifid at the base, which reach as far as the upper part of the tube of

---

<sup>1</sup> *Synonymes*. *C. pubescens*, Vahl. *C. ovata*, Flor. Peruv. *C. hirsuta*, Flor. Peruv. But Lambert affirms that it is totally different from the *hirsuta* of the Flora Peruviana. See *Illustration of the Genus Cinchona*, 4to. 1821. p. 4.



the corolla. The germen is tomentose, and changes to an oblong, narrow capsule, about one inch and a half in length, marked with ten striæ, of a reddish-brown colour, and crowned with the calyx.

The bark yielded by this tree is named *Quina amarilla*<sup>1</sup>, *Cascarilla de Loxa*, and *Cascarilla amarilla*; and is the yellow bark of the shops. It is brought to this country in chests containing about 90 to 100 pounds each; and consists of pieces about eight or ten inches in length, some quilled, but the greater part flat.<sup>2</sup> The quilled pieces are less rolled and thicker than the quilled pale bark; and the epidermis, which is of a tawny, greyish-brown colour, and covered with flat and stringy lichens, is more rough and chopped, easily separating, and often as thick as the bark itself, which is about one-eighth of an inch; while the interior is of a yellow colour, passing to orange. The flat pieces are generally without any epidermis, and considerably thicker than the quilled. Both are mixed in the same chest.

*Qualities.* — Yellow bark has nearly the same odour in decoction as the pale; the taste is more bitter, but less austere, and it does not afford any astringent feeling to the tongue when chewed. The internal colour is golden cinnamon, or subdued orange-yellow, becoming when moistened a lively orange. The fracture is woody and fibrous, presenting, when examined by a lens, the appearance of parallel, longitudinal, needle-like fibres, with a dry, agglomerated powder, in the interstices, of a yellow colour. It is easily reduced to fine powder, and the powder preserves the colour of the bark, but is brighter. The sediment, which the infusion lets fall in cooling, is of a brighter colour than the dry powder. The filtered aqueous infusion has a pale golden hue, with a shade of red; is clearer, and seemingly less mucilaginous than the former: it has all the bitterness of the bark; reddens litmus paper, and precipitates solution of galls; but the precipitate does not fall so instantaneously as in the infusion of the former species. With solution of isinglass a pinkish-yellow precipitate is produced: acetate of lead throws down a precipitate; and that with tartarized antimony is in yellowish white flakes and more copious than that which the pale bark affords. A solution of sulphate of iron changes its colour to a bluish-green, and, after many hours, gives a precipitate of the same hue. The ethereal tincture has a golden colour, affords resin when evaporated, and is affected by the same re-agents as that of the pale *Cinchona*; but

<sup>1</sup> *Yellow bark*; but the adjective signifies both yellow and pale, or wan. The name appears to be used in contradistinction to *naranjada*, orange colour, which is applied to the first officinal species.

<sup>2</sup> These are distinguished in commerce by the terms *with coat*, and *without coat*.



the water on which it is evaporated is less highly coloured. The alcoholic tincture appears to be in every respect the same as that afforded by the pale bark. It seems to agree in most of its properties with the first species examined by Vauquelin; which he states was brought to Spain in 1788, and, owing to its having been used for the royal family, got the name of royal Cinchona. Yellow bark owes its febrifuge properties to an alkali which has been termed Quinia, and which can be separated by a process similar to that employed for obtaining cinchonia. Quinia separates from the alcoholic solution in a viscid state, and when dried in the open air, in a moderate heat, becomes brown and brittle. When properly prepared it is white, bitter, insoluble in water, but very soluble in ether. It differs from cinchonia in containing oxygen as a proximate principle, in addition to carbon, hydrogen, and nitrogen.

According to Mutis and Zea, it is indirectly febrifuge only; but when genuine, all its varieties are excellent remedies.

The goodness of all the species of Cinchona depends on the proper age of the branches that are barked. The bark-collectors (*cascarilleros*) decide on the maturity of these in the following manner. They strip off from each branch a small piece of bark; and if it immediately redden on the inner side, they consider it sufficiently mature; but should the colour be not manifested in three or four minutes, it is rejected as being not yet in season.<sup>1</sup>

As Cinchona bark occasionally varies in its powers, and is often adulterated with other inferior barks, even by the bark-peelers (*cascarilleros*) who gather it; arising either from ignorance, or from a fraudulent desire of more quickly completing their contracts<sup>2</sup>; it is of importance to be able to distinguish good bark, and the best varieties from those of an inferior description. Mutis informs us, that the old trees furnish the best bark; and that the bark taken from the trunk and thicker branches, is superior to that from the younger branches. The following directions for choosing bark are those generally attended to in South America<sup>3</sup>: The essential characteristics are *colour, taste, and smell*; the secondary or accidental ones, are *exterior coat, fracture, weight, thickness, and quill*. The best bark of the first class is of an orange yellow colour; and the goodness decreases as the colour varies from this to a

---

<sup>1</sup> *Memoir on Quinquina, by M. Laubert, Chief Physician to the Spanish Army*: translated in *Lambert's Illustration*, 4to, 1821. p. 64.

<sup>2</sup> Humboldt says, "We saw at Peru the barks of two new species of *Weinmania* and *Wintera* mixed with those of *Cinchona*." *Personal Narrative*, vol. v. p. 769. *Trans.*

<sup>3</sup> Extracted from a MS. of *Don Felix Devoti*, a respectable physician at Lima, who had practised upwards of twenty-five years in South America.



very pale yellow. When of a dark colour, between red and yellow, it is always to be rejected; as this colour designates either that it is of a bad species, or that it has not been well preserved from the air and moisture. This dark colour, however, must not be confounded with a red colour in the inside, which constitutes a distinct species. The *taste* of bark should be bitter, but not nauseous nor very astringent, with a slight agreeable acidity just perceptible to the palate; and, when chewed, it should not appear in threads of much length. The *odour* of any of the barks is not very strong; but when they have been well cured and preserved, it is always perceptible; and the stronger it is, provided it be pleasant, the better may the bark be considered. The appearance of the *coat* or epidermis has led to many mistakes. It is, in many instances, merely accidental; depending on the variation in the height of the ground, and the exposure of the branches to the sun and air. Seven distinct appearances of the epidermis are remarked: 1. Negrilla, dark silver coat<sup>1</sup>; 2. Crespilla, short curled; 3. Pardo-obscura, dark open leopard grey<sup>2</sup>; 4. Pardo-clara, light open grey<sup>3</sup>; 5. Lagatijada, fine dark silver, lizard-coloured<sup>4</sup>; 6. Blanquissima, very pale<sup>5</sup>; and, 7. Cenicienta, ash-coloured. The three first are the best, and belong to bark produced on the highest mountains: the others rank in the order of their arrangement; the epidermis being always cracked and rough in proportion as the trees have been exposed to a scorching sun. With regard to *fracture*, some of the worst barks break even and clean as if cut with a knife, and some of the best have always a more or less splintery fracture.<sup>6</sup> The fibres of the fracture being sharp and short indicate the bark to have been gathered from mature branches; the long and thread-like from immature branches. The best barks are generally observed to be the heaviest. In point of *thickness*, very thin bark is inert, owing to the branches from which it was taken having been too young; and very thick bark, particularly if it break like common wood, argues that the tree must have been sickly; yet bark exceeding a line in thickness may be good; for al-

---

<sup>1</sup> This bark is occasionally found amongst the pale Cinchona sent to England. It is easily distinguished by its spotted surface. Ruiz says it must be ranked among those of a middling quality.

<sup>2</sup> This is found mixed with the pale bark of the shops. It is regarded as of middling quality.

<sup>3</sup> This is a very rare bark, and is that of the *C. fusca* of Ruiz. It is called *Asmonich* by the natives of Puzuzu and Muna, where it is found.

<sup>4</sup> The bark with this coat has the greatest affinity with the *yellow* bark of the shops. It is a good kind of bark.

<sup>5</sup> This bark is little valued in Spain, and is seldom met with in commerce.

<sup>6</sup> The idea of a resinous fracture being the characteristic of good bark originated when the virtue of bark was supposed to depend on the resin it contained.



though it is disapproved of at Cadiz, under the name of *quinon*, yet, excellent effects have resulted from much thicker bark in England. The moderately thick and firm bark is always preferred at Lima. The *quilling* of bark arises from the manner in which it is separated from the branches. This is effected by making a longitudinal incision in the branch, and passing under the bark a very fine knife. As the slip dries, it rolls up, owing to the internal surface shrinking more than the external: a feeble rolling, therefore, denotes that the bark is rather too old, or has been too slowly dried; too much quilling, that it is either too young or has been too hastily dried. The moderate *quill* of bark certainly denotes it to be of the best kind, and that it has been taken from branches of a proper age, and well dried; but the bark-collectors often produce this effect by fire, when there is a want of sun, as is frequently the case in some parts of the mountains. The fraud is known by the colour being much darker; and, when the bark is split, the inside exhibiting stripes of a whitish sickly hue. It should be preserved in cases, well secured from the air and humidity.

Analytical examinations of *Cinchona*, with the view of discovering on what principle its febrifuge properties depend, have been made by several chemists well qualified for the task. Vauquelin divided all the different species into three sections, according to their chemical properties.<sup>1</sup> The first comprised those which precipitate tannin, but not animal gelatin; the second, those which precipitate gelatin, but not tannin; and the third, those which precipitate at the same time tannin, gelatin, and tartar emetic. He conjectured, that on the principles producing these effects, particularly that which precipitates infusion of galls, the febrifuge properties of the barks depend, and that they are more or less febrifuge, in proportion to the quantity of these principles that are present. He asserted that the principle which precipitates tannin is of a brown colour and bitter taste; is less soluble in water than in alcohol; and that it also precipitates tartarized antimony, but not glue.<sup>2</sup> He imagined that it possessed some analogies with the resinous bodies, although it furnished ammonia on distillation; whilst the principle which, in some *Cinchonas*, precipitates glue, has a bitter and astringent taste; is more soluble in water than the principle which, in other

---

<sup>1</sup> He examined seventeen different kinds, but was not able to ascertain the names of the trees from which they were obtained.

<sup>2</sup> The effect of this principle was first noticed by Dr. Maton; and soon after by Seguin, who immediately concluded that it was gelatine; but this opinion was proved to be erroneous by Dr. Duncan, jun. who found that it was a principle *sui generis*, and named it *cinchonin*. Vide *Nicholson's Journal*, vii. 226.



kinds precipitates tan: it is also soluble in alcohol, and does not precipitate tartar emetic.<sup>1</sup> Fabroni conceived, that he was authorised in concluding from his experiments, that “the febrifuge virtue does not belong essentially and individually to the astringent, the bitter, or any other soluble principle, as the quantity of these increases by long boiling, while the virtues of the decoction decrease. Neither does the febrifuge virtue reside in that principle which destroys the emetic property of tartarized antimony, and precipitates iron, since the decoction contains more of it than the infusion, while its virtues are evidently less.”<sup>2</sup> It was concluded from these doubts, and many others that had been raised, that much was yet to be done before the principle of Cinchonas effective in the cure of fevers could be ascertained, and time has displayed the truth of this opinion.<sup>3</sup>

The latest analysis of the Cinchona barks, is that of MM. Pelletier and Caventou. The following are the components of the three officinal species: 1. In *pale bark* they found a peculiar acid, which had been previously found by Vauquelin, and named Kinic acid, combined with *Cinchonia*, forming an acidulous kinate of Cinchonia, a green fatty matter, a red nearly insoluble colouring matter, which they term red cinchonic, tannin, a yellow colouring matter, kinate of lime, gum, starch, and woody fibre. 2. In *yellow bark* they found that the alkaline base differs from cinchonia, in being incrySTALLIZABLE, very soluble in ether, and forming salts with the acids different from those formed by cinchonia, and which has been termed *Quinia*. The components of yellow bark are, an acidulous kinate of Quinia, a deep-yellow fatty matter, red cinchonic, tannin, yellow colouring matter, kinate of lime, starch, and woody fibre. 3. *Red bark* contains acidulous kinate of Cinchonia, kinate of Quinia, reddish fatty matter, red cinchonic, tannin, kinate of lime, yellow colouring matter, starch, and woody fibre.

We may venture to state the following as the known active

<sup>1</sup> *Annales de Chimie*, l. c.

<sup>2</sup> *Edinburgh Medical and Surgical Review*, ii. 338.

<sup>3</sup> In consequence of a chemical theory of the mode in which Cinchona acts on the living body, Fabroni made some curious experiments to ascertain the relative affinity of different Cinchonas to oxygen. In imitating his experiments with the three officinal species, I found that, when half a drachm of each of these barks in powder was separately mixed with half a fluid ounce of strong nitric acid, in similar vessels, the temperature of the atmosphere at the time being 70°, and that of the acid 71°, in the space of four minutes, the heat produced rose the mercury in the thermometer as follows:

Common pale bark,	— to 120°
—— yellow bark,	— to 123°
—— red bark,	— to 119°

The mixture in each vessel was gradually swollen as the heat increased, and nitrous fumes were given out, showing the evident decomposition of the acid.



constituents of Cinchonas; *Cinchonia* or *Quinia*, according to the species of bark employed, united with the Kinic acid<sup>1</sup>, *kinate of lime*, *resin*, *extractive*, *gluten* or *ferment*, *volatile oil*<sup>2</sup>, and *tannin*. I separated the *resin* in a pure state by evaporating the ethereal tincture on the surface of cold water; and the *gluten* Fabroni found was separable by water, occasioning the spontaneous fermentation of the decoction and infusion in summer, and decomposable by fermentation. MM. Alibert and Cabal demonstrated the presence of iron in Cinchona, by incinerating the bark, dissolving the ashes in nitric acid, and adding prussiate of potass, which precipitated prussiate of iron.

The two alkaline bodies, *Cinchonia* and *Quinia*, found in these barks, unite readily with the acids, particularly the sulphuric; and their sulphates are now frequently employed instead of the entire barks or their other preparations.

The *Sulphate of Cinchonia* is so little employed, on account of the expense of preparing it, that I shall merely describe its characters and composition. It crystallizes in short rhomboidal prisms, which are soluble in 6.5 of alcohol of sp. gr. 85., and in 54 parts of water at a temperature of 60° Fahr. According to M. Beup's experiments, these crystals consist of cinchonia 84.324, sulphuric acid 10.811, and water 4.865, or of one atom of cinchonia, one of acid, and two atoms of water.<sup>3</sup>

The *Sulphate of Quinia* is prepared by boiling, for half an hour, one pound of powdered yellow Cinchona bark in eight pints of distilled water, acidulated with one fluid ounce of sulphuric acid. After straining off the decoction, boil the residue in a similar quantity of acidulated water; then mix the strained decoction with the former, and stir into the mixture, by small portions at a time, four ounces of powdered lime. A reddish brown flocculent precipitate will fall down, which is to be separated, by throwing the whole on a linen cloth. Wash this precipitate with a little, cold, distilled water, and having dried it, digest it in rectified alcohol, in a moderate heat, for some hours. Decant off the spirit and repeat the digestions until the spirit cease to acquire a bitter taste; and, having mixed the spirituous solutions, distil the whole in a water-bath until three-fourths of the alcohol shall have passed over; and a brown viscid substance, covered with a bitter, alkaline liquid, remains in the retort. Pour off the alkaline liquid and saturate it with sulphuric acid; then having reduced it, by evaporation, to half the quantity, add a small portion of charcoal, and

---

<sup>1</sup> *Annales de Chimie*, lix. l. c. The name of the acid is derived from *kina*, *kina*, an old appellation of the bark. Dr. Duncan proposes to call it *cinchonic acid*, as the present name would lead to the supposition that it is procured from kino.

<sup>2</sup> Dr. Irwin first obtained a small portion of this oil.

<sup>3</sup> *Ann. de Chimie*, xxvii. p. 323.



after boiling for a few minutes, filter the fluid while hot and set it aside to crystallize. The brown viscid mass is to be boiled in a small quantity of water slightly acidulated with sulphuric acid, by which means a large portion of it will be converted into sulphate of Quinia. The crystals obtained from both the above parts are to be dried on bibulous paper. Sulphate of quinia thus prepared is in small, silky, acicular crystals, soluble in 740 parts of water at 55° Fahr., in 30 parts at 212°; and in 60 parts of alcohol of sp. gr. 85. It effloresces in the air, losing three-fourths of the water of crystallization; and should always be used in that state. It consists, in its crystallized state, of 61.644 parts of quinia, 13.698 of sulphuric acid, and 24.658 of water in 100 parts: or one atom of quinia, two atoms of acid and sixteen of water.<sup>1</sup>

*Medical properties and uses.* — Cinchona bark is a powerful and permanent tonic, possessing also antispasmodic and antiseptic powers; and is, undoubtedly, superior to all other remedies in counteracting febrile action, and restoring strength and vigour to morbidly weakened habits.

The stories which are related regarding the discovery of its febrifuge effects appear to be founded on fiction, and are unworthy of notice. The Peruvians, it has been supposed, were acquainted with its powers before the conquest of their country by the Spaniards, and from them the knowledge of it might have been acquired by their conquerors: but Humboldt renders this idea improbable, and says, that the use of the cinchona bark “is entirely unknown to the Indians in Loxa, Guaneabamba, and far around.”<sup>2</sup> They even regard it as poisonous; and “in Malacatis only, where many bark-peelers live, they begin to put confidence in the Cinchona bark.”<sup>3</sup>

The most probable history of the discovery of the febrifuge virtues of cinchona, is the following tradition, mentioned by Humboldt, in his Dissertation on the Cinchona Forests.<sup>4</sup> The Jesuits, at the felling of the wood, had taken notice of the considerable bitterness of the Cinchona, and, “there being always medical practitioners among the missionaries, it is said they had tried an infusion of the Cinchona in the tertian ague, a complaint which is very common in that part of the country;”

<sup>1</sup> *Ann. de Chimie.* xxvii. p. 323.

<sup>2</sup> *Humboldt on the Cinchona Forests*; in *Lambert's Illustration of the Genus Cinchona.* Lond. 1821, 4to. p. 22.

<sup>3</sup> Humboldt says that the present people of South America have the most inveterate prejudices against the employment of the different kinds of Cinchona: and in the very country where this valuable remedy grows, they try to cut off the fever by infusions of *Scoparia dulcis*, and hot lemonades prepared with sugar and the small wild lime, the rind of which is equally oily and aromatic. *Personal Narrative*, vol. v. p. 164. *Trans.*

<sup>4</sup> This salt was first noticed by M. Descamps, an apothecary at Lyons, who erroneously ascribed to it the febrifuge property of bark.



and having found it succeed in curing the disease, began to employ it as a febrifuge.

It was, nevertheless, little known by Europeans, until the countess of Chinchon, wife of Don Geronimo Fernandez de Cabrera Bobadella y Mendoza, count of Chinchon, viceroy of Peru, introduced it into Europe, on her return to Spain, in 1640. Its fame soon spread, and it was taken to Italy in 1649<sup>1</sup>, and through the means of Cardinal De Lugo and the Jesuits, was distributed over the continent.<sup>2</sup> It was in repute in England in 1658; but owing to its high price<sup>3</sup>, and some prejudices formed against it, it was very little used, till Talbot, an Englishman, again brought it into vogue by the many cures he performed with it in France, under the name of the *English remedy*. His secret of preparing and exhibiting it was purchased by Louis XIV. and made public. These circumstances throw light on the origin of some of the names by which it has been known: as, *Cortex* and *Pulvis Comitissæ*; *Cortex* and *Pulvis de Lugo*; and *Pulvis Jesuiticus* or *Pulvis Patrum*. It was called also, *Palos de calentura*, or fever wood, on account of its effects; and, from the place whence it was brought, *Peruvian bark*.

It was introduced into practice for the cure of intermittent fever. It was much opposed at first<sup>4</sup>, but soon acquired great reputation; and it still retains the reputation it acquired as a remedy for that disease; although, owing to peculiar idiosyncrasies and other accidental causes, it has occasionally failed in this country in agues, which were afterwards removed by other remedies, particularly arsenic. Some of these failures may perhaps have arisen from the kind of the bark employed: for notwithstanding the generally received opinion, that all the kinds of bark may be indifferently used, one for another, yet there is some reason for the assertions of the Spanish and American physicians, that they vary in other respects besides their degree of activity. By them the pale bark, *calisaya*, *quina*, *naranja-*

<sup>1</sup> The earliest printed treatise on the medical properties of Peruvian bark, with the exception of the *Schedion*, an Italian handbill printed in 1650, recommending it, and giving directions for exhibiting it, is that of *Joannus Jacobus Chifletius*, entitled "*Pulvis Febrifugus orbis Americani*." 4to. pp. 30. Anno 1653. Chifletius was physician to the Archduke Leopold William.

<sup>2</sup> Morton gives the above account on the authority of Bollus, a Genoese merchant, who had lived long in Peru, "*autor fide dignus*." *De Febribus Intermit.* c. vii.

<sup>3</sup> It was sold at first by the Jesuits for its weight in silver; yet Condamine relates that, in 1690, several thousand pounds of it lay at Piura and Payta for want of a purchaser. *Mémoires Acad. Roy.* 1738.

<sup>4</sup> Among others, it was opposed by Chifletius in the work already noticed, and also in a Tract, published in 1655, in defence of Chifletius, by Melippus Prolimus, against one Antimus Conygius, who had ventured to extol the bark.



*da*<sup>1</sup>, is considered as directly febrifuge, and the best adapted for the cure of ague; the yellow bark, *quina amarilla*, as only indirectly so, and better fitted for slow fevers and chronic debilities: while the red, *colorada*, *quina roxa*, is only fit to be used in cases of gangrene<sup>2</sup>, as its use is apt to be followed with disgusting nausea, severe vomiting, and insupportable colic. The differences of opinion with regard to the best time of giving it are now nearly settled. Boerhaave<sup>3</sup> and others recommended that the fever should be allowed to run on for some time before it was administered; but it is now generally agreed that the bark cannot be given too early after the stomach and bowels are cleared by an emetic and cathartic. Dr. Cullen recommended the exhibition of it in a large dose or doses immediately before the accessions<sup>4</sup>; but Morton's method of giving it directly after the hot stage of the paroxysm ceases, and repeating it in increased doses during the intermission, until the cold stage again returns, is now generally adopted. It may be safely given, however, during the paroxysm, as practised by Dr. Clarke of Newcastle, but many stomachs are apt to nauseate it at that time.

In remittent fevers, *Cinchona* is found equally efficacious; but the excitement, however, particularly in the remittents of warm climates, requires to be previously subdued by blood-letting, and the bowels to be kept open. It renders the remissions distinct, and by degrees checks altogether the febrile action. In other affections, depending on a similar state of habit, as, hemicrania, periodical pains, spasms, chorea, hysteria, epilepsy, passive hæmorrhagy, and in habitual, frequently returning coughs, it is also found useful: but it does not prevent the continuance of those paroxysms of ague which form one of the constitutional symptoms of stricture of the urethra, and some other local affections; and which can be cured only by removing the strictures and other sources of irritation.

In the low stage of continued fevers of the typhoid type, particularly when these are attended with symptoms of putridity, as in jail-fever, cynanche maligna, scarlatina maligna, confluent small-pox, and in putrid measles, the bark must be regarded as one of the most valuable remedies. The administration of it in pure typhus has been of late years judiciously

<sup>1</sup> According to Condamine, this was the bark first introduced into Europe. He says it yields by incision a yellow odorous resin; and that the Jesuits of La Paz (whence the best bark of this species is still obtained) used to gather it with care, and send it to Rome, where it was specific in agues. But the Loxa bark coming to Europe soon after, the three kinds were confounded together.

<sup>2</sup> Zea, *Annales de Hist. Nat.* l. c. Rushworth discovered the efficacy of the red bark in gangrene.

<sup>3</sup> *Aphorismi* &c. 767.

<sup>4</sup> *Mat. Med.* ii. 97.



delayed until the increased excitement is presumed to be subdued, and symptoms of great debility make their appearance; or until the morbid heat be carried off, and the skin opened. Several eminent modern physicians<sup>1</sup>, however, recommend it to be given early in the disease, and persevered in; but from our own experience we are inclined to consider the former the safer practice, and believe that the best effects will be produced from the Cinchona, when its use, in pure typhus, is not begun till the skin becomes moist, the tongue is in part cleaned, and the urine deposits a critical sediment. The best adjuncts in these cases are the diluted sulphuric, or the muriatic acids and aromatics, particularly the tincture of capsicum.

Cinchona was first conjectured to be useful in gout by Sydenham, and in some cases its efficacy is sufficiently evident. In rheumatism, also, Dr. Haygarth has lately strongly recommended it to be given, after the manner of Morton, Hulse, and Fothergill, from the commencement of the disease; the stomach and bowels being previously emptied by means of antimonial preparations. In my own practice, I have found it useful only after the liberal exhibition of calomel, tartarized antimony, colchicum and opium, when the pain has partly abated, or assumed an intermittent character, and the pulse has become softer. Its efficacy in this disease is much increased by the addition of spirit of turpentine.

In phthisis, bark is found beneficial when the accompanying hectic puts on more of the intermittent form than usual; when the debility is considerable, and blood is mixed in the sputa: and in several cases of pneumonia, when, after repeated large bleedings and evacuations, the pulse continued hard and thrilling, and the blood buffy, although the expectoration was free and the skin open, yet I have seen bark produce the happiest effects.

In various cutaneous diseases, as, lichen agrius, and lividus; in purpura<sup>2</sup>; in impetigo *erysipelatodes* and *scabida*; in some varieties of erysipelas, and in extensive ulcerations both from common inflammation and venereal affections<sup>3</sup>; in the termination of all acute dyseases after the urgent symptoms are subdued; and in dyspepsia, chronic debility, and nervous affections, Cinchona is found to possess great efficacy.

As a local remedy, bark is sometimes used in the form of gargle in malignant sore throat and aphthous affections; and as a wash to foetid gangrenous sores: but in these cases the red bark is to be preferred. Powerful effects, also, are said to have been produced upon the system by frictions with the extract, softened by saliva or oil, upon the thighs and other

<sup>1</sup> Clarke of Newcastle, Heberden, &c.<sup>2</sup> Willan.<sup>3</sup> Pearson.



parts of the body. It may be efficaciously administered per anum, when it cannot be taken into the stomach: but Denman says he found no advantage from its use as a clyster in the low state of puerperal fever, in which it has been highly extolled.

The *Sulphates of Cinchonia* and of *Quinia* have been successfully employed in all the cases in which the use of *Cinchona* is indicated. Sulphate of *Quinia* is much cheaper than that of *Cinchonia*; and on account of its being free from the lignin, resin, and other inert parts of the drug, as well as the smallness of the dose, is a preferable medicine to the bark. In too large doses it nauseates and excites head-ache. The best forms of exhibiting it are as pills made up with crumb of bread, or as a draught, formed with the acidulated infusion of the confection of roses. The dose of both salts is from one grain to four grains, which may be given three times a-day.

*Cinchona* bark is administered in a variety of forms. (See *Preparations and compositions.*) In substance it is reduced to the state of an impalpable powder; and although it loses some of its activity during the process of pulverization, yet, when it can be retained on the stomach, this is the best form of the remedy.<sup>1</sup> If it excite nausea or vomiting, or operate as a cathartic, or occasion costiveness, these inconveniences may in some degree be obviated by combining it with aromatics, opium, or a cathartic, as circumstances direct; or some of the lighter preparations, in which its active principles are supposed to be extracted, and free from the grosser parts, may be employed. The powder is given mixed in wine or in water; or, when the taste is an objection, in milk or syrup, or a solution of extract of liquorice, all of which effectually cover the taste, provided the dose be taken directly after it is mixed.<sup>2</sup>

<sup>1</sup> Fabroni says, “*Cinchona* loses its solubility, and consequently its activity, by long exposure to the air, and by pulverization long protracted with the view of rendering it as fine as possible. From  $\frac{12}{100}$  to  $\frac{6}{100}$  are obtained from bruised *cinchona*, which in fine powder yields only  $\frac{6}{100}$  or  $\frac{7}{100}$  to water.” Practitioners ought never to purchase bark in the state of powder, for in this state it is always found more or less adulterated. Dr. Paris (*Pharmacologia*) mentions that in a late officinal inspection of the shops of apothecaries and druggists, “the censors repeatedly met with powdered *cinchona*, having a harsh metallic taste.” This may arise from the admixture of a species of bark, lately introduced into Europe from Martinique, resembling the *Cinchona floribunda*; and which, by an analysis of M. Cadet (*Journ. de Pharm.* vol. ii. p. 54.), was found to contain iron. The *Cinchona floribunda* is both emetic and purgative; and if this new bark possess the same properties, it is unnecessary to add, that it must prove injurious when combined with good *Cinchona*. A less injurious, but equally fraudulent admixture is the powder of bark which has been employed in making the extract; and of very inferior bark, much of which, we have been informed, is imported for no other purpose.

<sup>2</sup> Mutis, conceiving that fermentation is the best method for extracting the active part of *Cinchona*, has proposed to make a beer of it, by fermenting one part of the bark in powder with eight parts of honey or sugar, and 80 or 100 of water. And Alibert having persuaded a brewer to make some beer with *Cinchona*, administered it to convalescents, weakened by protracted intermittents, with the best effects.



The dose of the powder is from grs. v. to ʒij. or more. In intermittents the full dose is sometimes given at first<sup>1</sup>; but in other diseases, grs. v. x. or xv. are sufficient to commence with, the dose being repeated every two, three, or four hours, and gradually increased, until one or two ounces, in some cases, be taken in twenty-four hours.

Officinal preparations. *Infusum Cinchonæ*, L. E. D. *Decoctum Cinchonæ*, L. E. D. *Extractum Cinchonæ*, L. E. *Extractum Cinchonæ resinosum*, L. D. *Tinctura Cinchonæ*, L. E. D. *Tinctura Cinchonæ composita*, L. E. D.

CINNAMOMI CORTEX. Vide *Laurus Cinnamomum*.

CINNAMOMI OLEUM. *Ibid.*

CITRUS. *Spec. Plant. Willd.* iii. 1426.

Cl. 18. Ord. 3. Polyadelphia Icosandria. Nat. ord. Pomaceæ, Linn. Aurantiæ, Juss.

G. 1391. *Calyx* five-cleft. *Petals* five, oblong. *Anthers* twenty, the filaments united into different parcels. *Berry* nine-celled.

*Species* 1. *C. medica*. The Lemon-tree. *Med. Bot.* 2d edit. 528. t. 189.

*Species* 4. *C. Aurantium*. The Orange-tree. *Med. Bot.* 523. t. 188.

1. CITRUS MEDICA.<sup>2</sup> Var.  $\beta$  *C. Limon*.

Officinal. LIMONES. LIMONUM CORTEX.—OLEUM, *Lond.* CITRI MEDICÆ CORTEX: OLEUM VOLATILE: SUCCUS, *Edin.* LIMON; FRUCTUS SUCCUS; EPIDERMIS, EJUSQUE OLEUM ESSENTIALE, *Dub.* Lemons: their rind, and its essential oil.

*Syn.* Citronier (*F.*), Citrone (*G.*), Limone (*I.*), Citri (*S.*), Lémōn (*Arab.*), Lému (*H.*), Jambéra (*Sans.*), Elimitchum pullum (*Tam.*), Jerook (*Malay.*), Usi (*Celebes.*).

The lemon-tree is a native of Assyria and Persia, whence it was brought into Europe; first to Greece and afterwards to Italy.<sup>3</sup> It is now cultivated in Spain, Portugal, and France, and is not uncommon in our green-houses.<sup>4</sup> It is a beautiful ever-green, of small growth, sending off numerous branches covered with a greyish bark. The leaves are alternate, of a shining pale green colour, ovate, acuminate, about four inches long and two inches broad, slightly indented at the edges, and supported on naked linear footstalks. The flowers, which appear the greater part of the summer, are odoriferous, large,

<sup>1</sup> This was always the case, on the first introduction of the remedy into European practice. In the *Schedion*, two drams in fine powder, are ordered to be macerated for three hours in a glass of white wine, and given at the commencement of the paroxysm. The patient was also ordered to abstain from all other medicine.

<sup>2</sup> Μηλεα μηδικη Theophrasti et Dioscoridis.

<sup>3</sup> Venit in Italiam post Virgilii et Plinii tempora, ante Palladii. *Willd. S. P.* iii. 1426. The fertility of the lemon-tree is proverbial in Italy. A wager was laid in 1812, by Signor Antonio Georgeri of Massa, with Marchese Calani of Spezia, that at Cresullo, half a mile from Massa, there was a lemon-tree which would mature, that year, fourteen thousand lemons. It exceeded the quantity. *Lander's Conversations*, vol. i. p. 122.

<sup>4</sup> It was first cultivated in Britain in the Oxford garden, about the year 1648.



and placed on simple and branched peduncles, arising from the smaller branches. The calyx is saucer-shaped, with the teeth pointed; the petals are oblong, concave, white, with a purplish tinge on the outside; the filaments, united at their base into four parcels, support yellow vertically-placed anthers; and the germen is superior, roundish, and having a simple style with a globular stigma. The fruit is an ovate berry, pointed at each end, rough, punctured, externally of a pale yellow colour, and internally divided into seven, nine, or eleven cells, containing four seeds in each, and filled with vesicles distended with a sharp acid juice. The rind is double: the exterior part thin, yellow, and chiefly made up of a great number of vesicles filled with a very fragrant oil; the interior thicker and whiter than the exterior, and coriaceous.<sup>1</sup>

Lemons are brought to England from Spain and Portugal packed in chests, and each lemon separately rolled in paper. The Spanish lemons are most esteemed.

*Qualities.*—Lemon *juice* is sharp, but very gratefully acid. It consists principally of the citric acid, mucilage, extractive matter, a small proportion of sugar and water. Before Scheele's process was known, many different unsuccessful plans were adopted for separating the citric acid; which is now obtained in a crystallized form, and admitted into the London and Dublin pharmacopœias.<sup>2</sup> The simple juice, although well depurated of its extractive matter, yet soon spoils; and, therefore, the crystallized acid dissolved in water is generally used in its stead.

The *rind* is warm, aromatic, and slightly bitter, qualities depending on the essential oil it contains, which is given out to water, wine, and alcohol. The *essential oil* obtained by distillation is extremely light, nearly colourless, and fragrant; and has the same taste as the rind, only in a greater degree. It is very volatile, yet does not readily rise with alcohol or with proof spirit.

*Medical properties and uses.*—Lemon *juice* is refrigerant and antiseptic. It is given diluted with water and sweetened, forming the beverage called lemonade, to quench thirst, and abate heat, in febrile and inflammatory diseases. Given alone to the extent of a table-spoonful for a dose, it allays hysterical palpitations of the heart; and in combination with carbonate of potass (f℥ss. of the juice to ℥j. of the salt), taken in a state of effervescence, it is used with great success to allay vomiting, and determine to the surface. A still more useful and pleasant effervescing draught is made by putting a table-spoonful of lemon juice, mixed with a small quantity of sugar, into a tumbler, and

<sup>1</sup> *Cærtner de Fructibus*, vol. ii. p. 189.

<sup>2</sup> For an account of this acid, vide *Acidum Citricum* among the Preparations.



pouring over it half a pint of aërated soda water. On account of its antiseptic powers, lemon juice is successfully used in scurvy: and for this purpose large quantities of it, in a concentrated state, are distributed in the navy; but the continued use of it is said to be hurtful to the general health of the men, and to hasten the progress of phthisis where it makes its appearance. The citric acid is likely to supersede its employment in the navy. Dr. Wright observes, that its powers are increased by saturating it with muriate of soda, and recommends such a mixture in remittent fever, dysentery, colic, putrid sore throat, and as being almost specific in diabetes and lientery. It is given also united with camphor, infusion of cinchona and wine, in the same cases; and mixed with ardent spirits and water with sugar it forms *punch*, which is a useful cordial in low fevers.

*Lemon peel* is added to stomachic tinctures and infusions, and is particularly applicable in dyspepsia, arising from irregularities in diet, and the inordinate use of ardent spirits.

The *essential oil* is chiefly used as a perfume, to cover the smell of sulphur in ointments compounded with it.

Officinal preparations. Of the juice, *Syrupus Limonis*, L. E. D. Of the rind, *Aqua Citri medicæ*. E.

## 2. CITRUS AURANTIUM.<sup>1</sup>

*Officinal*. AURANTII BACCÆ, CITRI AURANTII CORTEX, *Lond.* CORTEX, SUCCUS, *Edin.* FRUCTUS SUCCUS, CORTEX EXTERIOR, FRUCTUS IMMATURUS, ET FLORUM AQUA STILLATITIA, *Dub.* The fruit and outer rind of the Seville orange.

*Syn.* (*Immature fruit*), Fruits verts d'orange (*F.*), Unreife Pomeranzen (*G.*), Pomeransknopp (*Swed.*), Collungie pullum (*Tam.*), Jeroc manis (*Malay*), Jeruklegi (*Jav.*), Panneh dodang (*Cyng.*), Usi (*Celebes*)—(*Ripe fruit and its rind*), Orange, Ecorce d'orange (*F.*), Pomeranzen, Pomeranzenschuske (*G.*), Arancio (*I.*), Naranja (*S.*), Narenj (*H.*), Nágaranga (*San.*).

The orange-tree is a native of India and Persia, but it is now abundantly propagated in the south of Europe and the West India islands, and is also found in our green-houses. In its general appearance it resembles the lemon-tree, but the leaves, which are not so large as those of the lemon, and more pointed, are entire, smooth, and furnished with wings or appendages on the footstalks, by which it is particularly distinguished. The flowers, like those of the lemon, appear all the summer, are large, white, odorous, and arise from the smaller branches upon simple and branched pedicels. The parts of the flower resemble closely those of the lemon. The fruit is a globular berry, rough, and of a deep reddish yellow or orange colour; internally divided into nine cells, filled with a vesicular

<sup>1</sup> Aurantia forte a corticis colore, qui colore auri relucet, ut aurea mala vere nominari possunt: sive ab Arantia oppido dicta, veteribus ignota, insitione ad nos devenerunt, *Bauhin Pin.* p. 436.



pulp, and each containing from two to four seeds. The rind, like that of the lemon, is double: the exterior thin and glandular; the interior thick, whitish, and fungous. The China or sweet orange (*Citrus Sinensis*) is a variety of the same species as the Seville orange, and is medicinally employed for allaying thirst in febrile diseases. Both are imported chiefly from Spain, in chests, and packed in the same manner as lemons.

*Qualities*.—The *juice* of the Seville orange is a grateful acid liquor, with a slight degree of bitterness. It consists of nearly the same principles as the juice of the lemon; with a smaller portion, however, of citric acid. The *exterior rind* has a very grateful aromatic odour, and a warm bitter taste, depending on the essential oil contained in its vesicles. Both the bitter and aromatic parts are extracted by water and alcohol, and the essential oil can be obtained by distillation. The *unripe fruit*, named in common Curaçoa oranges, have the aromatic flavour of the rind with a greater degree of bitterness, and retain both when dried. They vary in size from that of a small pea to that of an acorn. The *distilled water* has the grateful perfume of the flowers.

*Medical properties and uses*.—The *juice* of the Seville orange is employed in the same diseases, and with the same intentions, as lemon juice, but it is not so generally used. The *rind* is a useful stomachic, carminative, and tonic, and is a common addition to bitter infusions in dyspepsia. In gout it is joined with magnesia and alkalies; and when the cinchona does not sit easily upon the stomach, it is a most useful adjunct to that remedy in whatever form administered. It has also been given alone in intermittents with seeming advantage.<sup>1</sup> The *oil* is only used as a perfume.

The dried *unripe fruit* (*Aurantium curassaventium*) is employed as an internal remedy in the same cases as the rind of the ripe orange. It is, however, more commonly used as a mechanical irritant in issues; for which purpose the smaller fruit is selected, and generally made round and smooth in the turning lathe. It is preferred for this purpose on account of its odour only; for the heat and moisture of the part in which the orange is lodged, swells it as much as the common pea; and, therefore, it requires to be renewed once in twenty-four hours.

The usual dose of the dried rind, and of the Curaçoa orange, is from grains xv. to ʒj., which are given three or four times a-day.

Officinal preparations. Of the juice, *Succus Cochlearice comp.* E. Of the rind, *Infusum Aurantii compositum*, L. *Tinctura Aurantii*,

---

<sup>1</sup> Murray's *App. Med.* vol. iii. p. 289.



L. D. *Syrupus Aurantii*, L. D. *Confectio Aurantii*, L. E. D. *Aqua Citri Aurantii*, E.

COC'CUS.<sup>1</sup> *Syst. Nat.* Gmelin, 2220.

Cl. 5. Ord. 2. Insecta Hemiptera.<sup>2</sup>

G. 229. Rostrum or Snout seated on the breast. *Antennæ* filiform. Abdomen bristled behind. Wings two, erect in the males; females apterous.

Species 22. *C. Cacti*. Cochineal Insect. *Reaum. Ins.* iv. t. 7. fig. 11, 12. *Phil. Trans.* lii. 661. pl. 21.

Officinal. COCCUS, *Lond.* COCCUS CACTI, *Edin.* COCCINELLA, *Dub.* Cochineal.

Syn. Cochenille (*F.*), Cochenille (*G.*), Cochenilje (*Dutch*), Konsienell (*Swed.*), Cuzzinel (*Dan.*), Coccinilia (*I.*), Cochinilla (*S.*), Cochineel poochie (*Tam.*).

This coccus is found in its wild state in Mexico, Georgia, South Carolina, and some of the West India islands, feeding on several species of cactus, particularly the common Indian fig, or prickly pear plant (*Cactus opuntia*):<sup>3</sup> but in Mexico, and some of the adjoining Spanish settlements<sup>4</sup>, where the insect is, at it were, domesticated and reared with great care, it feeds only on a species of cactus which was supposed to be the Cochineal Indian fig (*Cactus coccinifer*), but which, Humboldt says, is a distinct species, the fruit being internally white. It is cultivated for this purpose; and on it the insect attains to a greater size than in the wild state. It is a small insect, very seldom exceeding a barley-grain in magnitude; with the head, except in the males, scarcely distinct from the body, which is depressed, downy, and transversely rugose. The abdomen is of a purplish red colour, flat below and convex above; and the legs are six in number, short and black. The males, which are few in proportion to the females, there being one only to 150 or 200 females, are winged, slender and active, with the body of a red colour: the head is small, but very distinct from the neck, furnished with jointed feelers, and two long diverging white hairs, about five times the length of the body, which proceed from the tail: the body is elliptical, and furnished with white wings, which lie flat when the insect rests or walks, but are erected when it flies. The females have no wings and are sluggish, scarcely ever moving from the part of the plant where they fix themselves. The back is hemispherical, and crossed by nu-

<sup>1</sup> ΚΟΚΚΟΣ βαφικη Dioscoridis, is the Kermes or *Coccus Ilcis*, Linn. which was known, as a dye, by the Phœnicians, before the time of Moses; and was the *tola* of the Jews. *Beckman's Hist. of Inventions*, translation, vol. ii. p. 185.

<sup>2</sup> Cl. vii. *Ryngota*. Spec. 21. *Fabricii*.

<sup>3</sup> These plants have neither stem nor leaves, in the common acceptation of these words, but consist of roundish or oval compressed joints, that grow out of each other.

<sup>4</sup> "Oaxaca, Kascala, Chulula, Nueva Galicia, Chiapa, in New Spain; and Hambatio, Loja, and Tucuman, produce the greatest quantity." *Ulloa*, quoted by *Bancroft*.



merous wrinkles ; the body is of a much deeper red than that of the males : on the breast is an awl-shaped papilla, through which a fine thread is spun to form a web, with which the insect envelopes itself as soon as it is fully impregnated ; when it becomes torpid, and immediately after laying its eggs dies, and is a mere useless husk.

The wild cochineal is collected six times in the year, just before the females begin to lay their eggs ; a few being left on the plants to furnish a future supply. But the domesticated insect is collected thrice only in the same space of time, the domestication diminishing the number of broods to three in the year, owing to their propagation being suspended during the rainy seasons, whilst the downy covering of the wild species allow them to withstand the inclemency of these seasons. At the third gathering, branches of the plant, to which a certain number of females is left adhering, are broken off, and preserved with great care under cover during the rainy season ; and after this is over they are distributed over the out-door plantations of the cactus, where they soon multiply, and in the space of two months the first crop is fit to be gathered. The insects are detached from the plant by means of a blunt knife, then put into bags and dipped into boiling water to kill them ; after which they are dried in the sun : and although they lose two-thirds of their weight in this process, yet about 600,000lbs.<sup>1</sup> are brought annually to Europe.

Cochineal was used by the natives of Mexico, when the Spaniards arrived there in 1518 ; and was introduced into Europe about the year 1523. The domesticated kind, which is not only much larger, but yields a richer colour, and is consequently most esteemed, is known, in the language of the Spanish merchants, by the name *grana fina* ; the wild is one-half the size only of the other, covered with white down or powder, and is denominated *grana silvestra* ; but, as we receive them, both the kinds are often mixed together. They are imported in bags, each containing about two hundred weight, and have the appearance of small, dry, shrivelled, rugose berries or seeds, of a deep brown-purple or mulberry colour, with a white matter between the wrinkles. In this state they suffer no change from length of keeping. Dr. Bancroft directs that cochineal to be chosen as the best, which

---

<sup>1</sup> Each pound is said to contain 70,000 insects. The monopoly of cochineal is still in the hands of the Spaniards ; but attempts are making to propagate it in the East Indies, if the death of Mr. Anderson have not terminated them. Of the whole quantity brought to Europe, about 150,000lbs. may be considered as the present annual consumption of Great Britain (1812), which at 30s. per lb. cost 225,000*l.* sterling.



“ is large, plump, dry, and of a silver white colour on the surface.”<sup>1</sup>

*Qualities*.—Cochineal has a faint heavy odour, and a bitter austere taste. It is easily pulverized, affording a powder of a purplish red hue, which has been found to be composed of carmine, a peculiar animal matter, a fatty matter, phosphate and carbonate of lime, and muriate and phosphate of potass<sup>2</sup>: the colouring matter is taken up by water, alcohol, and solutions of the pure alkalis. The watery infusion is of a violet crimson, the alcoholic of a deep crimson, and the alkaline of a deep purple, or rather violet hue. The colour of the watery infusion is brightened by all the acids, except the oxymuriatic (*chlorine*), by which it is destroyed. It is brightened also by supertartrate of potass, and alum; and, at the same time, is partly precipitated. It is also precipitated by sulphate of iron of a brownish-violet colour, the liquid remaining a pale yellowish-brown; and by sulphate of zinc and acetate of lead of a purple-violet, the liquid being perfectly colourless. Hence, cochineal is incompatible as a colouring matter with these metallic salts.

*Medical properties and uses*.—Cochineal has lately been recommended as an antispasmodic and anodyne in whooping-cough. I have had no experience of its effects, and believe it to be better fitted for giving a fine colour to tinctures, and similar preparations.

COCHLEARIA.<sup>3</sup> *Spec. Plant. Willd.* iii. 448.

Cl. 15. Ord. 1. Tetradymania Siliculosa. *Nat. ord.* Siliquosæ, *Linn.* Cruciferæ, *Juss.*

G. 1228. *Silicle* emarginate, turgid, rugged; with gibbous, obtuse valves.

*Species* 8. *C. Armoracia*.<sup>4</sup> Broad Horse-radish. *Med. Bot.* 2d edit. 400. *Smith's Flora Brit.* ii. 690.

*Officinal*. ARMORACIÆ RADIX, *Lond.* COCHLEARIÆ ARMORACIÆ RADIX, *Edin.* RAPHANUS RUSTICANUS; RADIX, *Dub.* Horse-radish Root.

*Syn.* Cran; Raifort (*F.*), Murrettich (*G.*), Meerradys (*Belg.*), Peperrod (*Dan.*), Pepperrot (*Swed.*), Chrzan (*Polish*), Chren (*Russian*), Rafano rusticano (*I.*), Marvisco (*S.*), Rubao rusticano (*Portug.*), Morungy vayr (*Tam.*).

This plant is a perennial, growing wild in many parts of England in moist situations, and in waste ground, flowering in June; but it is generally cultivated for culinary and medicinal purposes. The root is long, tapering, and white, sending up many leaves, and a round, erect, branched stem, which rises about two feet in height. The radical leaves are

<sup>1</sup> *Philosophy of Permanent Colours*, 2d edit. i. p. 434.

<sup>2</sup> *Journ. de Pharm.* 1818. p. 526.

<sup>3</sup> Named from a fancied resemblance of the leaf to an old-fashioned spoon.

<sup>4</sup> *Ῥαφανὶς ἀγρία*, *Dioscoridis*.



petiolate, very large, lance-shaped, and waved, crenate and sometimes pinnatifid. Those of the stem are sessile, much smaller, lanceolate, sometimes divided at the edges, at other times entire. The flowers are in terminal clusters, numerous, and of a white colour. The leaves of the calyx are ovate, concave, spreading, and deciduous; the petals white, obovate, twice the length of the calyx, and inserted by narrow claws. The germen is heart-shaped, bearing a simple permanent style, crowned with an obtuse stigma, and changing into an elliptical bilocular pod, containing four seeds in each cell, which frequently prove abortive.

As the acrimony, on which its virtues depend, is lost in some degree by drying, it should be preserved in sand, in a cool place.

*Qualities.* — Horse-radish has a pungent odour, and a very hot, biting, acrid taste, with some degree of sweetness. When kept until it is quite dry, it loses more than two-thirds of its weight, and in time the whole of its pungency is dissipated.

Both water and alcohol extract its active principles. The infusion reddens litmus paper, and precipitates solutions of superacetate of lead and of nitrate of silver. Coction destroys its acrimony, which depends on a volatile oil that can be obtained separate when the mashed root is distilled with water. The oil is of a pale yellow colour, heavy, volatile at 60°, with an extremely pungent odour, and a sweetish, strong, acrid taste, exciting inflammation in the tongue and lips to which it is applied. Einhoff, who examined this root, says, the distilled watery liquid yields traces of sulphur.<sup>1</sup>

*Medical properties and uses.* — This root is stimulant, diaphoretic, and diuretic; and externally rubefacient. It is used, with advantage, in paralytic affections and chronic rheumatism, both internally and externally; and in dropsy, particularly when it follows intermittent fever, in which it was successfully employed by Sydenham. It has, also, been found efficacious in some cutaneous affections; and as a local remedy, a syrup made with an infusion of it, as recommended by Cullen<sup>2</sup>, removes hoarseness arising from relaxation. Horse-radish may be given in substance in doses of ʒss. or more, scraped, or in small pieces swallowed whole.

Officinal preparations. *Infusum Armoraciæ compositum*, L. *Spiritus Armoraciæ compositus*, L. D.

COCOS. *Spec. Plant. Willd.* iv. 400.

Cl. 21. Ord. 6. Monœcia Hexandria. *Nat. ord.* Palmæ.

<sup>1</sup> *Annales de Chimie*, lxx. 185.

<sup>2</sup> The syrup is made by infusing one drachm of scraped horse-radish in two ounces of boiled water, in a covered vessel, and adding double its weight of sugar. Of this syrup a tea-spoonful is to be swallowed leisurely, and repeated at intervals.



G. 1680. *Spathe* general, one-celled. *Spadix* branched.  
*Male flowers*. *Calyx* three-leaved. *Corolla* tripetalous.  
*Female* ——. *Calyx* two-leaved. *Corolla* six-petaled.  
*Style* none. *Stigma* hollowed. *Drupe* fibrous.

*Species* 3. *C. butyracea*. The Mackaw Tree. *Piso, Hist. Nat. lib. iv. p. 125.* (Pindova.)

*Officinal*. COCI BUTYRACEÆ OLEUM FIXUM, *Edin.* Palm Oil.

*Syn.* Huile de Cocobier du Bresil (*F.*), Olio di Cocco del Brasile (*I.*).

This species of palm is a native of Brazil, and is found in abundance near the mines of Ybaquenses. It is a lofty tree, with a rough bark, and the foliage forming a very dense shade. The fruit, which is collected throughout the year, is an obovate, one-celled, smooth, succulent drupe, of a yellow colour, with a point at the upper end; and, at the base, the hard persistent calyx. The nut has a cartilaginous skin and a fibrous pulp; and contains a cartilaginous hard kernel, having nearly the same taste as that of the common cocoa-nut.

This kernel yields the oil. It is first coarsely pounded, or ground in a mill, then macerated in hot water, until it parts with its oil, which, collecting on the surface of the water, concretes as it cools. It is afterwards purified by washing in hot water.

*Qualities*. — Palm oil has an agreeable odour, resembling that of violets, or of the Florentine Iris, and a slightly sweetish taste. It is of the consistence of butter, and has a light lemon-yellow colour; it becomes rancid by long keeping, loses its pleasant odour, and its yellow colour fades to a dirty white. It is said to be sometimes imitated with hog's lard coloured with turmeric, and scented with Florentine Iris root.

*Medical properties and uses*. — This vegetable butter is emollient; and as such is sometimes used externally in frictions.

COLCHICUM. *Spec. Plant Willd.* ii. 272.

*Cl.* 6. *Ord.* 3. Hexandria Trigynia. *Nat. ord.* Spathaceæ, *Linn.*  
*Junci, Juss.*

G. 707. *Spathe*. *Corolla* six-parted, with a rooted tube. *Capsules* three, connected, inflated.

*Species* 1. *C. autumnale*. Meadow-saffron. *Med. Bot.* 2d edit. 759. t. 259. *Smith, Flor. Brit.* 400. *English Botany*, 133.

*Officinal*. COLCHICI RADIX ET SEMINA<sup>1</sup>, *Lond.* COLCHICI AUTUMNALIS RADIX, *Edin.* COLCHICUM; RADIX, PRIMO VERE, FOLIIS JAM APPARENTIBUS. The root of Meadow-saffron.

*Syn.* Colchique (*F.*), Zeitlozen, Weissen saffron (*G.*), Tydeloosin (*Dutch*), Hundedöd (*Dan.*), Tidlösa (*Swed.*), Colchico Autumnale (*I.*), Zafran (*S.*), Rozsiad (*Pol.*).

This is an indigenous perennial plant, generally found growing in moist rich meadow grounds<sup>2</sup>, and flowering in

<sup>1</sup> All the Colleges have erred in using the word *radix* instead of *bulbus*.

<sup>2</sup> It is very abundant in Essex and Suffolk.



September. The bulb is solid, egg-shaped, and covered with a brown, membranous coat. The leaves, which appear in spring, are radical and spear-shaped, about five inches long, and one inch broad at the base. They wither away entirely before the end of summer, and are succeeded by the flower, which appears in autumn without any leaves.<sup>1</sup> It is, however, proper to state, that the bulb from which the flowers spring is the offset of that from which the leaves have decayed. There is no calyx. The corolla, which is of a pale, pinkish-lilac colour, springs directly from the bulb, and consists of a tube about five inches long, two-thirds of which are sunk in the ground, and a limb divided into six lanceolate keeled segments. The filaments are half the length of the segments of the corolla, subulate, united to the upper part of the tube, and supporting yellow erect anthers. The stigmas are revolute. The fruit is a three-lobed, three-celled capsule, on a thick, short peduncle. The impregnated germen remains under ground, close to the bulb, till the following spring, when it rises in its capsular form above the surface, accompanied by the leaves. The seeds are ripe about the end of June.

The thick, old bulb begins to decay after the flower is perfectly expanded, and the new bulbs, of which there are always two formed on each old bulb, are perfected in the following June; from which time until the middle of August they may be taken up for medicinal use. The bulbs when mature, on being cut transversely, yield a milky-looking acrid juice, which produces a beautiful, cærulean blue colour, if rubbed with the alcoholic solution of guaiac. To preserve the virtues of the plant, the bulb, as soon as possible after it is dug up, should be cut into transverse slices, not thicker than one-eighth of an inch, and dried by placing the slices on clean white paper, distinct from one another, without heat, or at a very low temperature. The test of the drug being good and properly dried, is the appearance of the blue colour, on rubbing it with a little distilled vinegar and the alcoholic solution of guaiac. The slices also should not appear deeply notched or panduriform; as this is the mark of the bulb having begun to empty itself for the nourishment of the young bulbs; and, consequently, to suffer in its medicinal powers, from the chemical change which, at this period, its contents must necessarily undergo for the nourishment of the offsets. It should be preserved in slices, in well stopped bottles.

*Qualities.*—The recent bulb of this plant has scarcely any

---

<sup>1</sup> From this circumstance, it is called "*naked lady*," in some parts of the country.



odour. When it is dug up at a proper season of the year, the taste is bitter, hot, and acrid, occasioning a warm sensation in the stomach, even when taken in a small quantity. At other seasons, however, and in some soils and situations, it possesses very little acrimony; and thence the contradictory opinions which authors have given of it. Its acrimony resides in a peculiar alkali, which can be separated from the other principles, and has been named *veratrine* by M. M. Pelletier and Caventou, who discovered it. The veratrine is obtained in form of a white powder: it combines with acids, but the neutral salts are not crystallizable.<sup>1</sup> The other components of the bulb are the following: a fatty matter, gallic acid, a yellow colouring-matter, gum, starch, inulin in great abundance, and lignin. Vinegar and wine are the best menstrua for extracting the active qualities of the bulb. A deposit forms in the wine, which Sir E. Home says is extremely acrid, exciting nausea and griping, and ought to be removed, as its removal does not alter the virtues of the medicine.<sup>2</sup>

*Medical properties and uses.* — Meadow-saffron possesses diuretic, purgative, and narcotic properties: and on the Continent, where it was recommended to notice by Baron Stoerck, it is a favourite remedy in dropsy, particularly hydrothorax, and in humoral asthma. But as it does not differ in its mode of action from squill, and is more uncertain in its operation, it has not been much used in that complaint in this country. In gout and rheumatism, however, its efficacy has been fully ascertained: and, in allaying the pain, it may be almost said to possess a specific property. It operates on the bowels chiefly, producing copious bilious evacuations; and on the nerves, diminishing the action of the arterial system. The petals of the flower and the seed possess the same medicinal properties as the bulb, but in a diminished degree. In the seed the veratrine exists in the testa or husk, and consequently the seeds should not be bruised in preparing the wine or tincture with them.

The dose in substance is from grs. iij. to grs. ix. of the dried bulb: and of the saturated vinous infusions, made by macerating ℥iss. of the dried bulb, or ℥j. of the dried petals, or of the seed, in f℥xij. of white wine, from ℥xxx. to ℥lxxx. may be taken whenever the patient is in pain.

Officinal preparations. *Acetum Colchici*, L. *Oxymel Colchici*, D. *Syrupus Colchici autumnalis*, E.

COLOCYNTHIDIS PULPA. Vide *Cucumis Colocynthidis*.  
CONIUM.<sup>3</sup> *Spec. Plant. Willd.* i. 1395.

<sup>1</sup> *Journ. de Pharm.* Aug. 1820.

<sup>2</sup> *Phil. Trans.* 1817, part ii.

<sup>3</sup> Κώνιον Dioscoridis. *Cicuta vulgaris major*, Park. 932. *Cicuta*, Dod. 461.



Cl. 5. Ord. 2. Pentandria Digynia. Nat ord. Umbellatæ.

G. 533. *Partial involucre* halved, three-leaved. *Fruit* nearly globular, five-streaked, notched on each side.

*Species* 1. *Conium maculatum*. Common Hemlock. *Med. Bot.* 2d. edit. 104. t. 42. *Smith Flora Britan.* i. 302.

*Officinal.* CONII FOLIA, *Lond.* CONII MACULATI FOLIA, *Edin.* CICUTA, *Dub.* The leaves and seed of Hemlock.

*Syn.* Cigue ordinaire (*F.*), Gefleckter Schierling (*G.*), Gevlakte Scheerling (*Dutch.*), Skarntyde (*Dan.*), Spräckligodört (*Swed.*), Swinia wesz (*Pol.*), Bo-ligolou (*Russian*), Cicuta Maggiore (*I.*), Conio manchado (*S.*).

Hemlock is a biennial, umbelliferous, indigenous plant, growing under hedges, by road-sides, and among rubbish, flowering in June and July. The root, which is fusiform, branching, whitish and fleshy, exudes, when cut, a milky juice. The stem rises erect about four or five feet in height, is branching and leafy, round, hollow, striated, smooth, shining, and maculated with brownish purple. The lower leaves are very large, above a foot in length, on large sheathing petioles, supradecomposed, and shining; the upper ones are bipinnate; the whole stand upon channelled footstalks, proceeding from the joints of the stem, are incised, smooth, of a deep-green colour on the upper surface, but paler underneath. The rays of the umbels are ten or twelve, those of the umbellules fifteen or sixteen. The involucre consists of from three to seven, short, turned-down, lancet-shaped leaflets, with white edges spread at the base; the involucre of three or four leaflets on one side only, and spreading. The flowers are very small; the petals white, the outer ones rather larger than the inner, cordate, inflected; the stamens the length of the petals, supporting white orbicular anthers; the styles two, filiform, diverging, and crowned with round stigmas. The fruit is ovate, striated, smooth, and brownish when ripe.

Hemlock is distinguished from other umbelliferous plants, with which it may be confounded, by its *large and spotted stem*<sup>1</sup>, the dark and *shining colour of its lower leaves*, and their *disagreeable smell*, when fresh and bruised, resembling in some degree the urine of a cat.<sup>2</sup>

For medical use, the leaves should be gathered about the end of June, when the plant is in flower; the small leaflets picked off, and the footstalks thrown away. The picked leaflets are then to be properly dried (*vide Powders*, Part III.); and as exposure to the air and light destroys the fine green colour of the plant, and injures its active qualities, the dried

<sup>1</sup> The *Chærophyllum bulbosum*, bulbous-rooted cow-parsley, has a spotted stem, but the joints are swelled, and the seeds rough.

<sup>2</sup> In Ray's Synopsis, *Conium maculatum* is named *Cicuta*; a name still retained by the Dublin College; and owing to which the water-hemlock, *Cicuta virosa*, has sometimes been confounded with it; and improperly used.



leaflets must be preserved in boxes completely filled, by gently pressing down the leaves, then covered with a closely fitted lid, wrapped in paper and sealed; or, if powdered, the powder may be preserved good, in closely stopped opaque phials, for many years.

*Qualities*.—The odour of properly dried hemlock leaves is strong, heavy, and narcotic, but not so disagreeable as that of the fresh leaves: the taste is slightly bitter and nauseous. They are easily pulverised; and the powder should retain the beautiful green colour of the leaves. The acrimony only of the fresh leaves is lost in drying; but the narcotic principle remains uninjured if the operation be well performed. The virtues of conium are extracted by alcohol and sulphuric ether. To the ether it communicates a very deep green colour; and when the tincture is evaporated on the surface of water, a rich dark green resin remains, in which the narcotic principle of the plant appears to reside; it contains the odour and taste in perfection; and half a grain produces head-ache, and slight vertigo. To this principle, which I discovered, Dr. Paris proposes to give the name of *conein*.<sup>1</sup>

*Medical properties and uses*.—Hemlock is a powerful narcotic, and is used as such both internally and as an external application. Stoerck, whose publications first brought it into general notice, rated its powers too high, and the multitude of discordant diseases which he enumerated as yielding to it, led many sober men to doubt its efficacy altogether. Hemlock is, nevertheless, a useful narcotic; and, if it have not succeeded in curing cancer in the hands of British practitioners, it has been advantageously used as a palliative in both scirrhus and open cancer, abating the pain, and allaying the morbid irritability of the system. It has also been found serviceable in chronic rheumatism, in scrophulous, syphilitic, and other ill-conditioned ulcers, and glandular tumors; in pertussis, and the protracted cough which often remains after pneumonic inflammation. An over-dose of it produces sickness, vertigo, delirium, dilatation of the pupils, great anxiety, stupor, and convulsions.<sup>2</sup> The best antidote is vinegar, after the stomach has been evacuated, and the cerebral excitement reduced by bleeding and purging.

The powder of the dried leaves, if well preserved, is the best form of this remedy. Hufeland recommends the fresh expressed juice from ℥ xij. to ℥ lx. for a dose. The dose of the powder is grs. iij., gradually increasing it every day, until a slight vertigo forbid its further increase.

Officinal preparation. *Extractum Conii*, L. E. D.

<sup>1</sup> *Pharmacologia*, p. 185.

<sup>2</sup>

*Pinguescere sæpe cicuta*

*Barbigeras pecudes, homini quæ est acre venenum. Lucretius.*



CONVOLVULUS. *Spec. Plant. Willd.* i. 844.

*Cl.* 5. *Ord.* 1. Pentandria Monogynia. *Nat. ord.* Campanaceæ, *Linn.*  
*Convolvuli, Juss.*

*G.* 323. *Corolla* bell-shaped, plaited. *Stig.* two. *Capsule* two-celled, each cell containing two seeds.

\* *Stem* twining.

*Sp.* 4. *C. Scammonia*. Scammony. *Med. Bot.* 2d edit. 243. t. 86.

*Sp.* 61. *C. Jalapa*. Jalap. *Med. Bot.* 2d edit. 246. t. 87.

1. CONVOLVULUS SCAMMONIA.<sup>1</sup>

*Officinal.* SCAMMONIÆ GUMMI RESINA, *Lond. Edin.* SCAMMONIUM; GUMMI RESINA, *Dub.* Scammony.

*Syn.* Scammonée (*F.*), Scammonium von Aleppo (*G.*), Het Scammoneum (*Dutch*), Scammonea (*I.*), Escamonea (*S. & Port.*), Sukmunya (*H. & Arab.*)

This plant is a native of Syria and Cochin China. It grows in abundance on the mountains between Aleppo and Latachea, and there the greater part of the Scammony of commerce is obtained.<sup>2</sup> The root, which is perennial, is tapering, from three to four feet in length, and from three to four inches in diameter, covered with a light grey bark, and contains a milky juice. It sends up many slender, twining stems, which extend from fifteen to twenty feet in length, adorned with arrow-shaped, smooth, bright green leaves, upon long footstalks. The flowers are in pairs upon the pedicels, consisting of a double calyx of four emarginated leaflets in each row; and a funnel-shaped, pale yellow, plaited corolla. The capsule is three or four-celled, containing small pyramidal seeds.

Scammony is obtained from the root of this plant<sup>3</sup>; and is collected in the beginning of June, in the following manner: The ground is cleared away from the root, the top of which is then cut off in a sloping direction, about two inches below the place whence the stalks spring; and the milky juice which flows from it is collected in a shell fixed at the most depending part. Each root yields a few drachms only, which are drained off in about twelve hours. "This juice from the several roots is put together, often into the leg of an old boot, for want of some more proper vessel, where, in a little time, it grows hard and is the genuine Scammony." "The Jews," Dr. Russel says, "buy the Scammony while it is soft, and mix it with the expressed juice of the stalks and leaves, with wheat flour, ashes, fine sand, or whatever else can answer their purpose." It is imported from Aleppo in what are called drums, which weigh from 75 to 125 lbs. each; and from Smyrna in cakes like wax, packed in chests. The former is light and friable, and is considered the best; that from Smyrna is more compact and ponderous, less friable, and fuller of impurities.

<sup>1</sup> Σκαμμωνία Dioscoridis.

<sup>2</sup> Russel's *Nat. Hist. of Aleppo*, ii. 246.

<sup>3</sup> No other part of the plant possesses any medicinal quality. Russel l. c.



*Qualities*.—Good Scammony is light, friable, and externally like a honeycomb. It has a peculiar, rather heavy, odour, not unlike that of old ewe-milk cheese; and a bitterish, slightly acrid taste. The colour is blackish or bluish grey, changing to dirty white, or lathering, when the surface of the mass is rubbed with the wet finger. The fracture is irregular but smooth, faintly shining, and the sharp edges of the shivers are of a lighter grey colour, and translucent. It is pulverulent; and the powder has a light grey colour. Its specific gravity is 1.235.<sup>2</sup> When it is of a dark colour, heavy and splintery, it should be rejected. When triturated with water, nearly one-fourth of it is dissolved, and the solution appears slightly mucilaginous, opaque, and of a greenish-grey colour. This solution is not affected by alcohol, solutions of superacetate and acetate of lead and sulphate of iron, nor precipitated by the acids; but with sulphuric acid it gives out the odour of vinegar. Solution of ammonia does not alter it, but that of potass occasions a yellowish precipitate, which is quickly re-dissolved on the addition of an acid. Ether takes up two parts in ten of scammony, and when evaporated, leaves a brownish semi-transparent resin. Alcohol dissolves two-thirds of its weight; but proof spirit is its best menstruum, taking up the whole except the impurities. Aleppo scammony contains, according to Bouillon La Grange and Vogel, 0.60 of resin, 0.20 of extractive, 0.03 of gum, and 0.35 of impurities. Smyrna Scammony contains 0.29 of resin, 0.08 of gum, 0.05 of extractive, and 0.58 of impurities. When these impurities consist of flour, sand, or ashes, they can be detected by dissolving the sample in proof spirit, as they sink and remain undissolved: but Scammony is sometimes also adulterated with the expressed juice of *Cynanchum monspeliacum*: and a fictitious Scammony is also sold for the real, consisting of jalap, senna, manna, gamboge, and ivory black.

*Medical properties and uses*.—Scammony is a drastic cathartic, operating, in general, quickly and powerfully. The ancients were acquainted with its purgative qualities; and also employed it as an external application for removing hard tumors, itch, scurf, and fixed pains; but for the latter purposes it is now never used. It is a good purgative, in the torpid state of the intestines, in leucophlegmatic, hypochondriacal, and maniacal subjects; in worm cases, and the slimy state of the bowels to which children are subject; and as a hydragogue cathartic in dropsy. Scammony has been regarded by some as a cathartic of so irritating a nature, as to require to be corrected by exposing it to the fumes of sulphur,

<sup>2</sup>*Brisson*.



defæcating it with lemon juice and other acids, and uniting it with demulcent mucilages; but, except in an inflamed or very irritable state of the bowels, it is a safe and efficacious purgative. It is however apt to gripe, on which account it is generally united with an aromatic, or a drop of some essential oil.

The dose of Scammony is from grs. v. to grs. xvj.; whether it be given in powder, or as a bolus, or in the form of mixture, triturated with almonds, gum, or extract of liquorice, and water.

Official preparations. *Confectio Scammonice*, L. D. *Pulvis Scammonice comp.* L. E. *Extractum Colocynthis comp.* L. *Pulvis Sennæ comp.* L.

## 2. CONVOLVULUS JALAPA.

*Official.* JALAPÆ RADIX, *Lond.* CONVULVULI JALAPÆ RADIX, *Edin.* JALAPA; RADIX, *Dub.* Jalap root,

*Syn.* Jalap (*F.*), Jalappenharz (*G.*), Jalapparot (*Swed.*), Scialappa (*I.*), Jalapa (*S.*).

This species of convolvulus is a native of South America, taking its name from Xalappa, a city of Mexico.<sup>1</sup> It grows in a dry sandy soil, and flowers in August and September. The root is perennial, of an irregular egg shape, and a dark almost black colour on the outside, ponderous, large<sup>2</sup>, and when fresh, abounding with a milky juice. It sends up many triangular, tuberculated, twining, twisted stems, which extend upwards of ten feet, with smooth petiolated leaves, of a bright green colour, varying in shape, some being cordate, others angular, and a few oblong and pointed. The flowers are on short, axillary peduncles that send off two pedicels, each bearing a large, bell-shaped, entire, plaited flower, of a reddish colour externally, and a dark purple within: with a calyx composed of five oval, concave, pale green leaves, somewhat indented at their points. The anthers are of a yellow colour, large, on slender short filaments; the style is shorter than the filaments, and the germen oval. The seeds are bristled.

The root of this plant, which is the Jalap of the shops, was first brought to Europe about the year 1609 or 1610.<sup>3</sup> The best comes from Vera Cruz in transverse slices, and also in egg-shaped, pointed, entire tubers, covered with a very thin, wrinkled, brown cuticle. That which is sliced is more liable to be adulterated, which is said to be sometimes done with slices of briony root; but the fraud is easily discovered by the

<sup>1</sup> It was cultivated in England by Mr. Miller in 1668; and a few years ago two specimens were in vigorous growth in Kew gardens, slips of the original plant, introduced there by Mons. Thouin in 1778.

<sup>2</sup> A root of jalap, which was carried by Michaux, junior, in 1803, from the botanic garden of Charlestown to Paris, and planted there in the garden of the Museum of Natural History, where it now grows, weighed 47 pounds and three quarters. *Memoirs de l'Institut*, tom. vi. 387.

<sup>3</sup> *Bauhin Prodromus*, 135.



spongy texture and whiter colour of the latter, and its burning less readily when applied to the flame of a candle.

*Qualities*. — Good Jalap root has a sweetish, heavy odour when broken, and a sweetish, slightly pungent taste. It is heavy, compact and hard, with a shining resinous fracture, which shows the internal part of a yellowish-grey colour, interspersed with deep brown, concentric circles. It is pulverulent, affording a powder of a pale brownish-yellow colour. Both water and alcohol, separately, extract a part; and when mixed, take up the whole of the active constituents of jalap. Ether dissolves three parts of ten submitted to its action; and affords, when evaporated over water, a transparent, insipid resin, and some extractive. Hence jalap appears to contain resin, starch, extractive, and ligneous matter.

*Medical properties and uses*. — Jalap is a stimulant cathartic, acting briskly on the bowels; and although occasionally griping severely, yet safe and efficacious. It is used in the same cases as scammony, whenever it is required effectually to evacuate the intestines; and as a hydragogue purgative it is supposed to possess singular efficacy. It has been asserted that it proves hurtful in hypochondriasis, bilious habits, and fevers, except of the intermittent kind; but Dr. Hamilton used it in all these instances, in typhus, and the exanthemata, with the best effects.<sup>1</sup> The watery extract purges moderately without griping, and is therefore well adapted for children; but the alcoholic scarcely at all purges, although it occasions the most violent tormina and gripings. It is frequently triturated with hard sugar, which renders its powder finer, and increases its activity; and with other cathartics, by which the action of both is reciprocally improved. In dropsical affections, the supertartrate of potass is a useful addition; and in the cachexia and worms, it is united with calomel, the operations of which it greatly quickens.

The dose is from grs. x. to 3 ss. in powder, pills, or bolus; with a drop or two of essential oil to prevent griping.

Officinal preparations. *Pulv. Jalapæ comp.* E. *Extractum Jalapæ*, L. E. D. *Tinctura Jalapæ*, L. E. D. *Tinct. Sennæ comp.* E.

CONTRAJERVÆ RADIX. Vide *Dorstenia Contrajerva*.

COPAIFERA. *Spec. Plant. Willd.* ii. 630.

Cl. 10. Ord. 1. Decandria Monogynia. Nat. ord. Dumosæ, Linn. Leguminosæ, Juss.

G. 880. Calyx none. Petals four. Legume ovate. Seed one, with an ovate arillus.

Species 1. *C. officinalis*. Copaiva tree. *Med. Bot.* 2d ed. 609. t. 216.

Officinal. COPAIBA, Lond. COPAIFERÆ OFFICINALIS RESINA, Edin. BALSAMUM COPAIBÆ, Dub. Copaiba Balsum.

<sup>1</sup> *Observations, &c. on Purgative Medicines*, 8vo. passim.



*Syn.* Beaume de Copahu (*F.*), Kopaiva balsam (*G.*), Hwit Indiansk Balsam (*Swed.*), Balsam Copayve (*Dutch*), Balsamo del Coppaiba (*I.*), Copayva (*S.*).

The Copaiba tree is a native of South America and the Spanish West India islands. It grows in great plenty in the woods of Tolu, near Carthagena, and in those of Quito and Brazil. It is a lofty, handsome tree, branching at the top, with a brownish ash-coloured bark. The leaves are large and pinnate, consisting of four pair of ovate pointed, alternate, ferruginous leaflets, with a terminal one, two, or three inches long, entire, shining, veined, narrower on one side than on the other, and placed on short petioles. The flowers are in terminal racemes, which are stiff, spreading, the length of the pinnæ, and loosely divided into eight alternate common peduncles, with the flowers, which are white, sitting closely on them. The petals are oblong, acute, concave, spreading: the filaments slender, incurved, bearing oblong, incumbent anthers; and the germen roundish, compressed, and on a short pedicel. The fruit is an oval, two-valved pod, containing a single egg-shaped seed, enveloped with a berried arillus.

The Copaiba balsam of the shops is procured by wounding or boring these trees to the pith, near the base of the trunk, when it flows abundantly<sup>1</sup>, in the form of a clear, colourless liquid; which is thickened, and acquires a yellowish colour by age. The operation is performed two or three times in the same year; and, from the older trees, the best balsam is obtained. It is brought to this country from the Brazils in small casks, each of which contains from one cwt. to one cwt. and a half of the balsam.

*Qualities.*—Genuine good Copaiba balsam has a peculiar but agreeable odour, and a bitterish, hot, nauseous taste. It is clear and transparent; its consistence is that of oil, the colour a pale golden yellow, and its specific gravity 0·950<sup>2</sup>; but when it is exposed with an extended surface to the action of the air, it gradually thickens, until at length it becomes solid, dry, and brittle like resin. It is insoluble in water, but is completely soluble in alcohol and ether. Sulphuric acid converts it into a brown, bituminous-like mixture, which gives out a strong odour of sulphur. Nitric acid, in the ordinary heat of the air, partially dissolves it, and renders it brown; but, at an increased temperature, the action is violent, the acid is decomposed, and nitrous fumes are copiously emitted. The muriatic and

<sup>1</sup> "Tanta quantitate distillat, ut spatio trium horarum ad lb. xij. effundat." *Piso. Nat. Hist.* 56.

<sup>2</sup> The adulterated balsam, which Lewis mentions, as being thick, white, and opaque, with a quantity of turbid, watery liquor at the bottom, is not now to be found.



acetic acids scarcely affect it. The pure alkalies form with it white, saponaceous compounds, which are soluble in water, forming opaque milky mixtures. It is soluble, also, in the expressed oils. In destructive distillation it yields some empyreumatic brownish-red oil, an acidulous water, carbonic acid gas, and olefiant gas, but does not yield benzoic acid. Hence, it approaches nearer in its nature to the turpentine than to the balsams. It is sometimes adulterated with mastic and oil, and occasionally with rape oil. Bucholz remarks, that if Copaiba does not dissolve completely in a mixture of four parts of alcohol and one of rectified sulphuric ether, its adulteration may be inferred.

*Medical properties and uses.* — Copaiba balsam is stimulant, diuretic, and gently purgative. It has been recommended in pulmonary complaints; but where the excitement is morbidly increased, or there is any degree of the inflammatory diathesis present, the heating and irritating quality of Copaiba renders it injurious. From its power of stimulating the urethra, it is more successfully used in gleet. It is equally efficacious in fluor albus, and in that state of the uterus sometimes occurring on the final cessation of the menses, which is accompanied with a sanious discharge, great bearing down, and many of the symptoms of incipient cancer. It certainly affords considerable relief in hæmorrhoidal affections; perhaps from its exciting the steady peristaltic motions of the intestines, at the same time that the determination of the blood to the hæmorrhoidal vessels is lessened, by the stimulant effect of the remedy on the kidneys. In too large doses, it excites inflammation of the kidneys, and its use should always be avoided when ulceration of these organs is suspected.

The dose of Copaiba is from  $\mathfrak{m}\text{x.}$  to  $\mathfrak{m}\text{xxx.}$ , twice or thrice a day, either dropped on sugar, or mixed with soft or distilled water, by means of mucilage or the yolk of an egg.

**CORIANDRUM.** *Spec. Plant. Willd.* i. 1448.

*Cl.* 5. *Ord.* 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

*G.* 552. *Corolla* radiate. *Petals* inflex-emarginate. *Involucre* universal, one-leaved. The partial ones halved.

*Species* 1. *C. sativum*.<sup>1</sup> Common Coriander. *Med. Bot.* 2d edit. 137. *t.* 53. *Smith Flor. Brit.* 320. *Eng. Bot.* 67.

*Officinal.* CORIANDRI SEMINA, *Lond.* CORIANDRI SATIVI SEMINA, *Edin.* CORIANDRUM, *Dub.* Coriander Seed.

*Syn.* Coriandre (*F.*) Koriandor saamen (*G.*), Koriander (*Dutch, Swed. & Dan.*), Koryander (*Pol.*), Coriandro (*Port.*), Coriandro (*I.*), Semilla de Clantro (*S.*), Cottamillie (*Tam.*), Mety (*Malay*), Kezereh (*Arab.*), Kitnuz (*Pers.*), D'hanya (*H.*), D'amyáca (*San.*).

This plant is an annual, a native of Italy; but is now found wild in some parts of this country<sup>2</sup>, owing to the abundant

<sup>1</sup> *Kopriavon* Dioscoridis. <sup>2</sup> About Ipswich, and some parts of Essex. *Smith.*



cultivation of it for medicinal purposes. It flowers in June, and ripens its seed in August. The stem is erect, about two feet in height, branching, divaricated, round, smooth, and obscurely striated. The leaves are compound; the lower ones pinnated, with gashed, wedge-shaped, somewhat roundish leaflets, and the upper thrice-ternate, with linear-pointed segments. Both the umbels and umbellules are many-rayed; with an involucre of one linear leaf, and involucels of three lanceolate narrow leaves, all on one side. The flowers are of a white or reddish colour. The calyx consists of five leaves: the petals are five also, oblong, and inflected at the tips, but those of the flowers of the circumference have the outermost petals larger, and not inflected. The fruit is globular, obscurely ribbed, and divisible into two concave hemispherical seeds; and this form of the fruit distinguishes coriander from all the other species of *Umbellatæ*. The whole plant, when green, has an abominably fœtid odour if bruised, which extends even to the fruit.<sup>1</sup>

*Qualities.*—The dried seeds have a grateful, aromatic odour, and a moderately warm, pungent taste; qualities which depend on an essential oil, that can be obtained separate by the distillation of the seeds with water. Their active principles are completely extracted by alcohol, but only partially by water.

*Medical properties and uses.*—These seeds are carminative and stomachic. They are sometimes used in flatulencies; but principally to cover the unpleasant taste, and correct the griping quality of some cathartics. The dose is ʒj. to ʒj., bruised.

CORNUA. Vide *Cervus Elaphus*.

CRETA. Vide *Calx*.

CROCUS. *Spec. Plant. Willd.* i. 194.

*Cl.* 3. *Ord.* 1. Triandria Monogynia. *Nat. ord.* Ensatæ, *Linn.* Irides, *Juss.*

*G.* 92. Corolla six-parted, equal. Stigma convoluted.

*Species* 1. *C. sativus*.<sup>2</sup> Common Saffron. *Med. Bot.* 2d edit. 763. t. 259. *Smith Flora Brit.* i. 39. *Eng. Bot.* t. 343.

*Officinal.* CROCI STIGMATA, *Lond.* CROCI SATIVI STIGMATA, *Edin.* CROCUS, *Dub.* The Stigmas of the Saffron.

*Syn.* Saffran (*F.*), Safran (*G. & Dan.*), Saffran (*Dutch & Swed.*), Zafferano (*I.*), Azafran (*S.*), Açafrão (*Port.*), Szafran (*Pol.*), Khoongoomapoo (*Tam.*), Zafran (*Arab.*), Abeer (*Pers.*), Safaron (*Malay*), Khohom (*Cyng.*), Cashmeerum (*Sans.*).

Common Saffron is a perennial bulbous plant, found wild in some parts of this country, which affords reason for supposing it to be indigenous; but it is probable, that it was

<sup>1</sup> Hence, Alston imagined that the name of the plant comes from *kopis*, a bug. *Mat. Med.* ii. 349.

<sup>2</sup> *Κροκος* Dioscoridis. Its English name is derived from the Arabic Sapharan. *Celsus.* See *Alston's Lectures*, ii. 119.



originally brought from Asia. It is cultivated for medicinal use, in great abundance, in Cambridgeshire and Essex. Formerly, it was chiefly grown at Saffron Waldon, but it is now confined to Stapleford. It flowers in September. The bulb is solid and depressed. The flower, which appears before the leaves, is sessile on the bulb, of a violet or lilac colour, and raised on a long, slender, white tube. The leaves are linear, a little revolute, of a deep rich green colour, with a white nerve in the centre, and inclosed with the tube of the flower in a membranous sheath. The corolla is parted into six nearly elliptical segments; the stamens are shorter than the corolla, and erect; and the style, which is the length of the corolla, hangs out at one side between the segments. The stigma is deeply three-parted, of a rich orange colour, and odorous; with the segments linear-involute at the margin, and crenate at the apex.

For the preparation of the Saffron, the flowers are gathered early in the morning, just as they are about to blow. They are then spread upon a table, and the stigmas, with a proportion of the style, carefully picked out of the flower, which is thrown away as useless. The stigmas are then dried upon a portable kiln of a peculiar construction, over which a hair cloth is stretched, and over it several sheets of white paper are laid; upon which the wet saffron is spread between two and three inches thick. It is now covered with other sheets of paper, and over them is laid a coarse blanket, five or six times doubled, which is pressed down with a board and a large weight after the fire is lighted. The first heat is strong, to make the saffron sweat; and, after an hour, when it is found formed into a cake, it is turned, and the same degree of heat continued for another hour. The fire is then reduced to a moderate heat, which is kept up for twenty-four hours, during which time the cake is turned every half-hour, so as to dry it thoroughly. It is then fit for the market.

In the shops is found Saffron from Sicily, France, and Spain, besides the English. The Spanish is generally spoiled with oil, in which it is dipped with the intention of preserving it; the Sicilian and French are better; but the English, as being fresher, more genuine and better cured, is always preferred. It is sometimes adulterated with fibres of smoked beef, the petals of the safflower (*Carthamus tinctorius*), and of officinal marigold (*Calendula officinalis*); or Saffron, from which tincture or infusion has been drawn, is mixed with a little good Saffron, and again pressed into a cake. These frauds are detected by infusing the suspected Saffron in hot water, when the expanded stigmas will be easily distinguished from the petals of the other flowers; and the deficiency of



the presence of colour and odour, or an unpleasant odour arising when the Saffron is thrown upon red-hot coals, will indicate the presence of the other fraudulent ingredients. It should be chosen fresh, in close, tough, compact cakes, moderately moist and possessing, in an obvious degree, all the undermentioned sensible qualities. The not staining the fingers, the making them oily, a musty flavour, and a whitish-yellow, or a blackish colour, indicate that it is bad, or too old.

*Qualities.*—Good Saffron has a sweetish, penetrating, diffusive odour; a warm, pungent, bitterish taste; and a rich, deep, orange-red colour. It yields its colour and active ingredients to water, alcohol, proof spirit, wine, vinegar, and, in a smaller degree, to ether. By distillation with water it affords a small quantity of a heavy, golden-yellow coloured essential oil. The watery infusion, which has the deep orange colour of the Saffron, is rendered of a very deep purple by strong sulphuric acid, the mixture emitting the smell of vinegar, and yielding a copious black precipitate when diluted with water: the oxymuriatic acid produces a copious yellow precipitate, the liquid retaining only a pale lemon colour. Hence, Saffron seems to contain chiefly extractive, which, according to Hermbstadt, is nearly pure<sup>1</sup>, and in the proportion in ten parts in sixteen of the vegetable; the remainder being chiefly ligneous fibre. I have found that it contains resin also; for sulphuric ether digested on saffron is coloured, and when evaporated on the surface of water, a pellicle of resin is left, whilst the coloured extractive, which is taken up with the resin, is dissolved in and colours the water.<sup>2</sup>

*Medical properties and uses.* — Saffron is regarded as a stimulant and antispasmodic; but, from the experiments of Dr. Alexander<sup>3</sup>, its powers appear to be inconsiderable. It was known to the ancients, who considered it as a remedy of great activity; in moderate doses exhilarating the spirits, easing pain, and producing sleep; but occasioning head-aches, coma, delirium, convulsive laughter, and even fatal effects, when

<sup>1</sup> This extractive, when pure, is named polychroite by Bouillon La Grange and Vogel, on account of the different colours it is capable of assuming. Chlorine and light destroy its colour in the watery infusion; sulphuric acid changes it to indigo, which gradually becomes lilac; and nitric acid gives it a green hue. Vide *Ann de Chim.* lxxx. p. 186.

<sup>2</sup> Chemists assert, that extractive is insoluble in ether; but I find that when resin also is present in any vegetable matter, ether is capable of taking up some extractive combined with the resin which it dissolves; and when the ethereal tincture is evaporated on the surface of water, these principles are separated, the resin remaining in the form of a pellicle on its surface, whilst the extractive is dissolved, colours the water, and forms, with the solution of muriate of tin, a brown flaky precipitate. Hence ether is a good test of these vegetable principles.

<sup>3</sup> *Experimental Essays*, p. 88.



given in large doses. In modern practice, however, it is scarcely ever given except as a cordial adjunct to more active remedies. The dose in substance is from grs. x. to ʒss; but it has been given in much larger doses without any sensible effect being produced.

Officinal preparations. *Syrupus Croci*, L. *Tinctura Croci sativi*, E. *Confectio aromatica*, L. D. *Pil. Aloes cum Myrrha*, L. *Tinct. Aloes comp.* L. E. D. *Tinct. Cinchonæ comp.* L. D. *Tinct. aloes*, D. *Tinct. Rhei comp.* L.

CROTON. *Spec. Plant, Willd.* iv. 531.

Cl. 21. Ord. 8. Monœcia Monadelphia. Nat. ord. Tricoccæ, Linn. Euphorbiæ, Juss.

G. 1718. Male. Calyx cylindrical, five-toothed. Corolla five-petalled. Stamens 10—15.

Female. Calyx many-leaved. Corolla none. Styles three, bifid. Capsule three-celled. Seed one.

Species 43. *C. Eluteria*.<sup>1</sup> *Elutheria. Med. Bot.* 2d edit. 633. t. 223.

—— 36. *C. Tiglium*. Purging Croton. *Flor. Zeyl.* 343. *Rumph. Amboyn.* iv. p. 98. t. 42. *Rheede Malab.* ii. p. 61. t. 33. *Ray. Hist. Plant.* 167, *Ainslie's Mat. Med. of Hindustan*, 4to. pp. 96. 291.

#### 1. CROTON ELUTERIA.

Officinal. CASCARILLÆ CORTEX<sup>2</sup>, *Lond. Dub.* CROTON ELUTHERIÆ CORTEX, *Edin.* Cascarilla Bark.

Syn. Cascarille (F.), Cascarillrinde (G.), Kaskerilla (Dutch), Kaskarille (Dan. & Swed.), Cascariglia (I.), Chacarilla (S.), Cascarilha (Portug.).

This tree is a native of the Bahama Islands, and has been also found in Jamaica by Dr. Wright. It is a small tree, seldom exceeding twenty feet in height, and branching thickly towards the top. The more tender branches, when broken, ooze out a thick balsamic liquor. The leaves are alternate on short petioles, ovate or cordate, lanceolate, and elongated towards the apex, which is blunt, entire and, on the upper surface, of a bright green colour. The flowers are in axillary and terminal racemes. The petals are whitish, oblong, obtuse, and spreading. The male flower has ten subulate filaments, supporting erect, compressed anthers: the female produces a roundish germen crowned with three bifid, spreading styles, with obtuse stigmas. The capsule is superior, trilocular, and contains a solitary shining seed.

Cascarilla bark is imported chiefly from Eleutheria, one of the Bahama Islands, packed in chests and bales. It consists of pieces about six or eight inches long, scarcely one-tenth of an inch thick, quilled, and covered with a thin, whitish epidermis.

Qualities. — Cascarilla bark has a pleasant, spicy odour, and

<sup>1</sup> It is the *Clutia Eluteria* of Linnæus.

<sup>2</sup> The London College erroneously refers this bark to the *Cascarilla* of Linnæus, the bark of which, however, has none of the sensible qualities of Cascarilla.



a bitter, warm, aromatic taste. The colour of the inside of the pieces is a reddish cinnamon hue, and their fracture close and short, of a dark reddish brown or purple colour. It is very inflammable, and is easily distinguished from all other barks by emitting, when burnt and extinguished, a fragrant smell, resembling that of musk, but more agreeable. Its active constituents are partially extracted by alcohol and water, and completely by proof spirit. Ether takes up one and a half in ten parts; and, when evaporated on the surface of water, leaves a thick pellicle of bitter resin; and dissolved in the water, a small portion of almost colourless, pungent extractive. According to Tromsdorff, who analysed it, 4696 parts yielded the following products: — Mucilage and bitter principle 864, resin 688, volatile oil 72, water 48, and woody fibre 3024 parts.<sup>1</sup> The ethereal tincture shows extractive also to be present, of a greenish yellow colour, very fragrant and pungent.

*Medical properties and uses.* — This bark is a valuable carminative and tonic. It was introduced into practice as such in 1690 by professor Stisser; and was afterwards much used in Germany, particularly by the Stahleans, as a substitute for cinchona bark, in the cure of intermittent and remittent fevers: but although they over-rated its virtues, yet it is an excellent adjunct to the bark in these diseases; rendering it, by its aromatic qualities, more agreeable to the stomach, and increasing its powers. It is successfully employed in dyspepsia, asthma, and flatulent colic; the latter stage of dysentery, and diarrhœa, particularly when occurring after measles: and in the gangrenous thrush peculiar to children.<sup>2</sup> The dose of the powdered bark is from grs. xij. to 3ss., three or four times a day.

Officinal preparations. *Infusum Cascarillæ*, L. *Tinctura Cascarillæ*, L. D. *Extractum Cascarillæ*, D.

## 2. CROTON TIGLIUM.

*Officinal.* TIGLII OLEUM, Lond. Oil of Croton, or Tiglium.

*Syn.* Huile de Croton (F.), — Nervallum cottay unnay (Tam.), Iummal Gota (Duk.), Nepala (Sans.), Naypulum vittilo noonay (Telingoo), Dund (Pers.), Batoo (Arab.), Beri (Malay).

The plant yielding the seed from which this oil is expressed, is a native of the Molucca islands, and of the greater part of the peninsula of India. It has an arboreous stem, covered with a soft, blackish bark. The leaves are ovate-acuminate, serrated and smooth, with two glands seated at the base; and are supported on petioles shorter than the expansion of the leaf. The flowers are in terminal racemes. The seeds, which

<sup>1</sup> *Annales de Chimie*, xxii. 219. and *Thompson's Chemistry*, 4th ed. v. 220.

<sup>2</sup> *Underwood, Diseases of Children*, 4th ed. i. 79.



are contained in trilocular capsules, are oblong, about the size of a large coffee bean, four-sided, flattish on two sides, and convex on the other, with four elevated ridges, running at equal distances from the base to the apex of the seed. The shell of the seed is black; but covered with a soft, pale yellowish-brown epidermis.

Croton seeds are imported into this country in cases; and, owing to the rubbing of the epidermis, when the cases are not completely filled, have generally a mouldy appearance. In this state they were, formerly, known in Europe under the name *Molucca grains*; but, as they were discarded from medical practice on account of their very drastic effects, arising from the imprudent manner in which they were exhibited, they ceased to be an article of commerce, until lately that the expressed oil has been introduced as a purgative: 100 parts of the kernels of the seeds when bruised yield 60 of acrid oil, and 40 of farinaceous matter. The acrid principle resides chiefly in the testa or skin of the cotyledons, and the corculum or embryo, and is mixed with the oil of the cotyledons in its expression. The goodness of the oil, therefore, depends on the seeds being shelled before they are bruised.

*Qualities.* — Croton oil is of a pale reddish-brown colour. Its taste is hot and acrid; and it leaves an uneasy feeling in the mouth and throat, which continues for many hours. Even a minute portion of the kernel of the seed when chewed leaves a hot, pungent sensation on the tongue, which remains for twenty-four hours. The oil is soluble in ether and oil of turpentine. Alcohol takes up two parts out of three, and the solution possesses the acrimony and the cathartic properties of the oil, whilst the undissolved portion is devoid of acrimony and inert when taken into the stomach. But much of what is taken up by the alcohol is fixed oil; and, from the experiments of Dr. Nimmo<sup>1</sup>, Croton oil is composed of 45 parts of an acrid principle, and 55 of fixed oil, resembling the oil of olives.

Dr. Nimmo has suggested the following means of detecting adulterations of Croton oil: pour into a phial, the weight of which is known, 50 grains of the oil; add alcohol, which has been digested on olive oil; agitate well; and, having poured off the solution, add more alcohol of the same kind until the dissolved portion is diffused in such a proportion of the alcohol that each half-drachm measure shall contain equal to one dose of the Croton oil for an adult; — by placing the phial near a fire, to evaporate what remains of the alcohol in the bottle, if the remainder be to that abstracted by the alcohol as 55 to 45, the oil is genuine. If it be adulterated with any

<sup>1</sup> *Journ. of Sciences*, vol. xiii. p. 66 — 69.



fixed oil, the residuum will be larger; if with castor oil it will be smaller than in the genuine oil.<sup>1</sup>

*Medical properties and uses.*—Croton oil is a powerful hydragogue purgative, operating in a very short time after it is taken. It has been given with great advantage in cases of obstinate constipation, convulsions, mania, apoplexy, and other diseases which require, along with the complete evacuation of the intestines, the lessening the circulating mass. The small doses in which this oil produces its effects, require the greatest caution to be observed in its administration, as it has occasionally induced the most dangerous hypercatharsis. In India, where it has long been used, ghee or butter, with orange or rice water or cold butter-milk, and the external effusion of cold water, are employed to counteract its too violent effects, when these occur. It is also used in India as an emenagogue with excellent effects, and as an external application in rheumatic affections.<sup>2</sup>

Croton oil is generally administered in doses of from one to five drops, made into pills with crumb of bread; or, combined with mucilage of gum, sugar and almond mixture, in the form of emulsion. Dr. Nimmo recommends the saturated alcoholic solution, in the dose of fʒss. rubbed up with simple syrup, and mucilage of gum, of each oz. ii., and oz. iv. of distilled water.<sup>3</sup>

CUBEBA. Vide *Piper Cubeba*.

CUCUMIS. *Spec. Plant. Wild.* i. 611.

*Cl.* 21. *Ord.* 8. Monœcia Monodelphia. *Nat. ord.* Cucurbitaceæ. *G.* 1741. *Male.* Calyx five-toothed. *Corolla* five-parted. *Filaments* three.

*Female.* Calyx five-toothed. *Corolla* five-parted. *Pistil* three-cleft. *Seeds* of the gourd argute.

*Sp.* 1. *C. Colocynthis*.<sup>4</sup> Bitter Cucumber. *Med. Bot.* 2d. ed. 189. t. 71.

*Officinal.* COLOCYNTHIDIS PULPA, *Lond.* CUCUMERIS COLOCYNTHIDIS PULPA, *Edin.* COLOCYNTHIS; FRUCTUS MEDULLA, *Dub.* The pulp of Coloquintida, or bitter Cucumber.

*Syn.* Coloquinte (*F.*), Koloquinten (*G.*), Koloquint (*Dutch, Dan. & Swed.*), Coloquintida (*I.*), Pepinero Coloquintida (*S. & Portug.*), Hunzil (*Arab.*), Indráini (*H.*), Indraváruni (*San.*), Makkul (*Ben.*), Peycoomutikai (*Tam.*), Dahuk (*Egyp.*).

This plant is an annual, a native of Turkey and Nubia<sup>5</sup>, flowering from May till August, and much resembling the cucumber in its herbage. The root is branching, and strikes

<sup>1</sup> *Journ. of Sciences*, vol. xiii. p. 66—9.

<sup>2</sup> *Mat. Med. of Hindostan*, 4to. Madras, 1813. The solution of the oil in oil of Turpentine produces a pustulous eruption when it is applied to the skin.

<sup>3</sup> *Journ. of Science.* vol. xiii. p. 69.

<sup>4</sup> *Κολυκυνθίς* Dioscoridis.

<sup>5</sup> Burckhardt, in his travels through Nubia, 4to. p. 184, says, “the ground was covered with the coloquintida, a plant very common in every part of this desert, (Wady Om-gat).”



deep into the ground. The stems are trailing, beset with rough hairs; the leaves are on long petioles, of a triangular form, variously sinuated, obtuse, of a fine green colour on the upper surface, and whitish and rough beneath. The flowers are solitary, axillary, and of a yellow colour. The calyx of the male flowers is bell-shaped; the corolla the same shape with the limb, divided into five-pointed segments; and the anthers, which stand on three short filaments, are long, erect, and adhere together on the outer side. The female flower is like the male, but the filaments have no anthers. The fruit is a round berry or pepo, the size of a small orange, yellow, and smooth on the outside when ripe; trilocular, each cell containing many ovate, compressed, whitish seeds, enveloped by a white spongy pulp.

When the fruit is ripe and yellow, it is peeled and dried in a stove; and in this state it is brought to this country. When it is larger than a St. Michael's orange, and has black, acute pointed seeds, it is not good.

*Qualities.* — Dried colocynthida is inodorous; but has an extremely bitter, nauseous taste, and the pulp feels mucilaginous when chewed. Independent of the seeds, it is altogether composed of a very light, easily torn, white, cellular matter. Ether, alcohol, and water, extract its virtues. The infusion in boiling water has a golden-yellow colour, gelatinizes as it cools, and resembles, except in colour and taste, mucilage of quince seed. This mucilage is soluble in cold water. Alcohol and all the acids coagulate the solution, which is precipitated by solutions of acetate and superacetate of lead, and nitrate of silver. Sulphate of iron strikes with it a deep olive colour. Its colour is rendered also greenish by solution of potass, which precipitates it; but the mucilage is dissolved by solution of ammonia. Ether, digested on the pulp, deposits, when evaporated on the surface of water, a white, opaque, bitter resin, and some extractive; from which the water acquires the bitter taste of the fruit, and precipitates solutions of potass, nitrate of silver, and acetate of lead. From these experiments colocynth pulp appears to consist chiefly of mucus, resin, the bitter principle, and some gallic acid. M. Vauquelin found that an alcoholic tincture of colocynth, when evaporated, yields a brittle, yellow substance, partially soluble only in water, the residue being a white filamentous mass, changing to yellow. This substance he has named *Colocyntine*; and he regards it as the active principle of the drug.<sup>1</sup>

*Medical properties and uses.* — The pulp of this fruit is a very powerful drastic cathartic. It was employed by the an-

---

<sup>1</sup> *Journ. de Pharm.* 1824. 416.



cients in dropsical, lethargic, and melancholic affections; but always with caution, on account of its violent effects. Orfila, from his own observations, asserts, that one or two drachms of it only, applied to the cellular tissue of the interior of the thigh of a man, produced death in the space of twenty-four hours.<sup>1</sup> When given alone, even in moderate doses, it purges vehemently, producing violent gripings, bloody dejections and, not unfrequently, convulsions and inflammation of the bowels. The watery decoction, or the infusion, is much less violent in its operation, and has been recommended in worm cases. It is scarcely ever given alone in any form, but is generally united with other purgatives to quicken their operation. The dose is from grs. iv. to grs. x. triturated with almonds, or gum, or some farinaceous matter.

Officinal preparations. *Extractum Colocynthis*, L. *Extractum Colocynthis comp.* L. D. *Pilulæ Aloes cum Colocynthide*, D.

CUMINUM. *Spec. Plant. Willd.* i. 1440.

Cl. 5. Ord. 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

G. 547. Fruit ovate, striated. *Partial umbels* four. *Involucre* four-cleft.

Sp. 1. *C. Cyminum*.<sup>2</sup> Cumin. *Med. Bot.* 2d ed. 143. t. 56.

Officinal. CUMINI SEMINA, *Lond.* Cumin Seed.

*Syn.* Cumin (*F.*), Ramischer Kumel (*G.*), Komyn (*Dutch*), Kummen (*Dan.*), Spis Kummin (*Swed.*), Semenza di Comino (*I.*), Semilla de Comino (*S.*), Kmin (*Polish*), Siragum (*Tam.*), Zereh (*Pers.*), Jantan (*Malay*), Kimoon (*Arab.*), Iirâ (*H.*), Iiraca (*San.*), Jeera (*Beng.*).

This plant is an annual, a native of Egypt, but cultivated in great abundance in Sicily and Malta; whence the seeds are brought to this country. It flowers in June. It rises eight or ten inches in height, on a slender, round, often procumbent, branching stem. The leaves are of a deep green colour, narrow, linear, and pointed; the flowers purple, in numerous four-rayed umbels; with umbellets having seldom more than four flowers. Both the involucre and involucels consist of three or four subulate unequal leaflets. The corolla is composed of five unequal petals, inflected, and notched at the apex: the filaments support simple anthers; and the germen is inferior, large, ovate, with two minute styles terminated by simple stigmas. The fruit consists of two oblong, striated seeds, united by their flat sides; of a pale brown colour, bristly on their convex surface.

*Qualities.* — Cumin seeds have a strong, peculiar, heavy odour, and a warm, bitterish, disagreeable taste. Water extracts little more than their odour; but alcohol takes up both odour and taste; and yields, when evaporated, an extract

<sup>1</sup> *Leçons faisant partie des Cours de Med. Legale*, de M. Orfila.

<sup>2</sup> *Κυμινον* Dioscoridis.



*Medicinal properties and uses.*—Cumin seeds are carminative and stomachic: but they are chiefly employed as an external stimulant in discussing indolent tumours.

CUPRUM. *Edin. Dub.* Copper.

Copper is a metal of a yellowish or brownish red colour, found very abundantly in many countries in both hemispheres of the globe. It is procured

i. Crystallized, (*Alloy*)      Sp. 1. *Native copper.*

*a.* and combined with iron. 3. *Purple copper.*

4. Grey copper.

5. *Copper pyrites.*

6. *Black copper.*

7. *White copper.*

iii. Oxidized. — 8. *Ruby copper.*

9. *Tile-red copper.*

———— 10. *Copper black.*

11. *Azure copper, or Mountain blue.*

— 12. *Malachite.*

— 13. *Emerald copper.*

14. *Octohedral arseniate of copper.*

15. *Hexahedral arseniate of copper.*

16. *Prismatic arseniate of copper.*

17. *Trihedral arseniate of copper.*

18. *Martial arseniate of copper.*

19. *Phosphate of copper.*

20. *Sandy copper.*<sup>1</sup>

<sup>1</sup> Vide *Aikin's Chemical Dictionary*, art. Copper.



cured chiefly in Cornwall.<sup>1</sup> The ore is first roasted to volatilize the sulphur, which is collected in chambers connected by flues with the kilns. It is then smelted, in contact with the fuel, in a large reverberatory furnace, to separate the iron; which, being less fusible than the copper, remains in the scoria, while the melted copper is drawn off through a plug-hole into earthen moulds. The copper, however, in this state is still very impure; and therefore it is re-melted and granulated; and lastly, refined, by being again melted with the addition of a little charcoal, which brings it to a state fit to bear the hammer, and to answer the various purposes of art.

Pure copper has a yellowish red colour; is sonorous, ductile, malleable, tenacious; has a styptic, disagreeable taste, and emits an unpleasant odour when rubbed. The specific gravity, when it is pure, and has been only fused, is 8.895. It has a granulated texture, and breaks with a hackly fracture; melts at a temperature equal to 27° of Wedgwood; is volatilized by a greater heat; and is oxidized, when heated in contact with atmospherical air, even at a temperature below that of ignition. When exposed to humidity and to air at the same time, it is tarnished, and a green crust is formed on its surface, which is a carbonate of copper.

Although copper in its metallic state was used as a remedy by the ancients, yet it is completely discarded from modern practice: and, notwithstanding, so much has been said of its deleterious effects, there is every reason for believing that clean copper, when taken into the stomach, exerts no action whatever on the system. Two cases of halfpence being swallowed by children, have come under my observation, in one of which the copper coin remained six months in the intestines, and in the other two months. Both were evacuated without having in the smallest degree injured the health, although the impressions on the coins were nearly effaced, and the metal was much corroded. But poisoning from the use of copper utensils in cookery, arises either from the formation of the green carbonate, mentioned above, owing to the vessels not being well cleaned, and the food being allowed to stand for some time in the pan exposed to the air, after it is taken from the fire; or from the formation of *verdegris*, when vinegar, used in making pickles, and other acid liquors intended for internal use, are boiled in brass or copper vessels. The salts of copper thus formed are poisons, exciting inflammation of the stomach; and many fatal accidents have arisen from the practices which produce them,

---

<sup>1</sup> The Parys mine in the isle of Anglesea is now nearly exhausted.



and from the equally dangerous mode of giving a fine green colour to vegetables by boiling halfpence with them; on which account copper utensils should be altogether banished from the kitchen, and also from the laboratory, where they are sometimes employed in making decoctions. The salts of copper may be detected in any suspected liquor, by placing in it a piece of clean polished iron, on which the copper is precipitated in a metallic state; or by dropping into the suspected fluid a solution of ammonia, which produces a beautiful blue colour, if any salt of copper be present. In cases of poisoning by any of the salts of copper, sugar is the best antidote.

The oxides of copper unite with acids, and form salts, which act very powerfully on the animal system; but of these the *subacetate* and the *sulphate* only are admitted into the list of *materia medica*.

### 1. SUBACETATE OF COPPER.

*Officinal.* *ÆRUGO*, *Lond. Dub.* SUBACETAS CUPRI, *Edin.* Verdegriis, or Subacetate of Copper.

*Syn.* Vert de gris (*F.*), Grunspan crystalle (*G.*), Verdegriise (*I.*), Cardenillo (*S.*), Vungala patchei (*Tam.*), Zungar (*Pers. & Duk.*), Sennang (*Malay.*), Zunjar (*Arab.*), Pitrai (*H.*), Pitalata (*San.*).

This salt is principally manufactured in the south of France, at Montpellier, and Grenoble.<sup>1</sup> In the former place, the *marc* of the grape, that is, the cake which remains in the wine-press after the juice is expressed, composed of the husks and stalks, is moistened with water, or with wine if poor, and disposed so as to excite in it the acetous fermentation. When this takes place it is spread in jars between well hammered plates of copper, heated over a pan of burning charcoal; a layer of fermented marc being placed between each plate of copper. The jars, each of which contains about 40 lbs. of copper, besides marc, are then loosely stopped with straw, and left at rest for ten, fifteen, or twenty days, at the end of which time the marc begins to whiten; and the copper is found to be covered with a green crust, interspersed with distinct silky green crystals. The plates are then moistened with water, and set up in racks, face to face, in a cellar: this is repeated once in seven days for six or eight times, until a thick coat of verdegriis is formed, which is scraped off; and the copper plates again subjected to the same process till they are completely corroded. When the plates are first used, the verdegriis is apt to be black, unless their surfaces be previously rubbed with a solution of verdegriis, which is suffered to dry before they are used.

Verdegriis in this rough state is sold by the makers, who are generally women belonging to the wine farms about Montpel-

<sup>1</sup> Vide Chaptal's Account of the Manufacture, *Phil. Mag.* vol. iv. 71.



lier, to commissioners, by whom it is further prepared. After being well beaten in wooden mortars, it is pressed down in bags of white leather, a foot in depth and ten inches wide, in which it is dried in the sun; and thus a loaf of verdegris is formed, which cannot be pierced with a knife.

In this process the copper is oxidized, and the oxide combined with a small portion of acetic acid, forming a subacetate, which is mixed with vegetable extractive matter and the stalks and husks of grapes. The Grenoble verdegris is a purer subacetate, being prepared by simply disposing plates of copper in a proper situation, and repeatedly moistening them with distilled vinegar until the surface be oxidized and changed into verdegris.

The Subacetate of Copper is imported into this country in the leather sacks, or bags, in which it is dried, each containing from fourteen to thirty pounds weight. But it is, also, now prepared in Great Britain.

*Qualities.* — Good Subacetate of Copper is inodorous. It seems at first nearly insipid, although exceedingly styptic; but leaves a strong metallic taste in the mouth. The mass is dry, not deliquescent, of a hard, pulverulent, foliaceous texture, and a beautiful, blueish-green colour. Distilled water at  $60^{\circ}$  dissolves 0.56 parts, while 0.44 remain in the state of a fine green powder, long suspended in the solution: that part which is dissolved is a superacetate of copper, the filtered solution reddening litmus paper; whilst the insoluble powder is a subacetate mixed with the impurities. When boiling water is used, the insoluble part is of a brown colour. Sulphuretted hydrogen gas decomposes the solution, precipitating a black sulphuret of copper. A small cylinder of phosphorus put into the solution is rapidly covered with a coat of metallic copper. According to Proust, verdegris consists of 43 parts of acetate of copper, 27 black oxide of copper, and 30 of water in intimate combination. Mr. Phillips, by a recent analysis, has ascertained that the quantity of the acetic acid in crystallized acetate of copper is 49.2, and that of the peroxide of copper 39.2, in 100 parts; the remainder being water<sup>1</sup>: while Dr. Ure asserts the following to be the composition of verdigris: — acetic acid 52.0, peroxide of copper 39.6, water 8.4.<sup>2</sup> By a more recent analysis of Vauquelin, that excellent chemist makes the proportion to be, acetic acid 51, oxide of copper 40, and water 9: or acetic acid 2 atoms, oxide of copper 1 atom, water 3 atoms.

<sup>1</sup> *Annals of Phil.* (new series), vol. i. p. 418.

<sup>2</sup> *Dict. of Chemistry*, art. Copper.



Besides the stalks and husks of grapes, verdegris is often adulterated with sand, and other earths. These are discovered by dissolving it in diluted sulphuric acid, which takes up the whole of the subacetate, and leaves the impurities. Or by boiling it in twelve or thirteen times its weight of distilled vinegar, allowing the undissolved part to settle, and ascertaining its amount. The addition of muriate of barytes will detect any admixture of the sulphate or the tartrate of copper.

*Medical properties and uses.* — Verdegris is tonic, and emetic. It has been used in epilepsy: and extolled as an emetic in cases which require that the stomach should be quickly evacuated, without weakening it, as in incipient phthisis: but its internal exhibition is always dangerous, and to be avoided. It is, however, a useful detergent and escharotic application to foul ulcers, and the callous edges of sores; and to consume fungus: but it is seldom used, although it is milder than the sulphate. It is also employed as a collyrium in chronic ophthalmia.<sup>1</sup>

The dose of verdegris to produce its tonic effect is under gr. ss.; and to operate as an emetic, from gr. j. to gr. ij. In overdoses it quickly proves fatal, acting both locally, and on the nervous system; and, on dissection, the coats of the stomach appear much thickened, and of a green colour. We formerly suggested the idea that fine filings of iron might precipitate the copper in its metallic state, and operate as an antidote: but this is now rendered unnecessary, as the experiments of Duval and others have proved that sugar is the antidote of cuprous poisons.<sup>2</sup>

Officinal preparation. *Ærugo preparata*, D. *Ung. Subacetatis Cupri*, E. *Linimentum Æruginis*, L. D.

## 2. SULPHATE OF COPPER.

*Officinal.* CUPRI SULPHAS, *Lond. Edin. Dub.* Sulphate of Copper.

*Syn.* Sulphate de cuivre (*F.*), Schwefelsaures Kupfer (*G.*), Kopper vitriol (*Belg.*), Blaae vitriol (*Danish*), Vitriuolo blo (*I.*), Caparosa (*S.*), Vitriolo de Cobre (*Portug.*), Zungbar (*Arab.*), Tuteya (*H.*), Tutt'ha (*San.*), Toorushco (*Tam.*), Neelatota (*Duk.*), Palmanicum (*Cyng.*):

A considerable part of this salt, which is the blue vitriol of commerce, is obtained by evaporation from the water of some copper mines. Its origin is derived from the natural sulphurets of copper, which suffering a chemical change from exposure to a moist atmosphere, are converted into the sulphate, and washed down by the rain and other water of the

<sup>1</sup> Dr. Paris (*Pharmacologia*) says, that it is the active ingredient in *Smellome's Eye Salve*, a nostrum in much repute.

<sup>2</sup> Vide *Traité des Poisons*, &c. par P. M. Orfila, tome i. p. 289.



mines.<sup>1</sup> It is also obtained by roasting copper pyrites, and exposing it to the action of air and moisture; in which case, as well as in the former, the compound is oxidized by attracting the oxygen of the surrounding atmosphere, at the same time that it changes the sulphur into sulphuric acid: so that by the gradual combination of these the sulphate is produced, and is then extracted by solution, and crystallized.

*Qualities.* — Sulphate of Copper is inodorous, and has a very harsh, acrid, styptic taste. It always reddens vegetable blues. It is in semitransparent crystals, which undergo a slight degree of efflorescence when exposed to the air: their form is that of a rhomboidal prism; and their colour a deep rich blue. Its specific gravity is 2·1943: and, according to Chenevix, it consists of 42·6 parts of hydrate<sup>2</sup> of copper, 33·0 of acid, and 25·4 of water of crystallization.

Sulphate of Copper is soluble in four parts of water at 60°, and less than two at 212°. It is insoluble in alcohol. The solution reddens litmus paper, the salt being truly a bi-sulphate. It is decomposed by the alkalies and alkaline carbonates, the sub-borate of soda, acetate of ammonia, the acetate and super-acetate of lead, and acetate of iron, nitrate of silver, oxymuriate of mercury, tartrate of potass, muriate of lime; and is precipitated by all the astringent vegetable infusions and tinctures; all which substances are therefore incompatible in prescriptions with this salt.

*Medical properties and uses.* — Sulphate of Copper is emetic astringent, and tonic, when taken internally. With a view to its emetic effect, it has been given in the early stage of phthisis, and where laudanum has been taken as a poison<sup>3</sup>; and as an astringent and tonic, in alvine hæmorrhages, intermittent fever, epilepsy, and some other spasmodic affections: but as the list of materia medica contains equally powerful and less injurious remedies, its internal exhibition ought to be altogether discontinued. Externally it may be employed to give a healthy stimulus to indolent foul ulcers, in which I have found it extremely beneficial; and as an escharotic, to consume fungus. Pledgets dipped in a weak solution of it are, also, sometimes used as a styptic in epistaxis, and other external hæmorrhages; and a still weaker solution is a useful collyrium in some kinds of ophthalmia. It forms the base of

---

<sup>1</sup> From this water at the Parys mine, a large supply of copper is obtained, by decomposing the sulphate, by throwing into the water old iron hoops.

<sup>2</sup> The hydrate consists of copper 25·6, oxygen 6·4, and water (which is intimately combined with the oxide) 10·6 parts. This water, as well as the water of crystallization, is expelled when the sulphate is decomposed by heat.

<sup>3</sup> See *Medico-Chirurgical Transactions*, vol. i.



a very unchemical preparation, Bate's *aqua camphorata*<sup>1</sup>, which the late Mr. Ware recommended, diluted with sixteen parts of water, in the purulent ophthalmia of infants.

As an emetic the dose is from grs. ij. to xv., in fʒij. of water; but as a tonic it should be given in the form of pill, beginning with gr.  $\frac{1}{4}$ ., and gradually increasing the dose to grs. ij.

Official preparation. *Solutio Cupri Sulphatis composita*, E. *Cuprum Ammoniatum*, L. E. D.

CURCUMA. Roscoe, *Linn. Trans.* viii. 354.

Cl. 1. Ord. 1. Monandria Monogynia. Nat. ord. Scitamineæ, Linn. Drymyrrhizæ, Juss.

Gen. Char. *Anther* double, two-spurred. *Filament* petal-like, three-lobed, bearing the anther in the middle.

Sp. 1. C. *Zedoaria*.<sup>2</sup> Zedoary. *Amomum Zedoaria*. Willd. *Spec. Plant.* i. 7.

Official. ZEDOARIA RADIX, Dub. Zedoary root.

Syn. Zeduire (F.), Zedoar wurzel (G.), Zedoaria (I.), Judwar (Arab.), Nirbisi (H. & San.), Banhaldi (Beng.).

This plant is a perennial, a native of the East Indies, growing in sandy open places in Ceylon and Malabar, where it is named *Acua* by the Brahmins; and flowering in April and May. The root is tuberous, oblong, and about the thickness of a finger; the leaves are palmated, broad, lanceolate, subsessile on their sheath; sericeous underneath; colour an uniform green.<sup>3</sup> The scape, which rises from among the leaves, is naked, and terminated by a lax, cylindrical, truncated, lateral spike of flowers.

The best Zedoary root comes from Ceylon, in firm, short, wrinkled pieces, of an ash colour externally. It should be heavy, and not worm-eaten.

*Qualities*. — The odour of Zedoary root is fragrant, and somewhat like that of camphor, the taste biting, aromatic, and bitterish, with some degree of acrimony. The pieces break with a short, close fracture, are pulverulent, and internally of a brownish-red colour. Its active principles are partially extracted by water, and more completely by alcohol. In distillation with water, a heavy, greenish blue essential oil is obtained, which deposits camphor. It seems to contain, independent of its aromatic and bitter principles, a large proportion of fecula.

<sup>1</sup> The following is the formula for Bate's preparation: — R *Cupri sulph. Boli gall. aa.* grs. xv. *Camphoræ* grs. iv. *Solve in aq. ferv.* fʒiv., *dilueque cum aq. frig.* O. iv. *ut fiat Collyrium*.

<sup>2</sup> The excellent reasons given by Mr. Roscoe for separating this plant from the genus *Amomum*, induce me to prefer his authority to that of Willdenow in this instance.

<sup>3</sup> Roxburgh; vide *Asiatic Researches*, 8vo. vol. xi. p. 165.



*Medical properties and uses.* — This root is tonic and carminative. It was much employed by Avicenna, and other Arabians, in vomitings, cholics, lientery, difficult menstruation, and as an antidote for venomous bites. It is certainly an agreeable stomachic, and useful in flatulent cholic, but it is scarcely ever used by modern practitioners. The dose of the powdered root may be from grs. viij to 3ss, two or three times a day.

CUSPARIA. Vide *Bonplandia Trifoliata*.

CYDONIÆ SEMINA. Vide *Pyrus Cydonia*.

DAPHNE.<sup>1</sup> *Spec. Plant. Willd.* ii. 415.

Cl. 8. Ord. 1. Octandria Monogynia. Nat. ord. Vepriculæ, Linn. Thymelææ, Juss.

G. 773. Cal. none. Cor. four-cleft, corollaceous, withering, inclosing the stamens. *Drupe* one-seeded.

\* *Flowers lateral*.

*Species* 1. *D. Mezereum*. Common Mezereon. *Med. Bot.* iv. 716. t. 68. *Smith, Flor. Brit.* 420.

*Officinal* MEZEREI CORTEX, *Lond.* DAPHNES MEZEREI CORTEX, *Edin.* RADICIS CORTEX, *Dub.*<sup>2</sup> The bark of the root.

*Syn.* Laureole gentile (*F.*), Kellerkals (*G.*), Pepperbompje (*Dutch*), Kielderhals (*Dan.*), Tibast (*Swed.*), Wyleze lyko (*Pol.*), Mezereo (*I.*), Mezereon (*S.*), Mezerao (*Port.*).

Mezereon grows wild in England and the north of Europe; but for medical use, and as an ornamental shrub, it is cultivated in gardens. Its flowers expand in March, before the leaves. It is a hardy plant seldom exceeding four feet in height, with a strong, woody, branching stem, covered with a smooth, grey cuticle and a tough fibrous inner bark. The root is of a fibrous texture, pale coloured, with a smooth olive coloured bark: the leaves which are protruded from the extremities of the branches are tender, pale green, deciduous, lanceolate, sessile, entire, and smooth: the flowers are of a pale rose colour, odorous, surrounding the twigs in clusters below where the leaves are sent off; they are sessile, two, three, and four clustered, with deciduous bractes at the base of each cluster; monopetalous, tubular, and the lip divided into four ovate, spreading segments. The stamens are alternately shorter; the four higher ones displaying their yellow anthers at the mouth of the tube: the germen is oval, supporting a flattish stigma on a very short style; and the fruit is a red, pulpy drupe, containing one round seed. There is a variety of the mezereon with white flowers and yellow fruit, but the medicinal effects of both are the same.

<sup>1</sup> Δαφνη Theophrasti et Dioscoridis.

<sup>2</sup> The Dublin College, in its Pharmacopœia, has quoted *Eng. Bot.* 119. erroneously, the plate referred to being that of *Daphne Laureola*, Spurge Laurel.



For medical use, the roots are dug up in the autumn, after the leaves have fallen. The cuticle of the dried root is corrugated, and the inner bark has a white, cotton-like appearance.

*Qualities.* — The inner bark of every part of this plant, when fresh, is very acrid, capable of producing inflammation, vesication, and a discharge of serum when applied to the skin; and when chewed, excites a considerable heat of the mouth and fauces, which continues for many hours afterwards. The fruit is equally acrid, acting as a corrosive poison, if eaten. The bark retains its acrimony when dried. It yields its virtues to water and vinegar. By digesting the bark in alcohol, then evaporating the liquid to separate the resin, and diluting the residual fluid with water, filtering, and adding acetate of lead, Vauquelin obtained a copious yellow precipitate, which, when freed from the lead by means of sulphuretted hydrogen gas, he found to be a vegetable principle, *sui generis*. He has given it the name of *Daphnin*.<sup>1</sup>

*Medical properties and uses.* — It operates as a stimulating diaphoretic, increasing the general arterial action, and determining powerfully to the surface; but is apt to disorder the primæ viæ, and occasion vomiting and purging. It was long externally employed, as a stimulus to ill-conditioned ulcers; and the recent bark macerated in vinegar and applied to the skin, is recommended in France for producing and keeping up a serous discharge in chronic local affections. To form the issue, the bark must be renewed every night and morning; and afterwards once in twenty-four hours, to keep open the drain. Dr. Withering employed it successfully as a local stimulant in a case of difficulty of swallowing occasioned by paralysis. Although the case was of three years' standing, the patient recovered the power of swallowing in about a month, by very frequently chewing thin slices of the root. For this purpose it should be sliced longitudinally, as the acrimony resides in the bark only, the woody fibre being nearly inert. Internally, a decoction of this bark has been used against chronic rheumatism, scrofulous swellings, lepra, and some other cutaneous diseases; and, till lately, it was considered an anti-venereal remedy of great efficacy. The dose in substance is gr. j. to grs. x.

Officinal preparation. *Decoctum Daphnes Mazerei*, E. *Decoctum Sarsaparillæ comp.* L.

DATURA. *Spec. Plant. Willd.* i. 1007.

Cl. 5. Ord. 1. Pentandria Monogynia. Nat. ord. Solanaceæ, Linn. Solaneæ, Juss.

---

<sup>1</sup> *Ann. de Chim.* lxxxiv. p. 174.



G. 377. *Corolla* funnel-shaped, plaited. *Calyx* tubular, angled, deciduous. *Capsule* with four valves.

*Species* 2. *D. Stramonium*. Thorn Apple. *Med. Bot.* 2d edit. 197. t. 74. *Smith, Flor. Brit.* 253.

*Officinal.* DATURÆ STRAMONII HERBA, *Edin.* STRAMONIUM; HERBA, *Dub.* The herbaceous part of the Thorn Apple plant.

*Syn.* Pomme Epineuse (*F.*), Stechapfel (*G.*), Dornappel (*Dutch*), Piigaeble (*Dan.*), Spikklubba (*Swed.*), Jondera (*Pol.*), Stramonio (*I.*), Estramonio (*S. & Port.*), Karoo Oomatie (*Tam.*), Dhetoora (*H.*), Jeuz massel (*Arab.*), Goozgneah (*Pers.*), Rctecubung (*Malay*), Kalu D'hatoora (*Beng.*), Kaloo Uttana (*Cyng.*), Kistna Datura (*Sans.*).<sup>1</sup>

This annual plant is a native of America, but is now naturalized to this country, and found growing on dunghills and by road-sides<sup>2</sup>, from the fruit ejected from gardens; flowering in July and August. It rises about two feet in height, with a round stem, branching and dichotomous above; spreading and leafy. The leaves, which spring from the forks of the stem on long round petioles, are large, of a dark green colour on the upper surface, and pale beneath; irregularly ovate-triangular in figure, sinuated, and unequal at the base. The flowers are, large, axillary, and solitary, on short erect peduncles. The calyx is about two inches in length, tubular, pentangular, and five-toothed: the corolla longer, of a white colour, funnel-shaped, and plaited: with the filaments, which support oblong, flat anthers, adhering to the tube; and the style filiform, terminated with a thick club-shaped stigma. When the corolla and its included parts drop, the calyx also separates, except the base, which remains, and, becoming reflex, enlarges with the receptacle as a support to the fruit. The fruit is a large, fleshy, ovate-roundish, four-cornered capsule, beset with sharp, awl-shaped spines; four-celled at the base, two-celled at the apex, and containing a great number of reniform compressed seeds. Both the leaves, capsule, and seeds, are medicinally used.<sup>3</sup>

*Qualities.*—The whole herb has a narcotic, foetid odour, producing head-ach; a bitterish, nauseous taste, and gives to the saliva a deep green tinge when chewed. According to Wedenberg<sup>4</sup>, it contains gum (*mucus*?) and resin, a volatile matter (which I find to be carbonate of ammonia), and a narcotic principle, which has lately been ascertained by M. Brandes to be an alkaline salt. He obtained it from the seeds, in which it is combined with malic acid, and named it *Daturium*. It is nearly insoluble in water and in cold alcohol, but boil-

<sup>1</sup> These oriental synonymes although inserted under this species, yet are those of *D. Fustuosa*.

<sup>2</sup> Very common about London.

<sup>3</sup> According to Dr. W. Ainslie, this species of *Datura* is not found in India; but the *D. fustuosa*, Dkétoora (*Hind.*), Dustura (*Sans.*), is well known and medicinally used. *Mat. Med. of Hindostan*, 4to. p. 42.

<sup>4</sup> *Dissertatio Medica de Stramonii Usu, &c.* Upsal. 4to.



ing alcohol dissolves it, and, on cooling, lets it fall in flocculi. It is crystallized with difficulty, but has been obtained in quadrangular crystals. It forms neutral salts with the acids.<sup>1</sup> The medicinal virtues of the herb are extracted both by water and alcohol. The watery infusion is transparent, with a very pale yellow hue, which is dissipated by acids, but very much deepened by the alkalies. It throws down whitish precipitates with acetate and superacetate of lead, and a black precipitate with nitrate of silver. Solution of sulphate of iron strikes a deep olive colour, and muriate of mercury renders it milky; but neither is precipitated till after a very considerable time.

*Medical properties and uses.* — Thorn-apple is narcotic and stimulant. Baron Stoerck first recommended it as an internal remedy, in cases of mania and epilepsy; but, as Cullen remarks, he was less violent in his commendations of it than of the other narcotic plants which he introduced.<sup>2</sup> It was afterwards tried by other continental physicians with unequal success; particularly by Greding, who made the greatest number of trials of it. But the most decided experiments in its favour have been made by Dr. Barton, of America, who regards it as a remedy of great efficacy. He found that, when the dose of the dried herb was gradually increased to thirty grains, it dilated the pupil, and produced paralysis of the eyelids; effects which were removed by a blister. Cataplasms of the bruised fresh leaves have been successfully used as an application to inflammatory tumours, and for discussing masses of indurated milk in the breasts of nurses: and an ointment made with the powdered leaves allays the pain of hæmorrhoids. Smoking the plant in the manner of tobacco, affords relief in the paroxysm of spasmodic asthma; a practice introduced into England from Ceylon. The inspissated, expressed juice of the leaves has been usually given; and the extract has been lately found almost specific in severe chronic pains. The root is given by the native practitioners in the Carnatic, in violent head-achs. Hufeland recommends the form of tincture. The dose of the extract, at first, should not exceed grs. ss. twice a day, increasing the quantity gradually, until grs. xii. be taken in twenty-four hours.

Several instances of the fatal effects of stramonium, when eaten by mistake, are recorded by authors.<sup>3</sup> It produces delirium, stupor, convulsions, furious madness, paralysis, cold sweats and death. As these effects depend on the narcotic principle, the best antidote is vinegar, after the stomach has been emptied.

<sup>1</sup> *Journ. de Physique*, xci. p. 144.

<sup>2</sup> *Materia Medica*, ii. 281.

<sup>3</sup> It is said to be sometimes used by the Turks instead of opium, or as a substitute for wine; and the Chinese infuse the seeds in beer. *Spratt's Hist. of the Roy. Soc.* 162.



DAUCUS. *Spec. Plant. Willd.* i. 1389.

*Cl.* 5. *Ord.* 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

*G.* 53. *Cor.* somewhat rayed. *Flor.* of the disk abortive. *Fruit* hispid with hair.

*Species* 1. *D. Carota*.<sup>1</sup> Common Carrot. *Med. Bot.* 2d edit. 130. t. 50. *Smith, Flor. Brit.* 300.

*Officinal.* DAUCI (*hortensis*) RADIX. — (*agrestis*) SEMINA, *Lond.* DAUCI CAROTÆ RADIX, *Edin.* DAUCUS SYLVESTRIS; SEMINA, *Dub.* The root of the cultivated Carrot, and the seed of Wild Carrot.

*Syn.* Carotte (*F.*), Karotte; Mohrrübe (*G.*), Karoot (*Dutch*), Derreurt (*Dan.*), Morot (*Swed.*), Marchew (*Pol.*), Carota (*I.*), Zanahoria (*S.*), Cenoira, (*Port.*), Jezer (*Arab.*), Gajer (*H.*), Garjara (*San.*), Carrot halung (*Tam.*), Zerdek (*Pers.*).

The carrot is a biennial indigenous plant. In its wild state it is found abundantly in pastures and on hills<sup>2</sup>; flowering in June and July. It is cultivated for culinary purposes and feeding cattle. The root is spindle-shaped, fleshy, and of a yellow colour; throwing up a round furrowed stem, which rises about two feet in height, and sends off long, erect, naked, floriferous branches. The leaves are large, petiolated, thrice-pinnate, cleft, and hairy. The flowers are in many-rayed compound umbels, flat on the top, and spreading, but after the flowering season they become condensed in a concave form. The involucre consists of several narrow trifid leaves; the involucrel is more commonly simple. The marginal flowers are white or yellow; the central, which are abortive, often of a dark blood colour. The seeds are in pairs, egg-shaped, convex, rough, and bristled on one side and flat on the other. These appearances are changed by cultivation, which also increases the size and the nutritious matter of the root.<sup>3</sup>

*Qualities.* — The sensible qualities of the *root* of the *cultivated carrot* are well known. It contains chiefly mucilage and sugar. The *seeds* of the *wild* variety have an aromatic odour, and a warm pungent taste; qualities depending on an essential oil, which may be separated by distillation with water.

*Medical properties and uses.* — The root of the garden carrot is emollient and antiseptic, and is successfully used, when boiled and beaten to a pulp, as a poultice to correct the discharge of fœtid and ill-conditioned sores, and to allay the pain of carcinomatous and phagedenic ulcers. The seeds are carminative and diuretic, and hence useful in flatulent cases; but they possess no efficacy in gravel, for which they have been extolled.<sup>4</sup>

<sup>1</sup> Σταφυλινος αγριος Dioscoridis.

<sup>2</sup> We have seen it in great abundance on the range of chalk hills which overlook Ryegate in Surrey.

<sup>3</sup> Carrots are seldom sufficiently boiled; but when well boiled, they are very digestible and nutritious.

<sup>4</sup> The red central flowers of the umbels have been extolled in epilepsy.



The dose of the bruised seed is from ʒj. to ʒj. or more.

DELPHINIUM. *Spec. Plant. Willd.* ii. 1226.

Cl. 13. Ord. 3. Polyandria Trigynia. Nat. ord. Multisiliquæ, Linn.  
Ranunculaceæ, Juss.

G. 1061. Cal. none. Pet. five. Nect. bifid, horned behind. Pods three or one.

\* \* *Three-capsuled.*

Species 13. *D. Staphisagria*.<sup>1</sup> Staves Acre. *Med. Bot.* 2d edit. 471. t. 168.

Official. STAPHISAGRIÆ SEMINA, *Lond. Dub.* DELPHINII STAPHISAGRIÆ SEMINA, *Edin.* Staves Acre seeds.

Syn. Staphisagre (*F.*), Stephanskraut laus kærner (*G.*), Luiskruid (*Dutch*), Lunsurt (*Dan.*), Staffans frö (*Swed.*), Gnidosz ziele (*Polish*), Stafisagria (*I.*), Piojera (*S.*), Alvarraz (*Portug.*).

This species of larkspur is a biennial plant, a native of the south of Europe, flowering from June to August. It is a handsome plant, from one to two feet in height, with a downy, erect, purplish, simple stem; and palmated leaves, the lobes of which are five or seven in number, of a pale green colour, oblong, ovate, and sometimes acutely indented. The flowers are of a blue or purplish colour, in an open, terminal spike, and supported on long flower-stalks; the uppermost petal projected backwards so as to form a hollow spur, which encloses two spurs of the superior leaflets of the nectary. The filaments are about twenty, and short, bearing large, yellow anthers; the germens three, close together, tapering, downy, and crowned with short filiform styles, having simple stigmas. The seeds are rough, brown, triangular, and contained in straight, oblong capsules. They are usually imported from Italy; for although the plant is occasionally reared in our gardens<sup>2</sup>, yet, it is difficult to preserve it through the winter, so as to enable it to perfect its seed.

*Qualities.* — Stavesacre seeds have very little odour, but that little is disagreeable; their taste is bitter, acrid, and hot. They are yellowish within and covered with a rough blackish cuticle. Their virtues are partially extracted by water and completely by alcohol. MM. Lassaigne and Feneulle discovered that their active properties depend on a peculiar alkaline principle, which they named *Delphine* or *Delphinia*. It is a white, crystalline powder; inodorous, acrid and bitter to the taste. Its acetate is a virulent poison.

*Medical properties and uses.* — These seeds are emetic and cathartic, but their operation is so violent that they are never internally administered. Owing to their stimulating powerfully the salivary gland, when chewed, they have been used

<sup>1</sup> *Σταφίς αγρία* Dioscoridis.

<sup>2</sup> It was first cultivated in England by Gerard in 1596.



as a masticatory in tooth-ach; but they are chiefly employed in powder, mixed with hair-powder for destroying pediculi of the head.<sup>1</sup>

DIANTHUS.<sup>2</sup> *Spec. Plant. Willd.* ii. 671.

Cl. 10. Ord. 2. Decandria Digynia. *Nat. ord.* Caryophyllei, *Linn.*  
Caryophylleæ, *Juss.*

G. 893. *Calyx* cylindrical, one-leaved, with four scales at the base.

*Petals* five, with claws. *Capsule* cylindrical, one-celled.

\* \* *Flowers* solitary, many on the same stem.

*Species* 9. *D. Caryophyllus*. Clove Pink, or Gillyflower. *Med.*  
*Bot.* 2d edit. 579. t. 205. *Smith, Flor. Brit.* 461. *Eng. Bot.*  
214.

*Officinal.* DIANTHI CARYOPHYLLI FLORES, *Edin.* CARYOPHYLLUM RUBRUM; FLORES, *Dub.* Flowers of the Clove Pink.

*Syn.* Giroflée musquée (*F.*), Gewürzhaft riechende Gartennelke (*G.*), Garofano (*I.*), Clavel (*S.*).

This is a perennial plant, a native of Italy, but found growing wild on ruined walls, as those of Rochester, Deal, and other old castles in England, flowering in July.<sup>3</sup> It is cultivated in gardens for medicinal use; in which case the flowers become full, and improve their native odour. The root is firm and fibrous. The stems, which rise from among tufts of channelled, linear, glaucous leaves, that are finely toothed a little above the base, but entire and smooth towards the apex, are erect, branched, and paniced, bearing many solitary flowers. The whole herb is glaucous. The calyx is striated, tubular, five-cleft, of a pale green colour, with four rhomboid, pointed scales at its base, about one-fourth of its length: the petals vary in colour from a pale flesh-red to the deepest carnation; are unequally crenated, smooth at the orifice, with long, narrow, whitish claws. The stamens are often short and abortive; the styles long, recurved, and downy on the upper side.

The varieties of this species of *Dianthus* produced by horticulturists are very numerous. For medicinal use, those should be chosen which have the richest colour and most spicy odour. The petals must be picked when the flower is fully blown.

*Qualities.* — The odour of the petals is fragrant and aroma-

<sup>1</sup> *Delphinia*, in a separate state, exerts violent poisonous properties, in very small doses; acting chiefly on the nervous system. To procure it, boil the powdered seeds in distilled water, and press through a cloth. Filter the decoction, and boil it for a few minutes with pure magnesia: re-filter, and boil the residuum left on the filter with alcohol; and, lastly, cautiously evaporate the solution. A white powder is obtained, which is *Delphinia*.

<sup>2</sup> From *Διος αυθος*, the flower of Jove; yet it was unknown to the ancients.

<sup>3</sup> Ray and Hudson suppose it to be an outcast of gardens, and not an indigenous plant of England. It is mentioned by Chaucer under the name "cloue gilofre;"

———— "to put in ale,  
Whether it be moist or stale."



tic, resembling that of the clove spice, the taste slightly bitter, and subastringent. Both water and alcohol extract their sensible qualities; and they yield an essential oil by distillation with water. The infusion strikes a black colour with sulphate of iron; acids redden its colour; and alkalies change it to green.

*Medical properties and uses.* — Notwithstanding the testimony of our forefathers in favour of the efficacy of these flowers in nervous affections<sup>1</sup>, modern practitioners value them merely for their sensible qualities, and employ them only to give an agreeable flavour and fine colour to a syrup, which is a pleasant vehicle for the exhibition of more active medicines.

Officinal preparations. *Syrupus Dianthi Caryophylli*, E. D.

DIGITALIS. *Spec. Plant. Willd.* iii. 283.

Cl. 14. Ord. 2. Didynamia Angiospermia. *Nat. ord.* Luridæ, *Linn.* Scrophulariæ, *Juss.*

G. 1155. *Cal.* five-parted. *Corolla* bell-shaped, five-cleft, belly-ing. *Capsule* ovate, two-celled.

*Species* 1. *D. Purpurea*. Purple Foxglove.<sup>1</sup> *Med. Bot.* 2d edit. 218. t. 78. *Smith, Flor. Brit.* 665. *Eng. Bot.* 1297. *Withering's Account of Foxglove*.

*Officinal.* DIGITALIS FOLIA, *Lond. Dub.* DIGITALIS PURPUREÆ FOLIA, *Edin.* Foxglove leaves.

*Syn.* Grande Digitalis (*F.*), Purpurrother Fingerhut (*G.*), Paarsch vingerhoed (*Dutch*), Fingerbölle (*Dan.*), Fingerborrsort (*Swed.*), Paluszniczek (*Pol.*), Digitale Porporina (*I.*), Dedalera purpurea (*S.*), Deda leira (*Portug.*).

Foxglove is an indigenous biennial plant, found growing generally on the sides of hills and roads, where the soil is dry, sandy, or gravelly; flowering from the middle of June to nearly the middle of August. The root is knotty and fibrous, sending up an erect stem about four feet in height; round, downy, and leafy. The lower leaves are in tufts, large, about eight inches in length and three in breadth, ovate and pointed, with bordered fleshy peduncles: the upper or stem leaves are alternate, sparse, and lanceolate; and both kinds have bluntly serrated, nearly crenate edges, and wrinkled velvety surfaces; with the upper surface of a beautiful deep green colour, and the under paler and more downy. The flowers, which are numerous, are attached on footstalks to one side of the upper part of the stem, so as to allow them to hang down and form a very elegant terminal spike. At the base

<sup>1</sup> Gerard says, that when they are made into a conserve, they are "exceeding cordiall, and wonderfully above measure comfort the heart."

<sup>2</sup> It was named *Digitalis* by Fuchsius, (*Plantarum omnium Nomenclaturæ*, 1541,) the first author who notices its medicinal properties from the German name Fingerhut, a finger stole. It had been previously described by Tragus under the name of *Campanula sylvestris*.



of each footstalk is a sessile pointed floral leaf. The uppermost segment of the calyx is narrower than the other four; the corolla is monopetalous, of an oblong bell-shape, and about the size of the little finger of an ordinary glove; bellying on the lower side, with a short, tubular base. The upper lip is lightly cloven, and more reflected than the under, which is larger. The corolla is guarded by long hairs at the mouth: its general colour is a bright pinkish purple, with the tube white, and the bellying part sprinkled on the inside with dark purple spots on a white ground, which give to the outside a speckled appearance. The filaments are white, curved, bearing large, oval, yellow anthers; the germen is pointed, supporting a simple style with the apex cloven. The seed-vessel, which is a pyramidal capsule, with a double partition produced by the inflected margins of the valves<sup>1</sup>, contains many small, ferrugineous, punctated seeds.

The leaves are the parts of the plant medicinally used. They should be gathered when the plant is in flower, and those only which are fresh selected. "The leaf-stalks and midrib should be rejected, and the remaining part be dried either in the sun-shine, or on a tin pan or pewter dish before the fire, or the plant be hung up, each leaf separate, in a warm kitchen." Practitioners ought annually to obtain a supply of the recent leaves, in the month of July, and dry them themselves; as in the herb-shops they are often so ill dried as to appear black, in which state they are useless. The powder should be kept in closely stopped, opaque phials.

*Qualities.* — Recent foxglove leaves are inodorous; but in the dried state they have a slight narcotic odour, and a bitter nauseous taste. Both water and alcohol extract their virtues. The watery infusion has a pale, olive-green colour, with the unpleasant odour and taste of the plant. It does not precipitate solutions of galls, tartarized antimony, nor sulphate of iron, which only deepens its colour, but it produces a yellowish precipitate with muriate of mercury, and a blackish-violet, very copious one with nitrate of silver. The dry powder, which should have a beautiful green colour, moistened and triturated with lime or calcined magnesia, and a glass rod dipped in muriatic acid held over it, exhibits copious white fumes, proving the presence of ammonia. The presence of ammonia is also apparent in the tincture; which is rendered milky by water.<sup>2</sup> Ten grains of the powder macerated in f ʒss of sulphuric ether lost

---

<sup>1</sup> *Gærtner de Fructibus*, &c. i. 247. t. 53. f. 6.

<sup>2</sup> Destouches, a French chemist, who analysed Foxglove, obtained also much carbonate of ammonia by distilling the aqueous extract. He obtained besides sulphate of potash and of lime, phosphate of lime, carbonates of lime, and of potash, and acetate of ammonia.



three grains of its weight, and yielded all its colour to the ether: and the ethereal tincture, on being evaporated on water, left a pellicle of dark-green, unctuous, resinous matter, whilst some yellowish extractive was dissolved in the water, and precipitated afterwards by oxymuriatic acid. From this imperfect analysis, foxglove appears to contain ammonia, extractive, and a pea-green resinous matter in which its narcotic power resides. More recent analysis has discovered a peculiar alkaline principle which has been termed Digitaline, or Digitalia. It is procured by digesting the leaves in ether, filtering and evaporating and dissolving the residuum in water, which is filtered and then treated with hydrated oxide of lead, which, being evaporated and the residuum digested in ether, yields up the Digitaline, which is obtained in the separate state by evaporation. It is a brown pasty substance, very bitter and very deliquescent.<sup>1</sup>

*Medical properties and uses.* — *Digitalis* is directly sedative,<sup>2</sup> and diuretic. It weakens the force of all the vital functions; and by a proper exhibition of it, the frequency of the pulse may be diminished any number of pulsations, and regulated at the pleasure of the practitioner; whilst at the same time it admits, to a certain extent, of the employment of such medicines as increase the firmness of the arterial action, and give tone to the habit. When given to the full extent of which the system can admit, the pulse intermits, and vertigo, indistinct vision, and nausea with vomiting, or purging occur; and if, after these indications, the quantity be still increased, or if any considerable portion of the recent herb be inconsiderately swallowed, it produces delirium, hiccough, cold sweats, convulsions, syncope, and death. It is supposed, notwithstanding its sedative effects are acknowledged, to increase the action of the absorbent system; but although the discharge of urine is very considerably increased during its use, and the load of water with which the body is oppressed in dropsies be thrown off, yet it is not easy to conceive how a direct sedative can operate as a stimulant; and therefore the *modus operandi* of foxglove, in producing its diuretic effect, may be regarded as still unexplained.<sup>3</sup>

As a *sedative* foxglove was early used in some acute diseases, but its powers were not understood. It is now efficaciously

---

<sup>1</sup> *Bib. Univ.* xxvi. p. 102.

<sup>2</sup> This term implies, as I understand it, any substance which diminishes the action of the heart and arteries, without first increasing it.

<sup>3</sup> Dr. Baildon observed a curious effect of posture in ascertaining the real effects of digitalis on the pulse. When, by gradually increased doses, he took it to the extent of grs. vj. in the day, the pulse fell to 40 from 110. But when it was actually at 40, the erect posture would raise it to 100; when sitting it was 72, and when lying down 40. He observed the same effect in several patients to whom he gave it. — *Edin. Med. and Surgical Journal*, iii. 271.



employed in inflammatory diseases; in active hæmorrhages, particularly from the uterine vessels, when the pulse is sharp, throbbing, and frequent; in mania; in scrofula; and in most cases of increased vascular action, or in which it is essential to lessen the usual impetus of the blood, as in aneurism. In mania it acts rather as a narcotic, soothing the nervous system and procuring sleep to the patient, than as a sedative.

Its beneficial effects in phthisis pulmonalis were known so early as 1710; and passing experience confirms the justness of the praises bestowed on it by a writer of that period.<sup>1</sup> Dr. Ferriar found its utility in this complaint much increased by combining it with myrrh and sulphate of iron.<sup>2</sup> Its use has also been extended to venereal ulcerations, chronic rheumatism, hooping-cough, and some spasmodic affections. As an external application it has long been used in Italy<sup>3</sup> to cleanse old sores; and Hufeland recommends it to be used in the form of fomentation for dispelling glandular swellings.

As a *diuretic* the use of foxglove was reintroduced by Dr. Withering in 1775.<sup>4</sup> He found that its beneficial effects in dropsies were more certainly obtained in those constitutions in which there is a laxity of fibre, pale countenance, feeble intermitting pulse, and cold skin; and where the swelling easily pits. But in florid habits, with great strength, tense fibre, and a hot dry skin, its sedative effect is perceived, but no diuresis follows. “If the belly in ascites be tense, hard, and circumscribed; or the limbs in anasarca be solid and resisting, we have but little hope.”<sup>5</sup> Experience has confirmed these judicious observations; and it is found that where this favourable state does not exist, it may be produced by bleeding, and the free use of neutral salts, calomel, and squill, all of which lower the system. The diuretic effect is checked when much nausea is present; and Withering says purging also checks it; but I have not found this to be the case. The kinds of dropsy in which its effects are most useful, are ascites, anasarca, hydrothorax, and that species of swelling which succeeds parturition, phlegmasia dolens, where the legs and thighs swell, become pale and semi-transparent, with pain in both groins. It has also been found of the greatest service when conjoined with nitrous acid, in the dropsy which occurs in broken-down constitutions, that have been long harassed by mercury.<sup>6</sup> *Digitalis* will not cure a dropsy attended with palsy, unsound viscera, or other complications

<sup>1</sup> Salmon. See *The Edin. Med. and Surg. Journal*, v. 303.

<sup>2</sup> *Essay on the Medical Properties of Digitalis*.

<sup>3</sup> Araldi (*Digitalis*) tutte pioghe salda.

<sup>4</sup> He was induced to try it from finding it the active ingredient in a family recipe for the cure of dropsy, regarding which his opinion was asked.

<sup>5</sup> *Withering's Account*, &c. 186.

<sup>6</sup> Carmichael on *Diseases which have been confounded with Shypilis*, 4to. p. 63-5.



of disease: but by allaying the urgency of the symptoms, it gains time for other medicines to act. No benefit has hitherto been obtained from its use in hydatids, and hydrocephalus.

Foxglove is administered in substance, or in decoction, or the watery infusion, or in tincture (see *Preparations*). When given in substance, it is frequently combined with aromatics, soap, or ammoniacum; and most advantageously with calomel and opium, when it is required only to produce its diuretic effect. It is always proper to begin with a dose not exceeding gr. j. of the powdered leaves, given in a pill twice a day; and gradually to increase it till its effects are apparent either on the kidneys, the stomach, the pulse, or the bowels. The medicine must then be discontinued; but in dropsy it may be repeated after an interval, if the whole of the water be not evacuated. During its employment diluents are useful and necessary; and immediately it is discontinued the strength should be recruited by generous food, steel, and cordial tonics. The deleterious effects of an overdose are to be counteracted by cordials, as, brandy, mint tea, and opium; and when these are not sufficient, by blisters.

Officinal preparations. *Decoctum Digitalis* D. *Infusum Digitalis*, L. E. *Tinctura Digitalis*, L. E. D.

DOLICHOS. *Spec. Plant. Willd.* iii. 1037.

Cl. 17. Ord. 4. Diadelphia Decandria, Nat. ord. Papilionacæ, Linn. Leguminosæ, Juss.

G. 1349. At the base of the standard, two oblong parallel scales, compressing the wings underneath.

\* *Twining.*

Sp. 16. *D. pruriens*. Cowhage. *Med. Bot.* 2d ed. 422. t. 153. *Chamberlaine's Practical Treatise on the Efficacy of Stizolobium or Cowhage.*

Offi. DOLICHI PUBES, *Lond.* DOLICHI PRURIENTIS PUBES, *Edin.* DOLICHES SETÆ LEGUMINUM, *Dub.* The hairs of the Dolichos pod.

*Syn.* Kiwách (*H.*), Capicach'hu (*San.*).

This is a perennial climbing plant, a native of America, and the East and West Indies. In Bengal, where it is named *Cadjuct*, it flowers in the cool months from September to March. The root is fibrous; the stem herbaceous, cylindrical, voluble, climbing, and branching; with ternate leaves, on footstalks from six to fourteen inches long, given off alternately at the distance of a foot from each other. The central leaflet is rhomboidal, the two lateral ones oblique, and all of them smooth on the upper surface, and hairy beneath. The flowers are papilionaceous, of a blood colour, peduncled, in pendulous solitary spikes, which hang from the axillæ of the leaves. The fruit is a coriaceous pod about four or five inches long, curved like the letter *f*, thickly covered with briskly short brown hairs; and containing three or five oval, compressed seeds.

The pods we receive are brought from the West Indies.



If incautiously touched, the spiculæ with which they are beset separate easily, and sticking in the fingers, occasion the most intolerable itching.

*Medical properties and uses.* — The spiculæ of dolichos pods operate as a mechanical anthelmintic. They have been found particularly useful in expelling the round worm, *lumbricus teres*; the spiculæ irritating, and aiding its expulsion, by wounding it, without affecting the intestines. The best mode of preparing the remedy, is to dip the pods in syrup or molasses, and then, with a knife, to scrape off the hairs along with the syrup. When the mixture attains the thickness of honey, it is sufficiently impregnated with the hairs, and is fit for use.

The dose of this mixture, for a child of three or four years old, is a tea-spoonful given in the morning for three days, and then followed by a brisk cathartic.

**DORSTENIA.** *Spec. Plant. Willd.* i. 682.

*Cl.* 4. *Ord.* 1. Tetrandia Monogynia. *Nat. ord.* Scabridæ, *Linn.* *Urticæ, Juss.*

*G.* 244. *Receptacle* common, one-leaved, fleshy, in which solitary seeds are nestled (or placed in sockets without attachment).

*Spec.* 5. *D. Contrajerva*, *Contrayerva*. *Med. Bot.* 2d edit. 705. 240.

*Officinal.* CONTRAJERVÆ RADIX, *Lond.* DORSTENIÆ CONTRAJERVÆ RADIX, *Edin.* *Contrajerva Root.*

*Syn.* *Contrajerva (F.), Giftwurz (G.), Contrajerva (Dutch), Contraherva (Port.), Contrajerva (I.), Contrahierba (S.).*

This is a perennial plant, a native of Peru, Mexico, and some of the West India islands. The root is fusiform, knotty, and branching, compact, furnished with many rough fibres; externally of a brown colour, and internally whitish. It sends up several leaves, which are about four inches in length and the same in breadth; of an irregular shape, but in general deeply lacinated into five or seven obtuse parts; and placed on long radical footstalks, winged towards the leaves. The fructification, which is remarkable, and on radical stalks or scapes which rise about four inches high, is a fleshy receptacle, shaped like an animal placenta, about an inch long, and three-fourths of an inch broad, placed vertically; and containing on its upper surface very small, scarcely conspicuous flowers, situated closely together, immersed in the receptacle, and occupying the whole of its disc. The capsule possesses an elastic power when ripe, by which the seeds are thrown out with considerable force.

Monardus is the first author who mentions this root, which, he says, is called *Contrajerva*<sup>1</sup> by the Spanish Indians, on account of its alexipharmic qualities. Dr. Houston<sup>2</sup>, however, asserted that the officinal *contrajerva* was the root of two other species of *Dorstenia*, the *D. Houstonia* and *D. Drakena* of

<sup>1</sup> The Spanish, for the English word *antidote*, is *contrahierba*. The Mexican name of the root is *Tuzpatlis*.

<sup>2</sup> *Phil. Trans.* No. 421. p. 195.



Willdenow; but the British Colleges follow the authority of Linnæus. It is brought to this country from the West Indies, packed in bales, in pieces of about two inches long.

*Qualities*. — Contrajerva root has a peculiar, but not unpleasant odour, and a bitterish, warm taste, leaving a pretty lasting impression on the tongue. It preserves its qualities when dried; and in the state of powder. Both water and alcohol, assisted by heat, extract its virtues. The watery decoction is of a dark, brownish-red colour, and exceedingly mucilaginous. — The alcoholic tincture reddens litmus paper, is not altered by a solution of sulphate of iron, but is precipitated by water.

*Medical properties and uses*. — This root is a stimulant sudorific and tonic. Huxham and Pringle first pointed it out as a remedy well suited to fevers of a typhoid type, and it is often employed in malignant, eruptive diseases, dysentery, and in some kinds of diarrhoea. It is also useful in atonic gout, chronic rheumatism, and the fever attending dentition in weak infants.

The dose of the powdered root is from grs. v. to ʒj; but it is seldom used alone.

Officinal preparation. *Pulv. Contrajervæ compositus*, L.

DRYOBALANOPS. *Asiatick Researches*, vol. xii. p. 539. Cl. 13. Ord. 1. Polyandria Monogynia. Nat. ord. Guttiferæ, Juss. *G. nova*. *Calyx* one-leafed, permanent; enlarged into a gibbous cup, with five ligulate long scariose wings. *Corolla* five-petalled. *Capsule* superior, one-celled, three-valved, seed solitary. *Embryon* inverse, without perisperm.

*Species*. D. *Camphora*. *Asiatic Res.* vol. xii. p. 539. Plate 4.

*Officinal*. CAMPHORA, *Lond. Edin.* CAMPHORA, RESINA, *Dub.* Camphor.

*Syn.* Camphre (*F.*), Der Kampferr (*G.*), Canfora (*I.*) Alcanfor (*S.*), Cafoor (*Arab.*), Carpoorum (*Tam.*), Carphura (*San.*).

That the camphor brought to Europe, from the islands of Sumatra and Borneo, is not the product of the *Laurus Camphora*, was remarked by Kœmpfer<sup>1</sup>; but no accurate description has yet been given of the tree which yields it. Mr. H. T. Colebrooke has, however, lately been enabled to determine the genus to which it belongs, and to which we have referred it, from the examination of some seeds in a very perfect state, sent from Tapanooly to Calcutta.

The *Dryobalanops Camphora* is a native of forests on the north-western coast of Sumatra, and especially in the vicinity of Tapanooly. It is stated by Mr. Prince to be found in abundance from the back of Ayer Bongey, as far north as Bacougan, a distance of 250 miles. It grows to a great height, and the *trunk*, which is arboreous and covered with

<sup>1</sup> “ Camphoram naturalem et crystallinam perquam pretiosam ac rarum imperitur arbor in Sumatra et Borneo insulis. Sed hæc arbor ex *Daphneo sanguine* non est.” *Amœn. Exot.* p. 773.



a brownish bark, often measures six or seven feet in diameter. The *leaves* are opposite below and alternate above, elliptical, obtusely acuminate, or rather beaked, parallel-veined, entire, smooth; 3.7 inches long, 1.2 broad, and supported on short *petioles*; with subulate, caducous *stipules* in pairs. The *perianth* is a one-leaved, five-parted, persistent capsule, which is superior, ovate, woody, fibrous, longitudinally furrowed, embraced at the base by the calycine hemispherical cup, and surrounded by its enlarged leaflets, which are converted into remote, foliaceous, spatulate, rigid, reflex wings: one-celled and three-valved. The seed is solitary, conforming to the cavity of the capsule: and has a strong terebinthinate fragrance.

The camphor forms in the heart of the tree, occupying portions of a foot and a foot and a half long, at certain distances: but the younger trees yield oil only, which has nearly the same properties as the camphor, and would ultimately be converted into the concrete substance. The natives, in searching for the camphor, make a deep incision in the trunk, about fourteen or eighteen feet from the ground, with a *billing* or Malay axe; and when it is discovered they fell the tree, and cut it into junks of a fathom long, which are again split. The camphor is found in a concrete state, and resembles whitish flakes in perpendicular layers, occupying a space the thickness of a man's<sup>1</sup> arm. A middling sized tree will yield nearly eleven pounds, and a large tree, double that quantity.

For the *qualities and medical properties of Camphor*, see *Laurus Camphora*.

**DULCAMARA CAULIS.** Vide *Solanum Dulcamara*.

**ERYNGIUM.** *Spec. Plant. Willd.* i. 1356.

*Cl.* 5. *Ord.* 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

*G.* 518. *Flowers* capitate. *Receptacle* paleaceous.

*Spec.* 6. *E. maritimum*, Sea Eryngo, or Holly. *Med. Bot.* 2d ed. 120.

*t.* 46. *Smith Flora Brit.* 289. *Eng. Bot.* 718.

*Officinal.* ERYNGIUM; RADIX, *Dub.* Eryngo Root.

*Syn.* Panicaut commun (*F.*), Eringio (*I.*), Eryngo (*S.*).

The sea holly is an indigenous perennial plant, growing abundantly on the sea shores, and flowering in July and August. The root is long, and creeping; sending up an erect, branching, round, obscurely furrowed, and leafy stem, nearly eighteen inches in height. The lower leaves are reniform, plaited, lobed, and petiolate; those of the stem are sessile; and the whole are smooth, rigid, toothed with sharp spines like those of the holly; and of a pale glaucous colour, with the nerves and spines white. The flower heads are terminal, conical, and supplied with paleæ; with an involucre of many,

<sup>1</sup> The Camphor thus found, is called by the natives *Se Tantung*.



spreading, pointed leaves. The paleæ, which separate the florets, are tricuspid, longer than the florets, and spinous. The teeth of the calyx are erect, and spinous; the corolla consists of five oblong blue petals, with inflected points. The filaments are longer than the corolla, the styles are filiform with simple stigmas; and the germen inferior and beset with hairs, becoming two oblong, connected seeds.

The root should be dug up for use, when the seed is ripe in autumn. It is white internally, and covered with a brown cuticle.

*Qualities.* — The root of eryngo has scarcely any odour; and only a soft, sweetish, slightly aromatic taste. Its virtues are extracted by water.

*Medical properties and uses.* — It is supposed to be diuretic and aperient: and has been recommended in gonorrhœa, jaundice, and visceral obstructions; but it is an inert, useless substance, and might with propriety be altogether rejected from the list of materia medica.

EUGENIA.<sup>1</sup> *Spec. Plant. Willd.* ii. 959.

*Cl.* 12. *Ord.* 1. Icosandria Monogynia. *Nat.ord.* Hesperidiæ, *Linn.* Myrti, *Juss.*

*G.* 972. *Calyx* four-parted, superior. *Petals* four. *Berry* one-celled, one-seeded.

*Species* 24. *E. caryophyllata*.<sup>2</sup> The Clove-tree. *Med. Bot.* 2d edit. 538. t. 193. *Journal de Physique*, tome xiv. 47. t. 1.

*Officinal.* CARYOPHYLLI. CARYOPHYLLI OLEUM, *Lond.* CARYOPHYLLUS AROMATICUS. *Floris germen, et ejus Oleum volatile*, *Edin.* CARYOPHYLLUS AROMATICA; CALYX, OLEUM ESSENTIALE, *Dub.* Cloves and Oil of Cloves.

*Syn.* Cloves. Clousele Girofles (*F.*), Gewürz nelken (*G.*), Kruidnagel (*Dutch*), Kryde nellike (*Dan.*), Kryddnegliker (*Swed.*), Gozdz do Potraw (*Pol.*), Gwasditschka (*Russian*), Clavo de espicia (*S.*), Cravo da India (*Port.*), Garofano (*I.*), Kerunful (*Arab.*), Laung (*H.*), Lavanga (*San.*), Craumboo (*Tom.*), Bruah Lawang (*Malay*). *The oil.* Huile de Girofle (*F.*), Nelkenöhl (*G.*), Olio di Garofano (*I.*), Azeyte de Clavos (*S.*), Craumbootylum (*Tam.*), Woorála tail (*Cyng.*).

The clove-tree is a native of the Moluccas, where it was originally abundantly found, particularly at Machian; but the narrow policy of the Dutch led them to destroy almost all the trees except those which they cultivated on the islands of Amboyna, Honimoa, Oma, and Nousalant, so as to give them a monopoly of the trade, which they have held since 1638. The French, however, obtained some plants, which they carried to the Isle of France, in 1770, and thence, in 1774, to Cayenne. In 1789 it was also introduced into the island of Dominica by William Urban Buêe, Esq.; and, in 1803, into the island of Sumatra, by Mr. William Roxburgh.

<sup>1</sup> This genus was named after Prince Eugene of Savoy.

<sup>2</sup> Καρυόφυλλα Græcorum.



At all these places it is now cultivated. It is a handsome tall tree, rising upon a stem of very hard wood, covered with a greyish smooth bark. The leaves are oblong, lanceolate, and pointed at both ends; firm, with many parallel nerves, on each side of the midrib; entire, sinuated, and supported on brown petioles about half the length of the leaf. The colour of the leaves is dull green, and when bruised their odour is strong, and aromatic. Its flowers are in terminal bunches, or corymbose panicles, which generally consist of 9, 15, or 21 flowers. The calyx is oblong, woody, and divided at the brim into four small, toothed segments. The corolla consists of four, roundish, notched, small petals; inclosing many slender filaments inserted into the calyx, bearing simple anthers. The germen is oblong, with a simple style: the fruit an inferior, coriaceous, bilocular berry.

The clove tree yields its first crop of cloves at the age of six years; and attains its highest state of bearing at twelve. The existence of the tree is limited to twenty-four years. Although the unopened flowers of this tree, and even the leaves, particularly their petioles, are extremely aromatic and odorous, yet the flowers are inodorous when they are fully blown; and the real fruit is not aromatic.<sup>1</sup> The cloves are the unexpanded flowers, which are first obtained when the tree is six years old. At Amboyna they are collected from October to December, when they begin to redden. They require to be dried quickly; on which account they are first immersed in boiling water, then exposed to smoke and a heat of 120° Fahr. till they begin to assume a brown hue; and afterwards the drying is finished in the sun. In the West Indies those cloves which are dried altogether in the sun are considered the best.

Cloves were first introduced into Europe by the Arabians, who brought them from India. They are imported into this country from the Dutch settlements; the best in chests, and an inferior kind in bags. The oil is brought in bottles; but a considerable quantity is drawn in this country. The best variety of the Amboyna cloves is smaller and blacker than the other varieties, very scarce, and as a mark of pre-eminence is named the *Royal* clove. The Dutch sometimes mix among the best cloves, those from which the oil has been drawn; and the fraud is not easily discovered, as the used cloves regain part of their flavour by this mixture. The oil is also much adulterated: and when it has a hot, fiery taste, and a great depth of colour, it may be suspected.

*Qualities.* — Good cloves have a strong, fragrant, aromatic

---

<sup>1</sup> *Journal de Physique*, l. c.



odour, and a hot, acrid, aromatic taste, which is very permanent. In form they resemble a small nail, scarcely exceeding half an inch in length; with a roundish conical head, and directly under it four sharp, spreading points, concave above. Their colour is deep reddish brown; the conical part of the head being lighter, and yellowish; and this part is very easily separated. To the touch they feel somewhat greasy. Water extracts their odour, but little of their taste: alcohol takes up both; and when evaporated, the extract is pungent and fiery, without any odour. Ether extracts completely their sensible qualities: and when the tincture is evaporated on water, a considerable portion of a very pungent, hot, unctuous resin, and some extractive remain.

Cloves yield by distillation in water one-sixth of their weight of a heavy, nearly colourless oil, which becomes yellow by age. It has the flavour of the cloves, but is comparatively milder. The Dutch oil is deeper, and of a reddish colour; and is extremely pungent and fiery; owing, it is supposed, to its containing in solution some of the resin of the cloves extracted by alcohol.<sup>1</sup>

*Medical properties and uses.* — Cloves are stimulant in a greater degree than any of the other aromatics. They are sometimes given alone in dyspepsia, when it is attended with a very languid state of the circulation, and a sense of coldness in the stomach; and in atonic gout: but they are chiefly used as corrigents to other medicines. The oil is used as a corrigent to griping extracts; and sometimes as a local application in tooth-ach. The dose of powdered cloves may be from grs. v. to grs. x.; that of the oil m̄ij. to m̄vi., triturated with sugar.

Officinal preparations. *Infusum Caryophyllorum*, L. *Spiritus Lavandulæ comp.* D.

EUPHORBIA.<sup>2</sup> *Spec. Plant. Willd* ii. 881.

Cl. 11. Ord. 3. Dodecandria Trigynia. *Nat. ord.* Tricoccæ, *Linn.* Euphorbiæ, *Juss.*

G. 959. *Corolla* four or five-petalled, fixed to the calyx. *Calyx* one-leafed, ventricose. *Capsule* tricoccous.

*Species* 7. E. *Officinarum*.<sup>3</sup> Officinal Euphorbium Plant. *Amcœnit. Acad.* vol. iii. p. 102. *Jackson's Morocco*, p. 81. *fig.*? *Bruce's Abyssinia*, vol. v. p. 41. *fig.*?

<sup>1</sup> Vauquelin obtained an oil resembling that of the clove from the leaves of *Agathophyllum ravenara*.

<sup>2</sup> Antonius Musa and Euphorbus were brothers; the former was physician to Augustus Cæsar, the latter to Juba, king of Lybia. Cæsar raised a statue to Musa: Juba named this plant after Euphorbus. "Ubi Jam Musæ statua? Periit! evanuit! Euphorbii autem perdurat, perennat, nec unquam destrui potest." *Crit. Bot.* 86.

<sup>3</sup> Δένδρον εὐφορβίον Dioscoridis.



*Officinal.* EUPHORBIAE GUMMI-RESINA, *Lond.* Euphorbium.

*Syn.* Euphorbe (*F.*), Euphorbium (*G.*), Winkel-euphorbium (*Dutch*), Pruskäda (*Swed.*), Euforbio (*I.*), Euphorbio (*S.*), Saynd kadood (*H.*), Ukcil Nefsch (*Arab.*), Nura-shÿ (*Beng.*), Shuddraykullu pomt (*Tam.*).

This is a perennial, succulent, shrubby plant, a native of Africa, where it grows in great abundance. The plant described and figured by Bruce under the name of *Kol-Quall*, and that which Jackson, in his Account of Morocco, says, the Arabs and Shellahs call *Dergmuse*, appear to be the same or varieties of the *E. Officinarum*. When arrived at maturity, it has a simple, erect, round stem, about five feet high; angled or furrowed with 18 or more longitudinal fissures. From the summit branches are thrown out in every direction, going off first horizontally, and then ascending, so as to give to the whole plant the appearance of the skeleton of a large goblet supported on a stalk or foot. The branches are about an inch in diameter, more distinctly angled than the stem, scolloped, and furnished with prickles everywhere double. It has no leaves, but, instead of them, tubercles adjoining to each pair of prickles. The flowers are sessile, on the extremities of the branches, of a crimson colour. The calyx is of one piece, persistent, with a four or five-toothed lip: the petals are four, turbinate, gibbous, thick, truncated, unequal in situation, and fixed by claws to the margin of the calyx. The filaments are more than twelve, threadlike, longer than the corolla, coming forth at different times, and carrying each two globular anthers: the germen is trigonous, with a simple, short style, crowned with three semibifid obtuse stigmas. The capsule is tricoccous, pedicellated, elastic; with round solitary seeds.

The succus proprius of all the species of *Euphorbia* is white, and concretes by exposure to the air into a solid substance. The *Euphorbium* brought to this country is said to be the product of some other species, besides the plant we have described: for instance, *E. antiquorum* and *E. canariensis* of Willdenow. Mr. Jackson says, that in the lower regions of Mount Atlas the inhabitants collect the concreted gum resin, which they call *furbiune*, in September. It is obtained by making slight incisions in the branches of the plant with a knife, from which a milk-like juice exudes, and forms into tears of an oblong or roundish form. The quantity yielded is so considerable, that the plants are cut once only in four years; the supply then obtained being sufficient for that space of time for all Europe. — The recent juice is so corrosive as to erode the skin wherever it touches; and the people who gather the gum are obliged to tie a cloth over the mouth and nostrils, to protect them from the acrid dust



of the withered branches, which induces the most violent sneezing.<sup>1</sup>

Euphorbium is imported in serons, each of which contains from 100 to 150 lbs. weight. It is in small, hollow, somewhat forked pieces, which appear as if the Euphorbium had concreted round the pedicels of the flowers; and it is often mixed with the tricoccous seeds, and other impurities.

*Qualities*.—It is inodorous; and, when first chewed, has little taste, but it soon gives a very acrid, burning impression to the tongue, palate, and throat, which is very permanent, and almost insupportable. Its specific gravity is 1.124. Water, when triturated with it, is rendered milky, but actually dissolves one-seventh part only of the quantity employed: alcohol dissolves one-fourth part, and affords a clear straw-coloured tincture, which is rendered milky by the addition of water: ether takes up six parts in ten, forming an opaline infusion. When the ethereal tincture is evaporated on water, it leaves on the side of the glass a pellicle of transparent resin, and on the water a cake of opaque adhesive whitish matter, which I found to consist of wax and resin, resembling an officinal plaster; while the water is rendered milky. The acrimony resides in the resinous matter. The analysis of Braconnot<sup>2</sup> makes 100 parts of Euphorbium to contain 37.0 of resin, 19.0 wax, 20.5 malate of lime, which was mistaken for gum, 2.0 malate of potass, 5.0 water, 13.5 woody matter, and 3.0 loss. He regards the resin as peculiar, from its being insoluble in alkalies, but soluble in the sulphuric and nitric acids.

*Medical properties and uses*.—Euphorbium possesses powerful cathartic, emetic, errhine, and rubefacient properties. It has been given as a hydragogue in dropsies: but owing to the violence of its effects its internal use is now exploded: neither as an errhine can it be used alone, for it occasions so much inflammation as to produce hæmorrhage from the nostrils, and swell the integuments of the head. When properly diluted, however, with starch or any other inert powder, and cautiously used, it is an effectual and excellent errhine in lethargy, deafness, palsy, amaurosis, and similar cases.

FARINA. Vide *Triticum hybernum*.

FERRUM. Iron.

*Syn.* Fer (*F.*), Ferro (*I.*), Eissen (*G.*), Yzer (*Dutch*), Jern (*Dan. Swed.*), Ferro (*Portug.*), Zelaco (*Polish*), Hierro (*S.*), Loha (*H. Duk. Sam.*), Ayas

<sup>1</sup> Bruce says, "When the tree (Kol-Quall) grows old, the branches wither, and in place of milk, the inside appears to be full of powder, which is so pungent, that the small dust which I drew upon striking a withered branch seemed to threaten to make me sneeze to death, and the touching of the milk with my fingers excoriated them as if scalded with boiling water." *Appendix*, 4to. p. 43.

<sup>2</sup> *Annales de Chimie*, lxxviii. 44.







and setting fire to the lowest stratum of coal. This part of the process expels any sulphur, water, or carbonic acid, with which the ore may be combined; and it is then smelted with coke in a conical furnace of the strongest masonry; the heat being raised to a very high degree by passing a blast of condensed air through the furnace, and to facilitate the separation of the melted metal, lime is used, as a flux. The scoria are drawn out through an opening towards the bottom of the furnace; and the melted metal, which is collected in a cavity at the bottom, is run off into moulds. In this state it is called pig, or cast-iron: and requires to be again fused and submitted to the action of the hammer, or passed between rollers, before it is sufficiently pure either for the majority of the purposes of art or of medicine.

Pure forged or bar iron is of a bluish white or grey colour, of a fibrous texture, and very brilliant in the fracture. It emits a peculiar odour when rubbed, and has a styptic taste. Its specific gravity varies from 7·6 to 7·8. It is attracted by the magnet, and becomes magnetic; properties which distinguish it from all other metals. It is very malleable, but less so than gold, silver, or copper; and is more ductile, tenacious, and elastic, than any other metal. Iron can be ignited by percussion, and melts at 158° of Wedgwood. Its surface is soon tarnished and oxidized when exposed to the air; and the oxidizement is much hastened by the presence of water, which it decomposes. Percussion at a high temperature separates from its surface oxidized scales; the sparks produced by its collision with flint are oxidized; and in the state of wire, when made red hot at one extremity, and introduced into a bottle of pure oxygen gas, it burns with great splendour, and is oxidized in globules. According to Proust, completely oxidized iron consists of 48 parts of oxygen and 52 of iron.

Iron is of all the metals the least injurious to the animal system, being in no respect poisonous. It was medicinally used by the ancients; for Dioscorides, we know, employed it quenched in wine as a remedy for dysentery; and its use was by no means unfrequent as an external application for the cure of malignant ulcers. The effects of iron, however, as an internal remedy, were very little understood until more modern times. It acts as a powerful tonic, increasing the general excitement, promoting the digestive powers and healthy secretions, giving a more florid hue to the blood, and augmenting in a great degree the energy of the muscular fibres. It answers the intentions for which it is prescribed more effectually, when it is given in small doses, minutely divided as it is found in chalybeate springs, and its use long continued. The diseases in which it is used are those which are dependent on, or attended with, a weak, languid, leucophlegmatic habit of body, as chlorosis, hysteria, dyspepsia, fluor albus,



gleet, passive hæmorrhagies, palsy, scrofula, rickets, and the last stage of phthisical affections; it is also beneficial in convalescence from almost all acute diseases, and has been lately recommended as a specific in cancer. The use of iron is contra-indicated whenever the inflammatory diathesis prevails, or there is any particular fulness of the vessels; or an increased secretion of bile, particularly in sanguineous habits. In these states of the system it occasions heat, thirst, headach, laborious respiration, and many other unpleasant symptoms; but, when given in a proper state of the body, few medicines are capable of producing more beneficial effects.

For the purposes of medicine, soft, malleable iron undergoes various preparations (see *Preparations and Compositions*): but at present we have to notice it only as it is mentioned in the lists of materia medica of the British pharmacopœias.

#### 1. METALLIC IRON.

*Officinal.* FERRI RAMENTA ET FILA, *Lond.* FILA ET LIMATURA, *Edin.* FERRI SCOBS, *Dub.* Iron filings and wire.

*Syn.* Limailles de Fer (*F.*), Gopulvertes Eissen (*G.*), Limatura di Ferro (*I.*), Limadura de hierro (*S.*), Cerumboo podie (*Tam.*), Sōw-ik (*Esquimaux.*).

These filings are obtained from the workers in iron; but as they are often mixed with copper filings and other impurities, it is necessary, in order to purify them, to draw them upwards through a sieve, or piece of coarse gauze, with a magnet.

*Medical properties and uses.* — Metallic iron exerts no action on the living system, unless it meets with acid in the stomach, in which case it becomes tonic. Iron filings, therefore, are not adapted for all the cases in which chalybeate remedies prove useful; and are chiefly suited to those cases of dyspepsia, hysteria, chlorosis, and general debility, which are accompanied with acidity in the first passages. When iron is oxidized by the assistance of watery fluids, hydrogen gas is evolved; hence, when the filings are rendered active in the stomach, foetid eructations are produced, and the fæces are coloured black; which, therefore, are evident symptoms of the medicine having taken effect. As an anthelmintic, iron filings may operate mechanically, and dislodge worms; but, even in worm cases, the oxidizement of it in the stomach renders it more useful.

Iron wire is used for pharmaceutical preparations, on account of the purity of the iron from which it is made; as the softest and purest iron only can be drawn.

The filings are given in the form of powder combined with some aromatic, or made into an electuary with honey, or in pills in combination with myrrh, ammoniacum, or some bitter extract. The dose may be from grs. v. to ʒss.

*Officinal preparations,* *Ferri Limatura Purificata*, *E.* *Oxidum Ferri Nigrum purificatum*, *E. D.* *Ferri Acetas*, *D.* *Subcarbonas Ferri præparatus*, *E. D.* *Carbonas Ferri præcipitatus*, *E. D.* *Ferri Sulphas*, *L. E. D.* *Ferrum tartarizatum*, *L. E.* *Tinctura Acetatis*



*Ferri, D. Liquor Ferri alkalini, L. Vinum Ferri, L. D. Murias Ammonicæ et Ferri, E. Sulphuretum Ferri, E. Rubrum Ferri Oxidum, E.*

## 2. OXIDIZED IRON.

*Officinal.* FERRUM; SQUAMÆ OXIDI, *Dub.* The Scales of the Oxide of Iron.

*Syn.* Bluettes de Fer (*F.*), Eissenoxyd (*G.*), Scaglio di ferro (*I.*), Escamas de hierro (*S.*).

These scales are detached by the hammer of the Smith from the surface of iron heated to redness in the forge, and hammered on the anvil.

*Qualities.* — They are inodorous and insipid, attracted by the magnet, sp. gr. 3·5, brittle, and reducible by trituration to a powder which is of a dull, greyish-black colour. This oxide is soluble in acids, without producing hydrogen gas; and appears to be the metal in a degree of oxidizement according to M. Berthier higher than the protoxide.<sup>1</sup> According to Lavoisier and Proust, they consist of 27 parts of oxygen, and 73 of iron: but M. Berthier makes the proportions to be iron 74·5, oxygen 25·5.

*Medical properties and uses.* — These scales are used in the same cases and in the same manner as the filings, and are preferable; for as they do not produce hydrogen gas when dissolving in the stomach, their use is unaccompanied by the distention and flatulence which the filings often occasion.

FERULA. *Spec. Plant. Willd.* i. 1411.

*Cl.* 5. *Ord.* 2. Pentandria Digynia. *Nat. Ord.* Umbellatæ.

*G.* 539. Fruit oval, compressed, plane, three streaks on each side.

*Species* 11. *F. Assafoetida*.<sup>2</sup> *Assafoetida.* *Kœmpfer, Amœnitates Exoticæ*, 535. *t.* 536.

*Officinal.* ASSAFÆTIDÆ GUMMI-RESINA, *Lond. Edin.* ASSAFÆTIDA, *Dub.* *Assafoetida.*

*Syn.* *Assafoetida* (*F.*), Stinkender Asand (*G.*), *Assafetida* (*I.*), *Asafetida* (*S.*), *Perungyum* (*Tam.*), *Angoo* (*Malay*), *Ungoozeh* (*Pers.*), *Duivelsdreck* (*Dutch*), *Dyrelsdraek* (*Dan.*), *Dyfelstrack* (*Swed.*), *Hilteet* (*Arab.*), *Hing* (*H.*), *Hinga* (*San.*), *Jo-eul* (*Chinese*).

This species of ferula is a native of the south of Persia, chiefly growing on the mountains in the provinces of Chora-saan and Laar, where it is named *hingisch*. The root is perennial, tapering, and ponderous; when fully grown, the size of a man's leg, covered with a blackish coloured bark, and near the top beset with strong rigid fibres. The internal substance is fleshy, white, and abounds with a thick, very fetid, milky juice. This stem is round, smooth and striated; rising erect to the height of nine feet, and is about seven inches in circumference at the base; surrounded with six or seven ra-

<sup>1</sup> *Annals de Chimie*, xxvii. 19.

<sup>2</sup> Σιλφιον μηδικον *Dioscoridis*. The plant described and figured by Dr. Hope of Edinburgh, in the 75th volume of the *Philosophical Transactions*, as that which yields the officinal assafoetida, is the *Ferula Persica* of Willdenow, and a native of the north of Persia.



dical leaves, nearly two feet long, bipinnate, with alternate pinnules, smooth, sinuated, lobed, or lanceolate; of a deep green colour, and fetid odour. The flowers are in plano-convex, terminal, compound umbels: the seeds oval, flat, foliaceous, of a reddish-brown colour, rough, with three longitudinal lines; and have a porraceous odour, and a sharp bitter taste.

When the root is four years old, it is fit to yield the assafoetida, which is procured in the following manner. At the season when the stem and leaves begin to decay, they are twisted off from the root, which is then exposed by digging away the earth that surrounds it. It is left in this state screened from the sun for forty days; then the top is cut off transversely; and after forty-eight hours the juice which has exuded is scraped off, and another transverse section is made. This operation is repeated three successive times, and then the root is allowed to remain untouched for eight or ten days, before another section is made. The root perishes after it is exhausted of the juice. The juice collected from a number of roots is put together and dried in the sun.

Assafoetida is brought into this country packed in cases, mats, and casks; that in the cases proving generally the best. It is in irregular masses, adhering to each other, externally of a brownish yellow colour, and containing many little shining tears of a whitish, reddish, or violet hue. The best is clear, and of a pale reddish colour, contains many of the white tears, and has the odour very strong.

*Qualities.*— Assafoetida has a strong, very disagreeable, alliaceous, fetid odour, and a bitter subacid taste; but these qualities, particularly the odour, on which much of the efficacy of the drug depends, are much injured by keeping.<sup>1</sup> It becomes brittle by exposure to the air; but is not easily reduced to powder, unless it be triturated with carbonate of ammonia. Its specific gravity is 1.327. It yields all its virtues to ether and to alcohol. It is diffused by trituration in water, forming a milky opaque mixture. The ethereal tincture, when evaporated on water, leaves a thick pellicle of brown, fetid resin, and gives the water a milky appearance. In distillation, either with water or alcohol, assafoetida yields an essential oil, on which its odour depends. The proportions of its components are gum 60, resin 30, and essential oil 10 parts, in 100; but Brugnatelli affirms that the part which has been regarded as gum is extractive.<sup>2</sup>

*Medical properties and uses.*— This gum resin is stimulant,

<sup>1</sup> Kœmpfer says: "Affirmare ausim drachmam unam, recens effusam, majorem spargere foetorem quam centum libras vetustioris, quem siccum venundant aromatarii nostrates." *Amœn. Exoticæ*, p. 535.

<sup>2</sup> *Compendio di Mat. Med.* p. 41. Pavia, 1817.



antispasmodic, expectorant, emmenagogue, and anthelmintic. It is more efficacious than any of the other fetid gums; producing its effects in a shorter space of time; and is therefore beneficially given as an antispasmodic in cases of hysteria, hyperchondriasis, dyspepsia, flatulent colic, tympanitis, and in nervous diseases: its expectorant powers have been found useful in asthma, and hooping-cough; and it ranks high as a remedy in chlorotic affections. We are informed that in India it is a successful native specific against the Guinea<sup>1</sup> worm. Its use is contra-indicated when the inflammatory diathesis is present; and owing to its stimulant quality, it is often combined with antimonials and nitre. It is used locally, in the form of enema in worm cases, flatulent colic, and in the convulsions attending dentition: and, sometimes, it is applied as a plaster for discussing tumours.

The dose is from grs. v. to ℥j, formed into pills, or diffused in water. Six drachms of assafoetida beaten with ʒss. of camphor forms a proper mass for a plaster.

Officinal preparations. *Mistura Assafoetidæ*, L. D. *Tinct. Assafoetidæ*, L. E. D. *Spiritus Ammoniacæ foetidus*, L. D. *Tinct. Castorei Comp.* E. *Pilulæ Assafoetidæ compositæ*, E. *Pilulæ Aloes et Assafoetidæ*, E. *Pil. Galbani comp.* L. *Enema foetida*, D.

FICUS. *Spec. Plant. Willd.* iv. 1131.

Cl. 23. Ord. 2. Polygamia Dioecia. *Nat. ord.* Scabridæ, *Linn.* Urticæ, *Juss.*

1931. Common receptacle turbinate, fleshy, converging, concealing the florets, either in the same or a distant individual.

Male. Calyx three-parted. Corolla 0. Stamens three.

Female. Calyx five-parted. Corolla 0. Pistil one. Seeds covered by a permanent, closed, somewhat fleshy calyx.

\* Leaves lobed.

Species 1. *F. Carica*.<sup>2</sup> The Fig-tree. *Med. Bot.* 2d edit. 714. t. 244.

Officinal. CARICÆ FRUCTUS, *Lond. Dub.* FICI CARICÆ FRUCTUS, *Edin.* The preserved fruit of the Fig.

Syn. Figue (*F.*), Feigen (*G.*), Vyge (*Dutch*), Figen (*Dan.*), Fiken (*Swed.*), Fico (*I.*), Higo (*S.*), Simie Attil pullumi (*Tam.*), Unjeer (*Pers.*), El Kermos (*A.*).

The fig-tree is a native of Asia, but it was introduced into Europe in the early ages. It flourishes in France, Spain, and Italy, and even sometimes ripens its fruit in England.<sup>3</sup> It flowers in June and July. It seldom rises above twelve feet in height, but sends off many spreading branches; and the trunk, which exudes a milky odorous fluid when wounded, is covered with an ash-coloured bark, and seldom exceeds seven inches in diameter. The leaves, which are annual in

<sup>1</sup> *Edin. Med. Journ.* ii. 304.    <sup>2</sup> *Συκη* of the Greeks, who termed the fruit *Συκα*.

<sup>3</sup> The first fig-trees introduced into England are still in the Archbishop's garden at Lambeth. They are supposed to have been planted by Cardinal Pole, and now bear excellent fruit.



Europe, but perennial between the tropics, are large, nearly a span in length, scabrous, and irregularly divided into three or five lobes; of a deep green colour on the upper surface, with a pale green longitudinal vein to each lobe; but on the under surface the whole is pale green, with the veins raised, reticulated, and downy: they are supported on round petioles. The fruit in its early stage serves as a common receptacle, and contains upon its inner surface both the male and female florets. It is turbinate, umbilicate at the top, in colour varied green and red, fleshy, soft, and hollow within.<sup>1</sup>

The fig-tree was very much cultivated by the ancients, who brought the fruit to perfection by a process which they termed caprification. They had observed that those figs which were perforated by an insect, the *Cynips Psenes* of Linnæus, always ripened better; and therefore, they tied a wild fig, on which this insect breeds, near the young cultivated figs, so as to cause the insects, when they issued from the wild fig, also to perforate them. The good effects arose from the crawling of the larvæ within the figs, scattering the pollen and thus forwarding the impregnation of the female florets. But the gardeners of Aleppo, ignorant of the cause of the benefit derived from the *Cynips*, imitate the process by pricking the figs with a needle dipped in oil, in order to procure early figs. The fruit when ripe is dried in ovens, to preserve it, and to destroy any of the larvæ of the *Cynips* that may remain; and then packed very closely in the small chests in which they are imported into this country.<sup>2</sup>

*Qualities.* — Dried figs have a sweet, peculiar taste. They are generally compressed; the cuticle is of a brownish colour, and crusted over with crystals of sugar; and within are numerous small, yellow, lenticular seeds in a sweet viscid pulp. They consist almost entirely of mucilage and sugar.

*Medical properties and uses.* — The dietetical use of figs is well known.<sup>3</sup> When eaten freely they are apt to occasion flatulent cholic and diarrhœa. They are used medicinally in demulcent decoctions, in pulmonary and other inflammatory complaints; and two ounces of them boiled in six fluid ounces of water, and strained, form a useful gargle in cynanche tonsillaris, when suppuration takes place. The figs themselves, roasted or boiled and split, form excellent cataplasms, when applied very hot to gum-boils, buboes, and other phlegmons.<sup>4</sup>

<sup>1</sup> *Gærtner de Fructibus*, ii. 66. t. 91.

<sup>2</sup> The most luscious dried figs in the world are those of Kalamata, in the Morea. They are dried upon rushes, with which the figs are pierced.

<sup>3</sup> Figs were the chief part of the food of the ancient *Athletæ*.

<sup>4</sup> The most ancient cataplasm on record was made of figs. It was used for the relief of Hezekiah, who lived 260 years before Hippocrates. "And Isaiah



FRAXINUS. *Spec. Plant. Willd.* iv. 1102.

Cl. 23. Ord. 2. Polygamia Dioecia. Nat. ord. Sepiariæ, Linn. Jasmineæ, Juss.

G. 1903. Hermaph. Calyx 0, or four-parted. Corolla 0, or four-parted. Stamens two. Pistil one. Capsule one-seeded, lanceolate. Female. Pistil one, lanceolate.

Sp. 15. F. *Ornus*.<sup>1</sup> Flowering ash. *Med. Bot.* 2d edit. p. 589. *Sibthorp Flora Græca*, t. 4.

Officinal. MANNA, Lond. Dub. Edin. Manna.

Syn. Manne (F.), Mannaesche (G.), Manna (Russian), Manna (I.), Mana (S.), Turenjeebeen (Arab.), Shirkhisht (H.).

This tree is a native of the south of Europe, growing abundantly in Calabria, Apulia, Sicily, and on the loftier mountains of Greece; and is cultivated in England as an ornamental tree, flowering in May and June. It seldom exceeds twenty feet in height, is very branching, and has a smooth grey bark. The leaves are deciduous, petiolate, opposite, and pinnate, composed of two or three pair of leaflets, with a terminal one: the leaflets are opposite, about one inch and a half long, and three-fourths broad, pointed at each end, obtusely and unequally serrated, smooth, and of a deep bright green colour. The footstalks vary in length, and are channelled; the gems are villous. The flowers, which grow in close panicles at the extremities of the young shoots, are supradecompound, opposite, scarcely the length of the leaves, smooth, and without bractes. The segments of the calyx are ovate, pointed, and nearly equal; the petals oblong and linear, obtuse, entire, attenuated at the base, spreading, twice the length of the calyx, and of a white colour. The filaments are two, spreading, white, smooth, and bearing yellow incumbent anthers. The germen is small, oval, and smooth, with a short, straight style, crowned with a notched stigma. The capsules droop, are lanceolate, notched, compressed, and bilocular at the base; with one cell generally abortive, while the other contains a cylindrical ferruginous seed.

Two other species of ash, the *rotundifolia* and *excelsior*, also produce manna.<sup>2</sup> It exudes in warm dry weather spontaneously from the stem and branches; and concretes into whitish tears which are scraped off and sold under the name of manna in the tear. The greater part of the manna, however, is obtained by longitudinal incisions about three inches

said, Take a lump of figs. And they took and laid it on the boil, and he recovered."

2 Kings, chap. xx. 7.

<sup>1</sup> *Μελια* Dioscoridis.

<sup>2</sup> It is also produced from the Tamarisk, and used as food by the Bedouin Arabs in the region of Mount Sinai. Burckhardt says, "Whenever the rains have been plentiful during winter, it drops abundantly. They gather it before sunrise, because if left in the sun it melts: they use it as we do sugar, principally in their dishes composed of flour. *Travels in Nubia*, 4to. 1819. Introd. p. lxxviii.



in length, made on one side of the tree only in the same season, and continued from the base of the trunk upwards as far as the branches, at the distance of an inch from each other. The manna flows at first in the form of a thick juice, which gradually concretes. It is collected in baskets, and known under the name of *manna grassa*, fat manna: and is in irregular masses of a reddish or brownish colour, often full of impurities. By making the juice to concrete on straws, and chips fastened near the incisions, a finer kind of manna is procured, which is called canulated or flaky manna, *manna in cannoli*. In Sicily this variety is received on the leaves of the prickly pear, *Cactus opuntia*. The collecting begins about the middle of June, and terminates in September.<sup>1</sup>

Manna is brought to Great Britain packed in chests. The different sorts are in separate packages, and are known by the names of Flake manna, Sicilian manna<sup>2</sup>, and Calabrian manna. The best is “in oblong pieces or flakes, moderately dry, friable, light, of a whitish or pale yellow colour, and in some degree transparent: the inferior kinds are moist, unctuous, and brown.”<sup>3</sup> The best flake manna bears the impression of the branch on which it had concreted on its inner surface. Manna is said to be occasionally counterfeited by a composition of honey or sugar, mixed with scammony or some other purgative<sup>4</sup>: but such frauds are now seldom attempted; and bad or counterfeit manna may be easily discovered by its colour, weight, transparency, and taste; which are different from those of real manna.

*Qualities.* — Manna has a slight peculiar odour, and a sweet taste, with some degree of bitterness, not very pleasant, and leaving a nauseous impression on the tongue. The finer pieces, which are often hollow, when broken and examined by the microscope, exhibit bundles of long beautiful spicular crystals: but the general texture of the pieces is granular. It is entirely soluble in water and alcohol; and the latter, when the solution has been assisted by heat, deposits on cooling five-eighths of the manna beautifully crystallized; and which may be regarded as pure manna, and an uncrystallizable mucilaginous extractive matter remains, on which probably the purgative quality of the drug depends. Fourcroy and Vauquelin suppose that the common manna of the shops contains four different ingredients. 1. Pure manna, constituting three-fourths of the whole: 2. A little common sugar: 3. A yellow

<sup>1</sup> Arcturius is the first Greek who notices manna. *Friend's Hist. of Med.* i. 271.

<sup>2</sup> The greatest produce of Sicilian manna is in the neighbourhood of Castillamare, Carini, Cefalu, and Caronia, where it yields an annual revenue of 40,000*l.* sterling. Smyth's *Sicily and its Islands*, 4to. 1824 p. 14.

<sup>3</sup> *Lewis.*

<sup>4</sup> *Alston's Mat. Med.* ii. 472.



nauseous-smelling substance, to which its purgative qualities seem owing: and, 4. Mucilage.

*Medical properties and uses.* — Manna is a very gentle laxative. It was extravagantly commended by some of the older physicians; but is now more justly regarded as a laxative fit for children only, and persons of very weak habits. When given in a dose sufficient for an adult, it is apt to occasion flatulence and griping; and therefore it is seldom used, except as an adjunct to senna, rhubarb, or solutions of neutral salts, with the view of covering their tastes.

The dose for children is from ʒj. to ʒiv.; and for adults from ʒj. to ʒij.

Officinal preparations. *Confectio Sennæ*, L. E. D. *Enema Catharticum*, D. *Enema fœtidum*, D. *Syrupus Sennæ*, D.

FUCUS. *Genera Plant.* Schreber.

Cl. 24. Ord. 3. Cryptogamia Algæ. Nat. ord. Algæ.

G. 1671. Male. Vesicles smooth, hollow, with villose hairs within, interwoven.

—— Female. Vesicles smooth, filled with jelly, sprinkled with immersed grains, prominent at the tip. Seeds solitary.

Species 2. *F. vesiculosus*.<sup>1</sup> Bladder-wrack. *Turner's Fuci*, ii. 44. *Eng. Bot.* 1066.

Officinal. FUCUS, *Lond.* QUERCUS MARINA; HERBA FRUCTIBUS PRÆSENTIBUS, *Dub.* Bladder-wrack, bearing the fruit.

This marine plant is a perennial, a native of the British shores; bearing the fructification in the spring. The root is an expanded, black, woody, callous disk. The frond is smooth and glossy, from one to four feet in length, and from half an inch to one inch and a half in breadth, of a dark olive green colour, becoming paler near the apices; every where linear, and dichotomous; and through its whole length furnished with a midrib of a blackish colour, as thick as a goose-quill at the base, but gradually growing pale and thin. In the membranous part are found immersed spherical vesicles, varying in size from a pea to a hazel-nut, always close to the midrib, their substance thin, and their cavity full of air. The fructification consists of compressed, turgid, solitary or twin receptacles at the ends of the branches; roundish, perforated, and filled with a tasteless pellucid mucus.

When the plant is dried in the usual manner it becomes brittle, and of a dull black colour; and sometimes is covered with a saline efflorescence: but, when the fresh plant is immersed in boiling water before drying it, the colour remains as vivid as in the fresh plant. It is much used in the manufacture of kelp.

*Qualities.* — The odour is slight, but peculiar; and the

<sup>1</sup> The ancients were unacquainted with this plant, which was first described by Clusius under the name of *Quercus marina*. *Hist.* i. 21.



taste nauseous, and similar to that of soda. When burnt, it yields charcoal and soda.

*Medical properties and uses.* — The burnt plant is considered as deobstruent; and has been exhibited in scrophulous affections and bronchocele: and Dr. Russel found the mucus of the vesicles an excellent resolvent when externally applied to scrophulous swellings.

Official preparation. — *Pulvis Quercus marinæ*, D.

GALBANI GUMMI RESINA. Vide *Bubon*.

GALLÆ. Vide *Quercus*.

GENTIANA.<sup>1</sup> *Spec. Plant. Willd.* i. 1331.

Cl. 5. Ord. 2. Pentandria Digynia. Nat. ord. Rotaceæ, Linn. Gentianæ, Juss.

G. 512. Corolla one-petalled. Capsule two-valved, one-celled; with two longitudinal receptacles.

\* Corollas five or nine-cleft, somewhat bell-shaped.

Species 1. *G. lutea*. Yellow-Gentian. *Med. Bot.* 2d edit. 273. t. 95.

Official. GENTIANÆ RADIX, Lond. GENTIANÆ LUTEÆ RADIX, Edin. GENTIANA RADIX, Dub. Gentian root.

Syn. Gentiane jaune (F.), Rother Enzian (G.), Gentiaan (Dutch), Entzianrod (Dan.), Baggsöta (Swed.), Genziana (I.), Jenciana (S.), Genciana Amarella (Port.).

This species of gentian is a perennial plant, found growing on the alps of Switzerland and Austria, the Apennines, the Pyrenees, and in North America. The root is thick, long, and cylindrical. The lower leaves are petiolate, large, spear-shaped, stiff, and having five large veins on the back, plaited, and of a yellowish green colour; those of the stem are concave, smooth, and egg-shaped, sessile, and almost embracing the stem, which rises three or four feet in height. The flowers are in whorls at the upper joints, large, yellow, peduncled and beautiful: the calyx, which is a membranous deciduous spathe, bursts on the side when the flower opens: the corolla is rotated, divided into five or eight narrow spreading segments, elliptical, and speckled with many thick dots. The filaments are shorter than the corolla, and furnished with long erect anthers: the germen is conical, crowned with two sessile reflected stigmas; and becomes a conical capsule, which contains numerous small seeds.

Gentian roots are brought to this country from Germany. They are in pieces of various lengths and thickness, twisted, wrinkled on the outside and covered with a brownish gray cuticle.

*Qualities.* — They have no particular odour, and the taste is intensely bitter without being nauseous. When cut transversely the pieces exhibit a yellow maculated heart, with thick

<sup>1</sup> Γεντιανη Dioscoridis. Said to have been named after Gentius, King of Illyria, who first discovered its medicinal properties 167 years before the birth of our Saviour.



bark verging to brown. The sensible qualities of Gentian root are extracted by ether, alcohol, and water. The two former extract a resin and a bitter extractive matter; and the latter some part of these and a considerable quantity of mucilage also, which occasions the infusion often to become ropy. Diluted alcohol is its proper menstruum. In the bitter extractive the virtues of the drug seem to reside.

*Medical properties and uses.*—Gentian root is tonic, stomachic, and in large doses aperient. Its use as a stomachic bitter is of a very ancient date; and it is still, perhaps, the most generally employed of this class of medicines. It has been found beneficial in dyspepsia, gout, hysteria, and jaundice; chlorosis, dropsy, and diarrhoea; and in all cases of general debility in which tonics are indicated. It is sometimes joined with the Cinchona in intermittents; and, according as the circumstances of the cases for which it is prescribed direct, it may be combined with orange-peel, chalybeates, aromatics, squill, mineral acids, and neutral salts. On account of its antiseptic effects on dead animal matter, its infusion has been used as an application to putrid ulcers. The forms in which it is generally given are infusion and tincture.

The dose in substance is from grs. x. to ℥ij.

Officinal preparations. *Extractum Gentianæ*, L. E. D. *Infusum Gentianæ compositum*, L. E. D. *Tinctura Gentianæ composita*, L. E. D. *Vinum Gentianæ compositum*, E.<sup>1</sup>

GEOFFROYA. *Spec. Plant. Willd.* iii. 1129.

Cl. 17. Ord. 4. Diadelphia Decandria. Nat. ord. Papilionaceæ, Linn. Leguminosæ, Juss.

G. 1362. Calyx five-parted. Drupe ovate. Nucleus compressed. Species 3. *G. inermis*. The Cabbage-tree. *Med. Bot.* 2d edit. 416. t. 151. *Phil. Trans.* lxxvii. 512. t. 10.

Officinal. GEOFFROYÆ INERMIS CORTEX, *Edin.* GEOFFRŒA; CORTEX, *Dub.* Cabbage-tree bark.

✓ *Syn.* Umari de la Jamaïque (*F.*), Geoffrunrinde (*G.*), Maskbark (*Swed.*), Geoffrea (*I.*).

This tree is a native of Jamaica, growing in the low savannahs. It is a lofty tree, branching towards the top; and covered with a smooth grey bark. The leaves are pinnate, composed of four or five pairs of lancet-shaped, pointed, smooth leaflets, in pairs on short footstalks, with a terminal one. The flowers are in clusters on large branched spikes. The calyx is bell-shaped, with five short obtuse teeth; the corolla papilionaceous, of a pale rose colour, consisting of a

---

<sup>1</sup> Dr. Paris (*Pharmacologia*) says, that the quack medicine known under the name of *Brodm's Nervous Cordial*, consists of the tinctures of *Gentian*, *Columba*, *Cardamom*, and *Bark*, with the compound spirit of *Lavender*, and *Wine of Iron*: and *Stoughton's Elixir*, of tincture of *Gentian*, with the addition of *Serpentaria*, *Orange peel*, *Cardamoms*, and some other aromatics.



roundish, concave *vexillum* notched at the apex; two oblong, obtuse, concave, somewhat shorter *alæ*; and an obtuse divided *carina*. The filaments, nine of which are united at the base, support simple, roundish anthers; the germen is oval, with a curved, tapering style, and hooked stigma. The fruit resembles a small plum, is pulpy, marked on each side with a longitudinal furrow; and contains a hard seed.

*Qualities*. — Cabbage-tree bark has a disagreeable, sweetish, mucilaginous taste. The pieces, as they are brought to this country, are externally grey; internally black, furrowed, and pulverulent, affording a powder resembling that of jalap. Its soluble components seem to be chiefly mucus, resin, extractive, saccharine matter, and a narcotic principle.

*Medical properties and uses*. — This bark is a powerful anthelmintic. Its properties, as such, were first noticed by Mr. Peter Duguid<sup>1</sup>, and have since been fully confirmed: but we are principally indebted to Dr. Wright for an accurate knowledge both of the plant and its virtues.<sup>2</sup> It is particularly useful in expelling lumbrici, and may be given in forms of powder, decoction, extract, and syrup; but the decoction is the form most commonly employed. (See *Preparations and Compositions*.) It operates as a cathartic, but has a narcotic effect also; and requires, therefore, to be given at first in small doses, “which may be gradually increased till nausea is excited, when the dose for that patient is ascertained.”<sup>3</sup> In over-doses, it is apt to occasion sickness, vomiting, fever, and delirium; and the same effects are produced if cold water be drunk during its operation. When such symptoms occur from either cause, they are generally removed by copious draughts of warm water, a dose of castor-oil, and plentiful dilution with lemonade or infusion of tamarinds. Owing to these deleterious effects of the remedy, it has not been generally used in this country.

The dose of the powder is from ʒj. to ʒss.;—and that of the extract, which is made by evaporating the decoction, grs. iij. The syrup, which is the decoction with a double portion of sugar added to it, may be taken in doses of from two to four spoonful.

Official preparations. *Decoctum Geoffroyæ inermis*, E.

GEUM. *Spec. Plant. Willd.* ii. 1113.

Cl. 12. Ord. 8. Icosandria Polygynia. *Nat. ord.* Senticosæ, *Linn.* Rosaceæ, *Juss.*

G. 1002. *Calyx* ten-cleft. *Petals* five. *Seeds* with a bent awn. *Species* 3. *G. urbanum*.<sup>4</sup> Common Avens, or Herb Bennet. *Med. Bot.* 2d edit. 502. t. 181. *Smith Flora Brit.* 554.

<sup>1</sup> *Physical and Literary Essays*, ii. 264.

<sup>2</sup> *Phil. Trans.* l. c.

<sup>3</sup> *Wright*, l. c.

<sup>4</sup> *Caryophyllata*, *Alston's Mat. Med.* vol. i. p. 402.



*Officinal.* GEUM URBANUM; RADIX, *Dub.* The root of Avens.

*Syn.* Bonoite officinale (*F.*), Nelkenwurzel (*G.*), Gemeen Nagelwortel (*Dutch*), Nellikerod (*Dan.*), Näglikerot (*Swed.*), Zarzyezka (*Pol.*), Cariofilata (*I. S.*), Guriofillata (*Port.*).

Avens is an indigenous perennial plant, common in woods and moist shady places, flowering from May to August. The root is fibrous, or rather composed of slender radicles, attached to a central root: the stem is about two feet high, nearly upright, roundish, hairy, leafy, branched at the top, and of a brownish hue on one side. The leaves are of a deep green colour, on channelled footstalks; those nearest the root pinnated and lyrate, with two pairs of unequal leaflets, and a larger wedge-shaped terminal one; the upper leaves are simple, trifid, and pointed; and all cleft, serrated, and hairy. The stem-leaves are accompanied with a pair of large, rounded, lobed, toothed stipules, resembling the smaller leaflets. The flowers are erect, terminal, and solitary, on peduncles. The calyx is externally brownish, internally pale green, of one piece, divided into five large pointed segments, and five intermediate ones much smaller, and lanceolate. The petals are bright yellow, small, with claws, and spreading: the filaments awl-shaped, nearly as long as the larger segments of the calyx, yellowish, inserted into the calyx; at first bent inwards, afterwards erect, and supporting yellow anthers. The germens are hairy, collected into a pear-shape, the styles hooked and jointed in the middle, with simple stigmas. The seeds are hairy, with purple naked awns, hooked, and kneed near the apex.

The root should be dug up in March; for the odour, which is then strong, is almost lost when the flowers appear. It should be dried in the air, with a moderate heat. The large roots are to be preferred. The cultivated are supposed to be superior to the wild.

*Qualities.* — Avens root has a fragrant odour, resembling that of cloves, and an aromatic, bitterish, austere taste. The pieces are of a dark brownish red colour on the outside, and internally white. Its sensible qualities are extracted both by water and by alcohol; and in distillation with water, it yields a small portion of heavy volatile oil. The watery infusion is of a reddish colour, froths when agitated, like a gummy solution, reddens litmus paper, and strikes a black colour with sulphate of iron.

*Medical properties and uses.* — The root of avens is astringent, tonic, and antiseptic. It is mentioned by Ray as a febrifuge, but has been very little used in Britain, and was altogether unnoticed, until Buckhave, in a work entitled *Observationes circa Radicem Gei urbani*, adduced numerous instances of its efficacy in intermittents. The Continental practitioners recom-



mend it in dysentery, chronic diarrhoea, and flatulent colic; and as a general tonic in all cases in which Cinchona is employed. The Dublin College has therefore judiciously inserted it in the list of materia medica; and, being an indigenous remedy, it deserves attention.<sup>1</sup> It may be exhibited in the form of powder; or in an electuary compounded with honey and rhubarb; or a decoction may be made with one ounce of the root and one pint of water; or a tincture with similar proportions of the root and alcohol. The dose of the powder is ʒss. or ʒj. four times a day; of the decoction fʒj. every hour; and of the tincture fʒss., properly diluted, three or four times a-day.

GLYCYRRHIZA.<sup>2</sup> *Spec. Plant. Willd.* iii. 1143.

*Cl.* 17. *Ord.* 4. Diadelphia Decandria. *Nat. ord.* Papilionaceæ, *Linn.* Leguminosæ, *Juss.*

*G.* 1366. *Calyx* bilabiate; upper lip three-cleft, lower undivided. *Legume* ovate, compressed.

*Spec.* 4. *G. glabra*. Common Liquorice. *Med. Bot.* 2d edit. 429. t. 152.

*Officinal.* GLYCYRRHIZÆ RADIX, *Lond. Dub.* GLYCYRRHIZÆ GLABRÆ RADIX: — EXTRACTUM, *Edin.* Liquorice root, and the extract.

*Syn.* Reglisse (*F.*), Sussholzwurzel (*G.*), Zoethout (*Dutch*), Lakris (*Dan.*), Lakrits (*Swed.*), Lakrycia (*Pol.*), Addimodrum (*Tam.*), Bikh-mehek (*Pers.*), Urat manis (*Malay*), Oyot manis (*Jav.*), Olinde (*Cyng.*), Legorizia (*I.*), Regaliza (*S.*), Alcacuz (*Port.*), Ussulussoos (*Arab.*), Iét'himad'h (*H.*), Yastimadhuca (*San.*).

The liquorice plant is a native of the south of Europe and Syria; but the greater part of what is used in Britain is the produce of its own soil by cultivation. The London market is supplied chiefly from Mitcham in Surrey.<sup>3</sup> It flowers in August. The root is perennial, running, when in its proper soil, a light sandy one, very deep; it is round; the thickness from that of a goose-quill to that of the thumb; long, thin, flexible; furnished with sparse fibres; covered with a brownish cuticle; internally fibrous, of a pale-yellow colour, and juicy. The stem rises four or five feet in height, is herbaceous and striated, with few branches. The leaves are alternate and pinnated, consisting of four or five pairs of ovate, retuse, petiolated leaflets, with a terminal one; of a pale green colour, and clammy on the under-side. The flowers are papilionaceous in long axillary sparse spikes, of a blue or purplish colour. The calyx is persistent, tubular, and divided above; the corolla consists of an ovate, lanceolate, obtuse, erect, concave *vexillum*; two ob-

<sup>1</sup> Augsburg beer, which is much prized on the Continent, is supposed to owe part of its excellence to the custom which prevails of putting a small bag of the root of *Avens* into each cask. *Nicholson's Journ.* vol. xxx. p. 80.

<sup>2</sup> Γλυκυρρίζα Dioscoridis. The name is derived from γλυκὺς, sweet, and ῥίζα, a root.

<sup>3</sup> Very little is now grown at Godalming, where it was formerly cultivated to some extent. Vide *Stevenson's Survey of Surrey*, p. 380. It was first cultivated in England, in 1558. *Stow*.



long, obtuse *alæ*, and a shorter *carina*. The filaments are ten, nine of them united at the base, bearing simple, roundish anthers: the germen is short, with a tapering style and blunt stigma. The legumes are ovate, flattened, smooth, acute, one-celled, containing two or three small kidney-shaped seeds.

When liquorice root is three years old, it is dug up for use in November. “The whole roots are then washed, the fibres cut off, and the smaller roots separated from the larger ones. The former termed the offal, are dried and ground to powder; the latter are packed up and sold to the druggists.”<sup>1</sup>

*Qualities*.—This root is inodorous, and the taste sweet and mucilaginous, leaving, when it is chewed without being peeled, a slight degree of bitterness in the mouth. The powder, if good, is of a brownish yellow colour, and has a rich sweet taste, more agreeable than that of the fresh root; but it is said to be often sophisticated with flour, and other substances not quite so wholesome, in which case it has a fine pale yellow colour.

The medical properties of the root depend on a saccharine matter, which approaches in its nature to *sarcocoll* and mucus: water, by coction, extracts both of these principles, but alcohol only the saccharine matter. For the properties of the extract, which is imported from Spain, see Part III. (*Preparations and Compounds*.)

*Medical properties and uses*.—Liquorice root is a pleasant demulcent; but on account of its bulk it is rarely used in substance.<sup>2</sup> The decoction of it, either alone or in combination with other mucilaginous vegetables, is often given in catarrh, and in hectic and phthisical cases. It is also administered in some cases of dyspepsia, where there is a deficiency of the natural mucus of the stomach, which is injured by the acrimony of ill-digested food, and a morbid state of its secreted fluids. The dose of the powder is from grs. x. to ʒj., that of the decoction a cupful frequently repeated.

Officinal preparations. *Extractum Glycyrrhizæ*, L. E. D. *Decoction Sarsaparillæ comp.* L. D. *Infusum Lini*, L. *Confectio Sennæ*, L. E.

GRANATI CORTEX. Vide *Punica*.

GRATIOLA.<sup>3</sup> *Spec. Plant. Willd.* i. 102.

Cl. 2. Ord. 1. Diandria Monogynia. Nat. ord. Personatæ, Linn. Juss.

G. 49. *Corolla* irregular, reversed. *Stamens* two, sterile. *Capsule* two-celled. *Calyx* seven-leaved; the two exterior leaves spreading.

<sup>1</sup> The price of the best roots is about 3*l.* per cwt. *Stevenson*, l. c.

<sup>2</sup> The ancients believed that chewing the root allayed thirst; but this opinion was founded on a mistake. *Cullen, Mat. Med.* ii. p. 407.

<sup>3</sup> The name means *Gratia Dei*—from the supposed virtues of the plant.



*Species* 1. *G. officinalis*. Hedge-Hyssop. *Med. Bot.* 2d edit. 359. t. 131. *Flora Danica*, t. 363.

*Officinal*. GRATIOLÆ OFFICINALIS HERBA, *Edin.* GRATIOLA; HERBA, *Dub.* The herbaceous part of Hedge-Hyssop.

*Syn.* Gratiolæ (*F.*), Gradenkraut (*G.*), Genadekruid (*Dutch*), Gudsnaades urt (*Dan.*), Jordgalla (*Swed.*), Kenjtrud (*Pol.*), Graziola (*I.*), Graciola (*S. Port.*).

This plant is a perennial, a native of the south of Europe, growing in marshy or moist pastures, and flowering in June and July. It is cultivated in Britain.<sup>1</sup> The root is creeping, cylindrical, fleshy, and fibrous; and sends up several upright, smooth, round stems, nearly a foot in height, with sessile leaves, in some degree sheathing, and opposite, in pairs. They are lanceolate, smooth, serrated towards the point; of a bright somewhat deep-green colour; nearly two inches long, and half an inch broad; punctured, and longitudinally veined beneath. The flowers are axillary and solitary, on slender reddish peduncles: the calyx is divided into five elliptical segments, with two lanceolate spreading bracteal leaves; the corolla is tubular, divided at the lip into four obtuse segments, the uppermost of which is much broader than the other three, and more reflected; the tube is yellowish, intermixed with reddish streaks; the limb pale purple. Two of the filaments only are furnished with anthers; and the style is tapering, erect, with a divided stigma. The capsule is oval, and contains many small seeds.

The sensible qualities of *Gratiola* are strongest when it is in flower, at which time, therefore, it should be gathered for use.

*Qualities*. — It has scarcely any odour; but the taste is very bitter and nauseous. Boiling water extracts its sensible qualities more perfectly than alcohol. The colour of the infusion approaches to that of Madeira wine: it slightly reddens litmus paper, and strikes an olive colour with a solution of sulphate of iron without occasioning a precipitate. When sulphuric acid is added to the unstrained infusion, it emits the odour of tamarinds; and when the infusion is filtered and slowly evaporated, spicular crystals are formed, which appear to be tartaric acid.

*Medical properties and uses*. — *Gratiola* is cathartic, diuretic, and emetic, producing, in very large doses, all the effects of an irritative poison. It has been much recommended by the German physicians in dropsy; and has also been used in jaundice and worm cases. Hufeland found it extremely efficacious in visceral obstructions, and scrofulous affections<sup>2</sup>: and we are even told that in the Vienna hospital it has cured the most confirmed cases of lues venerea!<sup>3</sup>

<sup>1</sup> It was first cultivated in Britain by Turner, in 1568.

<sup>2</sup> *Hufeland uber die Natur, &c. der Scrofula.*

<sup>3</sup> *Kostrzewski, Dissert. de Gratiola*, p. 64.



It is given either in the form of powder, or of infusion combined with aromatics. The dose of the powder is grs. xv. to ʒss; that of the infusion, made with ʒij. of the dried herb and Oss. of warm water, from fʒiv. to fʒj. three times a day.

**GUAIAACUM.**<sup>1</sup> *Spec. Plant. Willd.* ii. 538.

*Cl.* 10. *Ord.* 1. Decandria Monogynia. *Nat. ord.* Gruinales, *Linn.* Rutaceæ, *Juss.*

*G.* 819. *Calyx* five-parted, unequal. *Petals* five, inserted into the calyx. *Capsule* angular, three or five-celled.

*Species* 2. *G. officinale* Officinal Guaiacum. *Med. Bot.* 2d edit. 557. t. 200.

*Officinal.* GUAIACI RESINA ET LIGNUM, *Lond.* GUAIACI OFFICINALIS LIGNUM-RESINA, *Edin.* GUAIAACUM; LIGNUM, GUMMI-RESINA, *Dub.* The wood and resin of Guaiacum.

*Syn.* Guyac (*F.*), Guajakgummi (*G.*), Pokhout (*Dutch*), Franzos trace (*Dan.*), Franzosenholts (*Swed.*), Guajaco (*I.*), Guayaco (*S.*), Guajaco (*Port.*).

This tree is a native of Jamaica, Hispaniola, and the warmer parts of America.<sup>2</sup> It rises forty feet in height and is four or five in circumference, with many divided knotted branches. The bark of the trunk is of a dark grey colour, variegated with greenish or purplish specks, but that of the branches is ash-coloured, with fissures. The leaves are abruptly pinnate, consisting of two or three pair of smooth, shining, veined, obovate, dark green leaflets, almost sessile. The flowers are peduncled in a kind of umbels, which spring from the divisions of the smaller branches. The calyx consists of five, concave, oblong, blunt, spreading, unequal, deciduous leaves; the petals are five, of a rich blue colour, elliptical, concave, and spreading; the stamens are erect and villous, with yellowish hooked anthers; the germen is oval, with a short style and simple stigma; and the capsule sub-turbinate, on a short pedicel, smooth, and of a pale ferruginous hue, pentagonous, with ribbed angles, and five-celled; but two or three of the cells are often abortive.<sup>3</sup> The seeds are solitary, and angled.

All the parts of this tree possess medicinal qualities; but the wood and the peculiar substance afforded by it are the only parts used: the virtues of the wood depend altogether on the peculiar matter it contains. This is spontaneously exuded from the tree, and is called native gum: it concretes in tears, which are semipellucid and very pure; but the greater part of it is obtained by making incisions into the trunk, or, as it is termed, jagging the tree. This operation is performed in May; and the juice, which flows copiously, is concreted by the sun. It is also obtained by sawing the

<sup>1</sup> The Spanish name *Guayaco*, is derived from the Caribbee. *Humboldt*.

<sup>2</sup> The tree was cultivated in this country by the Duchess of Beaufort, 1699.

<sup>3</sup> *Gært. de Fructibus*, ii. 148. t. 113. fig. 1.



wood into billets, and boring a hole longitudinally through them; so that, when one end of a billet is laid on a fire, the guaiac melting, runs through the hole from the opposite end, and is collected in a calabash. Boiling the chips or raspings in salt and water, also separates the guaiac, which as it rises to the surface may be collected by skimming.

The wood is brought to this country either in large solid pieces which weigh from four to five cwt. each, and are covered with a yellowish alburnum: or it is already rasped. The guaiac, or gum as it is improperly termed, arrives in casks and mats; the former containing from one to four cwt., the latter generally less than one cwt. each.

*Qualities.* — The *wood* of *Guaiacum* is inodorous, but when heated it emits an aromatic odour; and the taste is bitterish, subacid, and biting. It is very hard, heavier than water, externally yellowish, and internally of a blackish brown colour mixed with green streaks. The *resin*, or *Guaiac*, has a fragrant odour, with scarcely any taste, but occasions, when swallowed, a sensation of heat in the throat. It has a resinous aspect; is of a greenish brown colour externally, and internally presents a mixture of greenish, reddish, and brownish tints. It is somewhat translucent, breaks with a vitreous fracture, and is easily reduced to a powder, which is grey at first, but becomes green in a short time when it is exposed to the air and light; a change which appears to depend on the absorption of oxygen.<sup>1</sup> The specific gravity of *Guaiac* is 1.2289. It was generally regarded as a gum resin, till Mr. Brande's experiments showed it to be a substance *sui generis*, differing from both gum and resin.

When *Guaiac* is digested in *water* a little extractive only is dissolved, in the proportion of 9 parts in 100, and the infusion has a greenish brown colour and a sweetish taste. *Alcohol* dissolves readily 95 parts in 100, and the solution is decomposed by the mineral acids, affording precipitates which assume various tints of colour (see *Tinctura Guaiaci*). *Sulphuric ether* dissolves four parts in ten of *Guaiac*, and when the solution is evaporated on water, it leaves a tough, pellucid, pale brown pellicle, which appears to be pure *Guaiac*. It becomes green after some time; and a small portion of extractive remains dissolved in the water. The *alkaline solutions* and their carbonates dissolve it readily: and the solutions are precipitated

---

This effect of light and air was first noticed by Dr. Wollaston, who found that the most refrangible produced this change; and subsequent experiments of Mr. Brande clearly proved it to arise from oxygen. I found that the change takes place in an hour, when the powder is exposed to sunshine. It appears to be again deoxidized if exposed to the least refrangible rays only, according to Dr. Wollaston's experiments.



by the diluted sulphuric, the nitric, and the muriatic acids. *Sulphuric acid* dissolves it with scarcely any effervescence, affording a solution of a rich claret colour, which, when fresh prepared, deposits a lilac-coloured precipitate on the addition of water; and when heated separates some charcoal. *Nitric acid* dissolves it with a strong effervescence and a copious extrication of nitrous fumes; and when the solution is evaporated, it yields a large portion of oxalic acid. The diluted acid converts it into a brown resinous substance. *Muriatic acid* dissolves a small portion only, and affords a solution of a brown colour. I found that, during the solution of Guaiac in these acids, the heat which was evolved raised the thermometer in the following proportions; in the sulphuric 44; in the nitric 120; and in the muriatic 8 degrees. Nothing comes from the distillation of Guaiac in water; but Mr. Brande obtained from 100 parts of it, distilled *per se* in close vessels, the following products: acidulous water 5·5, thick brown oil 24·5, thin empyreumatic oil 30·0, charcoal remaining in the retort 30·5, and 9·5 of gases, which were chiefly carbonic acid and carburetted hydrogen.<sup>1</sup> From these experiments it is clear that Guaiac differs from resin; and we also learn that the mineral acids are incompatible in prescriptions with it.

It is sometimes adulterated with common resin and Manchinal gum. The former is detected by the turpentine emitted when the suspected Guaiac is thrown on hot coals; and the latter by adding to the alcoholic solution a few drops of sweet spirit of nitre, and diluting with water; the guaiac is precipitated, but the adulteration floats in white striæ.

*Medical properties and uses.*—Both the wood and the guaiac are stimulant, diaphoretic, diuretic, and purgative. The wood was introduced into Europe by the Spaniards as a remedy for lues venerea in 1508, and gained much celebrity, from curing Van Hutten; but it had long before been used for the same purpose by the natives of St. Domingo. It obtained so much reputation, that the exhibition of mercury was discontinued for a considerable length of time<sup>2</sup>, and even in the eighteenth century its specific powers over this disease were maintained by Boerhaave: but frequent disappointments and more correct observations have shown that it possesses no powers of eradicating the venereal virus; and that it is useful only after a successful mercurial course for repairing the strength and vigour of the system: “and where a thickened state of the ligaments, or of the periosteum, remains, or where there are

---

<sup>1</sup> *Philosophical Trans.* 1806: and *Phil. Mag.* xxv. 107.

<sup>2</sup> It was then sold for seven gold crowns a pound.



foul indolent ulcers<sup>1</sup> ;” or in suspending the progress of some of the secondary symptoms for a short time, as ulcers of the tonsils, eruptions, and nodes. The decoction of the wood has been found more useful in cutaneous diseases, scrofulous affections of the membranes and ligaments, and in ozæna. The Guaiac itself is an efficacious remedy in chronic rheumatism and arthritic affections<sup>2</sup>, as well as those diseases for which the decoction of the wood is usually given; and in every respect it may be regarded as the active ingredient of the wood. Its sensible effects are a grateful sense of warmth in the stomach, dryness of the mouth, and thirst, with a copious flow of sweat, if the body be kept externally warm, or, if the guaiac be united with opium and antimonials: but when the body is freely exposed, instead of producing diaphoresis, it augments considerably the secretion of urine. It may be exhibited either in substance or in tincture. The dose is from grs. x. to ʒss, in the form of pills or of bolus: or made into an emulsion with water by means of mucilage or yolk of egg. Larger doses purge.

Officinal preparations. Of the wood — *Decoctum Guaiaci comp.*, E. of the Guaiac — *Mistura Guaiaci*, L. *Tinctura Guaiaci*, L. E. D. *Decoctum Sarsaparillæ comp.*, L. D. *Tinctura Guaiaci ammoniata*, L. E. D. *Pulvis Aloes comp.*, L. D.

HÆMATOXYLON.<sup>3</sup> *Spec. Plant. Willd.* ii. 547.

Cl. 10. Ord. 1. Decandria Monogynia. Nat. ord. Lomentaceæ, Linn. Leguminosæ, Juss.

G. 830. *Calyx* five-parted. *Petals* five. *Capsule* lanceolate one-celled, two-valved, with the valves boat-shaped.

*Species* 1. *H. campechianum*. The Logwood-tree. *Med. Bot.* 2d ed. 455. t. 163.

*Officinal.* HÆMATOXYLI LIGNUM, *Lond. Edin. Dub.* Logwood.

*Syn.* Bois de Campeche (*F.*), Kampesch-holz; Blauholz (*G.*), Campechehout (*Dutch*), Campeschetrace (*Dan.*), Campechetrad (*Swed.*), Campeggio (*I.*), Palo de Campeche (*S.*), Pao de Campeche (*Portug.*).

This tree is a native of South America, and attains to great perfection at Campeachy, in the bay of Honduras. It was introduced into Jamaica in 1715, and from its quick growth now abounds in a degree which much incommodes the landholders in the neighbourhood of Savannah la Mar;

<sup>1</sup> *Pearson's Observations on the Effects of various Articles of the Materia Medica in the cure of Lues venerea*, p. 10.

<sup>2</sup> *The Chelsea Pensioner*, a nostrum, by which Lord Amherst was cured of Rheumatism, is composed of Guaiac ʒj., Pulv. Rhei ʒij., Supertart. Potassæ ʒj., Sulph. ʒij., Nucismyrist. j. in pulv. trit. et per opem mellis misce ut fiat Electuarium. Two large spoonful to be taken night and morning. *Jesuit Drops* also consist of Guaiacum, Balsam of Copaiba, and Oil of Sassafras, made into a tincture by spirit. *Pharmacologia*, p. 257.

<sup>3</sup> From αἷμα, blood, and ξυλον, wood. *Millar's Dictionary*. The trivial name *Campechianum*, and the English term *Campechi wood*, originated from *Palo Campechio*, the name imposed by the Spaniards who first discovered the wood.



flowering in March and April.<sup>1</sup> The stem and branches are generally crooked, the former is seldom above twenty inches thick; and the tree scarcely ever rises more than twenty-four feet in height. It is covered with a dark-coloured, rough bark; and has many smaller ramifications, which are close, prickly, and beset with strong spines. The leaves are abruptly pinnate, each composed of four or five pairs of sessile, obcordate, obliquely nerved leaflets. The flowers are in terminal spicular clusters; the calyx consists of brownish-purple coloured, oblong, obtuse segments; the petals are spreading, obtusely lanceolate, and of a reddish-yellow colour. The stamens are downy, tapering, shorter than the corolla, with small oval anthers; the fruit is a double-valved pod, containing five or six small, flat, reniform seeds.

Logwood is brought to this country in logs, which are afterwards chipped. Those pieces which have a deeper colour are to be preferred. It is much employed as a dye-wood.

*Qualities.* — This wood is inodorous, but has a sweet astringent taste. It is hard, compact, heavy, and of a deep-red colour, which it gives out both to water and alcohol. The recent infusions made with distilled water are yellow, but those with common water have a reddish purple colour, which is deepened by the alkalies, and changed to yellow by the acids. They form precipitates with the sulphuric, nitric, muriatic, and acetic acids; solutions of alum, sulphates of iron and of copper, acetate of lead, and tartarized antimony<sup>2</sup>; which are, therefore, incompatible in prescriptions with these infusions and decoctions. The colour of the precipitates varies; those with the acids are reddish-brown, with alum and tartarized antimony violet, with sulphate of iron blueish-black, sulphate of copper purplish-blue, acetate of lead reddish-black, and sulphate of magnesia, purple. According to Chevreul, logwood contains a volatile oil, tannin, two kinds of colouring matter, one of which is soluble both in water and alcohol, the other soluble in alcohol only; acetate of lime and of potass<sup>3</sup>; and a peculiar substance, which is procured in small brilliant crystals of a reddish-white colour, and a slightly astringent, bitter, and acrid taste; and which he named *hematin*.<sup>4</sup>

---

<sup>1</sup> It was cultivated in this country by Mr. Miller in 1739; but is now seldom found in our hot-houses.

<sup>2</sup> When an infusion or decoction of logwood is kept for some time, it becomes capable of producing a precipitate with gelatine; but when recent, no such effect takes place either with glue or isinglass. This was first observed by Dr. Bancroft.

<sup>3</sup> *Annales de Chimie*, lxvii. 254. *Thomson's Chemistry*, v. 206.

<sup>4</sup> To obtain hematin, digest logwood rasped in water of the temperature 125°, filter and evaporate to dryness. Digest the residue for a whole day in alcohol of sp. gr. 0.837, filter and concentrate by evaporation; then add a small portion of water, evaporate a little farther, and leave it to itself. Crystals of hematin are formed in abundance.



*Medical properties and uses.*—Logwood is supposed to be astringent: but this is a questionable opinion, for although it produces an ink with sulphate of iron, it possesses no acerbity, and does not produce a precipitate with gelatine.<sup>1</sup> It is employed in diarrhœa, and in the latter stage of dysentery; but the extract is more usually ordered. It has the advantage of giving tone to the general system, and thus obviates the lax state of the intestines. When taken into the stomach, it imparts a red colour to the urine. The decoction may be taken in doses of two or three fluid ounces, frequently repeated.

Official Preparation. *Extractum Hæmatoxyli*, L.

HELLEBORUS. *Spec. Plant. Willd.* ii. 1335.

Cl. 13. Ord. 6. Polyandria Polygynia. *Nat. ord.* Multisiliquæ, *Linn.* Ranunculaceæ, *Juss.*

G. 1089. *Calyx* none. *Petals* five or more. *Nectaries* bilabiate, tubular. *Capsules* many seeded, nearly erect.

*Species* 3. *H. niger*. Black Hellebore. *Med. Bot.* 2d ed. 473. t. 169.

*Species*. 6. *H. fœtidus*. Fœtid Hellebore, or Bear's foot. *Med. Bot.* 2d ed. 477. t. 170. *Smith Flor. Brit.* 598. *Eng. Bot.* t. 613.

#### 1. HELLEBORUS NIGER.

*Official.* HELLEBORI NIGRI RADIX, *Lond.* HELLEBORI NIGRI RADIX, *Edin.* —; (MELAMPODIUM<sup>2</sup>) RADIX, *Dub.* The root of Black Hellebore.

*Syn.* Hellebore (*F.*), Schwartz Niesswurzel (*G.*), Swart Prustrot (*Swed.*), Ellebro negro (*I.*), Helleboro negro (*S.*), Kadugaroganic (*Tam.*), Kherbeksiya (*Pers.*), Kāli Koothie (*H.*), Kherbec usivud (*Arab.*).

Black hellebore, so named from the dark colour of the root, is a native of Austria, the Apennines, and Italy; flowering from December till March; whence it has been called Christmas rose, and has obtained a place in our gardens.<sup>3</sup> The root is perennial, transverse, rough, knotted, externally black, internally whitish, and sends off many depending fibres. The leaves, which are deep green, spring directly from the root on long, maculated petioles; and are composed generally of five leaflets, pedate, two being supported on one partial petiole on each side, and one terminal: the leaflets are ovato-lanceolate, smooth, shining, and coriaceous, with the upper half of each sparsely serrated. The flower-stalks are scapes, six or eight inches long, erect, round, somewhat tapering, sheathed, variegated with red, and bearing one or two flowers. The floral leaves supply the place of a calyx, are oval, and indented at the apex. The corolla consists of five large, roundish, concave, spreading petals, at first white, with a tint of red, deepened by age, but finally changing to green, after the pollen is shed, and the seed impregnated. The nectaries are greenish yellow, tubular, two-lipped, the upper lip longer and slightly emarginate, the lower

<sup>1</sup> Vide *Bancroft, Phil. of Permanent Colours*, 2d ed. vol. ii. p. 395.

<sup>2</sup> Named from Melampus, a soothsayer, who first gave it as a purgative.

<sup>3</sup> It was cultivated in Britain by Gerarde in 1596.



finely notched. The filaments are numerous, and thread-like, with yellow anthers. The germens, which vary in number from four to eight, become beaked pods, containing many oval, black, shining seeds.

This plant has been supposed to be the *ελλεβορος μελας* of Hippocrates, but there is every reason for agreeing with Willdenow, that his fifth species, *Helleborus orientalis*, is the drug of the ancients. It was found by Bellonius and Tournefort<sup>1</sup> growing in plenty about Mount Olympus, and the island Anticyra, which was formerly celebrated for its production. Sometimes the roots of *Helleborus viridis*, *Adonis vernalis*, *Trollius europæus*, *Actæa spicata*, *Astrantia major*, and *Aconitum neomontanum* are, either ignorantly or fraudulently, substituted for those of black hellebore. These are distinguished chiefly by their colour being paler than the roots of the hellebore.

*Qualities.* — The fibres of the roots, which are the parts used in medicine, are about the thickness of a straw, from four inches to a foot in length, corrugated, of a deep brown black on the outside, and internally white or yellowish. They have an unpleasant odour; and a nauseous, bitterish, acrid taste, benumbing the tongue, and leaving upon it an impression, “as when it hath been a little burnt with eating or supping any thing too hot.”<sup>2</sup> The acrimony is impaired by keeping; and appears to depend on a volatile matter, as water distilled from the root has an acrid taste. Both alcohol and water extract its medicinal properties; and, as the spirituous preparation is the most active, these appear to depend on its resinous part. By coction with water it yields a very considerable portion of gummy matter and some resin. (See the extract.)

*Medical properties and uses.* — Black hellebore root is a drastic cathartic, and on this property probably depends its emmenagogue and hydragogue powers. In smaller doses it is supposed to act as an alterative. It has been much celebrated in mania, melancholia, dropsy, scabies, and worms; but does not appear to possess any particular advantages over the other resinous purgatives, which act with less virulence. As an emmenagogue it is useful in plethoric habits, when preparations of iron are contra-indicated. When black hellebore is taken in too great quantity, it occasions violent vomitings, inflammation of the stomach, vertigo, tremblings, convulsions, cramps, and death. These effects are to be obviated by evacuating the stomach, by drinking copiously of mild mucilaginous fluids, and then employing the most powerful antiphlogistic measures. (See *Extract*.) It is seldom prescribed in substance;

<sup>1</sup> *Bellonii Obs.* l. iii. c. 41. *Tournefort, Voyage*, ii. let. 21. p. 189. Grew.



but either in the form of tincture, or of extract, or of decoction made with two drachms of the root to a pint of water.<sup>1</sup> The dose of the root is from grs. x. to ℥j., which purges strongly; but to produce its other effects two or three grains are sufficient. Of the decoction f℥j. may be given every four hours.

Official Preparations. *Tinctura Hellebori nigri*, L. E. D. *Extractum Hellebori nigri*, E. D.

## 2. HELLEBORUS FÆTIDUS.

*Official.* HELLEBORI FÆTIDI FOLIA, *Lond.* HELLEBORASTER; FOLIA, *Dub.* The leaves of Fœtid Hellebore.

*Syn.* Hêllebore fœtide (*F.*), Stinkende Niesswurzell (*G.*), Elleboro fetido (*I.*), Helleboro hediondo (*S.*).

This is a perennial, indigenous plant, growing under hedges and in shady places, on a chalky soil, flowering in March and April. The root is small and bent, with a great many slender, dark-coloured fibres. The stem is about eighteen inches high, round, strong, naked, divided and subdivided into branches, and compressed towards the top. The leaves, which stand upon long channelled footstalks, surrounding the middle of the stem, are of a deep, lurid-green colour, and pedate: the leaflets long, narrow, lanceolate, obscurely serrated, generally nine in number, four united at their bases on each side, and one terminal. At each ramification of the flower-stem is a stem-clasping, broad stipule, or scaly leaf, trifid at the place where the stem first branches, then bifid, and at the ultimate subdivisions entire, oval, and pointed; those on the flowering stem are of a much paler green than the proper leaves. The flowers are numerous, terminal, peduncled and pendent: the petals are five, two of which are oval, and the other three heart-shaped; the whole concave, persistent, of a pale-green colour tipped with a claret-purple: the nectaries are five or eight, similar to those of the black hellebore. The stamens are numerous, and flat at the base, the length of the petals, with white anthers; the germens three or more, becoming beaked pods, containing round, black seeds.

These roots were long used as domestic worm medicine, before they were introduced into the list of materia medica by the London College.<sup>2</sup>

*Qualities.* — The odour of the recent plant is fœtid, the taste of the leaves, when chewed, bitterish, biting, and so acrid as to excoriate the mouth. The stipules possess these qualities in a greater degree than the proper leaves.

*Medical properties and uses.* — The leaves of fœtid hellebore are strongly cathartic and emetic; and in overdoses prove highly deleterious. They have been successfully, although

<sup>1</sup> Wintringham, *Thesaurus Med.* p. 87.

<sup>2</sup> “Probably upon the authority of Dr. Bisset.” *Med. Bot.* 478.



they are now rarely, used as an anthelmintic against the lumbricus teres, for which they were strongly recommended by Dr. Bissett; and Woodville tried them with advantage upon a girl of twenty years of age, in the Middlesex Dispensary. They are given dried, in the form of powder; or a decoction made by boiling ʒij. of the recent leaves, or ʒss. of the dried, in fʒviij. of water for fifteen minutes; or a syrup made with the expressed juice of the recent leaves moistened with vinegar, which is supposed to correct the violent effects of the drug.

The dose of the powder is from grs. vj. to ʒj.; of the decoction from fʒj.; and of the syrup a tea-spoonful at bedtime, and one or two in the morning, to children betwixt two and six years of age, on two or three successive days.

HERACLEUM. *Spec. Plant. Willd.* i. 1421.

Cl. 5. Ord. 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

G. 541. Fruit elliptical, emarginate, compressed, striated, margined.

Corolla difform, inflex, emarginate. Involucre caducous.

Species *H. gummiferum*. Gum-bearing Heracleum. *Willd. Hortus Berolin.* i. t. 53, 54.

Officinal. AMMONIACUM, *Lond. Edin. Dub.* Ammoniac.

Syn. Gomme Ammoniaque (*F.*), Ammoniak (*G.*), Gomma Ammoniaco (*I.*), Goma Amoniaco (*S.*), Ushok or Feshook (*Arab.*), Semughbilsheeren (*Pers.*).

The plant which yields this gum-resin is a native of Africa, and the East Indies. It has not been scientifically described by any one who has seen it growing in its native soil<sup>1</sup>, and the description which is about to be given, is that of a plant which Willdenow reared from seed found in the ammoniacum of the shops, and has named *Heracleum gummiferum*. The London College, on his authority, has admitted it as the ammoniacum plant; it flowers in June and July. The root is tapering, a span in length, fleshy, whitish, and twice divided at the apex. The stem rises three feet in height, is branched, erect, about an inch thick at the base, deeply furrowed, and sparsely furnished with hairs. The branches are opposite and divaricated. The radical leaves are a span in length, cordate, three-lobed, toothed, pubescent on the under surface, and supported on roundish, channelled, furrowed, petioles: the stem-leaves are opposite, somewhat cordate, three or four inches long, toothed, on petioles, the margin of the base of which is leafy, ventricose, and sheathing. The umbels are large and many-rayed, composed of many-flowered convex umbellules. The involucre is polyphyllous, with the leaflets linear, lanceolate, and deciduous; as are also the involucels, which, however, are persistent. The marginal flowers are hermaphrodite and rayed,

<sup>1</sup> Mr. Jackson has seen and described the plant, of which he has also given a figure, but neither can be regarded as scientifically correct; and of course not authority to be relied on. *Account of Morocco*, p. 83.



the central hermaphrodite without the germen. The margin of the calyx is obsolete. The *marginal* flowers have a pentapetalous unequal corolla, the two outer petals being large, dilated, and somewhat cordate; and the three inner inflex, cordate, and half the size only of the others. The *central* flowers are pentapetalous, equal, with inflex cordate petals. The filaments are five, capillary, with roundish anthers. There is no germen in the central flowers, but in the marginal it is oblong, inferior, and crowned with two short styles inserted into a glandular body, with capitate stigmas. The fruit is oblong, slightly emarginate, consisting of two striated seeds, convex on one side, and on the other plane.

Willdenow could not obtain any of the gum-resin from this plant; but he has no doubt of its being the plant from which it is obtained. Mr. Jackson, in his account of Marocco, informs us, that the ammoniacum plant, which in the Arabic is named *feshook*, resembles the fennel<sup>1</sup>, is ten feet in height, and one inch thick in the thickest part of the stem. The plant grows at El-araiche and M'Sharrah Rumellah; and neither bird nor beast is seen near the spot, but it is attacked by a horned beetle which perforates the stem with its horn, and the juice runs out at the wound. The ammoniacum is, however, procured by incisions also, and allowed to drop on the ground, where it hardens by the air and sun; on which account that from Barbary is mixed with a red earth, and is not saleable in the London market. The best ammoniacum is brought from the East Indies, packed in cases and chests. It is in large masses composed of small round fragments or tears: or in separate dry tears, which is generally considered a sign of its goodness.

*Qualities.* — Ammoniacum has a peculiar faint but not ungrateful smell; and a bitter, nauseous, sweet taste. The tears are yellow on the outside, and white within; brittle, and break with a vitreous fracture. Their specific gravity is 1.207. Ammoniacum is adhesive in the warm hand, softens by heat, but does not melt; and is partially soluble in water, alcohol, ether, solutions of alkalies, and vinegar. When triturated with water the solution is milky, but after some time it lets fall a resinous matter; which is the part of the ammoniacum that is taken up by ether and alcohol. Water or alcohol, when distilled off ammoniacum, bring over nothing from it. According to the analysis of Braconnot, it is composed of 70.0 parts of resin, 18.4 gum, 4.4 glutinous matter, and 6.0 water, in 100.0 parts; 1.2 parts being lost in the analysis.<sup>2</sup> I find that sul-

<sup>1</sup> Both Dioscorides and Pliny describe ammoniacum as the juice of a species of *Ferula* growing in Libya. *Dioscor.* l. iii. c. 98. *Plin.* l. xii. c. 23.

<sup>2</sup> *Annales de Chim.* lxxviii. 69. *Thomson's Chemistry*, v. 143.



phuric ether takes up six grains in ten of ammoniacum, and when evaporated, leaves a yellowish white resin<sup>1</sup>, which is long of hardening, and is insipid, although it possesses the odour of the gum-resin: the taste resides in the gum, which in other respects possesses the properties of acacia-gum. Water, therefore, is the proper menstruum for ammoniacum.

*Medical properties and uses.*—Ammoniacum is a stimulating expectorant, deobstruent, and antispasmodic; and is in large doses purgative. Externally it is discutient and resolvent. It is prescribed with advantage in asthma, chronic catarrh, and some other pulmonary affections; but, on account of its stimulating properties, its use must be avoided where any inflammatory action of the chest is going forward. As a deobstruent it is useful in visceral obstructions, hysteria, and chlorosis: and in that peculiar state of the bowels often accompanying hypochondriasis and dyspepsia, in which there is an almost constant degree of colic, particularly after taking food, and which appears to arise from a viscid mucus lodged in the intestines, a combination of ammoniacum and rhubarb is singularly efficacious. As an antispasmodic, Cullen properly considers it as the least powerful of the fetid gums. It may be combined with tartrized antimony, squills, assafoetida, and ipecacuanha, to promote its expectorant powers; and with myrrh, iron, and bitters, when its deobstruent properties are required. It is given either in substance, or diffused in water in the form of emulsion. Externally, it is applied under the form of plaster to scirrhus tumours and white swellings of the joints. (See *Preparations and Compositions*.)

The dose of ammoniacum is from grs. x. to grs. xxx.

Officinal preparations. *Mistura Ammoniacy*, L. D. *Emplastrum Ammoniacy*, L. *Emplastrum Ammoniacy cum Hydrargyro*, L. *Emplastrum Gummosum*, E. *Pilulæ Scillæ comp.* L. E.

HIRUDO.<sup>2</sup> *Syst. Nat. Gmelin*. i. 3095.

Cl. 6. Ord. 1. Vermes intestina.

G. 280. Body oblong, truncated at both extremities, cartilaginous, moving by dilating the head and tail.

Sp. 2. *H. medicinalis*. The Medicinal Leech. *Amœnit. Academ.* vii. 40.

Officinal. HIRUDO MEDICINALIS, *Dub.* The Leech.

*Syn.* Sang sue (*F.*), Blutiul; *Ægle*; Lake (*G.*), Sanguisuca; Mignatta (*I.*), Sanguijuela (*S.*), Khéruheen (*Arab.*), Jonc (*H.*), Jelauca (*Sanc.*), Utter (*Tam.*), Patchet (*Malay*), Zeloo (*Pers.*).

This species of leech is common throughout Europe, inhabiting lakes and stagnant pools. The body is about three

<sup>1</sup> Nitric acid converts this resin into a yellow matter, which imparts a permanent yellow colour to silk.

<sup>2</sup> Named from *haurio*, expressive of its well known peculiar action. — *Johnson's Treatise*, p. 40.



inches long, tapering towards the head, composed of rings, and capable of being very much lengthened and contracted. The colour of the back is dark olive, divided by four yellow or buff-coloured, longitudinal lines, two of which are lateral, with a black line running through their centres; and the other two, which are on the upper part of the back, dividing it into three, nearly equal parts, are broken with black. Within these lateral and upper lines are two others, which appear like chains of black and yellow. The belly is pale olive, thickly maculated with black or very dark blue, irregular spots. The mouth is triangular, placed in the centre of a horse-shoe sucker which is under the head: and at the anal extremity there is a broad, circular sucker, by which it attaches itself to different bodies.

Leeches are oviparous. All the ova are discharged in one involucre, near the surface and the margins of pools, and are hatched by the heat of the sun. They do not cast the skin, as has been generally supposed; but at certain times throw off a tough, slimy substance from their bodies, apparently the production of disease; and from which they get disincumbered by drawing themselves through between the moss and the matted roots of rushes.<sup>1</sup> During winter they remain almost torpid, hid amongst the thick network of aquatic roots which surround the pools.

Norfolk supplies a great part of the leeches which are brought to the London market: some are taken also in Suffolk, Hampshire, Kent, Essex, and Wales, but many are imported from Bourdeaux and Lisbon.<sup>2</sup> They are caught in spring and autumn by people who wade into the pools and allow them to fasten on their limbs; or more generally, the catchers beat, as they wade in, the surface of the water with poles, which sets the leeches in motion, and brings them to the surface; when they are taken with the hand and put into bags. As they come to the surface just before a thunder storm, this is regarded as a good time for collecting them. They are best preserved in vessels half filled with soft water, kept in an equal and moderate temperature (50° Fahr.), and covered over with a coarse cloth, so as to admit the air. The water should be changed once a week; and all the dead or sickly leeches removed from the general stock, for they are subject to much disease and great mortality. Leeches which have been used should not be returned to the stock till they appear to have completely

---

<sup>1</sup> I give this on the authority of Mr. Dickson of Covent Garden, who has made many curious observations on the œconomy of the leech.

<sup>2</sup> These differ from English leeches chiefly in having the belly of one uniform colour.



regained their health and vigour, which is known by their feeling hard and firm when handled. As we are ignorant of their proper and natural food, it is useless to attempt to feed them<sup>1</sup>; but in winter it would perhaps be advantageous to put some moss into the vessel in which they are preserved.

*Medical uses.* — Leeches appear to have been first used by Themison. They are applied in cases where local blood-letting is necessary, as in ophthalmia, and particularly to places where cupping-glasses cannot be applied. In some habits, where there is a disposition to erysipelatous inflammation, their bites, which are triangular, occasion a considerable degree of irritation, and œdematous swellings follow, which are exceedingly troublesome; but in general they easily heal, and occasion no inconvenience. It is sometimes exceedingly difficult to make them bite, which they never will do when they are sick. The best mode of applying them, is to take them out of the water for some minutes before they are to be used, and to dry them well with a very soft cloth directly before they are applied. The part should also be well cleaned with soap and water, then washed with a little pure water, and made very dry. If there be any hairs on the spot, these must be close shaved. I have found this method preferable to that of wetting the part with milk and sugar, blood, or any other matter. When they, nevertheless, will not readily fix, or when it is wished to apply them very exactly on a particular spot, as, for instance, close to the angle of the eye in ophthalmia, Dr. Johnson recommends to puncture the part with a lancet; but I find that putting them into a large quill cut at both ends, and applying the end at which the head of the animal lies to the part, with the finger over the other end, is an excellent mode of making them bite. The quill is withdrawn after they are firmly fixed. They drop off spontaneously whenever they have gorged themselves with blood; and they may be separated at any time by sprinkling a little salt on the head. Very few leeches can draw more than half a fluid ounce of blood; and, therefore, it is necessary, in order to increase the quantity, to keep the orifices bleeding, by bathing them with hot water. It has been recommended to cut off the tail of the leech, so as to allow the blood to be discharged as fast as it is sucked, the leech continuing to suck notwithstanding this mutilation. After leeches drop off, the application of a very little salt makes them disgorge all the blood they have sucked; and if they be immediately thrown into clean water, and this re-

---

<sup>1</sup> Dr. Johnson says they live by adhering to and sucking the fluids of fish, frogs, &c.; but they take no kind of solid food. *Treatise on the Medicinal Leech*, p. 61.



peatedly changed for three or four times, they soon recover their health and vigour. Dr. Johnson advises the use of vinegar instead of salt, which is not apt to blister the lips of the leech as salt does, preventing it from sucking for some considerable time.

**HORDEUM.** *Spec. Plant. Willd.* i. 472.

*Cl.* 3. *Ord.* 2. Triandria Digynia. *Nat. ord.* Gramineæ.

*G.* 151. *Calyx* lateral, two-valved, one-flowered, three-fold.

*Species* 3. *H. distichon*. Common Barley. *Viborg-Cereal.* 35. t. 3.

*Officinal.* HORDEI SEMINA, *Lond.* HORDEI DISTICHI SEMINA,

*Edin.* HORDEUM DISTICHUM; SEMINA, *Dub.* Barley.

*Syn.* Orge (*F.*), Gerstengraupen (*G.*), Gemeenegerst (*Dutch*), Byg (*Dan.*), Korn (*Swed.*), Jeezmien (*Pol.*), Orzo (*I.*), Cebada (*S.*), Cevada (*Port.*), Iow (*H.*), Bārlee Arise (*Tam.*), Dhourra (*Arab.*).

Barley is asserted by Reidesel to be a native of Tartary, but the fact is not well ascertained.<sup>1</sup> It is an annual plant, and cultivated in almost every country of Europe. This species, which is the most generally cultivated in Britain, has a long flat spike or ear, with a double row of defective or male florets on each flat side, and a single row of fertile florets at each edge. The valves of the calyx, or outer chaff, are linear, and one half shorter than the corolla or inner chaff, which terminates in a straight, serrated awn or beard, sixteen times its own length. When ripe, the husk is coriaceous, angular, and continues close about the grain, which, when freed from it, is ovate, grooved, and angular.

Barley is used as an article of food, but less so than it was in former times; and it is now chiefly cultivated for the purpose of forming malt liquors and ardent spirits. It is formed into pearl barley by two different operations; the barley is first spread out and moistened; and then, in this state, by means of machinery, is denuded of the cuticle, or shelled. It is afterwards rounded in a mill, which at the same time polishes the little granules into which it is formed.

*Qualities.* — Pearl barley is inodorous, and has a slightly sweetish taste. It consists of roundish granules of a pearly whiteness, composed almost entirely of starch, with some gluten, mucilage, and saccharine matter<sup>2</sup>, which are dissolved in

<sup>1</sup> Carden asserts that it is a native of Athol, in Scotland. Diodorus Siculus refers it to Egypt, where, he says, Osiris found it wild, and first cultivated it.

<sup>2</sup> Einhof, who analysed barley both in the unripe and ripe state, found that 3840 parts of barley, in grain, afforded 430 of a volatile matter, 720 husk, and 2690 of meal; and from the same quantity of barley-meal, he obtained 360 of volatile matter, 44 albumen, 200 saccharine matter, 176 mucilage, 9 phosphate of lime, with some albumen, 135 gluten, 260 husk, with some gluten and starch, and 2580 of starch; 76 parts were lost in the analysis. When this meal is macerated in alcohol it yields a yellow-coloured acrid thick oil, which is supposed to give the peculiar flavour to spirits from raw grain, and to be lost in malting. *Thomson's Chemistry*, v. 254.



boiling water. The decoction very soon runs into the acetous fermentation. Barley is never used medicinally in substance.

Officinal preparations. *Decoctum Hordei*, L. E. D. *Decoctum Hordei compositum*, L. D.

HUMULUS. *Spec. Plant. Willd.* iv. 769.

Cl. 22. Ord. 5. Dioecia Pentandria. *Nat. ord.* Scabridæ, Linn. Urticæ, Juss.

G. 1795. Male. Calyx five-leaved. Corolla none.

———— Female. Calyx one-leaved, obliquely spreading, entire.

Corolla none. Styles two. Seed one, within a leafy calyx.

Species 1. *H. Lupulus*. The Hop. *Eng. Bot. t.* 427. *Smith's Flor. Brit.* 1077.

Officinal. HUMULI STROBILI, *Lond. Edin.* The Strobiles of the Hop.

*Syn.* Houblon grim pant (*F.*), Hopfen (*G.*), Luppolo (*I.*), Hoblon (*S.*).

The hop is an indigenous perennial plant, growing in hedges, and flowering in July. It is very abundantly cultivated in Kent, Essex, Surrey, and Suffolk, and the strobiles are picked about the end of August or the beginning of September.<sup>1</sup> The root sends up many long, striated, angled, rough, flexible stems, which support themselves by twining round upright bodies in a spiral direction from left to right. The leaves are opposite, in pairs, petiolate, heart-shaped, serrated, entire, or lobed, and of a dark green colour on the upper disc. Both the leaves and petioles are scabrous, with minute prickles; and at the base of each leaf-stalk are two interfoliaceous, entire, reflected, smooth stipules. The flowers are axillary, and furnished with bractes: the males are yellowish white, in panicles, and drooping: the females, which are on distinct plants, are in solitary cones or strobiles, ovate, and pendulous; composed of membraneous scales of a pale greenish colour, tubular from being rolled in at the base, and two-flowered, each containing one round, flattish seed, of a bay-brown colour, surrounded with a sharp rim, and compressed at the tip.

At the proper season, while the strobiles are yet scarcely ripe, the plants are cut about three feet from the ground, the poles on which they are twined pulled up, and the strobiles carefully picked off, one by one. Those that are over-ripe or defective are separated from those that are ripe enough, and both kinds are carried to the kiln as soon as possible after they are picked. The heat of the kiln requires to be regulated

---

<sup>1</sup> The culture of the hop plant was introduced into England from Flanders in 1524, and the strobiles were first used for preserving English beer in the latter part of the reign of Henry VIII.; but the prejudice against them was very considerable, and the city of London, a hundred years afterwards, petitioned the parliament to prevent their use. There are now, however, severe penalties inflicted on brewers who use any other bitter for preserving their beer.



with great nicety; and in order to prevent them from drying too fast, many kilns have two floors, on the uppermost of which the greener hops are laid, and gradually dried before being brought to support the heat of the lower floor.<sup>1</sup> Charcoal is the fuel usually employed; other kinds of fuel injuring the flavour of the hops. The strobiles are considered sufficiently dried when they become crisp; but they acquire a degree of toughness and tenacity before they are bagged, from being laid in heaps in the store-houses. Five pounds of moist or under-ripe hops make one pound only when taken from the kiln. The best hops are brought to market in fine canvass sacks called "pockets," each of which contains about  $1\frac{1}{4}$  cwt. of hops.

*Qualities.*—Hops have a strong, peculiar, fragrant, sub-narcotic odour, and a very bitter, aromatic, astringent taste. They have a pale greenish yellow hue, appear like thin transparent veined leaves; and although not tough, yet are difficult to pulverize. Some late experiments by Dr. A. W. Ives prove, that the active properties of the strobiles reside in a substance which forms one-sixth part only of their weight, and which is easily separated, by merely sifting in a fine sieve. Dr. Ives has named it *lupulin*. He found in 120 grains of lupulin, five grains of tannin, ten of extractive, eleven of bitter principle, twelve of wax, thirty-six of resin, and forty-six of woody fibre (lignin). Hops, from which all the lupulin is separated, yield an extract, which possesses none of the virtues of the hop.<sup>2</sup> The virtues of the strobiles are extracted by boiling water, or alcohol, or ether. The watery infusion has a pale straw colour, is rendered muddy by the mineral acids; alkalies deepen its colour; it strikes an olive with sulphate of iron; is precipitated by alcohol, solutions of superacetate of lead, nitrate of silver, and tartarized antimony: and when rubbed with magnesia or lime, a rod dipped in muriatic acid discovers the presence of ammonia. The ethereal tincture, when evaporated on water, leaves a pellicle of greenish intensely bitter resin, and deposits some extractive. By distillation in water, hops yield a volatile aromatic oil. From these experiments they appear to contain resin, extractive, volatile oil, tannin, an ammoniacal salt, and what has been termed the bitter principle.

*Medical properties and uses.*—Hops are narcotic, tonic, diuretic; and, externally applied, anodyne and discutient. Their use as a preservative of beer has been long known. They are also said to possess the power of procuring sleep in the delirium of fever, and in mania, when used as a pillow; and owing to this effect having been confirmed in the case of the

---

<sup>1</sup> This is the case at Farnham in Surrey. See *Stevenson's Survey*, 563.

<sup>2</sup> *Annals of Phil.* p. 194.



late king, George the Third, their efficacy as a general narcotic, when introduced into the stomach, has been investigated.<sup>1</sup> Dr. Maton observed that, besides allaying pain and producing sleep, the preparations of hops reduce the frequency of the pulse, and increase its firmness in a very direct manner. One drachm of the tincture and four grains of the extract given once in six hours, reduced the pulsations from ninety-six to sixty in twenty-four hours.<sup>2</sup> He found the extract exceedingly efficacious in allaying the pain of articular rheumatism: but our own experience has not afforded us sufficient proof of its utility as a sedative; and Dr. Bigsby's<sup>3</sup> experiments have lessened very much the confidence practitioners were disposed to give to it; indeed, we are rather surprised that the Edinburgh College has adopted it into their list of *Materia Medica*. An ointment compounded with the powder of the hop and lard is recommended by Mr. Freake as an anodyne application to cancerous sores. We have seen a fomentation of it afford much relief in painful swellings and tumours. It may be given in the form of powder, infusion, tincture, or extract. The dose of the powder is from grs. iij. to ℥j; that of the infusion, which is made with ℥ss. of the hops and Oj. of boiling water, f℥jss. with f℥ss. of cinnamon-water, twice or thrice a day.

Officinal preparation. *Extractum Humuli*, L. *Tinctura Humuli*, L. E.

#### HYDRARGYRUM.<sup>4</sup> Mercury or Quicksilver.

*Syn.* Mercure (*F.*), Quicksilber (*G.*), Kwitzilver (*Belg.*), Quicksolv (*Dan.*), Quicksilver (*Swed.*), Mercurio (*I.*), Azogue (*S. & Portug.*), Abue (*Arab.*), Pára (*H.*), Sodum (*San.*), Rassum (*Tam.*), Parah (*Duk.*), Seemab (*Pers.*), Rassa (*Malay*).

This metal is found in Spain, Germany, and Hungary; Siberia, the Philippines, China and Peru. The most productive mines are those of Idria, Carinthia, and the Palatinate; Almaden, near Cordova, in Spain<sup>5</sup>; and Guanica Velica, near Potosi, in Peru.<sup>6</sup> It is procured,

##### A. In its metallic state:

- |                          |                               |
|--------------------------|-------------------------------|
| i. Unalloyed.            | Sp. 1. <i>Native mercury.</i> |
| ii. Alloyed with silver. | 1. <i>Native amalgam.</i>     |

<sup>1</sup> *De Roche's. De Humuli Lupuli Viribus medicis.*

<sup>2</sup> *Observations on the Humulus Lupulus, &c. by A. Freake.*

<sup>3</sup> *Vide London Medical Repository, vol. v. p. 97.*

<sup>4</sup> Ὑδράργυρος Græcorum.

<sup>5</sup> This is the oldest and the richest mine of Cinnabar in Europe. It was wrought by the Romans two thousand years ago; and yields about 6000 quintals of fluid mercury, annually.

<sup>6</sup> These were discovered in 1566 and 1567, by Henry Garces, a Portuguese. Garces was a native of Porto, and went to Peru in the Spanish service. Examining one day the red earth which the Indians used as paint, and called *limpi*, he observed it was native cinnabar; and knowing that mercury was extracted from cinnabar in Europe, he began to work the Peruvian mines.







tin. When the metal quickly loses its lustre, is covered with a film, or is less fluid and mobile than usual, leaves a stain on a delft plate, or does not readily divide into round globules, but into those with tails, it may be suspected. Lead is discovered by dissolving a portion of the suspected mercury in nitric acid, and adding to the solution, water saturated with sulphuretted hydrogen gas, which gives a brown precipitate if it be present, and by this means one part of lead may be detected in 15,260 of mercury. Bismuth is detected by pouring the above-mentioned nitric solution into distilled water, when the bismuth will appear as a white precipitate. Exposing the mercury to heat detects zinc; and tin is discovered by a weak nitro-muriatic solution of gold, which is precipitated purple by tin. It is purified by distillation with iron filings, or by agitation in diluted sulphuric acid until the acid cease to become turbid; and then, after washing and drying the globules into which it has been divided, passing them through a pin-hole in the bottom of a funnel of writing paper.<sup>1</sup>

*Qualities.*—Pure mercury is inodorous, insipid, and of a bright white or silver colour. Its specific gravity is 13.568.<sup>2</sup> It is always fluid at the ordinary temperature of the atmosphere, but becomes a solid malleable metal in a degree of cold sufficient to sink the thermometer to 39° below 0 of Fahrenheit.<sup>3</sup> It boils at 656°, and is volatilized unchanged in close vessels, but is not capable of combustion.<sup>4</sup> Mercury is oxidized by the air at its usual temperature, when subject to agitation, and is fully saturated with oxygen in a continued heat of 600°. It is oxidized by, and combines with the sulphuric, nitric, and oxymuriatic acids; and its oxides also enter into combinations with the other acids. It unites with sulphur and phosphorus, and combines with many other metals, forming what are called amalgams.

*Medical properties and uses.*—Mercury in its metallic state exerts no action on the animal system. It has nevertheless been administered in doses of a pound or more with the view of operating mechanically, and overcoming by its weight the obstruction of the intestines which exists in ileus: but, as it cannot act by its gravity on the ascending part of the bowels, it is not easy to conceive how it should have been ever recommended; and the events of the cases in which it has been given have sufficiently proved the futility of the practice.

Mercury, however, when prepared for medicinal use, is a

---

<sup>1</sup> This method was invented by Professor Brunchi, of Pisa. Vide *Phil. Mag.* iv. p. 548.

<sup>2</sup> Cavendish.

<sup>3</sup> Crichton. *Phil. Mag.* xiv. 49.

<sup>4</sup> *Thomson's Chemistry*, i. 175. If, however, the galvanic fluid be passed through it, the beautiful luminous stars in which it is dispersed seem to prove its combustibility.



remedy of the most extensive application. It is a powerful and general stimulant, but its effects are certainly different from those of other articles which are ranked in the same class. It enters into the circulation, quickens the vascular action, and excites powerfully the whole of the glandular system; increasing all the secretions and excretions, and has even been detected in the urine.<sup>1</sup> It has been supposed, that it is peculiarly determined to the salivary glands; but if, as there is every reason to suppose, these glands are endowed with more irritability<sup>2</sup> than the rest of the habit, it is easy to conceive that the same degree of stimulus, which is operating on the whole system, will produce a greater effect on them in a direct ratio according to their greater susceptibility. But although its general action is stimulant, yet the various preparations of it produce different effects, operating sometimes as stimulants, sometimes as cathartics, or emmenagogues, and locally as errhines: and hence the great variety of diseases in which it has been found useful: as, febrile affections, spasms, cachetic diseases, glandular obstructions, cutaneous eruptions, and membranous inflammation. (See *Preparations and Compositions*.)

But the most important effect of the preparations of mercury is their specific operation in syphilis. They were used, and their effects, when accumulated in the habit, were known so early as the 13th century; and the writings of Theodorick<sup>3</sup> contain cautions against catching cold during the course; but the first notice of mercury as a remedy in Lues Venerea is contained in a tract by Jo. Almenar, a Spaniard, published in 1516; who recommends it after the manner of the Arabians, but condemns pushing the remedy so as to promote salivation. In a tract on Syphilis by Laurentius Phrisius, published in 1532,<sup>4</sup> four formulæ of ointments in which mercury is an ingredient are given under the title of *Unguenta Emporicorum*; and it does not appear that it was then used even as an external application by regular practitioners. Physicians, however, did not venture to give mercury internally, till Paracelsus broke the fetters of ancient authority, and proved that it might be exhibited not only with safety, but with advantage. Since his time, a period of nearly 300

---

<sup>1</sup> Mem. della Reale Acad. di Torino, tom. xxix. p. 228.

<sup>2</sup> That the salivary glands and their excretories are very excitable, is evident from the flow of the saliva being much increased by affections of the mind, as the thinking of any kind of food which is particularly grateful to the taste.

<sup>3</sup> He was a friar, afterwards bishop of Cervia, and died between the years 1270 and 80. See *Friend's History of Physic*, ii. 360.

<sup>4</sup> *Epitome opusculi de curandis Pusculis, ulceribus et doloribus morbi Gallici, mali Fruntzoss appellati, autore Laurentio Phrisio, artium et medicinæ doctore.*



years, experience has fully sanctioned its use; but, although, as Mr. Pearson justly observes, “not one medicine besides derived from the animal, vegetable, or mineral kingdom, has maintained its credit, with men actually employed in extensive practice, during a tenth part of that period<sup>1</sup>:” yet, it is remarkable, that in the present day its utility in syphilis has been questioned.<sup>2</sup> Many various theories of the operation of mercury have been advanced; the most satisfactory of which is that of Mr. Hunter, who supposed that the stimulant operation of the mercury induces and maintains an action which is incompatible with the morbid action produced by the venereal virus, until the poison is either destroyed, or evacuated from the body by the excretories. But whatever may be the principles on which it operates, its efficacy in this disease is certain, when it is judiciously and cautiously administered. The mode of giving it, and the morbid effects which it produces under certain circumstances, shall be mentioned when its preparations are described: it is only necessary to observe further in this place, that although men of the first medical talents have, occasionally, declaimed against its use,<sup>3</sup> and although much mischief may have of late years arisen from its indiscriminate employment by the speculative and the ignorant, yet, in the hands of judicious and cautious practitioners, it will continue to rank as one of the most useful of the articles of the materia medica.

Officinal preparations.<sup>4</sup>

- I. By distillation to purify the metal.
  1. *Hydrargyrum purificatum*, L. E. D.
- II. By trituration; (suboxidized.)
  - a. With animal fat.
    2. *Unguentum Hydrargyri fortius*, L. *Ung. Hydrargyri*, D.
    3. ————— *Hydrargyri*, E.
    4. ————— *mitius*, L. D.
    5. *Linimentum Hydrargyri*, L.
    6. *Emplastrum Ammoniaci cum Hydrargyro*, L. D.
    - *Hydrargyri*, L. E.
  - b. With saccharine substances.
    7. *Pilulæ Hydrargyri*, L. E. D.
  - c. With carbonate of lime.
    8. *Hydrargyrum cum Cretâ*, L. D.
  - d. With carbonate of magnesia.
    9. *Hydrargyrum cum Magnesia*, D.
- III. By the action of heat and air; (oxidized.)
  10. *Hydrargyri Oxydum rubrum*, L. *Oxydum Hydrargyri*, D.

<sup>1</sup> Observations, &c. p. 97.

<sup>2</sup> Medico-Chirurg. Trans. vol. ix.

<sup>3</sup> Saunders — Observations on the Hepatitis of India, &c.

<sup>4</sup> In forming this table, we have been much assisted by the excellent table drawn up by Dr. Duncan, jun. in the Edinburgh New Dispensatory; and we trust our alterations will render it more practically useful.



## IV. By the action of acids.

## a. With sulphuric acid ; (suboxidized.)

11. *Subsulphas Hydrargyri flavus*, E. *Oxydum Hydrargyri sulphuricum*, D.

## b. With nitric acid ; (suboxidized.)

12. *Unguentum Hydrargyri nitratis*, L. E. *Unguentum Super-nitratis Hydrargyri*, D.

13. *Unguentum Nitratis Hydrargyri mitius*, E.

————— (oxidized.)

14. *Hydrargyri Nitrico-oxydum*, L. *Oxydum Hydrargyri rubrum per Acidum nitricum*, E. *Oxydum Hydrargyri nitricum*, D.

15. *Unguentum Hydrargyri nitrico-oxydi*, L. *Unguentum Oxidi Hydrargyri rubri*, E. *Unguentum Subnitratis Hydrargyri*, D.

## c. With muriatic acid.

† sublimated ; (oxidized.)

16. *Hydrargyri Submurias*, L. *Submurias Hydrargyri mitis*, E. *Submurias Hydrargyri sublimatum*, D.

17. *Pilulæ Hydrargyri submuriatis*, L.

————— (oxidized and acidified.)

18. *Oxymurias Hydrargyri*, L. *Murias Hydrargyri corrosivus*, E. *Murias Hydrargyri corrosivum*, D.

19. *Liquor Hydrargyri Oxymuriatis*, L.

†† precipitated ; (oxidized.)

20. *Submurias Hydrargyri præcipitatus*, E. D.

## d. With acetous acid ; (suboxidized.)

21. *Acetis Hydrargyri*, E. *Acetas Hydrargyri*, D.

## V. By precipitation with earths and alkalies from acid solutions.

## a. By lime-water from the nitric solution ; (suboxidized.)

22. *Hydrargyri Oxydum cinereum*, L.

## b. By ammonia from the nitric solution ; (suboxidized.)

23. *Oxydum Hydrargyri cinereum*, E. *Pulvis Hydrargyri cinereus*, D.

## c. By ammonia from the muriatic solution ; (oxidized.)

24. *Submurias Hydrargyri ammoniatum*, D. *Hydrargyrus Præcipitatus albus*, L.

25. *Unguentum Submuriatis Hydrargyri ammoniati*, D. *Ung. Hydrargyri Præcipitati albi*, L.

## VI. Combined with sulphur.

## a. By trituration.

26. *Sulphuretum Hydrargyri nigrum*, E. D.

## b. Sublimated.

27. *Hydrargyri Sulphuretum rubrum*, L. D.

HYOSCYAMUS.<sup>1</sup> *Spec. Plant. Willd.* i. 1010.

Cl. 5. Ord. 1. Pentandria Monogynia. Nat. ord. Luridæ, Linn. Solaneæ, Juss.

G. 378. Corolla funnel-shaped, obtuse. Stamens inclined. Capsule covered with a lid, two-celled.

<sup>1</sup> Ὕος κνᾶμος, Hog-bean.



*Spec.* 1. *H. niger*. Common Henbane. *Med. Bot.* 2d edit. 204. t. 76. *Smith, Flor. Brit.* 598. *Eng. Bot.* 591.

*Officinal.* HYOSCYAMI FOLIA ET SEMINA, *Lond.* HYOSCIAMI NIGRI HERBA; SEMINA, *Edin.* HYOSCYAMUS; HERBA, *Dub.* The leaves and seed of Henbane.

*Syn.* Jusquiamé (*F.*), Bilsenkraut (*G.*), Bilsenkruud (*Dutch*), Fandensnosser (*Dan.*), Giusquiamo nero (*I.*), Beleno (*S.*), Yosciamo (*Port.*), Belmort (*Swed.*), Bielun (*Pol.*), Khorassānie Ajooan (*H.*), Sickran (*Arab.*), Khorasanie onsum (*Tam.*), Buzirulbury (*Pers.*), Adas-pedas (*Malay*), Adas (*Jav.*)

Common henbane is an indigenous annual, frequent on waste grounds, and at the sides of roads, particularly on a calcareous soil, flowering in July. The root is long, tapering, compact, and fibrous; the stem erect, woody, round, and branched, rising about three feet in height. The leaves are alternate, sessile, and embracing the stem; large, the lower ones being above a foot in length; deeply sinuated, undulated, woolly, and of a sea-green colour. The flowers are in terminal, recurved, leafy simple spikes; and each is simple and erect. The calyx is permanent, pitcher-shaped, with a regular, five-cleft border reticulated with veins; the corolla straw-coloured, and beautifully pencilled with a net-work of purple veins. The filaments are inserted into the tube of the corolla, tapering, downy at the base, and supporting purple anthers. The style is purplish, with a blunt round stigma. The capsule is globular, invested with the body of the calyx, bilocular, and closed with a convex smooth lid. It contains numerous small irregular brown seeds.<sup>1</sup>

The whole of the plant is covered with soft, white hairs, feels clammy and slightly adhesive, and is poisonous when eaten.

*Qualities.*—The odour of the recent leaves is strong, somewhat foetid and narcotic,<sup>2</sup> and the taste mucilaginous, and slightly acrid; but, when dry, they have scarcely either odour or taste. The virtues of the plant are completely extracted by diluted alcohol. The watery infusion is of a very pale-yellow colour, and insipid; and has the narcotic odour of the plant. It is not altered by the acids. The alkalies change the colour to a deep greenish-yellow, which, on the addition of an acid, disappears, and a brownish, flocculent precipitate is produced. It is copiously precipitated by solutions of superacetate of lead white: and by nitrate of silver black. Sulphate of

<sup>1</sup> The seeds abound with oil, and may be eaten with impunity. The roots, which resemble parsnips, have occasionally been eaten, and fatal effects produced. These roots, when moulded in the turning loom and strung in the form of beads, are sold under the name of *Anodyne Necklaces*: and are worn by infants to allay the irritation of teething; a humiliating instance how deeply error is rooted in the human mind!

<sup>2</sup> In the recent state the odour of the leaves occasions stupor and delirium.



iron strikes with it a pale olive colour, and a dark precipitate is slowly formed. Thence Henbane appears to contain resin, mucus, extractive, a peculiar alkaline salt, and gallic acid. M. M. Miessner and Brandes have examined the nature of this alkaline salt, which they have named *Hyociamia*; and have ascertained, that on it depends the peculiar virtues and the poisonous properties of the plant. It crystallizes in long prisms, and forms neutral salts with the acids.

*Medical properties and uses.* — Henbane is narcotic. Its operation is very similar to that of opium, increasing at first the strength of the pulse, and producing some sense of heat; effects which are followed by proportional diminution of excitement and sleep. In some habits it occasions diaphoresis, or diuresis, and, sometimes, a pustular eruption; at other times it purges; and, in overdoses, produces sickness, stupor, dimness of sight, hard pulse, delirium, and coma, with dilatation of the pupils; until, the pulse gradually becoming weak and tremulous, petechiæ make their appearance, and death ensues. Dissections show the effects of inflammation in the stomach, bowels, and the membranes of the brain. After an emetic is given, and the stomach fully cleared, vinegar is the best antidote.

The effects of Henbane as an anodyne were known to the ancients;<sup>1</sup> but, as those were ill understood, and its use was almost completely relinquished till the time of Baron Stoeck, he may be regarded as having introduced it. It may be employed in all the cases in which the use of opium is indicated, where the latter disagrees with the habit, or where its constipating effect is wished to be avoided. In painful and spasmodic affections, hysteria, rheumatism, and gout, much benefit has resulted from its use; and we have found, that it is particularly serviceable, when united with colocynth or other powerful cathartics, in colica pictonum. It is used externally to lessen and allay the irritation of very sensible parts: thence fomentations of the leaves have been found serviceable in scrophulous and cancerous ulcers, hæmorrhoids, and other painful swellings; and Hufeland recommends the leaves and marshmallow flowers boiled in milk, with the addition of a few grains of acetate of lead, as a topical application in scrophulous ophthalmia. Smoking the leaves, like tobacco, is said to allay the pain of tooth-ach. Its effects in dilating the pupil, when an infusion of it is dropped into the eye, are similar to those of Belladonna, and thence it is also employed as a preparative to the operation for cataract. It is used,

---

<sup>1</sup> “*Hyoscyamus in potu cibove sumptus, qualem ebriorum mentis alienationem infert.*” *Dioscorid. Alexif.* . xv. 407.



generally, in the forms of extract and tincture only; but should not be prescribed in combinations with alkalies; as these destroy its narcotic powers in twenty-four hours.

Officinal preparations. *Extractum Hyoscyami*, L. E. D. *Tinctura Hyoscyami*, L. E. D.

HYSSOPUS, *Spec. Plant. Willd.* iii. 47.

Cl. 14. Ord. 1. Didynamia Gymnospermia. Nat. ord. Verticillatæ, Linn. Labiatae, Juss.

G. 1096. Corolla, lower lip three-parted, with a small intermediate subcrenate segment. Stamens straight, distant.

Species 1. *H. officinalis*. Common Hyssop. *Med. Bot.* 2d edit. 318. t. 113.

Officinal. HYSSOPI OFFICINALIS HERBA, *Edin.* HYSSOPUS; FOLIA, *Dub.* The herbaceous part and leaves of Hyssop.

Syn. Hyssope (*F.*), Isop (*G.*), Yzoop (*Dutch*), Isop (*Dan. & Swed.*), Isopo (*I.*), Hysopo (*S.*), Hyssopo (*Port.*), Zufāiy yeābus (*Arab.*).

This is a perennial plant, a native of Siberia and Austria; cultivated in our gardens,<sup>1</sup> and flowering from June to September. The root is knobbed, woody, and fibrous: the stalk about two feet in height, obscurely quadrangular, erect, shrubby, and branching. The leaves do not exceed an inch in length, and one-third of an inch in breadth, are sessile, of a somewhat glaucous, deep-green colour, elliptical, entire, punctured, and stand in pairs. The flowers are produced on one side, in long, half-verticillated, terminal spikes, and intermixed with leaves. The calyx is persistent, nearly tubular, divided at the edge into five, acute teeth, striated, and of a purplish colour at first, but afterwards green: the corolla is violet-coloured, with a long, whitish tube; the upper lip short, round, and notched at the apex; the lower one separated into three segments, the undermost of which is inversely ovate. The filaments are crowned with simple anthers; the style is slender and bifid; and the seeds are four, at the bottom of the calyx.

*Qualities*.—The leaves of hyssop have an agreeable, aromatic odour, and a bitterish, moderately warm taste; qualities that appear to depend on a volatile oil of a yellow colour, which can be obtained separate by distillation with water. It is elevated by alcohol also; but soon exhales, and the spirit loses the odour it had when newly distilled.

*Medical properties and uses*.—Hyssop is stimulant and tonic. It has been recommended in hysteria; and was formerly employed in catarrhal and other pulmonary affections, with the view of promoting expectoration; but the stimulant

<sup>1</sup> It was first cultivated in England by Gerarde in 1596. It is not the esof of the Hebrews, nor the ὕσσωπος of the Greeks. It has been supposed to be the *xife* or *cyfe* of the Arabians. *Alston's Mat. Med.* ii. 152.



properties of hyssop render its use doubtful in these diseases; and, as a tonic, it scarcely merits the least attention.

INULA. *Spec. Plant. Willd.* iii. 2089.

Cl. 19. Ord. 2. Syngenesia Superflua. *Nat. ord.* Compositæ discoideæ, *Linn.* Corymbiferæ, *Juss.*

G. 1489. Receptacle naked. Pappus simple. Anthers ending in two bristles at the base.

Species. I. *Helenium*. Elecampane. *Med. Bot.* 2d edit. 64. t. 26. *Smith, Flora Brit.* 890. *Flora Danica*, t. 728.

Officinal. HELENII. ENULA CAMPANA; RADIX, *Dub.* Elecampane root.

*Syn.* Inula Laleniere (*F.*), Alantwurz (G.), Alantswortel (*Dutch.*) Aland (*Dan. & Swed.*), Omanowy (*Pol.*), Enula Campana (*I.*), Enula Campana (*S.*), Usululrasan (*Arab.*), Beykzunjebulshamee (*Pers.*).

This species of *Inula* is an indigenous perennial, found occasionally in pastures and rich moist soils,<sup>1</sup> flowering in July and August, and ripening its seed in September. The root is thick, branched, externally of a brown or grey colour, and internally white. The stem, which rises about three feet in height, is leafy, round, and furrowed; branched near the top and villous. The leaves are large, ovate, serrated, veined, of a deep-green colour on the upper surface, and on the under reticulated, tomentose, and whitish: the radical ones are petiolate, but those of the stem sessile and embracing. The flowers are terminal, solitary, large, and of a golden colour. The calyx is scaly; the exterior scales are large, ovate, imbricated, and externally tomentose; the interior are narrow, linear, equal, and chaffy. The florets of the ray are numerous, spreading, twice the length of the calyx, linear, with the apex tridentate. The anthers end in two bristles at the base. The seeds are quadrangular, smooth, slightly curved, and furnished with a somewhat chaffy pappus. The receptacle is reticulate and papillous.

The roots of elecampane found in the shops, are generally obtained from garden plants. They are fit for use in the second year of their growth; and at this age are preferable to the older roots, which become stringy and woody. They should be dug up in autumn.

*Qualities.* — Elecampane root when dry has an aromatic, yet slightly foetid odour; and when chewed, the taste is at first disagreeable, glutinous, and in some degree resembling that of rancid soap; then aromatic, bitter, and hot. Both water and alcohol extract its virtues: the tincture possessing more of the bitterness and pungency of the root than the watery infusion. The decoction, after standing some hours,

<sup>1</sup> Ἑλένιον *Dioscoridis*. It is not unfrequent in *Essex*. *Hudson*. Between Worcester and Ludlow, and Bishop's Castle and Newton. *Smith*. I have seen it near Ewell, *Surrey*.



deposits a white powder resembling starch in appearance; but its properties show it to be a distinct principle; and it has, therefore, been named *Inulin*.<sup>1</sup> In distillation with water, this root yields a concrete, flaky substance, which seems to hold an intermediate place between camphor and volatile oil in its nature.<sup>2</sup>

*Medical properties and uses.* — Elecampane is usually ranked as a tonic; and supposed to possess deobstruent, diuretic, and expectorant properties. It was formerly regarded as a remedy of great efficacy in dyspeptic affections, flatulencies, palsy, dropsies, uterine obstructions, and pulmonary complaints; but Cullen observed, that its diuretic powers were trifling; and could not discover that it possessed any expectorant properties.<sup>3</sup> It is scarcely ever used by the regular practitioner. The dose of the powdered root may be from ʒj. to ʒj.

IRIS.<sup>4</sup> *Spec. Plant. Willd.* i. 224.

Cl. 3. Ord. 1. Triandria Monogynia. *Nat. ord.* Ensatae, Linn. Iridae, Juss.

G. 97. Corolla six-parted; the alternate segments reflected. Stigmas petal-like.

\* *Bearded with ensiform leaves.*

*Species* 7. *I. florentina*.<sup>5</sup> Florentine Iris. *Med. Bot.* 2d edit. t. 262. *Sibthorp Flora Græca*, 28. t. 39.

*Officinal.* IRIDIS FLORENTINÆ RADIX, *Edin.* The root of Florentine Iris.

*Syn.* Iris de Florence (*F.*), Violenwurzel (*G.*), Florentynse Iris (*Dutch*), Violrot (*Swed.*), Violrod (*Dan.*), Iyrsa (*Arab.*), Irva (*H.*), Ireos (*I.*), Lirio de Florencia (*S.*), Lirio de Florença (*Port.*).

This species of Iris, which is found in a wild state in Carniola, the island of Rhodes, Laconia, and other places of the south of Europe, is cultivated in our gardens;<sup>6</sup> flowering in May and June. The root is tuberous, horizontal, somewhat jointed, and sends off many fibres from the under part. The leaves spring directly from the root, spreading in opposite directions, are sheathing, sword-shaped, vertical, nerved, curved inwards at the apex, and of a sea-green colour, yellowish at the base. From amidst them, the stem rises upwards of a foot in height, erect, simple, naked, round, and commonly bearing two flowers. The flowers are large, of a pale, whitish-blue colour, erect, terminal, and inodorous, bursting from a ventricose, nerved, floral leaf: the petals are alternate, three larger and three smaller: the larger have thickish claws

<sup>1</sup> This substance, which was first noticed and its properties investigated by Rose, was named by Dr. Thomson, *System of Chemistry*, 4th edit. iv. 697.

<sup>2</sup> *Newman's Chem. by Lewis*, 2d edit. ii. 216.

<sup>3</sup> *Mat. Med.* ii. 459.

<sup>4</sup> "Iris a cœlestis arcus similitudine nomen obtinuit." *Dioscorides*,

<sup>5</sup> *Ἰρις* Theophrasti.

<sup>6</sup> It was cultivated by Gerarde in 1596.



about an inch long, bordered with a thin edge, green on the outside, and bearded within with yellow-tipped, white hairs: the border is an inch in width and longer, reflected, whitish, and striated near the flexure: the smaller are whitish-blue, stand erect, are bent inwards with a reflected margin; and have thick, attenuated, greenish claws. The anthers are white, covered by the stigmas, which have the colour of the corolla, and are cleft at the apex into two, acute, serrated, upright segments. The capsules are three-celled, containing many seeds horizontally placed.

The roots of the Florentine iris are brought in a dry state from Leghorn, packed in large casks. They are in irregular knobbed pieces, with the cuticle pared off; of a dirty yellowish white colour, and full of small holes, which mark the places whence the radical fibres issued. The best pieces break with a rough but not fibrous fracture.

*Qualities.* — These roots, when recent, have a bitterish, nauseous taste, and are very acrid; but this acrimony is lost by drying. In their dry state they are brittle, easily pulverized, have a sweetish, bitter taste, with a slight degree of pungency, and the agreeable odour of the violet, for which they are chiefly valued. When chemically examined, they appear to consist principally of fecula, with a portion of mucilage and saccharine matter: and to contain malic acid, as their infusion strikes a brown colour with sulphate of iron.

*Medical properties and uses.* — The fresh root is cathartic, and has been recommended in dropsies; the dry is nearly inert; but, in neither state, does it merit a place in the list of materia medica.

**JUNIPERUS.** *Spec. Plant. Willd.* iv. 851.

*Cl.* 22. *Ord.* 13. Diœcia Monodelphia. *Nat. ord.* Coniferæ.

*G.* 1841. Male. *Amentum* ovate. *Calyx* a scale. *Corolla* none.

*Stamens* three. Female. *Calyx* three-parted. *Petals* three.

*Styles* three. *Berry* three-sided, irregular, with the three tubercles of the calyx.

*Species* 6. *J. Sabina*. Savine. *Med. Bot.* 2d edit. 10. t. 5.

*Species* 10. *J. communis*. Common Juniper. *Med. Bot.* 2d edit.

13. t. 6. *Smith, Flora Brit.* 1085. *Eng. Bot.* 1130.

#### 1. JUNIPERUS SABINA.<sup>1</sup>

*Officinal.* SABINÆ FOLIA, *Lond.* JUNIPERI SABINÆ FOLIA, *Edin.*

SABINA; FOLIA, *Dub.* Savine leaves.

*Syn.* Sabinne (*F.*), Sadebaum (*G.*), Sevenboom (*Dutch*), Sevebom (*Dan.*), Sabina (*I.*), Sabina (*S.*).

This shrub is a native of the south of Europe and the Levant; but has been long cultivated in our gardens, flower-

---

<sup>1</sup> Βραχὺς Dioscoridis. There are two varieties of Savine; the variety β is our plant.



ing in May and June. It seldom rises above three feet in height; is covered with a brown bark, and divided into numerous, subdivided branches; which are completely invested with very small, erect, firm, opposite, pointed leaves, of a bright green colour, which lie over one another, and terminate the branches in sharp points, giving the whole shrub a very lively aspect. The *male* and *female* flowers are on different plants. The *male* catkin consists of three opposite flowers placed in a triple row, and a tenth flower at the end: and at the base of each flower is a broad scale fixed laterally to a columnar pedicel. There are filaments in the terminal flower only; tapering and united at the base, with simple anthers, which are sessile in the lateral flowers. In the *female* flowers, the calyx is three permanent scales; the petals are stiff, sharp, and also permanent; and the germen supports three styles with simple stigmas. The fruit is a spurious fleshy berry of a blackish-purple colour; marked with tubercles, the vestiges of the calyx and petals, and containing three, small, hard seeds.

*Qualities.* — The leaves and tops of Savine have a strong, heavy, disagreeable odour, and a bitter, hot taste, with a considerable degree of acrimony. These qualities depend on an essential oil, which is obtained in considerable quantity by distillation with water. Both water and alcohol extract the active principles of Savine; and Lewis found that “on inspissating the spiritous tincture, there remains an extract consisting of two distinct substances, of which one is yellow, unctuous or oily, bitterish, and very pungent; the other black, resinous, tenacious, less pungent, and subastringent.”<sup>1</sup>

*Medical properties and uses.* — Savine is a powerful stimulant, possessing diaphoretic, emmenagogue and anthelmintic properties. It has certainly a considerable effect on the uterine system; but, on account of its stimulating properties, is suited to those cases only of amenorrhœa which are unattended by fever, and in which the circulation is languid. In plethoric habits its use should be preceded by repeated bleedings<sup>2</sup>; and at all times its internal exhibition requires caution. It has been given in gout, and worm cases also, but is seldom used. As an external local stimulant or escharotic, the dried leaves in powder are applied to warts, flabby ulcers, and carious bones; and the expressed juice diluted, or an infusion of the leaves, as a lotion to gangrenous sores, scabies, and tinea capitis; or mixed with lard and wax as an issue

---

<sup>1</sup> *Mat. Medica.*<sup>2</sup> Home. *Clinical Experiments*, 387.



ointment. The dose of the powdered leaves is from grs. v. to grs. x. two or three times a-day.

Officinal preparations.—*Oleum volatile Juniperi Sabinæ*, E. D. *Extractum Sabinæ*, D. *Ceratum Sabinæ*, L.

## 2. JUNIPERUS COMMUNIS.<sup>1</sup>

Officinal. — JUNIPERI BACCÆ ET CACUMINA, *Lond.* JUNIPERI COMMUNIS BACCÆ, *Edin.* JUNIPERUS; BACCÆ. *Dub.* Juniper berries.

*Syn.* Genevrier ordinaire (*F.*), Wachholderbeeren (*G.*), Sevenboom (*Dutch*), Sevebom (*Dan.*), Ginepro (*I.*), Embro (*S.*), Zimbrow (*Portug.*), Caw-caw-ynew muna (*Cree Indian*).

The common juniper is indigenous, growing on heaths and chalky hills, and flowering in May. It is a low, very branching, rigid, smooth, evergreen shrub; but, when planted in a good soil, it rises to fifteen feet in height. The leaves are very numerous, narrow, entire, sharply pointed, channelled, of a glaucous colour on the upper surface, and sessile, standing in ternaries. The catkins are axillary, sessile, solitary, ovate, small, and furnished with bractes; the male flowers yellow at first, and afterwards brown, with great abundance of pollen; the female smaller, and of a yellowish-green colour. The berry is globular, in colour blackish-purple with a glaucous bloom, composed of the scales of the amentum, which become fleshy and coalesce. The seeds are three, and angular.<sup>2</sup>

The berries require to remain two years on the tree before they are fully ripe. The greater quantity of those which are used in Britain are brought from Germany, Holland, and Italy. The Italian berries are less shrivelled, and have a fresher and more beautiful bloom upon them than the German, and are therefore generally preferred. They are imported in bags.

*Qualities.*—Juniper berries have a peculiar aromatic odour, and a sweetish, pungent, bitterish taste, when chewed. In distillation with water they yield a volatile, terebinthinate oil of a greenish colour, on which their virtues depend.<sup>3</sup> Both water and alcohol extract their active properties. Their principal constituents are mucus, saccharine matter, and volatile oil.

*Medical properties and uses.*—Juniper berries are diuretic and cordial. They have been long known as a remedy in hydropic affections; but they cannot be depended on alone, although they form an excellent adjunct to foxglove and squill. The tops are also used: and as the virtues of the

<sup>1</sup> Ἀρκευθος μικρα Dioscoridis.

<sup>2</sup> The resinous substance known by the name of Sandarach, which is brought from Morocco, exudes from the stem of the juniper in warm climates.

The flavour and diuretic properties of Hollands depend on this oil. English gin is flavoured by oil of turpentine.



berries depend on the essential oil, which is found in the woody part also of the plant, they must be equally efficacious. They have been recommended in scorbutic and cutaneous affections; and Rosenstein asserts that a strong decoction of them soon clears the hands in scabies. The berries are sometimes given in substance, triturated with sugar or some neutral salt; but the best form is that of infusion, made with ℥iij of the berries bruised, and Oj of boiling water. The dose of the first preparation is from ℥j. to ʒss.; that of the infusion, a teacupful every three or four hours.

Officinal preparatons. *Oleum Juniperi*, L. E. D. *Spiritus Juniperi compositus*, L. E. D.

KINO. *Vide* PTEROCARPUS.

KRAMERIA. *Spec. Plant. Willd.* i. 693.

Cl. 4. Ord. 1. Tetrandria Monogynia. *Nat. ord.* Rosaceæ.

G. 253. *Calyx* none. *Corolla*, four petals: the superior *Nectary* three-parted; and inferior two-leaved.<sup>1</sup> *Berry* dry, echinated, and containing one seed.

*Spec.* 1. *K. triandra*. Triandrous *Krameria*. *Flor. Peruv.* tom. i. p. 61. *Icon.* xciii.

*Officinal.* KRAMERIÆ RADIX, *Lond.* *Krameria* or *Ratanhy* Root.

*Syn.* *Ratanhie* (*F.*), *Ruiz para los dientes* (*S.*), *Ratanhia* (*Huanaco*), *Mapato* (*Tarma*).

This plant is a native of Peru, growing on the argillaceous, sandy, and arid acclivities of the mountains in the provinces of Huanuco, Tarma, Canta, Xauxa, Caxtambo, and Huamalis, and very abundantly near the city of Huanuco. It was also found by Humboldt in the province of Guancabunba in Peru. It flowers throughout the year; but is in the height of blossom in October and November. It is a shrub, with very long, much branched, spreading roots, of a blackish-red colour exteriorly, red interiorly, and having an intensely bitter, styptic taste. The stem is procumbent, round, and divided into numerous branches, which when young are white and silky, but as they grow they become naked below, and acquire a black colour. The leaves are sparse, sessile, oblong-obovate, pointed, entire, and covered with a white, silky pubescence on both surfaces. The flowers are terminal, solitary, and pedunculated. The corolla, for there is no calyx, is subpapilionaceous, consisting of four lake-coloured petals, the inferior larger than the others, sericeous externally, but internally smooth and shining: the nectary is tetraphyllous, the two upper leaflets being spathulate, the two lower roundish, concave, and scale-like. The stamens are

<sup>1</sup> This part of Willdenow's character applies solely to *K. Ixina*; the *pentaloba* of the *Flora Peruviana*, the only species which he describes. The name *Ratanhia*, signifies trailing plant.



three, each composed of a flesh-coloured filament, inserted between the germen and the superior leaflets of the nectary, and an urceolate anther, terminated with a pencil of very short hairs, and perforated with two holes at the apex. The style is red, awl-shaped, supporting a simple stigma, and seated on an ovate germen, which changes to a dry, hirsute drupe.

Ratanhy root is collected for medicinal purposes after the rains. As imported, it consists of pieces of various sizes ; but seldom exceeding half an inch in thickness. The root breaks short, exhibiting in the fracture a woody centre, and an easily separable, fibrous, dark-red bark.

*Qualities.* — The bark of Ratanhy root, when chewed, tastes bitter, astringent, and at first nauseous ; but the impression left in the mouth is sweetish and astringent, not unlike that produced by catechu. The woody centre is nearly insipid, and perfectly inert as a remedy. Ratanhy root yields its properties to boiling water, affording a dark-brown infusion, which emits an odour not unlike that of a raw potatoe, tastes astringent and very bitter, and leaves the same impression in the mouth as the bark of the root. All the mineral acids throw down copious precipitates when added to the infusion, but no precipitate is caused by acetic, citric, or oxalic acid. The pure alkalies produce no precipitate, but deepen the colour of the infusion to a rich claret-brown. Lime-water throws down a very copious pinkish precipitate, which is soluble in muriatic acid. Solution of sulphate of iron strikes a black colour with infusion of Ratanhy root ; that of acetate of lead, throws down a pale-brown precipitate, leaving the infusion nearly colourless and limpid ; and that of iodine, a copious fawn-coloured precipitate. Alcohol produces no effect on the infusion. Solution of isinglass separates tannin.

Ratanhy root digested in alcohol yields a deep, reddish-brown tincture, which, when evaporated, leaves a deep red, brittle resin. When this tincture is poured into water, it throws down the resin of a pink colour. In ether the tincture is less deep coloured, and when the ethereal tincture is evaporated on water, it leaves a pelicle of dark red resin on the surface, and a small quantity of extractive is diffused through the water, colouring it a light-brown. From these experiments we may conclude, that the bark of Ratanhy root contains a large proportion of tannin, some gallic acid, gum, fecula, and resin. From the effects of the mineral acids on the infusion, they may be regarded as incompatible in prescriptions with this root. Vogel states, that he found the constituents of 100 parts of the root, to be 40·00 of a pecu-



liar principle, 1·50 of mucilage, 0·50 starch, 48·00 fibrine, and 10·00 of water and loss.

*Medical properties and uses.* — Ratanhy root is powerfully astringent. It has been long esteemed in Peru as a remedy in dysentery, attended with bloody stools: as a detergent in ulceration of the gums, and a stomachic corroborant. It is also employed for fixing the teeth, when they become loosened by the receding of the gums;<sup>1</sup> and for giving a fine red colour to the gums and lips. It is powerfully styptic when applied to wounds, and on this account has been used in internal hæmorrhages, particularly hæmaturia. Alibert states that it has been used with success in France, in cases of leucorrhæa. It is little known in Great Britain as a medicine, although it has been long known to those who manufacture port wine; and large quantities of its extract are prepared solely for this purpose in South America. It is certainly likely to prove a valuable addition to the Materia Medica, in intermittents, diarrhœas, hæmorrhages, and all cases in which astringents are indicated. It has, also, been found useful in chronic rheumatism; in gastrodynia, attended by dyspepsia, head-ach and vertigo; and in all diseases of the digestive organs, in which the powers of the stomach are impaired. When there is great debility of the nervous system, it operates as powerfully and more immediately than the Cinchona bark; whilst, in cases of general asthenia, its invigorating effects are very evident. Ratanhy root may be exhibited in substance, or in the form of extract, or in infusion and decoction. The dose in substance is from gr. x. to ʒss.: of the infusion made with ʒss. of the bruised root to fʒvj. of boiling water, from fʒx. to fʒij.: and of the decoction, made with ʒij. of the bruised root, and 0j. of distilled water, from fʒj. to fʒij. On the continent it is exhibited in the form of tincture, made by digesting for twelve days ʒiij. of the powdered root with ʒij. of orange peel, ʒss. of serpentaria root, and ʒj. of saffron, in 0ij. of rectified spirit of wine. The extract is also much used.

LACTUCA. *Spec. Plant. Willd.* iii. 1523.

Cl. 19. Ord. 1. Syngenesia æqualis. *Nat. ord.* Compositæ semiflosculosæ, *Linn.* Cinchoraceæ, *Juss.*

G. 1404. *Receptacle* naked. *Calyx* imbricate, cylindrical, with a membranous margin. *Pappus* simple, stipitate. *Seed* even.

*Species* 1. *L. Sativa*. Garden Lettuce. *Blackwell*, t. 8.

*Species* 12. *L. virosa*. Strong-scented Lettuce. *Med. Bot.* 2d edit. 75. t. 31. *Smith, Flora Brit.* 819.

1. LACTUCA SATIVA.<sup>2</sup>

*Officinal.* LACTUCA, *Lond.* LACTUCÆ SATIVÆ HERBA; LACTU-

<sup>1</sup> An excellent toothpowder may be composed by mixing one part of finely powdered Ratanhy root with three parts of powdered charcoal. *T.*

<sup>2</sup> Ὠριδακίνη Theophrasti.



CARIUM, *Edin.* The herbaceous part and inspissated juice of the Garden Lettuce.

*Syn.* Laitue (*F.*), Lattich (*G.*), Lataw Gewoone salade (*Dutch*), Lattuca (*I.*), Lechuga (*S.*), Khasky (*Arab.*).

This species of lettuce is cultivated almost generally over Europe. The root is fibrous; and sends up a corymbose stem, which sometimes rises three feet in height. The leaves are roundish, obovate or cordate; shining, crisped, rugose, irregularly plaited, and of a yellowish green colour; but the plant is so well known as to require no description. When it is in flower, the slightest touch on the pedicels occasions the exudation of drops of a white, opaque, milky-looking fluid. The leaves and stem, immediately under the cuticle, contain a secreted juice, which is pellucid and colourless when in the vessels of the plant, but becomes milky when first exposed to the air, and afterwards acquires a brownish colour, resembling that of East Indian opium. This is the *Lactucarium*<sup>1</sup> of the Edinburgh college. The best method of procuring it, as first suggested by Mr. John Young, surgeon in Edinburgh, is to cut off the top of the stem, when it is in flower, about a foot above the ground, and to absorb the milky juice that exudes by means of a moist sponge, from which it can be again compressed into a proper vessel to be inspissated. But, as the cut surface soon ceases to bleed, another slice must be taken under the first, and this may be repeated as long as the fresh-cut surface will yield the juice. The process may be repeated two or three times a day.

*Qualities.* — *Lactucarium* has the colour and, in some degree, the taste and odour of opium. Distilled water dissolves the greater portion of it; and the clear solution, which is of a deep brown colour, when treated in the same manner as opium, discovers the presence of that principle detected in opium by Serturmer, and denominated Morphia; on which its narcotic property depends. It contains extractive, resin and mucilage; and Dr. John states, that caoutchouc also is one of its components.

*Medical properties and uses.* — *Lactucarium* has been proposed as a substitute for opium by Dr. Cox, of Philadelphia; but its value as a narcotic has been more lately examined by Dr. Duncan, senior, who conceives it to be particularly well adapted for allaying the cough in phthisis pulmonalis: and his opinions have been confirmed by the experience of many other respectable practitioners. It may, undoubtedly, prove useful as a soporific<sup>2</sup>, where, from peculiar idiosyncrasy, or

<sup>1</sup> This name was imposed by Dr. Duncan, sen. who first suggested its use as a narcotic. Vide *Observ. on Pulmonary Consumptions*, by A. Duncan, M. D. Appen. p. 162.

<sup>2</sup> I have always found that, when I eat lettuce to supper it acts as a soporific.



other causes, opium cannot be taken. The dose is from gr. i. to grs. vj. in the form of a pill: or, of a tincture made with one ounce of lactucarium and a pint of diluted alcohol, from ten to sixty drops may be taken.

Officinal preparations. — *Succus spissatus Lactucæ Sativæ*. E.

## 2. LACTUCA VIROSA.<sup>1</sup>

Officinal. — LACTUCÆ VIROSÆ HERBA, *Edin.* Strong-scented Lettuce leaves.

*Syn.* Laitue vireuse (*F.*), Wilder lattich (*G.*), Stinkende salade (*Dutch*), Stinkende laktuk (*Dan.*), Lattuca Salvatica (*I.*), Alface brava (*Portug.*).

This is an indigenous biennial plant, found growing on the banks of ditches, and borders of fields, flowering in July and August. The stalk rises about three feet in height, erect, slender, prickly below, smooth above, round, paniced, and not very leafy. The leaves are rather smooth and toothed, the lower ones numerous, obovate, undivided; those of the stem smaller, often lobed, amplexicaule, with the midrib beset with prickles on the under side. The bractes are cordate and pointed. The flowers are numerous, compound, of a sulphur-yellow colour, on short peduncles, furnished with small scaly leaves, and one at the base of each. The calyx is oblong, and composed of small lanceolate scales; and the corolla consists of florets scarcely longer than the calyx. The seeds are elliptical, compressed, striated, black, and furnished with a stipitate, scabrous pappus.

The leaves and stem contain a white opaque juice under the cuticle, that abounds more copiously when the plant is in flower; at which time, therefore, they should be gathered, and the juice immediately expressed.

*Qualities.* — The odour of the leaves is heavy and foetid, resembling in some degree that of opium; their taste is bitter and acrid: qualities depending on their milky juice. When triturated with water, the solution never clears; and the morphia, which is separated by treating it like opium, cannot be freed from the mucus of the extract.

*Medical properties and uses.* — The expressed juice is narcotic and diuretic (see *Preparations* and *Compositions*.) The leaves themselves are not used.

Officinal preparations. — *Succus spissatus Lactucæ virosæ*, E.

LAURUS. *Spec. Plant. Willd.* ii. 477.

Cl. 9. Ord. 1. Enneandria Monogynia. *Nat. ord.* Oleraceæ, *Linn.* Lauri, *Juss.*

G. 798. *Calyx* none. *Corolla* calycine, six-parted. *Nectary* of three two-bristled glands, surrounding the germen. *Filaments* interior, glanduliferous. *Drupe* one-seeded.

Sp. 1. *L. Cinnamomum*. The Cinnamon Tree. *Mat. Med.* 2d edit. 670. t. 223. *Percival's Account of Ceylon*, 4to. 346—350.

<sup>1</sup> Οπίδαξ ἀγρία Dioscoridis.



*Sp. 2. L. Cassia.* The Cassia Tree. *Carua*, *Rheede Hort. Malabar*, i. p. 107. t. 59. *Herb. Amb.* ii. 65. t. 14.

*Sp. 3. L. Camphora.* The Camphor Laurel. *Med. Bot.* 2d edit. 681. t. 236. Michaux, *North American Sylva*, vol. i. pl. 83.

*Sp. 10. L. nobilis.* Common Sweet Bay. *Med. Bot.* 2d edit. 678. t. 235.

*Sp. 34. L. Sassafras.* Sassafras Laurel. *Med. Bot.* 2d edit. t. 234. *North American Sylva*, vol. ii. fol. 61.

#### 1. LAURUS CINNAMOMUM.<sup>1</sup>

*Officinal.* — CINNAMOMI CORTEX. CINNAMOMI OLEUM, *Lond.*

LAURI CINNAMOMI CORTEX, *Edin.* CINNAMOMUM; CORTEX, OLEUM ESSENTIALE, *Dub.* Cinnamon, and Oil of Cinnamon.

*Syn.* Canelle (*F.*), Kanohl (*G.*), Kaneel (*D.*), Caneel (*Dan.*), Ackta Canel (*Swed.*), Canella (*I.*), Canela (*S.*), Dárchini (*H.*), Dárcasita (*San.*), Karruwa Puttay (*Tam.*), Canella vulgar (*Port.*), Kayu-manis (*Malay*), Darsini (*Arab.*).

The cinnamon tree is a native of Ceylon<sup>2</sup>, growing in great abundance in many parts of the island, particularly near Colombo. It also grows plentifully in Malabar, Cochin China, Sumatra, and the eastern islands. It has been cultivated in the Brazils, the Mauritius, and other places. France is partly supplied from Guiana. The soil in which it thrives best is nearly pure quartz sand. That of the cinnamon garden, near Colombo, was found by Dr. J. Davy to consist of 98.5 of silicious sand, and 1.0 only of vegetable matter in 100 parts. "The garden is nearly on a level with the lake of Colombo; its situation is sheltered; the climate is remarkably damp; showers are frequent, and the temperature is high and uncommonly equable."<sup>3</sup> The tree seldom rises above thirty feet in height; has a slender branching trunk covered with a brown ash-coloured cuticle, and from the root spring a number of suckers, which form a bush round the trunk. The leaves, which stand in opposite pairs in short, slightly channelled petioles, are from six to nine inches in length, oblong, pointed, trinerved; when young reddish, but afterwards of a bright-green colour; and have a spicy odour, and a hot taste when rubbed and chewed. The flowers, which appear in January, are white and inodorous, in axillary and terminal panicles. The petals are oval, pointed, concave, and spreading, longer than the filaments, which are in ternaries, flattish, erect, and

<sup>1</sup> *Κιννάμωμον* Dioscoridis. The Malays call cinnamon kayu-manis, which is sometimes pronounced as if it were written kaina-manis, which Mr. Marshal supposes to have been the original of the ancient Greek name kinnamomon; which, however, Scaliger (*Not: in Garz.*) derives from the Hebrew *kinamon*.

<sup>2</sup> Notwithstanding the jealousy of the Dutch, the cinnamon tree, long before the British obtained possession of Ceylon, was cultivated at the Isle of France, in several parts of India, Jamaica, and some other of the West India islands. Mr. Miller first cultivated it in this country in 1768; and a plant of it has regularly flowered and ripened seed in the hot-house of the Bishop of Winchester at Farnham for several years past.

<sup>3</sup> *Davy's Account of the Interior of Ceylon*, 4to. p. 39.



the three innermost glanduliferous at the base: and the anthers are double. The fruit is an oval berry, resembling a small acorn, but with the apex depressed, and the pulp fleshy, with a terebinthine odour; and a taste not unlike that of the juniper berry.

There are several varieties of the Cinnamon tree known at Ceylon. Seba enumerates ten, but the four following only are said to be barked: 1. Honey, sharp, sweet, or royal Cinnamon, (*Rase Curundu*, in the language of the natives,) which is the finest sort; 2. Snake Cinnamon, (*Nai Curundu*,) similar to the first; 3. Camphorated Cinnamon, (*Capura Curundu*,) so named from its having the odour of camphor, and the root yielding camphor by distillation; and 4. Bitter astringent Cinnamon, (*Cahatte Curundu*,) which has smaller leaves than the former varieties.<sup>1</sup> The trees that grow in the valleys, in a white sandy soil, are fit to be barked when four or five years old, but those in a wet soil, or in shady places, require to be seven or eight years of age. The bark is good for nothing if the tree be older than eighteen years. The tree was formerly propagated by a species of pigeon, that ate the fruit, and voided the seed; but since Falck, one of the Dutch governors, about the middle of the eighteenth century, raised it from berries sown in his garden, it has been regularly cultivated.

The barking, particularly in the vicinity of Negombo and Matura, commences early in May, and continues until late in October. The *chaliahs*, or people who perform it, are under native officers called cinnamon *moodeliars*<sup>2</sup>, who are answerable for the quantity barked. Branches of three years old are selected, and lopped off with a pruning knife, or bill hook called a *ketta*. To remove the bark, a longitudinal incision is made through it on both sides of the shoot, so that it can be gradually loosened, and taken off entire, forming hollow cylinders. The bark in this state, tied up into bundles, is allowed to remain for twenty-four hours, by which a fermentation is produced that facilitates the separation of the epidermis, which with the green pulpy matter under it is carefully scraped off. The bark now soon dries, contracts, and assumes the quilled form, after which the smaller pieces are put within the larger.<sup>3</sup>

<sup>1</sup> The other sorts mentioned by Seba are: Sandy Cinnamon, *Welle Coronde*, which feels gritty when chewed; Glutinous Cinnamon, *Sewel Coronde*; Insipid and inodorous Cinnamon, *Nicke Coronde*; Drum Cinnamon, *Dawel Coronde*, so named because the natives make drums of the wood; Prickly Cinnamon, *Catte Coronde*; flowering Cinnamon, *Mael Coronde*, the tree being always in bloom; and Three-leaved Cinnamon, *Toupat Coronde*. *Phil. Trans.* xxxvi. 97—105.

<sup>2</sup> Under the Moodeliars are inferior officers, named *Mohandrums* and *Aratchays*; and in 1811 General Maitland appointed a superior, who is named *Maha-Moodeliar*.

<sup>3</sup> Prior to the 15th century, all the Cinnamon used in Europe was imported by the Arabs, and passed through the hands of the Venetians; after this the Portuguese



The Cinnamon, when dry, is tied up in bundles of 30 lbs. weight, and carried to the government store-house, where the quality is determined by inspection of the bundles. It was formerly chewed, and the surgeons who used to be thus employed, had their mouths so excoriated, as to be unable to continue the process longer than two days together: but tasting is now seldom had recourse to.

Cinnamon is brought home in bags or bales, weighing 92 lbs. each<sup>1</sup>: and, in stowing it, black pepper is mixed with the bales to preserve the Cinnamon. According to Mr. Marshall's account, the annual quantity of Cinnamon sold at the East India Company's sales, taken on an average of the last eight years, up to 1810, is 318,258 lbs.; at an average price of six shillings per pound.<sup>2</sup> But much Cinnamon, of an inferior kind, reaches Europe through private merchants, particularly from China.

The *oil of Cinnamon* is prepared by macerating the<sup>3</sup> bark in sea-water for two days, then distilling with a slow fire, and separating the oil from the water with which it comes over. It is generally adulterated with alcohol or expressed oil. Eleven pounds of Cinnamon are required to procure one ounce of the oil. Cinnamon is sometimes intermixed with Cinnamon from which the oil has been drawn, and with Cassia. The former is detected by the weakness of its odour and taste; and the latter by its thickness, smooth fracture, and remarkably slimy taste.

*Qualities.*—Cinnamon has a very pleasant, fragrant odour, and a pungent, aromatic, sweetish taste; but, when it is very hot, without sweetness, and leaves a mawkish taste in the mouth, it is of an inferior quality. The best is rather pliable, but breaks in splinters: is as thin as paper, and of a light yellowish colour: thickness and a dark or brown colour are marks of inferiority. What is called Chinese Cinnamon is darker coloured, rougher, denser and breaks shorter. The taste is harsher, more pungent and ligneous, without the sweetness of the Ceylon Cinnamon. These qualities depend

became the sole importers, and continued to be so until 1645, when their trade was divided with the Dutch, who obtained entire possession of it in 1658; and were the principal cinnamon merchants until 1796, when Ceylon fell into the power of the British.

<sup>1</sup> The bags are made of cloth of the cocoa-nut bark.

<sup>2</sup> *Annals of Phil.* vol. x. p. 358.

<sup>3</sup> The bark of the roots yields an aromatic essential oil, denominated oil of Camphor, which is used in Ceylon as a rubefacient in painful affections of the joints, and in sprains. Oil of Cinnamon has always been dear. Alston says, that he saw the finest oriental sold in Holland, in 1721, at 50 stivers the drachm, "or the common price of the pound of Cinnamon at Amsterdam." *Lect. on Mat. Med.* vol. ii. p. 7.



on the *essential oil*, which may be separated by macerating the bark in alcohol, and distilling the tincture; in which process the oil does not rise with the spirit, but remains in the retort. From 3xvj. of the bark, Neumann obtained only two scruples and a half of oil.<sup>1</sup> It has a pale-gold colour, is heavier than water, perfectly soluble in alcohol, and has the odour and taste of the Cinnamon concentrated. But both a heavy and a light oil is obtained from Cinnamon, 80 lbs. yielding about five ounces of the former, and two of the latter.

*Medical properties and uses.* — Cinnamon *bark* is astringent, cordial and tonic. Hence it is found to be efficacious in alvine fluxes, proceeding from a weakened and languid state of the intestines, dyspepsia, and chronic, nervous debility; and, when given in the form of watery infusion, it removes nausea, and checks vomiting. But the principal use of Cinnamon is to cover the nauseous state of other remedies. The *oil* is a powerful stimulant and stomachic; and is used as such in cramps of the stomach, flatulent colic, hiccough, and nervous languors. It is, sometimes, inserted into the hollow of a decayed tooth to allay the pain of tooth-ach.

The dose of the bark in powder is from grs. x. to ℥j.; that of the oil from m.j. to m.ij. on a lump of sugar, or rubbed down with yolk of egg and syrup, so as to form a mixture with water.

Officinal Preparations. *Aqua Cinnamomi*, L. E. D. *Spiritus Cinnamomi*, L. E. D. *Tinct. Cinnamomi*, L. E. D. *Tinct. Cinnamomi comp.*, L. *Pulvis Cinnamomi compositus*, L. E.

## 2. LAURUS CASSIA.<sup>2</sup>

*Officinal.* LAURI CASSIÆ CORTEX; FLOS NONDUM EXPLICITUS, *Edin. Dub.* The bark and flower buds of the Cassia tree.

*Syn. Of the bark:* Casse (*F.*), Casia (*G.*), Cannellina (*I.*), Louranga puttay (*Tam.*), Kayu manis (*Jav.*), Kayu legi (*Malay*), Sing Rowla (*Nepaul*), Seleckheh (*Arab.*), Tej (*H.*), Twacha (*San.*). *Of the buds:* Fleur de la Cannelle (*F.*), Cassia Bloemen (*D.*), Tejpatka konpul (*H.*), Sirnāgāpoo (*Tam.*).

The Cassia tree is a native of Malabar, Ceylon, Sumatra, and Java, and has been generally supposed to be rather a variety of the Cinnamon than a distinct species of *Laurus*; although Marsden's description of the plant<sup>3</sup>, and Gærtner's of the fruit<sup>4</sup>, afford some reason for thinking that it is properly marked as a different species. It rises fifty feet in height, and gives out, almost from the bottom, large, spreading, horizontal

<sup>1</sup> Neumann's *Chemistry*, ii. 188.

<sup>2</sup> *Κασσία* Dioscoridis. It is this Dawul Kurundu of the Cingalese, the Cannella Matto of the Portuguese, and the Wilde Canule of the Dutch. *Marshall, Phil. Trans.* 1817.

<sup>3</sup> *History of Sumatra*, 125.

<sup>4</sup> *De Fructibus*, ii. 69. t. 92. If Gærtner be correct, the fruit of the Cassia is depicted instead of that of the Cinnamon, in the plate of the Cinnamon plant, in Woodville's *Medical Botany*.



branches: the leaves are from four to six inches long, elliptical, narrow, pointed, entire, smooth, longitudinally nerved, of a deep green colour above, and pale grey beneath: the flowers are in axillary clusters, six together on slender flower-stalks: they are monopetalous, white, small, and divided into six stellated points: the fruit is an ovate, oblong, black berry with a mucronated apex, standing in a bell-shaped, coriaceous, angled, unequally five or six-toothed calyx; it contains a bitterish pulp, and when dried is insipid and inodorous.

Like the Cinnamon, those trees which grow in a dry soil and high exposed situation yield a superior bark to those in a moist soil and shaded spot. The larger branches and the trunk are said to be the parts of the tree barked: and the cuticle only appears to be scraped off, the cellular integument being left, which, as the bark is taken from the larger branches, is thick, spongy, and full of a slimy mucus. This plant is never decorticated at Ceylon. According to Mr. Marshall<sup>1</sup>, the *Cassia bud* of commerce is the hexangular fleshy receptacle of the seed of the *L. Cinnamomum*, and not the *L. Cassia*, as supposed by the Dublin College. They are not prepared at Ceylon, but come chiefly from China, through Calcutta, Madras, and Bombay.

Cassia is imported in chests, half chests, and occasionally in quarter chests.

*Qualities.* — The odour of Cassia bark is similar to that of Cinnamon, but fainter; and the taste is more pungent, but less agreeable; appearing slimy when much chewed. It is of a cinnamon colour, in pieces more or less quilled, about one-tenth of an inch in thickness; which break with a short, close fracture, and show it to consist of two parts, the inner darker and of a fine texture, and the outer paler and somewhat spongy. When these are separated, the inner part has all the sensible qualities of real Cinnamon, only more pungency, whilst the outer has scarcely either flavour or taste: and I am of opinion that the allowing this cellular integument, from which the Cinnamon is freed, to remain in the Cassia, constitutes the chief cause of the difference between these two barks.<sup>2</sup> *Cassia buds* have the same odour and taste as the Cinnamon bark. They are of a brown colour, and resemble a nail, with a round head, surrounded with the hexangular calyx, which gradually terminates in a point. Both the bark and buds yield in distillation with water an essential oil, similar to that of cinnamon, on which their qualities depend.

---

<sup>1</sup> *Annales of Phil.* vol. x. p. 245.

<sup>2</sup> The rejected or third sort of Cinnamon, prepared in Ceylon, has been imported into England, and sold as Cassia.



*Medical properties and uses.*—Cassia bark and buds are stimulant cordials; and are used in the same cases, and in the same manner as Cinnamon bark.

Officinal preparation. *Aqua Lauri Cassiæ distillata*, E.

### 3. LAURUS CAMPHORA.

Officinal. CAMPHORA, *Lond. Edin.* CAMPHORA; RESINA,<sup>1</sup> *Dub.* Camphor.

*Syn.* Camphre (F.), Kampfer (G.), Canfora (I.), Alcanfor (S.), Cafoor (Arab.), Kaafur (Malay), Cafur (H.), Curfura (San.), Carpoorum (Tam.).

The species of laurel here designated, is a native of North America, China, and Japan. It yields Camphor; but I have already stated, that the camphor which comes from Sumatra, that is, the greater part of what is brought to Europe, is the produce of the *Dryobalanus Camphora*, a tree belonging to a different genus altogether from the laurel. The Camphor laurel<sup>2</sup> rises to a considerable height, is much branched, and covered with a smooth greenish bark. The leaves, which stand on long footstalks, are acuminate at both extremities, entire, smooth, ribbed, of a pale yellowish-green colour on the upper surface, alternate on the under glaucous, and two or three inches in length. The flowers are small, white, pedicellated, in roundish close clusters, which terminate long, axillary peduncles. The corolla consists of six, small, ovate, unequal petals, enclosing a tuberculated, bristled nectary, which surrounds the germen; the filaments are shorter than the corolla, and support round anthers; the germen is roundish, with a simple style and obtuse stigma. The fruit, which resembles that of the Cinnamon<sup>3</sup>, is a red, oval berry seated in a small, yellow cup supported in pairs on a long peduncle.

The roots, wood, and leaves of this tree have a very strong odour of Camphor; and, from the roots and smaller branches, it is obtained by distillation. They are cut into chips, which are suspended in a net, within a kind of still or iron pot, the bottom of which is covered with water, and an earthen head fitted to it: heat is then applied, and the steam of the boiling water, penetrating the contents of the net, elevates the camphor into the capital, where it concretes on rice straws with

<sup>1</sup> This is an error of the Dublin College, chemists being now agreed that Camphor is not a resin, but a proximate vegetable principle, *sui generis*.

<sup>2</sup> Specimens of it are common in our hot-houses; but they rarely flower.

<sup>3</sup> Camphor is not the production of those plants merely from which that known in commerce is obtained, but has also been procured from the roots of the Cinnamon, Cassia, and Sassafras laurels; from those of Galangale, Zedoary, Ginger; and from Cardamom seeds and long Pepper. The essential oils of Lavender, Sage, Thyme, Peppermint, Rosemary, and several other labiated plants, yield it: and an artificial Camphor is prepared by passing muriatic acid gas through oil of turpentine. The varieties of Camphor thus obtained, however, differ in some respects from common Camphor. The *Shorea robusta* of Roxburgh (*Car. Pl.* vol. iii. p. 212.) is said to yield a very fine species of camphor.



which this part of the apparatus is lined.<sup>1</sup> But, as we have already stated, the greater part of the Camphor brought to Europe is obtained in Sumatra, where the trees which yield it are cut and split, and the Camphor which is found concentered in the heart of them is picked out, and washed in a ley of soap. It is imported into this country in chests, drums, and casks; and is in small, granular, friable masses, of a dirty white, or greyish colour, very much resembling in appearance half-refined sugar. It often contains earth and other impurities.<sup>2</sup>

Camphor was introduced into Europe by the Arabians. Formerly all the crude Camphor brought to Europe was purified by the Venetians, and afterwards by the Dutch, who kept the art secret; but it is now practised to a considerable extent in this country. It is sublimed in glass vessels, after being mixed with one-twentieth of its weight of quick lime; and is afterwards fused, either “by increasing the heat suddenly when the sublimation is almost ended, without transferring the Camphor to different vessels, or by melting the sublimed flowers in a vessel for that purpose.”<sup>3</sup> Thus refined, it is in large round cakes, about two or three inches thick, concave on one side, convex on the other, and generally perforated.

*Qualities.*—Pure Camphor has a strong, peculiar, fragrant, penetrating odour; and a bitter, pungent, aromatic taste. It is white, transparent, unctuous to the touch, and friable, breaking with a shining, foliated or tabular fracture, which displays crystalline texture: and although brittle, yet it is, also, in some degree ductile, and therefore not easily pulverized. It swims on water, its specific gravity being 0.9887:<sup>4</sup> and is so volatile, that if it be not kept in well-stopt vessels, it loses a very considerable proportion of its bulk and weight by evaporation, particularly in a moist atmosphere. It melts at a temperature of 88°, boils at 400°, and sublimes in close vessels, crystallizing unchanged in hexagonal plates.<sup>5</sup> It is readily ignited, and burns with a brilliant flame, giving out

<sup>1</sup> According to Kæmpfer, the process is carried on chiefly by the peasants of Satzuma. *Amæn.* 779.

<sup>2</sup> Zea describes a variety of Camphor which is procured in South America, from a tree, the botanical characters of which are not yet known; but which is termed *caratte* by the natives. The Camphor exudes from the bark in the form of tears.

<sup>3</sup> *Aikin's Dictionary of Chemistry*, art. *Camphor*. Professor Robison, who saw the process as it was conducted in Holland, says, that the Camphor is in a liquid state in the subliming vessel. *Black's Lectures*, ii. 351.

<sup>4</sup> Brisson.

<sup>5</sup> Mr. W. Phillips says, that the native crystal of Camphor is a flat octohedron. *Paris, Pharmacologia*, 5th edit. vol. ii. p. 117.



much smoke. When triturated with water, very little is dissolved<sup>1</sup>, although it communicates to the water its odour and pungency; but the addition of carbonic acid gas augments very much the solvent power of water over Camphor. Alcohol, ether, the fixed and volatile oils, the sulphuric and nitric acids a little diluted, and the muriatic, the strong acetic, and the fluoric acids, dissolve Camphor, which is again separated unaltered from these solutions by the addition of water. Concentrated sulphuric acid decomposes it, forming artificial tannin; and by repeatedly distilling it with nitric acid, it is converted into camphoric acid. Alkalies exert scarcely any action on Camphor: but it unites with, and converts into a soft tenacious mass, the hardest resinous substances. Camphor, when mixed with clay, and distilled in close vessels, is decomposed, and resolved into a volatile oil and charcoal; in the proportion, according to Buillon la Grange, of 45·856 of the former, and 30·571 of the latter: thence, as a chemical compound, it appears to differ from the essential oils, only in containing a larger proportion of carbon. Dr. Thomson says its ultimate components are carbon 6·375, hydrogen 1·250, oxygen 1·000.

*Medical properties and uses.* — Camphor is stimulant, narcotic, and diaphoretic, but its stimulant powers are very transitory, and followed by sedative effects. It acts chiefly on the nervous system. The Arabians appear to have first used Camphor as a medicine<sup>2</sup>, and by them it was regarded as refrigerant; an opinion which, even in more recent times, has been the subject of much controversy. In moderate doses it operates as a cordial, increasing the heat of the body, and exhilarating; besides softening, and rendering fuller the pulse, and promoting diaphoresis: in larger doses, it allays irritation and spasm, abates pain, and induces sleep. But in immoderate doses Camphor produces vomiting, vertigo, delirium, convulsions, and other deleterious effects.

As a stimulant, Camphor is beneficially used in all fevers of the typhoid kind, cynanche maligna, malignant measles, confluent small-pox, and as an adjunct to bark and opium to check the progress of gangrene; and in spasmodic affections, as, hysteria, epilepsy, chorea, asthma, and painful menstruation. Its narcotic and anodyne effects being produced with very little increase of pulse, it has been successfully employed for allaying pain and irritation even in some inflammatory diseases, as, pneumonia, acute rheumatism, gonorrhœa, small-

---

<sup>1</sup> Cadet asserts, that one French pint of water dissolves about sixteen grains of camphor, which are again precipitated by pure potass. *Ann. de Chimie*, lxii. 132.

<sup>2</sup> They called it *Canfur* as well as *Caofoor*. *Clusius Exot.* 245, quoted by *Alston*.



pox, when attended with convulsions, gout, and in the delirium of mania, and inflammatory fevers. But in these cases its use should be preceded by evacuations; and the remedy itself combined with nitre, or antimonials: and in maniacal cases with opium.<sup>1</sup> Camphor is also given internally to obviate the irritating effects of some other medicines, as, meze-reon, cantharides, the saline preparations of mercury, and drastic purgatives; to correct the nauseating property and prevent the irritation which squill is apt to produce on the coats of the bladder.

Camphor may be administered in the solid form; but as in this state it is apt to occasion nausea, it is generally ordered in a state of minute division, suspended in fluids by means of mucilage or the yolk of eggs; sometimes by magnesia, which, assisting its division, and rendering it smooth as starch, admits of its combination with acids; and as several of the gum resins, when triturated with it, form a soft, uniform, soluble mass, they also may be employed for diffusing it in water.<sup>2</sup> It may be advantageously united with ammonia, aromatics, opium, bark, and other tonics, in low fevers and diseases of debility; with calomel, antimonials, digitalis, and neutral salts, in inflammatory diseases; with the foetid gums and other narcotics, in spasms and convulsive affections; and with squill and ipecacuanha, in pulmonary complaints.

As a local anodyne, Camphor is used in frictions, dissolved in oils, alcohol, or acetic acid, for allaying rheumatic and muscular pains; and with the addition of laudanum we have found it of great efficacy when rubbed on the abdomen, in flatulent colic, dysentery, and inflammations of the viscera. In collyria it is useful in ophthalmia; and dissolved in oil, as an injection, in ardor urinæ; and as an enema in the tenesmus occasioned by ascarides, or other irritations of the rectum.<sup>3</sup> A pill of Camphor and opium, or a solution of Camphor in oil of turpentine, put into the hollow of a carious tooth, affords almost immediate relief in tooth-ach. Twenty or thirty grains of Camphor added to a common poultice and applied to the perinæum allays chordee in gonorrhœa.

The dose of Camphor is from grs. ij. to ℥j., repeated at shorter or longer intervals according to the extent of the dose. It may be rubbed up with mucilage and almond emulsion, so

<sup>1</sup> In some experiments upon dogs with Camphor, M. Defermon found that contraction of the spleen takes place, producing a rugose appearance of its surface, and a movement throughout the whole viscus.

<sup>2</sup> *Murray's System of Mat. Med. and Pharm.* ii. 157.

<sup>3</sup> In some constitutions it must be exhibited in this form with caution; "two scruples of it given to a woman in a glyster, proved so irritating as to bring on pains resembling those of labour." *Heberden, Med. Trans.* vol. i. p. 473.



as to suspend it in water; and this form is preferable to that of pills or bolus. The bad effects of an overdose are most effectually obviated by opium.

Officinal preparations. *Mistura Camphoræ*, L. D. *Emulsio camphorata*, E. *Spiritus Camphoræ*, L. E. D. *Tinctura Camphoræ composita*, L. E. D. *Acidum acetosum camphoratum*, E. D. *Linimentum Camphoræ*, L. E. D. *Linimentum Camphoræ comp.*, L. *Linimentum Saponis*, L. E. D.

#### 4. LAURUS NOBILIS.<sup>1</sup>

*Officinal.* LAURI BACCÆ ET FOLIA, *Lond.* LAURI NOBILIS FOLIA; BACCÆ; OLEUM FIXUM, *Edin.* Laurel berries and leaves, and the fixed oil of the berries.<sup>2</sup>

*Syn.* Baies de Laurier (*F.*), Lorbeeren (*G.*), Bobek drzewo (*Pol.*), Bacchi di Lauro Riccio (*I.*), Bayas (*S.*).

This tree is a native of Italy and the south of Europe; but is cultivated in this country, and is not uncommon in our gardens, flowering in April and May. It is a handsome ever-green; and although it appears as a shrub in England, yet in its native soil and climate it rises twenty or thirty feet in height. The bark is smooth, and of a green olive colour. The leaves are lanceolate, about three inches long, and an inch and a half broad, on short petioles, smooth, entire, veined, often waved at the margin, of a firm texture, and a deep-green colour. The flowers are male and female on different plants, in short racemes, and of an herbaceous or yellowish-white colour. The corolla is divided, in both descriptions of flowers, into four oval segments. The berry is superior, of an oval shape, fleshy, and of a dark purple almost black colour.

Laurel berries, and the oil which is obtained by boiling the berries in water, are imported from the Streights. The simple expressed oil is insipid.

*Qualities.* — Both the *leaves* and the *berries* have a sweet, fragrant odour, and an aromatic, astringent taste; and the *oil*, which is of a yellowish green colour, has a stronger but similar odour and taste. Water distilled from the leaves shews traces of prussic acid; and it is probably on this component that their medicinal and poisonous properties depend.

*Medical properties and uses.* — Bay leaves, berries, and oil, are narcotic and carminative. They were formerly given in coughs, flatulent colic, hysteria, and obstructed menstruation, but their internal use is now abandoned; and, as an external application, they are generally compounded with other stimulants. Having found great advantage from the use of prussic acid, largely diluted, as a local application in impetigo, I have lately employed infusions of bay berries, with nearly the same beneficial results.

<sup>1</sup> Δαφνῆ Dioscoridis.

<sup>2</sup> Δαφνιδες Dioscoridis.



## 5. LAURUS SASSAFRAS.

*Officinal.* SASSAFRAS; LIGNUM ET RADIX, *Lond.* LAURI SASSAFRAS LIGNUM, RADIX, *Edin.* SASSAFRAS; LIGNUM, CORTEX, RADIX, *Dub.* The wood, root, and bark of Sassafras.

*Syn.* Sassafras (*F.*), Sassafras (*G.*), Sassofrasso (*I.*).

This species of laurel is a native of the southern parts of North America and Cochin China. It is cultivated in Jamaica; and withstands the cold of our climate so as to be frequently reared in gardens as an ornamental shrub. The flowers appear in May and June. In America the plant rises twenty, thirty, and even fifty in height, with the trunk about twelve inches in diameter, covered with a rough, furrowed, grey bark, and brownish towards the top. The leaves are of different shapes and sizes; some being oval, entire, and about four inches long and three broad; and others, the most numerous, lobed, about six inches long, and nearly as broad; they are of a lucid green colour, downy on the under surface, petiolate, and alternate. The flowers, which appear in spring immediately under the leaves before they begin to be evolved, are small, and produced in pendant panicles; and at the base of the pedicels are linear bractes. The corolla is divided into six, narrow, convex, yellowish, or greenish white segments, inclosing, in the male flowers, nine stamens supporting yellow anthers. The hermaphrodite flowers, which are on a separate plant, have six stamens only, and a simple style. The berry, which is oval, and, when ripe, of a deep-blue colour, is contained in a small red cup, supported on a peduncle from one to two inches in length.

The sassafras laurel was discovered by the Spaniards, immediately after their conquest of Florida, in 1538, under Ferdinand de Soto, and termed by them cinnamon wood, on account of its odour.<sup>1</sup> It is imported in what are termed logs; which are straight and branched pieces, light, of a spongy texture, and covered with the thick, rough bark. The bark is separated, and the wood then cut into chips, as is also the root.

*Qualities.*—Sassafras wood, root, and bark have a fragrant odour, and a sweetish aromatic taste. The wood is of a brownish-white colour; and the bark ferrugineous within, spongy, and divisible into layers. Their sensible qualities and virtues depend on an essential oil, which can be obtained separate by distilling the chips or the bark with water. It is very fragrant, hot, and penetrating to the taste, of a pale-yellow colour, and heavier than water. Water extracts the virtues of sassafras partially; alcohol completely; and when

---

<sup>1</sup> *Savary's Dictionary*, ii. l. 487.



the tincture is evaporated, it leaves an extract which contains the whole virtue of the plant.

*Medical properties and uses.* — Sassafras is a stimulating diaphoretic and diuretic. It has been employed in cases of scurvy, chronic rheumatism, gout, and in cutaneous affections. It was once regarded as serviceable in lues venerea, but it has no pretension whatever to the character of an antisymphilitic. Its effects are very uncertain; and even the diaphoresis which it is supposed to occasion may rather be ascribed to the guaiac, and other more powerful medicines, with which it is generally combined. An infusion of the chips taken as tea, is a common domestic remedy in the above complaints; but I know instances in which it has been taken regularly every morning for a couple of years without any perceptible benefit. The infusion, however, is the best form of giving the remedy, as much of the oil is dissipated in making the decoction.<sup>1</sup> The oil is sometimes given with the same intentions as the infusion.<sup>2</sup>

Officinal preparations. *Oleum Sassafras*, L. E. D. *Decoctum Sarsaparillæ compositum*, L. D. *Decoctum Guiaici*, L. E. D. *Aqua Calcis comp.*, D.

LAVANDULA. *Spec. Plant. Willd.* iii. 60.

Cl. 14. Ord. 1. Didynamia Gynospermia. Nat. ord. Verticillatæ.

G. 1099. *Calyx* ovate, somewhat toothed, supported by a bracte.

*Corolla* resupine. *Stamens* within the tube.

*Species* 1. L. *Spica*. Lavender. *Med. Bot.* 2d edit. 221. t. 114.

*Officinal.* LAVANDULÆ FLORES<sup>3</sup>, *Lond. Dub.* LAVANDULÆ SPICÆ FLORES, *Edin.* The flowers of Lavender.

*Syn.* Lavande (F.), Lavendelblumen (G. Dutch, Dan. Swed.), Lawanda (Pol.), Lavanda (I.), Alhuzema (S.), Alfazema (Port.).

This plant is a perennial, a native of the south of Europe, but commonly cultivated in our gardens<sup>4</sup>, flowering from June to September. It is a much-branched shrub, rising in its proper soil often six feet in height; the woody part of the stem being covered with a rough brown bark, while that of the shoots, which are four-cornered, is of a pale glaucous colour. The leaves of the most common variety are glaucous, narrow, nearly linear, and entire, the lower petiolate, and

<sup>1</sup> Dr. Paris (*Pharmacologia*) has given the following formula, as that by which much of the nostrum called Godfrey's Cordial is prepared: "Infuse ʒix. of Sassafras, and of the seeds of Coriander, Carraway, and Anise, of each ʒj., in six pints of water; simmer the mixture until it is reduced to four pints; then add lbvj. of Treacle, and boil the whole for a few minutes: when it is cold, add fʒiij. of tincture of opium.

<sup>2</sup> In Virginia a beer is made by boiling the young shoots of the Sassafras in water, adding molasses to the decoction and fermenting it.

<sup>3</sup> Ἰφύων Theophrasti.

<sup>4</sup> It was cultivated in England so early as 1568, according to Turner.



the upper ones sessile. The flowers are produced on the young shoots, in terminal spikes, which consist of interrupted whorls. The corolla is blue, tubular, and labiate, the upper lip larger and bifid, the lower divided into three segments. The filaments are within the tube, and support small simple anthers; the style, which is slender, and crowned with a bilobed stigma, rises from the centre of four naked seeds at the bottom of the tube.

There are two other varieties <sup>1</sup> of this species; but they are more rare, and do not differ in their sensible and medicinal qualities. The flowers are cut in dry weather, when they begin to blow.

*Qualities.* — Lavender flowers have an agreeable fragrant odour, and warm bitterish taste. Alcohol extracts their virtues completely, and elevates in distillation all their odorous parts; water acts less completely. The oil, however, on which their virtues depend, is obtained separate in distillation with water; in the proportion, according to Lewis <sup>2</sup>, of one ounce of oil from sixty ounces of the flowers.

*Medical properties and uses.* — Lavender is stimulant, and tonic. The oil extracted by alcohol enters into several compositions. The dried leaves in powder were used formerly as a sternutatory; but they are now neglected.

Officinal preparations. *Oleum Lavandulæ*, L. E. D. *Spiritus Lavandulæ*, L. E. D. *Spiritus Lavandulæ compositus*, L. E. D.

LEONTODON. *Spec. Plant. Willd.* iii. 1544.

Cl. 19. Ord. 1. Syngenesia Æqualis. *Nat. ord.* Compositæ Semiflosculosi, *Linn.* Cichoraceæ, *Juss.*

G. 1407. Receptacle naked. Calyx double. Pappus stipitate, hairy.

*Species* 1. *L. Taraxacum*. <sup>3</sup> Dandelion. *Med. Bot.* 2d edit. 39. t. 16. *Smith Flor. Brit.* 822. *Eng. Bot.* 510.

*Officinal.* TARAXACI RADIX, *Lond.* LEONTODI TARAXACI HERBA, RADIX, *Edin.* TARAXACUM; (DENS LEONIS) RADIX FOLIA, *Dub.* The root and leaves of common Dandelion.

*Syn.* Dent de Lion; Pissenlit (*F.*), Lowenzahn wurzel (*G.*), Paapenkriud (*Dutch*), Lövetand (*Dan.*), Lejentand (*Swed.*), Papawa ziele (*Pol.*), Tarassaco (*I.*), Cardillos tagarnina (*S.*), Dente de Leão (*Port.*).

This is one of our most common indigenous plants, flowering from April to September. The root is fusiform, and externally of a dark colour. The leaves are all radical, in general runcinate, but in very moist situations nearly entire<sup>4</sup>, toothed, smooth, and of a pleasant green colour. The flower-stem is an erect, one-flowered simple scape, naked, smooth, fistulous, fragile, and abounding with a milky bitter juice.

<sup>1</sup> β *L. angustifolia* flore albo. γ *L. latifolia*. <sup>2</sup> *Mat. Med.* 371.

<sup>3</sup> Ἀφάκη Græcorum.

<sup>4</sup> It must be distinguished from *Leontodon palustre*; Marsh Dandelion.



The flower is terminal, large, of a golden colour and closes in the evening: the calyx is smooth, with the exterior scales loosely turned down: the florets are very numerous, ligulate and toothed at the extremities. The receptacle is spheroidal and punctured. The seeds are obovate, furrowed, of a pale olive colour; and furnished with a radiated pappus, on a large stipe.

The herbaceous part of this plant is blanched and used on the continent as a salad; but, in this country, although it is designated by the Edinburgh and Dublin colleges, yet it is very seldom used, the root possessing much more of the principle on which the medicinal powers of the plant depend. The recent, full-grown root only should be used. It is white, and covered with a brown cuticle.

*Qualities.* — Dandelion is inodorous, but has a bitter, somewhat sweetish, acidulous taste. The milky juice reddens the vegetable blues, owing, according to Hermbstadt<sup>1</sup>, to the presence of tartaric acid. Water extracts its virtues better than alcohol; and scarcely any thing is taken up by ether: yet Dr. John detected caoutchouc in it. The decoction is precipitated by infusion of galls and solutions of nitrate of silver, muriate of mercury and superacetate of lead. Sulphate of iron strikes with it a pale olive colour and, after some time, throws down a precipitate. Hence it is probable, that the active principles of taraxacum are extractive, gluten, a bitter principle which does not appear to be resinous, and tartaric acid either free or as a supertartrate. The above mentioned re-agents are incompatible with the decoction.

*Medical properties and uses.* — Dandelion is aperient and diuretic. It has been long used on the continent as a remedy in jaundice, dropsy, pulmonic tubercles, hepatic obstructions, and some cutaneous diseases.<sup>2</sup> In this country it has been lately tried: and although its powers appear to have been over rated by the German physicians, yet it certainly possesses some efficacy in these diseases: and Dr. Pemberton affirms, that he has seen great advantage result from using the extract in chronic inflammation and incipient scirrhus of the liver, and in chronic derangements of the stomach.<sup>3</sup> Dr. W. Phillip considers it well adapted for cases in which bile is deficient without an impaired state of stomach. It may be given in the form of extract, or of infusion, made by boiling ʒij. of the sliced root in Oij. of water down to a pint, and to the strained fluid adding ʒiij. of supertartrate of potass: f ʒij. may be given for a dose three or four times a day.

Official preparations. *Extractum Taraxaci*, L. D.

<sup>1</sup> Thomson's Chemistry, 4th ed. v. 641.

<sup>2</sup> Bergius, Mat. Med. ii. 649.

<sup>3</sup> Diseases of the Abdominal Viscera, 42.



## LICHEN.

Cl. 24. Ord. 5. Cryptogamia Algæ. Nat. ord. Algæ.

Generic. Char. Male. Scattered warts.

————— Female. Smooth shields or tubercles, in which the seeds are embedded.

Species 1. *L. Islandicus*. Iceland or Eryngo-leaved Liverwort.

Eng. Bot. 1330. Flor. Danica, 155. Regnault, *Observations on Pulmonary Consumption*.

Species 2. *L. Rocella*. Dyer's Lichen, or Orchall. Eng. Bot. 211.

## 1. LICHEN ISLANDICUS.

Officinal. LICHEN, Lond. LICHEN ISLANDICUS, Edin. Dub. Iceland Liverwort.

Syn. Lichen d' Islande (*F.*), Isländisches Moos (*G.*), Yslandsch moos (*Dutch*), Islands moos (*Dan.*), Islands mossa (*Swed.*), Liquen Islandico (*S.*), Musgo Islandico (*Port.*), Lichene Islandico (*I.*).

This species of Lichen is an indigenous perennial. It is very abundantly found in Iceland, and in the north of Germany; and is more or less common on all the heaths and mountains of the north of Europe.<sup>1</sup> It grows to the height of two or three inches only, and has a rugged, bushy aspect. The frond is dry, coriaceous, lobed and laciniated, the lobes being subdivided and notched, resembling in appearance a buck's horn; but concave above and convex beneath; their surface is smooth, shining, and blistered; the margins beset with short, very minute, rigid, parallel hairs: and the colour of the whole is greenish-yellow, or greyish-brown.

This plant is used in Iceland and Lapland as an article of diet; being boiled in broth or milk after being freed from its bitter by repeated maceration in water; or dried and made into bread. It has of late years been brought in considerable quantity to this country for medicinal purposes.

*Qualities.*—The dried Lichen differs very little in its appearance from the recent plant. It is inodorous, and has a bitter mucilaginous taste; is neither very tough nor very brittle, but is not easily pulverized. When macerated in water it absorbs more than its own weight of the fluid, and the blisters appear like little white, opaque, glands, while the other parts of the plant are diaphanous. If the water employed in the maceration be warm, it acquires a strong, bitter taste, very similar to that of an infusion of Quassia. The macerated Lichen boiled in water affords a yellow-coloured inodorous decoction, which thickens as it cools and becomes a tremulous jelly, resembling starch, but without any viscosity. After some time this jelly cracks, separates from the watery part, and dries into semi-transparent masses, which are not soluble in cold water, but soluble in boiling water; and from

<sup>1</sup> It grows abundantly in the Asturias. *Journ. de Physique*, 1806.



which it is again precipitated by infusion of galls. According to the analysis of Proust, 100 parts of Lichen afford 64 parts of a substance insoluble in hot water, somewhat resembling vegetable gluten, 33 parts of a matter soluble in hot water, resembling starch, and three parts of a bitter extractive principle.<sup>1</sup>

*Medical properties and uses.*—Iceland Liverwort is tonic, and demulcent. From some remarks of Linnæus, made in 1737 in the *Flora Lapponica*, it would appear that the Danish physicians had long before that time employed this lichen, and found it efficacious in hæmoptysis, and pulmonary complaints; but it did not excite the attention of even the continental physicians, till after Scopoli's observations on it, in 1769, were published; and very few years have passed since it was known as a remedy in this country. Its virtues for the cure of phthisis have been very highly extolled; but experience has not altogether confirmed the truth of the praises which have been lavished on it.<sup>2</sup> Its supposed specific effects are said to depend on the combination of its tonic bitter and its demulcent properties. As a demulcent it is certainly superior to the mucilages; and owing to the bitter principle it contains, its decoction affords all the good effects that can be obtained from the other demulcents, and the mucilages, without loading the stomach. It allays the tickling cough, and relieves the oppressed breathing; involves the acrid matters contained in the stomach and bowels which often induce diarrhœa, and renders more bland the whole mass of animal fluids, so as to mitigate hectic fever, while, at the same time, it tends to invigorate the digestive organs. Still, however, its efficacy in phthisis is very circumscribed; but the circumstances above enumerated ought not to be over-looked, nor the Iceland Lichen regarded, as it often is, as a demulcent not more worthy of notice than the other articles of the same class. Besides phthisis, it has been also found useful in debilities after acute diseases, and in emaciations, particularly those arising from the great discharge of ulcers; in diarrhœas, dysentery, and hooping-cough.

It is generally exhibited in the form of decoction; (see *Preparations and Compositions*;) but as the bitter proves hurtful where the lungs or other viscera are actively inflamed, that part must be therefore separated. This is effected by cutting or pounding the Lichen, macerating it in several waters, and then, after boiling it for ten minutes, and decanting off the

---

<sup>1</sup> *Journal de Physique*, 1806.

<sup>2</sup> It still possesses a high reputation as a remedy for phthisis by the natives of Iceland. *Mackenzie's Travels*, 4to. *Appendix*, p. 411.



water, boiling it to the form of a mucilage in a fresh portion of water.

Officinal preparation. *Decoctum Lichenis*, L. D.

## 2. LICHEN ORCELLA.<sup>1</sup>

Officinal. LITMUS; LACMUS TINCTORIUS, *Dub.* Litmus.

*Syn.* Orchel, Orseille (*F.*), Oricello (*I.*), Orciglia (*S.*).

This is an indigenous Lichen, found in Portland island: but as an article of commerce it is obtained from the Levant; and also the Canary islands<sup>2</sup>, which produce annually 2600 quintals. It is a small species, seldom exceeding two inches in height, and is firmly fixed to the rocks by a solid base. From this base rises a tuft of worm-like stems, round, acutely pointed, often curved, more or less branched, smooth, of a white-grey, or brownish hue, and studded about their upper part with scattered tubercles, replete with white powder, which have been thought the seeds; but the fructification of this species is not well understood.

From this Lichen is prepared the *argol* or *archil* of commerce. It was known to the ancients, being the Λεικην of Dioscorides and the *Phycos thalassion* of Pliny. Its use as a dye was, however, lost; till it was again accidentally discovered by a merchant of Florence, in 1300<sup>3</sup>, observing that urine gave the lichen a fine violet colour. The preparation was long a secret, and was confined to Florence and Holland: but it is now known in England, and large manufactories of it are carried on in London and Liverpool. The Lichen, after being dried and cleaned, is reduced to powder in a mill resembling an oil-mill.<sup>4</sup> It is then mixed in a vat with one half of its weight of pearl-ash, and moistened with human urine: fermentation soon succeeds, and is kept up by stirring, and by successive additions of urine, until the colour of the material changes first to red and then to blue. In this state it is mixed with a third of its weight of good potass, and spread out to dry.<sup>5</sup> Chalk is sometimes added to it, but with no other view than to increase the weight.<sup>6</sup> It is generally sold in the form of cakes, but sometimes in that of a moist pulp.

*Qualities.* — Prepared argol has a slight violet odour,

<sup>1</sup> Vide *Dillenius' Hist. Muscorum*, 4to. p. 120. tab. 17. fig. 39., where it is figured under the name of *Coralloides corniculatum fasciculare tinctorium, fusci teretis facie*.

<sup>2</sup> The ancients named the Canaries the Purple Isles, from the abundance of orchilla which they yielded. *Mem. de l'Acad. des Inscriptions*, tom. iv. p. 457.

<sup>3</sup> *Thomson's Chemistry*, 4th ed. vol. v. p. 284. *Bancroft on Philosophy of Colours*, 2d ed. vol. p. 292.

<sup>4</sup> Sometimes it is not ground, but prepared in the entire state.

<sup>5</sup> *Nicholson's Journ.* 4to. ii. 311.

<sup>6</sup> Archil is chiefly used by the dyers, and in times of scarcity the Lichen has been sold at 1000*l.* sterling the ton. It is often mixed with the Lichen *fuciformis*.



arising from orris-root, with which it is always mixed, and a mawkish taste, leaving some degree of pungency in the mouth. When moist, the form of the Lichen is evident in the pulp. It communicates to water and to alcohol a beautiful violet colour, which, however, is very evanescent: all acids and salts, with an excess of acid, change it to red, which is again destroyed, and the blue restored by the addition of alkalies; and even exposure to the air of a room, in which many people are assembled, reddens the watery infusion. The tincture is least liable to change when kept, if it be reddened by an acid, and kept in close vessels.

*Use*.—This species of Lichen is said to have been “administered medicinally, with an intention of allaying the tickling attendant on phthisis, and in hysterical coughs;” but we must suppose the recent Lichen is meant, or before it has undergone any preparation as a colouring matter. We know of no other use of the prepared Lichen than as a dye stuff, or a chemical test of the presence of acids; and it is certainly the most delicate.

LINUM.<sup>2</sup> *Spec. Plant. Willd.* i. 1533.

*Cl.* 5. *Ord.* 5. Pentandria Pentagynia. *Nat. ord.* Gruinales, Linn. Caryophylleæ, Juss.

*G.* 590. *Calyx* five-leaved. *Petals* five. *Capsule* five-valved, ten celled. *Seed* solitary.

\* *With alternate leaves.*

*Species* 1. *L. usitatissimum*. Common Flax. *Med. Bot.* 2d edit. 566. *t.* 202. *Smith. Flora Brit.* 342. *Curtis Lond. fasc.* 5. *t.* 22.

\*\* *With opposite leaves.*

*Species* 26. *L. catharticum*. Purging Flax. *Smith, Flora Brit.* 344. *Eng. Bot.* 382.

#### 1. LINUM USITATISSIMUM.

*Officinal.* LINI USITATISSIMI SEMINA, *Lond. Edin.* LINUM; SEMINA, *Dub.* Linseed, and Linseed oil.

*Syn.* Grains de Lin (*F.*), Leinsaamen; Flachsamen (*G.*), Vlasch (*Dutch*), Hør (*Dan.*), Lin (*Swed.*), Len (*Pol.*), Alleverei (*Tam.*), Tokhemkutan (*Pers.*), Bidgierammu (*Malay*), Semi di Lino (*I.*), Laxor (*S.*), Linhaca (*Port.*), Buzruk (*Arab.*), Tisi (*H.*), Atasci (*San.*).

The common Flax is an annual plant, flowering in July. It is supposed to have been originally brought from those parts of Egypt which are annually inundated by the rising of the Nile; but it is now found growing wild in this country, and is cultivated in most parts of Europe. The root is simple and fibrous. The stem is erect, round, smooth, slender, and leafy; branched, with a panicle at the summit, and rising about two feet in height. The leaves are small, lanceolate, entire, obscurely three-nerved, smooth, sessile, standing nearly

<sup>1</sup> Translation of the Dublin Pharmacop. p. 165.

<sup>2</sup> Λινον Dioscoridis.



upright, and alternate on both the stem and branches. The flowers are petiolate: the calyx persistent, composed of five, sharp-pointed, keeled, tri-nerved, ovate leaflets; and the corolla consists of five notched, oblong, sky-blue, streaked petals, which spread into funnel-formed blossoms. The filaments are white, dilated, and slightly united at the base; the germen is ovate, and crowned with five blue, thread-like, spreading, reflected stigmas. The capsule is globular, the size of a common pea, crowned with a sharp spine, formed by the junction of the spines of the valves in one point, and containing in each cell an elliptical shining seed.<sup>1</sup>

Although this plant is extensively cultivated in Britain, yet the greater part of the linseed used here is brought from the Baltic. The seed ripens in September; and the plant is then pulled up as soon as the heads begin to change brown and hang downwards, otherwise the seeds are soon scattered.

*Qualities.* — These seeds are inodorous, and have an oily, mucilaginous, sweetish taste. They are small, flat, oval, and covered with a smooth, shining, brown-coloured cuticle, which abounds with a mucus, that can be extracted pure by infusion in boiling water. By expression, they yield one-sixth of their weight of fixed oil. The mucus of linseed is colourless, insipid, inodorous, and resembles in its viscosity mucilage of acacia gum; but differs from it in the following particulars: Alcohol precipitates it in white flocks, but the liquid remains clear; superacetate of lead throws down a dense precipitate; but oxy-sulphate of iron and silicated potass produce no sensible effect. For the qualities of the oil, see *Preparations*.

*Medical properties and uses.* — Linseed is emollient and demulcent. The mucus obtained by infusion is a cheap and very useful demulcent in catarrh, pneumonia, diarrhœa, and dysentery: visceral inflammations, calculus, gonorrhœa, ardor urinæ; and during the exhibition of oxymuriate of mercury. When the seeds are boiled in water, the mucus is obtained in union with a portion of the oil; forming a useful local remedy when given in the form of enema in abrasions of the intestines and tenesmus, particularly in the advanced stage of puerperal fever, when the offending matter in the bowels stimulates to frequent and involuntary stools: but the portion thrown up must be small in quantity.<sup>2</sup> The seeds,

<sup>1</sup> The partitions of the cells are singular. Gærtner thus describes them: “Dissepimenta membranacea, conduplicata, laminis suis extrorsum partitis, ita valvularum marginibus inserta, ut, cum hæ dehiscunt, illæ corii follii adinstar explicentur:” and adds, “Dissepimentorum in Radiola atque Lino fabrica, hactenus sine pari est, et essentialem hujus generis præbet characterem.” *De Fructibus*, ii. 147.

<sup>2</sup> *Denman's Midwifery*, ii. 251.



ground into powder or meal and simply mixed with boiling water, form an excellent poultice; valuable on account of the facility with which it is made.

Officinal preparations. *Infusum Lini*, L. *Oleum Lini*, L. E. D.

## 2. LINUM CATHARTICUM.<sup>1</sup>

*Officinal.* LINUM CATHARTICUM, *Lond.* LINUM CATHARTICUM; HERBA, *Dub.* Purging Flax.

*Syn.* Lin. Purgatif (*F.*), Purgier flachs (*G.*), Purgeervlasch (*Dutch*), Liden Vildhör (*Dan.*), Villhör (*Swed.*), Lino purgativo (*I.*), Cantilagua (*S.*), Linho purgante (*Portug.*).

This is an indigenous annual plant, found on dry and hilly pastures, flowering from June to August. The root is small, and sends up several delicate, leafy, erect, smooth stems, simple at the base, but dichotomous above, many-flowered, and from three to nine inches high. The leaves are opposite, of a sub-elliptical, lanceolate shape, obtuse, entire, green on the upper surface, and glaucous beneath. The flowers are small and white; nodding before they open, after which they rise and stand erect. The leaves of the calyx are pointed, serrated, and one-nerved; in the corolla, the petals are obovate, acute, white, and spreading; the filaments are united, forming a circle round the lower part of the germen, which is furnished with capitate stigmas. The seeds are yellow and shining.

*Qualities.*—Purging Flax, whether in the recent state or the dried, is nearly inodorous, and has a bitter, sub-acrid taste. Water extracts the virtues of the plant, which communicates to it, besides its sensible qualities, a yellow colour. Macerated in ether, it affords a green tincture, which deposits, when it is evaporated on the surface of water, a green, bitter resin, and an extractive matter, on which the virtues of the plant seem to depend.

*Medical properties and uses.*—This species of Flax was celebrated as a purgative by Gerarde. It may be given in the form of infusion, made with ʒij. of the dried plant, and ʒj. of boiling water, of which fʒij. is a dose: of the dried plant in powder, ʒj. may be taken for a dose. But it possesses no particular advantages, and only swells unnecessarily the list of purgatives.

LYTHRUM. *Spec. Plant. Willd.* ii. 865.

*Cl.* 11. *Ord.* 1. Dodecandria Monogynia. *Nat. ord.* Calycanthemæ, *Linn.* Salicariæ, *Juss.*

*G.* 951. *Calyx* twelve-toothed. *Petals* six, inserted into the calyx. *Capsule* two-celled, with many seeds.

*Species* 1. *L. Salicaria*.<sup>2</sup> Loosestrife, or Purple Willow Herb.

*Med. Bot.* 2d edit. *Smith Flora Brit.* 510. *Eng. Bot.* 1061.

*Officinal.* LYTHRUM SALICARIA; HERBA, *Dub.* Loosestrife.

*Syn.* Salicaire (*F.*), Brauner Wiederich (*G.*), Partike (*Dutch*), Salicaria (*I.*).

<sup>1</sup> Λινοκάριπος Græcorum.

<sup>2</sup> Λυθρον Dioscoridis.



This is an indigenous, perennial plant, found wild in almost every part of Europe, in marshes and on the banks of rivers, flowering from July till September. It is an elegant plant. The root is woody, branched, and extended; sending up an erect, leafy, slender, reddish, downy stem, about three feet in height, quadrangular, and sometimes hexagonous. The leaves are opposite, sessile, lanceolate, and cordate at the base; smooth on the upper surface, but pubescent beneath, and at the margin. The flowers are in the axillæ of the leaves, forming a leafy spike of a verticillated aspect: the calyx is red, hairy, and the segments of different shapes; six being awl-shaped and erect, and six small, ovate, concave, and bent inwards; the petals are oblong, undulated and of a purple colour. The stamens are alternately longer and inflected. The capsule is elliptical and small.

*Qualities*.—Loosestrife, in the dried state, is inodorous and has an herbaceous, sub-astringent taste. In coction with water, it renders the fluid mucilaginous; and the decoction strikes a black colour with sulphate of iron.

*Medical properties and uses*.—This plant is astringent and tonic. It is recommended by De Haen, and has long been celebrated in Ireland as a remedy in diarrhœa: it has also been found useful in dysentery. It is always proper to give a purgative prior to its use being begun. The best form of giving it is that of decoction, made by boiling  $\mathfrak{z}$ i of the recent root with  $\mathfrak{O}$ j of water. The dose of the dried herb, in powder, is from  $\mathfrak{z}$ ss to  $\mathfrak{D}$ iv., that of the decoction of the root  $\mathfrak{f}$  $\mathfrak{z}$ i, repeated every third hour.

MAGNESIÆ SUBCARBONAS. *Lond.* Subcarbonate of Magnesia.<sup>1</sup>

The greater part of the Subcarbonate of Magnesia found in the shops has long been manufactured on a great scale; and, therefore, the London College has, properly, given it a place in the list of the Materia Medica.

This salt, under the name of Magnesia alba, was formerly prepared from the bitter liquor called *mother of nitre*, which remains after the preparation of nitre, when no more of that salt can be obtained by evaporation from the water which is

---

<sup>1</sup> A carbonate of magnesia exists ready formed in the serpentine rocks in Moravia. The colour of this mineral is a cream-grey: the lustre dull; and the fracture flat conchoidal, with splintery edges. It is opaque, adheres strongly to the tongue; and has sp. gr. 2.915. It acquires positive electricity when rubbed on woollen cloth. According to Bucholz, it consists of carbonic acid 51 parts, and magnesia 46.59 in 100 parts of the mineral. — *Ann. de Chim.* t. lxxiv. p. 76. A native carbonate of magnesia is also found in Hindostan; and, according to Dr. Thomson's analysis, consists of carbonic acid 51.627, magnesia 47.566, deutoxide of Manganese 0.807, in 100 parts. *First Princip. of Chem.* vol. ii. p. 34.



washed out of the earth of nitre beds.<sup>1</sup> Of late years, however, it has been prepared, on a large scale, from the bittern, or liquor which remains after the crystallization of common salt, muriate of soda, from sea-water, which is chiefly a solution of muriate of magnesia. The bittern is heated to  $212^{\circ}$ , and decomposed by impure carbonate of potass, or of ammonia, which is stirred into it the moment it reaches the boiling point, and the fire withdrawn. A double exchange takes place, the muriatic acid leaving the Magnesia, unites with the alkali of the carbonate, forming a muriate of potass or of ammonia which remains in solution, whilst the disengaged carbonic acid of the carbonate combines with the magnesia, and forms an insoluble subcarbonate which is precipitated. The supernatant liquor is next drawn off, and the precipitate, after being repeatedly washed, is dried upon chalk stones. It is brought to market in square masses, which, when the article is good, are very light, and smooth to the touch. It is sometimes adulterated with chalk, and occasionally with gypsum. The chalk is detected by putting a portion of the mass into diluted sulphuric acid, which converts the Magnesia into a soluble sulphate, but produces an insoluble salt with the lime of the chalk: the gypsum is detected by boiling a portion of the Magnesia in distilled water, and adding to the solution muriate of barytes, which produces an insoluble precipitate, if gypsum be present. When it is prepared by decomposing the Sulphate of Magnesia by means of carbonate of potass, if not exceedingly well washed, it sometimes contains sulphate of potass, which may be detected by dissolving it in diluted nitric acid, and adding nitrate of barytes to the solution; an insoluble sulphate will be thrown down if sulphate of potass be present.

This salt is, in correct chemical language, a hydrated carbonate, and consists of 52.4 parts of carbonic acid, and 47.6 of Magnesia, in 100 parts: or, admitting that one atom of the carbonate is in weight = 42, its constituents will be one atom of the acid = 22, and one atom of the earth =  $20^2$ : but according to Berzelius the proportions are of acid 36.263, magnesia 43.956, water 19.781. By diffusing it in water, and passing carbonic acid through the mixture, it is converted into a bicarbonate, which crystallises in transparent, hexagonal prisms, each terminated by a hexagonal plane, about six lines in length and two in breadth, and consisting of one atom of acid, 2.75, one atom of Magnesia, 2.5, and three atoms of water,  $3.875^3$ .

<sup>1</sup> *Hoffmani. Opera*, t. iv. p. 479. *Black on Magnesia alba*, p. 2.

<sup>2</sup> *Phillip's Trans. of Pharm.* 1824. But according to Fourcroy and Kirwan, the constituents are acid 50, magnesia 25, water 25, in 100 parts.

<sup>3</sup> Berzelius. *Afhandlingar*, vj. 21.



A solution of this salt, with a great excess of carbonic acid, has been lately sold under the name of aerated magnesian water.

*Qualities.* — Carbonate of Magnesia is insipid, inodorous, perfectly white, and unalterable in the air. Its specific gravity is 0.2941. It is soluble in 480 parts of water at 60°, and is less soluble in hot water. It is decomposed by a strong heat, which drives off the carbonic acid.

*Medical properties and uses.* — For these, see Part iii. 7718

*Officinal preparations.* *Magnesia*, L. E. D.

**MAGNESIÆ SULPHAS**, *Lond.* **SULPHAS MAGNESIÆ**, *Edin.* **SULPHAS MAGNESIÆ**; *olim Sal catharticum amarum*, *Dub.* Sulphate of Magnesia. Bitter purging Salt.

*Syn.* Sulphate de Magnesie (*F.*), Schwefelsaure Magnesia (*G.*), Englesk laxesalt (*Dan.*), Sale amaro; Ossisolfato di Magnesia (*I.*), Sal Amargo (*S.*), Sal Cathartico amargo (*Portug.*).

This salt is found native in a pure state;<sup>1</sup> but it is more commonly combined with gypsum<sup>2</sup> and other salts, and in solution in sea-water, and several mineral springs. It was first artificially obtained in England in 1675, by Dr. Grew, from the evaporation of the water of the Epsom spring: whence it was named Epsom salt: and in 1700 it was made in considerable quantity from two springs at Shooter's-hill in Kent<sup>3</sup>; but the discovery of it in bittern, or the residual brine after the crystallization of sea-salt, soon opened a more copious source from which it might be obtained at all times; and for many years past, almost all the Sulphate of Magnesia used in this country has been manufactured from bittern. This substance consists chiefly of muriate of magnesia, muriate of lime, some common salt, and a small portion of sulphate of lime; and therefore it is probable, that the Sulphate of Magnesia is obtained by decomposing the muriate by means of sulphate of iron, or sulphuric acid in some form, although some affirm that the bittern is only boiled down to a high point of concentration, when the Sulphate of Magnesia forms, and is purified by a second solution and crystallization. Much of the Sulphate of Magnesia, however, now sold, is prepared from magnesian limestone, by a process invented by Dr. Henry of Manchester; and as it contains no muriate of magnesia, it is

<sup>1</sup> In the mercury mines of Idria it is found crystallized, and named by the Germans *Haarsalz*. According to Klaproth, it contains 1 per cent. of oxide of iron. *Analyt. Ess.* 80.

<sup>2</sup> It is found in the gypsum quarries of Piedmont; and, as Proust relates, it abounds so much in Spain, that in Andalusia large tracts are covered with an efflorescence of it after floods. *Journ. de Physique*, xxxiii. 312.

<sup>3</sup> It is also made in Bohemia from the mineral water of Sedlitz: and in the neighbourhood of Genoa from a pyrites found there; by first roasting the mineral and then exposing it to the air under a shade for six months, occasionally watering it, and then lixiviating.



much less disposed to attract moisture from the atmosphere and deliquesce, than that prepared from bittern. Much of the sulphate found in the shops contains some muriate of magnesia, which renders it deliquescent; and consequently, it requires to be preserved in close covered jars. It is often adulterated with glauber salt, which is made to resemble Epsom salt, by stirring it briskly, when it is about to crystallize. It may be detected, by precipitating the magnesia by pure ammonia, aiding by heat; filtering, and evaporating the filtered fluid to dryness, by a heat sufficient to volatilize the sulphate of ammonia: if it contain glauber salt, the soda will remain fixed. Or it may be detected by no precipitation ensuing, on adding carbonate of potass to the solution. Muriate of lime is detected by the oxalic acid.

*Qualities.* — Sulphate of Magnesia is inodorous, and has a very bitter, nauseous, saline taste. It is usually in small needle-like crystals, but the form of its regular crystal is a quadrangular prism, acuminate by four planes. When pure it effloresces; and is soluble in its own weight of water at 60°, increasing the volume of the fluid rather more than 4-tenths, or a solution of ℥j of Sulphate of Magnesia in f℥j of water, measures eleven fluid drachms and a quarter. Heat expels its water of crystallization; and the mass is melted, but not decomposed; it loses merely its water of crystallization, and a minute portion of its acid. According to Bergman, 100 parts consist of 29.35 of sulphuric acid, 17 of magnesia, and 53.65 of water of crystallization.<sup>1</sup> Its specific gravity is 1.66. It is decomposed by the alkalies, and their carbonates, but the bicarbonates do not decompose it; neither is it precipitated by the carbonate of ammonia. It is decomposed by lime-water, the muriates of ammonia, of barytes and lime, nitrate of silver, and acetate and super-acetate of lead, which are therefore incompatible with it in prescriptions.

*Medical properties and uses.* — This salt is purgative and diuretic. It operates readily without griping; and, notwithstanding its nauseous taste, is generally retained by the stomach when almost all other things are rejected, especially when it is administered in small, repeated doses, largely diluted, or united with acidulated infusion of roses. In these forms it is a useful purgative in hypochondriasis, colica pictonum, ileus, puerperal fever, and in all acute diseases. In small doses, combined with compound spirit of ammonia, and dis-

<sup>1</sup> According to Dr. Henry, the composition is, acid 38, magnesia 18, and water 44 parts, in 100 of the salt. — Mr. Phillips (*Translation of the Pharmacopœia*) states the components to be sulphuric acid 32.52, magnesia 16.26, water 51.22. Dr. Thomson (*First Prin. of Chem.*) makes it to consist of 1 atom sulphuric acid = 5, 1 atom of magnesia = 2.5, and 7 atoms of water = 7.875.



solved in the infusion of Quassia or Gentian, it has been found very useful in dyspepsia accompanied with costiveness. It is also used as an adjunct to stimulating clysters. By moderate exercise in the open air while taking this salt, the purgative effect is diminished, and its diuretic property increased. The dose is from ʒss to ʒij dissolved in water, gruel, or any other vehicle; and taken either at once, or in divided doses frequently repeated.

Officinal preparations. *Enema Catharticum*, D. *Enema foetidum*, D.

MALVA. *Spec. Plant. Willd.* iii. 774.

Cl. 16. Ord. 6. Monadelphia Polyandria. Nat. ord. Columniferae, Linn. Malvaceae, Juss.

G. 1290. *Calyx* double, the exterior three-leaved. *Capsules* numerous, one-seeded.

\* \* *With angular leaves.*

*Species* 43. *Malva sylvestris*.<sup>1</sup> Common Mallow. *Med. Bot.* 2d edit. 554. t. 199. *Smith Flor. Brit.* 740. *Eng. Bot.* 671.

*Officinal.* MALVA, Lond. MALVÆ SYLVESTRIS HERBA, FLORES, Edin. Mallow.

*Syn.* Mauve (F.), Kasepappel (G.), Malva (I.), Malvas (S.).

This is a perennial, indigenous plant, common over all Europe, growing on waste grounds and at the sides of roads; and flowering from May till August. The root is fusiform, branching, and of a whitish colour. The stem frequently erect, branched, round, hairy, and many flowered. The leaves are alternate, petiolate, cordate, divided into seven lobes, plaited, somewhat rough, and crenate; the upper ones are almost palmate. At the base of each footstalk are two stipules. The flowers, which stand on slender, hairy peduncles, are large; composed of five, inversely cordate, purple petals, three times longer than the calyx, which is hispid. The capsules are from ten to fifteen in number, of a rounded kidney form, crustaceous, brittle, close all round, of a dark straw colour, excavated, and wrinkled on the back. The seeds are kidney-shaped, ash-coloured, and furnished with an arillus which opens inwardly.

*Qualities.*—Common Mallow is inodorous, and has a weak, herbaceous, mucilaginous taste. The decoction has a mawkish disagreeable taste; is precipitated by acetate of lead; and is little more than a simple solution of vegetable mucus.

*Medical properties and uses.*—This herb is more demulcent and emollient than the root. Its decoction is employed in dysentery, ischuria, strangary, and nephritic complaints, but is in every respect inferior to that of Althea root. It is chiefly used in the form of enema in tenesmus, and nephritic

<sup>1</sup> Μαλαχη Græcorum. Malva, quasi molva, quod alvum moliat.



colic: and in that of cataplasms and fomentations in phlegmonous inflammation.

### MANGANESIUM. Manganese.

This is a brittle, greyish-white, brilliant metal, somewhat resembling iron in its external aspect, of a granular texture and not possessing ductility or malleability. It has not been discovered native in its metallic state<sup>1</sup>, but its ores are found in most of the countries of Europe both in primitive and transition mountains. It has neither odour nor taste: is softer than cast iron, and is not magnetic. Its specific gravity is 8.013.<sup>2</sup> It rapidly attracts oxygen from the air, loses its lustre, and progressively becomes violet, brown, and ultimately black. It rapidly decomposes water. It is very abundantly found in the state of the grey oxide. Manganese in the ore, both in primitive and transition mountains, is found

#### A. United with oxygen.

##### i. oxidized

##### Sp. 1. *Gray manganese ore.*

Var. *a.* Radiated. *b.* Foliated.  
*c.* Compact. *d.* Earthy.

##### 2. *Black manganese ore.*

##### 3. *Sulphuret of manganese.*

##### 4. *Phosphate of manganese.*

##### 5. *Silicate of manganese.*

— *a.* and combined with }  
sulphur.

— *b.* ————— with }  
phosphoric acid, and }  
iron.

— *d.* ————— with }  
Silica and iron.

Of these species the first only has been introduced into the list of materia medica.

*Officinal.* MANGANESE<sup>3</sup>, *Dub.* Manganese, or more properly Black Oxide of Manganese.

*Syn.* Manganese (*F.*), Braunstein (*G.*), Manganese (*I. S.*).

Under the name of black oxide of manganese are implied all the varieties of the first species. It was discovered in England by Boyle, in the beginning of the 17th century, but was regarded as a modification of iron ore, till the separate experiments of Scheele and Bergman, published in 1774, proved it to be an oxide of a peculiar metal; which Gahn afterwards succeeded in obtaining in its metallic state. It is found in Great Britain, Germany, Switzerland, the north of Italy, and France.

The greater part of the black oxide of manganese used in

<sup>1</sup> La Perouse suspected that he had found manganese in a metallic state; but his opinion was not confirmed.

<sup>2</sup> John, vide *Gehlen's Journ.* iij. p. 460.

<sup>3</sup> This term is improperly used by the Dublin College: for although the black oxide was originally named *manganese*, and is still so named in commerce; yet in a professedly scientific work, more accuracy of nomenclature is required.



England is obtained near Exeter in Devonshire, in Cornwall, and at Howth, near Dublin. It occurs crystallized and amorphous; and is generally in combination with small portions of oxyde of iron, carbonate of lime, silex and barytes.

*Qualities.* — Black oxide of manganese differs in its external characters. Its usual colour varies from iron-grey to black: when crystallized it is shining, but when amorphous devoid of lustre. Its texture is radiated, foliated, compact or earthy. None of the varieties are very hard; all of them are brittle, and several of them soil the fingers. Their specific gravity varies from 3·5 to 4·7. One hundred parts of the black oxide consist of 71·23 of metallic manganese, and 28·67 of oxygen. Exposed to the heat of ignition, all the varieties afford oxygen gas; and when mixed in powder with sulphuric acid, they afford it at a low temperature. It converts muriatic acid into oxymuriatic acid<sup>1</sup>; or more properly, according to the theory of Sir H. Davy, the hydrogen of the muriatic acid is attracted by the oxygen of the oxide of manganese, and its chlorine is evolved.

*Medical properties and uses.* — This metallic oxide is only used for procuring oxygen gas; and for fumigation in cases of infection. To procure oxygen gas, a portion of the oxide is put into an iron retort, fitted with a long curved tube, the extremity of which being placed under an inverted jar filled with water in a pneumatic trough, the retort is put into a common fire, and exposed to a full red heat. The caloric at this high temperature weakens the affinity between the manganese and the oxygen with which it unites, and causing it to assume a gaseous state, the oxygen gas is transmitted through the water, and collected in the jar. From the necessity of oxygen for carrying on the process of animal respiration, much benefit was expected from the breathing oxygen gas in disease; but experience has not confirmed the high expectations which were formed of its powers. It certainly increases the force and velocity of the pulse; and has been exhibited with seeming advantage in asthma, chlorosis, scrofula, typhoid fevers, and other diseases of debility. Diluted with from ten to twenty parts of atmospheric air, one or two quarts of it may be breathed at intervals in the course of the day.

But a more certain benefit is obtained from the use of this oxide of Manganese in fumigations. Medicine is indebted to Morveau for the discovery of this mode of destroying infection, and the numerous instances in which it has proved beneficial have fully established its use. For a fumigation the following

---

<sup>1</sup> The greatest consumption of black oxide of manganese is for the formation of the oxymuriatic acid, as employed in the art of bleaching.



ingredients are required: common salt  $\text{℥iv}$ , oxide of manganese in powder  $\text{℥iv}$ , sulphuric acid  $\text{℥viij}$ , and water  $\text{℥iv}$ : the water and acid must be previously mixed together, and then after the mixture is cold, poured over the other ingredients in a China basin. The doors and windows of the room which is to be fumigated, must be closely shut for two hours after the charged basin has been placed in it; then thrown open, and a current of air allowed to pass through the room. Chlorine is extricated by this process, and neutralizes the infectious and contagious matter, floating in the atmosphere and attached to the walls and floors of the infected rooms. All metallic furniture should be previously removed.

MANNA. Vide *Fraxinus Ornus*.

MARRUBIUM. *Spec. Plant. Willd.* iii. 109.

Cl. 14. Ord. 1. Didynamia Gymnospermia. Nat. ord. Verticillatæ, Linn. Labiatæ, Juss.

G. 1111. Calyx salver shaped, rigid, ten-streaked. Corolla, upper lip bifid, linear and straight.

\* \* With ten-teethed calyces.

Species 8. *M. vulgare*.<sup>1</sup> White Horehound. *Med. Bot.* 2d edit. 332. t. 118. *Smith Flora Brit.* 636. *Eng. Bot.* 410.

Officinal. MARRUBIUM, Lond. MARRUBII VULGARIS HERBA, Edin. MARRUBIUM ALBUM; FOLIA, Dub. Horehound leaves.

Syn. Marrube blanc (F.), Weisser andorn (G.), Witte andoorn (Dutch), Ilvidmarru (Dan.), Andorn (Swed.), Szanta biala (Pol.), Marrubio (I.), Marubio blanco (S.), Marroyo blanco (Port.).

White Horehound is an indigenous, perennial plant, growing in waste grounds, and flowering in July. The root is fibrous, sending up numerous stems about eighteen inches high, quadrangular, erect, and very downy. The leaves are in pairs, upon broad footstalks, rounded, crenate, wrinkled, hoary, and woolly on the under surface. The flowers are white, in crowded, axillary whorls, sessile, villous and furnished with setaceous, awned bractes. The calyx is tubular, furrowed, and divided at the margin into ten, narrow segments, which are hooked at their points: the corolla is tubular, compressed, opening at the mouth into two lips, the upper of which is narrow and cloven; the under broader, reflected, and three-cleft, with the middle segment broad and scalloped. The filaments are two long and two short, with simple anthers, within the tube; and the style is slender, with a cloven stigma. The seeds are four, at the bottom of the calyx.

Qualities.—Horehound dried has an aromatic odour, which however is soon lost by keeping; and a bitter taste. Both water and alcohol extract its virtues. The infusion reddens

---

<sup>1</sup> Πρασιον Dioscoridis. Lemery says the name is derived from the Hebrew word *Marrob*, which means a bitter juice.



tincture of litmus, gives a deep olive green precipitate with sulphate of iron, a brown with nitrate of silver, and a pale yellow with muriate of mercury: acetate and superacetate of lead do not affect it. The active principles of Horehound, therefore, appear to be a bitter extractive, volatile oil, and gallic acid.

*Medical properties and uses.*—Horehound is tonic, diuretic, and laxative. It was formerly much used in pulmonary affections, and is still a popular remedy for asthma and obstinate coughs. It loosens the belly when taken in large doses, and was consequently recommended in jaundice, cachexies, menstrual obstructions, and hysteria; and although its powers are not found by modern practitioners equal to the account which the ancients gave of them, and therefore it is very seldom prescribed, yet, we have seen decided advantage from its exhibition in phthisis. The dried herb may be given in powder, in doses of from ʒss to ʒj; or of the expressed juice of the fresh plant from fʒss to fʒjss may be taken twice or thrice a day. It is also used in the form of infusion.

MASTICHE. Mastich. See *Pistacia Lentiscus*.

MATONIA. *Maton*, *Trans. of the Linnean Society*, vol. x. part 2d.

Cl. 1. Ord. 1. Monandria Monogynia. *Nat. ord.* Scitamineæ, *Linn. Spec. M. Cardamomum*.<sup>2</sup> The Cardamom Tree. *Smith in Rees Encyclop. Van Rhee* (Elettari.) *Hort. Malabar.* vol. ix. t. 4, 5. *Linnean Trans.* vol. x. part 2d. *Roxburgh. Ind.* p. 3. N. 226. *Asiatic Res.* vol. xi. p. 335.

*Officinal.* CARDAMOMI SEMINA, *Lond.* AMOMI REPENTIS SEMINA, *Edin.* CARDAMOMUM MINUS; SEMINA, *Dub.* Lesser Cardamom Seeds.

*Syn.* Petit Cardamome (*F.*), Kleine Kardamomen (*G.*), Kardamom (*Dutch*), Kadamome (*Dan.*), Kardemumma (*Swed.*), Amomo minore (*I.*), Cardamomo (*S. & Port.*), Kardamome (*Russian*), Kákula (*Arab.*), Purbi and Guzrate Cláchi (*H.*), Ela (*San.*), Kapol (*Javanese*), Yaylersie (*Tam.*), Capulaga (*Malay*).

The plant which produces these seeds is a native of India, growing on the mountains above Cochin and Calicut; in shady places on the declivities, and in the valleys. The cultivated plant does not flower till it is four years old. It rises twelve feet in height. The root is oblong, jointed, tortuous, of a whitish colour, and sending off numerous fibres. The stems which emerge from the root, are simple stripes, like a reed, round, smooth, and the thickness of the human thumb. The leaves are alternate and sheathing, about four spans in length; broad, green, and striated with parallel veins; and have a strong subacrid, aromatic taste and odour. The midrib of the leaf on the upper surface is pale green; on the under a much deeper green. The flowers are in racemes, which are sent off



from the root, and creep along the ground; and are furnished with oblong leaflets like capsules. The calyx is monophyllous, inferior, small, and divided into three obtuse teeth at the margin: the corolla is monopetalous, tubular, and four-cleft, the three outer segments being long, narrow, and of a straw colour, and the central one large, broad, concave, and irregularly oval. The filament is broad, slightly grooved, supporting a large, double, emarginate, crestless anther, having a deep fissure between its lobes, to receive the style, which is slender, and bearing a funnel-shaped ciliated stigma. The capsule is berried, and trilocular.

The ripe fruit is gathered in November; and the capsules, which are dried over a gentle and slow fire, change as they dry from green to a whitish-straw colour, and become thinner in the bark: whilst the permanent calyx and footstalk is detached by rubbing them between the hands.<sup>1</sup>

Three species of this genus are known; but that which is above described yields the officinal cardamoms. They are brought to this country in the Bengal ships in cases, each containing about 120lbs. weight. For the purpose of preserving them, they are kept in the capsules, which are small, triangular, striated, and of a pale, clear, straw colour.

*Qualities.* — Cardamom seeds have an agreeable aromatic odour, and warm spicy taste. They are easily separated from the capsule; and are of a brown colour, angular, corrugated, and pulverulent. Water, alcohol and ether extract their virtues; the two latter most completely. The watery infusion has a turbid appearance, and lets fall a flocculent precipitate, on the addition of alcohol, the acids, solutions of sulphate of iron, muriate of mercury, and acetate of lead; but the sulphate of iron does not alter its colour. The alcoholic tincture is rendered milky by water. The ethereal has a yellowish-green hue, and, when evaporated on the surface of water, leaves neither resin nor extractive, but a considerable portion of essential oil, which has the flavour and taste of the seeds in perfection. Cardamoms therefore seem to be entirely composed of fecula, mucus, and essential oil.

*Medical properties and uses.* — Cardamom seeds are carminative and stomachic. They are less stimulating than pepper, and are, therefore, used, united with rhubarb and magnesia, in the flatulent colic of children; and as a grateful addition to bitters in dyspeptic, and gouty affections of the stomach: but they are principally employed to give warmth to other remedies.

---

<sup>1</sup> In gathering the fruit, the fruit panicles are plucked up by the roots; and the pods being stripped through the fingers, are sorted into three classes; 1. Valli Kai, or head fruit; 2. Nadu Kai, middle fruit; 3. Poulo Kai, abortive fruit. *Linn. Trans.* x. p. 229.



The dose in powder is from grs. vj to ʒj.

Officinal preparations. *Extractum Colocynthidis compositum*, L.D. *Tinctura Cardamomi*, L.E.D. *Tinctura Cardamomi composita*, L.D. *Tinct. Cinnamomi comp.*, L. E. *Tinct. Gentianæ comp.*, L. *Tinct. Rhei*, L.E.D. *Tinct. Rhei cum Aloe*, E. *Tinct. Sennæ*, L.D. *Spiritus Ether. aromat.* L. *Vinum Aloes soccot.*, E. *Confectio. aromat.* L. *Pulvis Cinnamomi comp.* L. E. D. *Pilulæ Scilliticæ*, E. *Infusum Sennæ*, D.

MEL. *Lond. Edin.* Honey.

*Syn.* Miel (*F.*), Gemeiner Honig (*G. Dutch*), Honning (*Dan.*), Honung (*Swed.*), Miod (*Pol.*), Med (*Russian*), Tayn (*Tam.*), Shahid (*Pers.*), Mele (*I.*), Miel (*S.*), Mel (*Port.*), Ussub (*Arab.*), Medhú (*H. and San.*).

Honey is collected by bees from the nectaries<sup>1</sup> of flowers, in which it is abundantly secreted; but it probably undergoes some change within the insect before it is excreted by it, and deposited in the comb. That it does not, however, undergo the process of digestion as food is likely, for the honey or sugar on which bees are fed during winter is not again excreted as honey; and the flavour of honey varies according to the nature of the flowers from which it is collected. Thus the honey of Minorca, Narbonne, and England are known by their flavours; and the honey prepared in different parts even of the same country differs.<sup>2</sup> It is separated from the comb by dripping, and by expression: the first method affords the purest sort; the second separates a less pure honey; and a still inferior kind is obtained by heating the comb before it is pressed. When obtained from young hives, which have never swarmed, it is denominated virgin honey. It is sometimes adulterated with flour, which is detected by mixing it with tepid water; the honey dissolves, while the flower remains nearly unaltered.

*Qualities.* — Honey has a peculiar saccharine aromatic odour; and a sweet acidulous sharp taste. In colour it varies from white or yellowish-white, to a pretty deep shade of amber or golden-yellow; in consistence, from the fluidity of limpid oil to the stiffness of soft suet: and when the more limpid kind is kept, partly crystallizes into little irregular concretions. It evidently contains sugar, mucilage, wax, and an acid; and occasionally some essential oil, as in the perfumed honey of

<sup>1</sup> The nectary is a glandular organ of the corollas of flowers. In many flowers it forms part of the petals themselves; in others it is a distinct organ. It is not easy to assign the use of honey in the vegetable œconomy.

<sup>2</sup> In some parts of Asia and America a poisonous honey is met with, which probably owes its deleterious properties to the flowers on which the bees feed. It is supposed that the honey extracted from the *Azulea pontica* and some species of the genera *Kalmia*, *Andromeda*, and *Rhododendron* is poisonous. In the island of Bourbon, honey of a green colour, and very fragrant, is procured; and bears a high price in India, to which it is chiefly exported. But bees do not sip the honey secreted in all flowers: thus they refuse that of the Crown Imperial, *Fritillaria imperialis*, and of the Oleander, *Nerium oleander*, which kills thousands of flies.



the Crimea. Honey is soluble in water, and partially in alcohol; and, like sugar, passes into the vinous and acetous fermentation. When heated over a slow fire it throws up a scum; and if the heat be continued so as to produce evaporation, the vapour is inflammable; and the honey becomes brown, and acquires an unpleasant flavour, which is strong in proportion to the degree of temperature employed. Lowitz found that the addition of charcoal to a solution of honey deprives it of odour, taste, and colour; but the colour again returns when the solution is evaporated. Cavezzali separated the sugar by first melting the honey, then adding carbonate of lime (egg-shells) in powder as long as any effervescence appeared; and, after separating a scum which forms by rest, filtering it, and setting it aside to crystallize. The crystals he purified by washing them with alcohol.<sup>1</sup> Proust separated it from a ready granulated honey by the action of alcohol.<sup>2</sup> Nitric acid converts honey into oxalic acid.

*Medical properties and uses.*—Honey is laxative, and externally detergent and stimulant. Simple honey is seldom ordered as an internal medicine; indeed, when freely eaten as food, it passes off quickly by stool, and induces colic in some habits; on which account, simple syrup should perhaps be preferred in all cases for forming medicinal preparations for internal use. As a local stimulant, it is employed in glysters; and forms an excellent adjunct to gargles in cynanche, and aphthous ulceration of the mouth and fauces. It is also a useful detergent to foul ulcers.

*Officinal preparations.* *Mel despumatum*, L. D. *Mel Boracis*, L. *Mel Rosarum*, L. D. *Oxymel Colchici*, D. *Oxymel Scillæ*, L. D.

MELALEUCA. *Spec. Plant. Willd.* iii. 1428.

*Cl.* 18. *Ord.* 3. Polyadelphia Icosandria. *Nat. ord.* Hesperideæ, *Linn.* Myrti, *Juss.*

*G.* 1392. *Calyx* five-cleft, half superior. *Corolla*, petals five. *Filaments* numerous, connate in five bodies. *Style* one. *Capsule* half-covered, three-celled.

*Species Nova.* *M. Cajuputi*.<sup>3</sup> *Cajuputi Melaleuca*, *Rumphius* (*arbor alba minor*). *Herbar. Amboinense*, ii. lib. 2. cap. 26. t. 17.

*Officinal.* CAJUPUTI OLEUM, *Lond.* MELALEUCÆ LEUCADENDRI OLEUM VOLATILE, *Edin.* OLEUM CAJUPUT, *Dub.* *Cajuputi oil.*

*Syn.* Cajeput (*F.*), Kajeputohl (*G.*), Cajeput (*I.*), Cajuputa (*Malay*), Kyä-pootie tylum (*Tam.*), Kynpootie ka tail (*Duk.*).

<sup>1</sup> *Annales de Chimie*, xxxix. 110.

<sup>2</sup> *Journal de Physique*, lix. 428.

<sup>3</sup> As the specimens of the tree which yields the true *Cajuputi* oil, which were sent home by Mr. Christopher Smith, differ from the *M. Leucadendron*, which was formerly supposed to yield it, and agree with the *Arbor alba minor* of Rumphius; Dr. Maton and Sir J. E. Smith have fixed this as a new species under the name of *M. Cajuputi*.



The tree which yields this oil is a native of Amboyna, Java, and the south part of Borneo, where it grows very abundantly in dry arid places. It is named *Cajuputa*<sup>1</sup> in the Malay language; and also by the natives *Daun Kitsjil*, and *Caju-Kilan*. It is a small tree, in some situations rather a shrub than a tree with a running root, often arched, and half above the ground. The stem is covered with a rough, pale, lamellated bark. The leaves are alternate, on short petioles, not unlike those of the willow, about three inches long, and little more than half an inch broad, lanceolate, and somewhat falcated; entire, smooth, three-nerved, firm, dry, fragile, of a pale yellowish-green colour, and having a very grateful odour. The flowers are white, sessile, and accompanied with minute, ovate bractes. The calyx is tubular, five-toothed, and one half deciduous; the petals are roundish, and concave; and the bundles of the filaments, which are long, filiform, and bearing small ovate anthers, are fixed within the tube of the calyx. The germen is inferior, roundish, crowned with a simple slender style longer than the filaments; and becomes a three-celled capsule, containing many small, oblong, angular seeds.<sup>2</sup>

To prepare the oil, the leaves are collected in a hot dry day, and put into thoroughly dry bags; in which nevertheless they soon spontaneously heat and become moist, as if macerated in water. They are then cut in pieces, infused in water, and left to ferment for a night; after which they are distilled. The quantity of oil they yield is very small, scarcely more than three fluid drachms being obtained from two bags of leaves.<sup>3</sup> When newly drawn it is very limpid, pellucid, and volatile; and Rumphius says, smells strongly of cardamoms, but is more pleasant. It is generally imported in copper flasks or canisters; but lately some has been brought home in quart glass bottles. On account of the high price of real *Cajuputi* oil, it is said to be often adulterated with oil of turpentine, and coloured with resin of milfoil.

*Qualities.*—The odour of this oil, as it is brought to us, is at first powerful, and similar to that of a mixture of oil of turpentine and camphor, but it soon becomes extremely fragrant and agreeable: the taste is pungent, and resembles very much that of camphor. It is limpid, transparent, and gene-

<sup>1</sup> Kayu-puti means in English, *white wood*, thence Rumphius terms it *Arbor alba*. *Mat. Med. of Hindostan*.

<sup>2</sup> The natives of the Moluccas macerate the leaves and flowers in fresh oil, and afterwards impregnate the oil with the smoke of benzoin. This preparation they call *Minjac Money*, or odorated oil, and use it as an unguent for the head. *Rumphius Herb. Amboin.* l. c.

<sup>3</sup> *Rumphius*.



rally of a blueish green colour, which is said to be partly derived from the copper of the flasks; but Mr. Brande says, "none of the samples which I have examined contain copper."<sup>1</sup> When dropped on the surface of pure water, it diffuses itself over it, and very soon completely evaporates, which is a good test of its purity; and it burns rapidly, without leaving any residuum. Like other volatile oils, it is entirely soluble in alcohol, which is not the case when it is adulterated with oil of turpentine. It is partially soluble in water.

*Medical properties and uses.*—Cajuputi oil is a highly diffusible stimulant, antispasmodic, and diaphoretic.<sup>2</sup> When taken into the stomach, it produces a sensation of heat, fills and quickens the pulse; and soon afterwards a copious sweat breaks out. It is efficaciously given in dropsy, chronic rheumatism, palsy, hysteria, flatulent colic, and other spasmodic and nervous affections. As a local and external stimulant, it is employed diluted with olive oil, as an embrocation to allay the pain of gout and rheumatism, and to restore vigour to joints after sprains. When put into a carious tooth, it lulls the pain of tooth-ach; and we have seen much benefit derived from rubbing it on the temples, in defective vision from a weakened state of the eyes. The dose is three or four drops on a lump of sugar.

MELISSA. *Spec. Plant. Willd.* iii. 146.

Cl. 14. Ord. 1. Didynamia Gymnospermia. *Nat. ord.* Verticillatæ, *Linn.* Labiatæ, *Juss.*

G. 1118. *Calyx* dry, nearly flat above: with the upper lip subfastigiate. *Corolla*, upper lip somewhat arched, bifid; lower lip with the middle lobe cordate.

*Species* 1. *Melissa officinalis*.<sup>3</sup> *Officinal or Common Balm. Med. Bot. 2d edit.* 335. *t.* 119.

*Officinal.* MELISSÆ OFFICINALIS FOLIA, *Edin.* Balm leaves.

*Syn.* Melisse (*F.*), Melisse (*G.*), Melissa (*L.*), Balsamina (*S.*).

Balm is a perennial plant, a native of the south of Europe, growing in mountainous situations, and flowering from July to September. It is cultivated in our gardens.<sup>4</sup> The root is fibrous, and sends up annual stems, which rise about two feet high, and are branched, quadrangular, and smooth. The leaves are opposite in pairs, of a bright green colour, ribbed, deeply serrated, and cordate; the lower ones on long footstalks, and the upper nearly sessile. The flowers, which are in small axillary bunches forming semi-whorls, stand on

<sup>1</sup> *Manual of Pharmacy*, p. 45.

<sup>2</sup> "Hujus olei binæ guttæ cum cerevisia vel vino propinatæ sudores excitant vehementes, cui fini apta medicamenta India exhibet perpauca." *Rumphius*.

<sup>3</sup> Μελισσοφυλλον *Dioscoridis*, bees being very fond of it.

<sup>4</sup> It was cultivated by Gerarde in 1596.



slender peduncles, at the base of which are small, oblong, notched, hairy bractes. The calyx is tubular and pentangular; the upper lip tridentate; the lower shorter, and cut into two acute teeth. The corolla, which is tubular, of a yellowish-white colour, with the upper lip shorter, and notched, and the lower three-cleft, encloses the anthers: the seeds are four, ovate, and placed at the bottom of the calyx.

For medicinal use, the herb should be cut before it flowers.

*Qualities*.—The recent plant has the agreeable odour of lemons, which is lost in drying; and an austere, slightly aromatic taste. In distillation with water, it yields a small portion only of a yellow essential oil, on which its odour depends. The watery infusion tastes rough; reddens slightly litmus paper; and affords with oxysulphate of iron a deep olive, with nitrate of silver a deep brown, and with acetate of lead a copious greenish-white precipitate.

*Medical properties and uses*.—Balm is stimulant, stomachic, and diuretic. It was formerly prized as a corroborant in nervous affections; but it is now used only in infusion, as a diluent in fevers.

**MENISPERMUM.** *Spec. Plant. Willd.* iv. 824.

(*Cocculus*. De Candolle, *Syst. Nat.* i. 515.)

*Cl.* 22. *Ord.* 10. Diœcia Dodecandria. *Nat. Ord.* Menispermæ. *G.* 1826. *Male*. *Calyx* two-leafed. *Petals* four or six exterior, eight, interior. *Stamens* sixteen.

— *Female*. *Corolla* similar to that of the male. *Stamens* eight, sterile. *Germens* two or three. *Berries* one-seeded.

*Sp.* 4. *M. palmatum*. Palmated Menispermum. *Cocculus palmatus*. De Cand. *tom.* i. p. 522. *Berry Asiatic Res.* 10. p. 385. *t.* 5.

*Officinal*. **CALUMBA**, *Lond.* **COLOMBÆ RADIX**, *Edin.* **COLOMBO RADIX**, *Dub.* Calumba Root.

*Syn.* Colombe (*F.*), Kolumbowurzel (*G.*), Colomba (*I.*), Kalumb (*Mozambique*), Columboo vayr (*Tam.*).

The London College has, now, properly referred this root to the *Cocculus palmatus* of De Candolle, the *Menispermum palmatum* of Willdenow, which, for the sake of uniformity, we have placed as the title of the article, referring our readers, however, to De Candolle's work.

This species of *Cocculus* is a native of the eastern part of southern Africa, growing in great abundance in the forests of Mozambique, between Oibo and Mozambo. The roots are dug up by the natives in the month of March, and transported to Tranquebar, where it is a staple article of export with the Portuguese.<sup>1</sup> An entire root was taken to Madras by Mons. Fortin, in 1805, and a plant raised from it

---

<sup>1</sup> The root was formerly erroneously supposed to be named from the principal town in the island of Ceylon, which was regarded as its place of export.



there by Dr. Anderson, from a drawing of which it was ascertained to belong to the natural order Menispermæ; but as it was a male plant only, the genus and species were undetermined until they were fixed by De Candolle. Dr. Berry drew up the following character of the male plant, which has been adopted by De Candolle; but the female plant is yet undescribed. The root is perennial, ramose, and bears fusiform tubers. The stems are annual, withering at the end of seven months; voluble, simple, round, hairy, about the thickness of a goose quill, bearing distant alternate, five-lobed, five-nerved leaves: with entire acuminate lobes; and supported on round hairy petioles, shorter than the leaves. The male flowers are in axillary, solitary, compound racemes, hairy, and shorter than the leaves: bearing partial, alternate peduncles, with sessile flowers; and lanceolate, ciliated, deciduous bractes. The calyx is hexaphyllous, with three exterior leaflets, and three interior, equal, oblong, obtuse and glabrous. The corolla consists of six minute, oblong, wedge-shaped, concave, fleshy, obtuse petals. The stamens are six, a little longer than the corolla, the anthers four-lobed and four-celled: there is no pistillum. The roots are dug up in March; but the offsets only are taken; each offset being a sessile tuber.

The dried root is brought to this country packed in bags, and sometimes in cases. It is in transverse sections, generally about one-third of an inch in thickness, and one or two inches in diameter. The bark is thick, and easily detached, internally bright yellow, and covered with a wrinkled olive brown cuticle. The interior part of the root is of a pale brownish colour, and has a spongy texture, with darker converging rays, which are the remains of sap-vessels. The pieces are frequently much perforated, evidently by worms, and not, as has been supposed, by stringing to facilitate its drying. Those pieces which have the fewest worm-holes, the brightest colour, and are solid and heavy, are the best. It is said that the root of white bryony, tinged yellow with the tincture of *Calumba*, has been fraudulently substituted for this root.

*Qualities.* — *Calumba* root has a very slight aromatic odour, and a bitter taste. It breaks with a starchy fracture, and is easily pulverized. Water at 212° takes up one-third of its weight; and the infusion has all the sensible qualities of the root. These are also extracted by alcohol; but proof spirit is its best menstruum. The infusion is not altered by solutions of sulphate of iron, nitrate of silver, muriate of mercury, and tartarized antimony; but a copious precipitate is produced by the infusion of galls, and yellow *Cinchona* bark, by acetate



and superacetate of lead, oxymuriate of mercury, and lime-water. Hence Calumba root appears to contain cinchonia. M. Planche found it to contain a large proportion of a peculiar animal substance, a yellow, bitter, resinous matter and one-third of its weight of starch. By repeated distillation, he also obtained a volatile oil; and, from the residue, malate of lime and sulphate of lime.

*Medical properties and uses.*—Calumba root is an useful antiseptic and tonic.<sup>1</sup> It was first brought into notice by Fr. Rede, in 1685. It is frequently employed with much advantage in diarrhœas arising from a redundant secretion of bile, and in bilious-remittent fever, and cholera, in which it generally checks the vomiting. It also allays the nausea and vomiting which accompany pregnancy; and, according to Percival, it is equally serviceable in stopping the severe diarrhœa and vomiting which sometimes attend dentition.<sup>2</sup> Denman found it more useful than the Cinchona in the low stage of puerperal fever.<sup>3</sup> As a tonic, unaccompanied with astringency and possessing little stimulus, it has been recommended in phthisis and hectic fever, to allay irritability, and strengthen the digestive organs: and in dyspepsia. It may be given combined with aromatics, orange peel, opiates, and alkaline, or neutral salts, as circumstances require. We have found the powder, in combination with rhubarb and sulphate of potass, exceedingly serviceable in mesenteric fever.

The dose of the powdered root is from grs. xv. to ʒss, repeated three or four times a-day.

Officinal preparations. — *Infusum Calumbæ*, L. E. *Tinctura Calumbæ*, L. E. D.

MENTHA. *Spec. Plant. Willd.* iii. 74.

Cl. 14. Ord. 1. Didynamia Gymnospermia. *Nat. ord.* Verticillatæ, *Linn.* Labiatæ, *Juss.*

G. 1102. *Corolla* not quite equal, four-cleft; the broader segment emarginate. *Stamens* upright, distant.

\* *Spiked.*

*Sp.* 7. *M. viridis*. Spearmint. *Smith (spec. 3.) Flora Brit.* 612. *Med. Bot. 2d edit.* 338. t. 121.

\* \* *Capitate.*

*Sp.* 13. *M. piperita*. Peppermint. *Smith (spec. 4.) Flora Brit.* 613. *Med. Bot. 2d edit.* 336. t. 120. *Eng. Bot.* 461.

\* \* \* *Verticillate.*

*Sp.* 20. *M. Pulegium*. Pennyroyal. *Smith (spec. 12.) Flora Brit.* 624. *Med. Bot. 2d edit.* 342. t. 122.

<sup>1</sup> The Africans of Mozambique esteem it as a remedy for venereal affections, and the Chinese employ it as an aphrodisiac.

<sup>2</sup> *Medical and Experimental Essays*, vol. ii.

<sup>3</sup> *Introd. to Midwifery*, ii. 524.



1. MENTHA VIRIDIS.<sup>1</sup>

*Officinal.* MENTHA VIRIDIS, *Lond.* MENTHA SATIVA; FOLIA, *Dub.* Spearmint.

*Syn.* Baume verte (*F.*), Frauenmurze (*G.*), Menta Romana (*I.*), Menta (*S.*).

This is an indigenous, perennial plant, growing in marshy places, and flowering in August. For medicinal purposes, it is cultivated. The root is creeping; the stem quadrangular, and foliaceous, rising about two feet in height, erect, smooth, and branching. The leaves are opposite, nearly sessile, lanceolate, about two inches and a half long, and an inch broad; of a deep-green colour above, paler beneath, pointed, serrated, smooth, and sometimes a little hairy underneath: the flowers are supported on smooth, partial flower-stalks, verticillated, in long, pointed, paniced spikes; furnished with setaceous, ciliated, lanceolate bractes, longer than the flowers: the calyx is cylindrical and furrowed, with five, nearly regular teeth: the corolla funnel-shaped, tender, smooth, and of a purple colour: the stamens are varying in length, with roundish anthers: and the style, which is filiform, with a bifid divaricated stigma, rises from a four-cleft germen: the seeds are four, small, and generally abortive, owing to the viviparous nature of the roots.

For medicinal use, spearmint is generally cut just as the flowers appear; but for obtaining the essential oil, the flowering plant is preferred. It should be cut in very dry weather.

*Qualities.* — Spearmint has a strong aromatic odour, and a warm slightly bitter taste; neither of which qualities is impaired by drying. Both alcohol and water extract its virtues.

*Medical properties and uses.* — Spearmint is stomachic and carminative. The infusion is serviceable in allaying sickness and vomiting in a weakened state of the stomach.

*Officinal preparations.* *Aqua Mentha viridis*, L. D. *Infusum Menthae*, comp. D. *Oleum Menthae viridis*, L. D. *Spir. Menthae viridis*, L.

## 2. MENTHA PIPERITA.

*Officinal.* MENTHA PIPERITA, *Lond.* MENTHÆ PIPERITÆ HERBA, *Edin.* MENTHA PIPERITIS; HERBA, *Dub.* Peppermint.

*Syn.* Menthe poivrée (*F.*), Pfeffermünze (*G.*), Peperminthe (*Dutch*), Peparmynta (*Swed.*), Menta piperita (*I.*), Verbal nena de sabor de Pimienta (*S.*), Hortelaã apimentada (*Port.*).

Peppermint is an indigenous, perennial plant, growing in moist places, and flowering in August and September. It is generally cultivated for medicinal use; particularly about Mitcham in Surry<sup>2</sup>, whence the London market is chiefly sup-

<sup>1</sup> *Μινθη* Hippocratis. Hæc species dignoscitur pedicellis semper glaberrimis. *Smith, Flor. Brit.* 613.

<sup>2</sup> Considerably more than one hundred acres of this herb are grown in the parish of Mitcham; but the greater part of the peppermint is made into a *liqueur* which is sold as a dram in London. *Stevenson's Survey*, pp. 377, 378.



plied. There are three varieties of peppermint, the first of which is the officinal plant. The root is creeping: the stem quadrangular and channelled, nearly upright, and about two feet high, branching, purplish, and rather hairy, with the hairs bent backwards: the leaves are of a dark-green colour, opposite; petiolate, ovate, rather pointed, serrated, the upper side smoother and less pubescent than the under, which is paler, with white and purple veins: the flowers are in terminal spikes, solitary, almost capitate, interrupted beneath, with the lower whorl more remote, and on a footstalk: the bractes are lanceolate and ciliated: the calyx is furrowed, tender, studded with glandular points, the base entirely naked, very smooth, and five-cleft, with the teeth of a blackish purple colour, and ciliated: the corolla is purple; and conceals within its tube the anthers, which are on short filaments: the germen is four-cleft, with a filiform style longer than the corolla, and furnished with a bifid stigma.

Dr. Smith supposes that this plant was discovered by Doctor Eales; and on examining the Linnean Herbarium, now in his possession, he found that the *Mentha piperita*, described by Linnæus, was not our officinal plant, but merely a variety of the *M. hirsuta*, with the odour of peppermint. It was, however, first described by Petiver. “The cultivators of the plant observe, that to keep up its quality, the roots must be transplanted every three years: otherwise it degenerates into the flavour of spearmint.”<sup>1</sup> If the plant be cut in wet weather it changes to black, and is little worth.

*Qualities.* — The odour of both the recent and dried plant is penetrating and grateful, in some degree resembling camphor: and the taste pungent, warm, glowing and bitterish, followed by a sensation of coldness in the mouth. These qualities depend on an essential oil and camphor. The oil can be obtained separate by distillation in water, is of a yellowish colour, and holds the camphor in solution. One pound of the recent plant should yield from fʒj. to fʒjss of the oil.

*Medical properties and uses.* — Peppermint is tonic, antispasmodic, and carminative. It is chiefly used to allay nausea and griping, to relieve flatulent colic, and in hysteria: or, as a vehicle, to cover the nauseous taste of other medicines; but to many palates it is extremely disagreeable. It may be given under the form of watery infusion; but the distilled water and the essential oil are generally preferred.

Officinal preparations. *Aqua Menthæ piperitæ*, L. E. D. *Oleum Menthæ piperitæ*, L. E. D. *Spiritus Menthæ piperitæ*, L. E.

### 3. MENTHA PULEGIUM.<sup>2</sup>

Officinal. PULEGIUM, Lond. MENTHÆ PULEGII HERBA, Edin. PULEGIUM; HERBA, Dub. Pennyroyal.

<sup>1</sup> *Linnean Transactions*, v. 176.

<sup>2</sup> Γλήχων Dioscoridis.



*Syn.* *Menthe peuliot* (F.), *Polei* (G. Dutch, Dan.), *Puleja* (Swed.), *Poley* (Pol.), *Puleggio* (I.), *Poleo* (S.), *Poejo* (Port.).

This is an indigenous perennial plant, growing on heaths and in moist meadows, and flowering in September. Like the other mints, it is cultivated for medicinal purposes; and becomes more luxuriant and erect. The stem is obtusely quadrangular, trailing, branching and somewhat hairy: the leaves are petiolate, small, obtuse, bluntly serrated and, in a small degree, hairy underneath; the flowers, which are supported on stalks covered with short thick hairs, are in sessile whorls, numerous, and many-flowered; and without bractes: the calyx is of a purplish-green colour, tender, furrowed and covered with thick short hairs: five-cleft, with the teeth pointed, unequal and ciliated: the corolla is twice the length of the calyx, purple, four-cleft, with the base white and externally villous: the stamens are erect, and longer than the corolla: and the germen similar to that of the former species.

*Qualities* — The odour is similar to that of spearmint, but less fragrant; the taste aromatic and pungent, with a slight flavour of camphor. These qualities reside in a very volatile essential oil, which rises in distillation with water.<sup>1</sup>

*Medical properties and uses.* — Pennyroyal was formerly regarded as emmenagogue, expectorant, and diaphoretic; and was in repute for promoting the uterine evacuation, and relieving hysteria, whooping-cough, and asthma; but it is now justly considered of no value and seldom used in regular practice.

*Officinal preparations.* *Aqua Pulegii*, L. E. D. *Oleum Pulegii*, L. D. *Spiritus Pulegii*, L.

**MENYANTHES.** *Spec. Plant. Willd.* iii. 810.

*Cl.* 5. *Ord.* 1. Pentandria Monogynia. *Nat. ord.* *Preciæ*, Linn. *Lysymachiæ*, Juss.

*G.* 299. *Corolla* hirsute. *Stigma* cloven. *Capsule* one-celled.

*Sp.* 4. *M. trifoliata*.<sup>2</sup> Buckbean. *Med. Bot.* 2d edit. t. 97. *Smith Flor. Brit.* 225. *Eng. Bot.* 405.

*Officinal.* **MENYANTHES**, *Lond.* **MENYANTHIS TRIFOLIATÆ FOLIA**, *Edin.* **TRIFOLIUM PALUDOSUM**, *Dub.* The leaves of Buckbean.

*Syn.* *Menianthe* (F.), *Bitterklée* (G.), *Water-drieblad* (Dutch), *Vandklever* (Dan.), *Vattenklofver* (Swed.), *Troylist* (Pol.), *Trifolio fibrino* (I.), *Menyanthes de tres en rama* (S.), *Trevo d'agua* (Port.).

This is one of the most beautiful of our indigenous plants. It is a perennial not uncommon in watery situations, in a black, boggy soil, flowering in June and July. The root is long, round, fibrous, and of a black colour; the stems are

<sup>1</sup> "1 cwt. of fresh penny-royal affords an average produce of 1lb. of essential oil." *Brande's Manual*, p. 146.

<sup>2</sup> *Μήνανθος* Theophrasti.



spreading, branched, and clothed with sheathing footstalks, each of which supports a ternate leaf, formed of three obovate, smooth, bluntly-toothed leaflets, of a beautiful green on the upper surface and pale beneath: the flower-stalk, which springs from within the sheath of a leaf, is longer than the leaves, erect, smooth, round, bearing a thyrsus of about ten flowers, accompanied by small ovate entire bractes: the calyx is obtusely five-toothed: the corolla a funnel-shaped petal, cleft into five deep segments, which are white, tipped with rose-colour, and closed within with long, fleshy, shaggy fibres on their upper side: the anthers are sagittate, and of a red colour: the germen round: and the stigma cloven and notched, on a slender style twice the length of the stamens.

*Qualities.* — The leaves of buckbean have a faint, disagreeable odour, and an intensely bitter nauseous taste, which is extracted by infusion with water.

*Medical properties and uses.* — Buckbean is tonic, diuretic, and purgative. It has been used with seeming benefit in remittent and intermittent fevers, rheumatism, arthritic affections, and in cachectic and cutaneous diseases. In large doses it is apt to excite vomiting. The dose of the dried leaves powdered, is from ℥j. to 3j.; or of an infusion made with ℥ss. of the dried leaves, and boiling water Oss., from f℥j. to f℥jss., may be taken three or four times a day. It is adviseable to unite some aromatic with either of these forms.

MEZEREI CORTEX. Mezereon Bark. Vide *Daphne*.

MOMORDICA. *Spec. Plant. Willd.* iv. 601.

Cl. 21. Ord. 8. Monœcia Monadelphica. Nat. ord. Cucurbitaceæ. G. 1739. Male. Calyx five-cleft. Corolla five-parted. Filaments five.

———— Female. Calyx five-cleft. Corolla five-parted. Style trifid. Gourd opening elastically.

Species 13. *M. Elaterium*.<sup>1</sup> Squirting cucumber. *Med. Bot.* 2d edit. t. 72.

*Officinal.* ELATERII PEPONES, *Lond.* ELATERIUM, *Edin.* ELATERIUM; FRUCTUS, *Dub.* The fruit of the wild cucumber: Elaterium.

*Syn.* Concombre sauvage (*F.*), Esselsgurken (*G.*), Ezelskomkommers (*Dutch*), Cocomero salvatico (*I.*), Cohombrillo amargo (*S.*).

This species of momordica is a perennial native of the south of Europe, flowering in June and July. It is cultivated in England<sup>2</sup>, but does not survive the severity of our winters. The root is large and fleshy, sending forth several thick, rough, trailing stems, which branch and extend three or four feet every way: the leaves are on long petioles, large, rough, of a greyish green colour, and cordate: the flowers are axillary,

<sup>1</sup> Ἐλατήριον Dioscoridis. Σικυς αγριος Antiquorum.

<sup>2</sup> It was cultivated by Gerarde in 1596.



similar in appearance to those of the common cucumber, but smaller, of a pale yellow colour, with a greenish base: the male flowers stand on short peduncles, but the female sit on the germen, which is inferior; the fruit into which it swells has the appearance of a small oval cucumber, of a greyish colour, and covered with prickles. When fully ripe it quits the peduncle, and casts out the seed and juice, with great force, and to a considerable distance, through the hole in the base where the footstalk is inserted.

For medicinal use, the fruit is gathered in September, just before it is ripe; and the clear juice which runs from it, and that obtained by the expression of the fruit, are inspissated, and form the elaterium of the shops.

*Qualities.* — The juice is nearly inodorous, and has a very slightly bitter taste. It deposits by rest a considerable portion of feculent matter, containing a peculiar principle, to which Dr. Paris has given the name of *Elatin*, and on which the active properties of the fecula depend. For an account of its nature, see *Extractum Elaterii*, among the Preparations.

*Medical properties and uses.* — This fruit is a very violent cathartic. It was much employed by the ancients, who regarded every part of the plant as purgative; but Dr. Clutterbuck has proved that this is an error.<sup>1</sup> It is also probable that the term *Elaterium* was given by the ancients to very different substances, and Hippocrates applies it to any violent purgative. Dioscorides extolled the fruit as highly efficacious in melancholic and maniacal attacks. It is still frequently prescribed in dropsies; but when incautiously given, it may bring on a dangerous hypercatharsis. The juice, which is lodged in the centre of the fruit, directly around the seeds, and which is improperly termed the extract, is always employed in preference to the recent fruit.

Officinal preparation. *Extractum Elaterii*, L. E. D.

MORUS. *Spec. Plant. Willd.* iv. 368.

Cl. 21. Ord. 4. Monœcia Tetrandria. Nat. ord. Scabridæ, Linn. Urticæ, Juss.

G. 1664. Male. *Calyx* four-parted. *Corolla* none.

—— Female. *Calyx* four-leaved. *Corolla* none. *Calyx* becoming a berry. *Seed* one.

*Species* 5. *M. nigra*.<sup>2</sup> Common mulberry tree. *Med. Bot.* 2d edit. 712. t. 243.

*Officinal.* MORRI BACCÆ, *Lond.* Mulberries.

*Syn.* Murier noir (*F.*), Schwarze Maulbeeren. (*G.*), Morone o Gelso (*I.*), Moras (*S.*).

This species of mulberry is a native of Persia, whence it was brought to Italy, and gradually spread over Europe. It

<sup>1</sup> *Lond. Med. Repository*, xii. 67.

<sup>2</sup> Συράμνος The ophrasti.



is now abundantly cultivated in this country, flowering in June and ripening its fruit in September. The tree, which seldom exceeds thirty feet in height, is covered with a brownish-grey bark; the leaves are numerous, on short footstalks, cordate, serrated, veined, about three inches long and nearly as broad; rough on the upper surface, which is of a deep green colour and covered with minute warts; and paler and villous underneath. The male flowers, which are on the same tree as the female, are in close roundish catkins, composed of caducous florets, which consist of four concave, oval, erect, calycinal leaves, inclosing four filaments bearing simple anthers; the female flowers contain a roundish germen, crowned with two divaricated styles furnished with simple stigmas, inclosed in a calyx of four ovate, concave, erect leaves; which, after flowering, swell, become succulent and coloured, investing the seed; and, many of them being crowded together on one peduncle, form a spurious compound berry, that has the appearance of a real, succulent berry, composed of a number of smaller berries.<sup>1</sup>

*Qualities.* — Mulberries are inodorous, have a sweet, acidulous taste and abound with a deep, blood-red juice. Hermbstadt found that their acidulous quality depends on the presence of tartaric acid, and I ascertained that they contain also jelly and mucus.

*Medical properties and uses.* — This fruit is cooling and laxative; and, when not too ripe, allays thirst and proves exceedingly grateful in febrile diseases. It is seldom, however, used medicinally. When eaten too freely as an article of food, it is apt to occasion diarrhœa.

Officinal preparation. *Syrupus Mori*, L.

MOSCHUS. *Syst. Nat. Gmelin.* i. 172.

Cl. 1. Ord. 5. Mammalia Pecora.

G. 28. Horns none. Fore teeth eight in the lower jaw. Tusks one on each side in the upper jaw, projecting out of the mouth.

Species. *M. moschiferus*. The Musk Deer. *Pennant Quadr.* 56. t. 10. f. 1.

Officinal. MOSCHUS, *Lond. Edin. Dub.* Musk.

Syn. Musc (*F.*), Bisam (*G.*), Muskus (*Dutch*), Mysk (*Swed.*), Denner (*Dan.*), Muschio (*I.*), Almizcle (*S.*), Almisca (*Port.*), Meshk (*Arab.* and *H.*), Castorie (*Tam.*), Jebat (*Malay*), Mesk (*Pers.*), Rutta Oorrola (*Cyng.*).

The animal from which musk is obtained, is an inhabitant of the alpine mountains of the east of Asia, particularly of the Himālā mountains, which divide Thibet from India, where it is known by the name of *Custeru*. It is a solitary animal,

---

<sup>1</sup> It is a curious fact, that two mulberry trees growing near one another, bear generally more fruit, and that of a finer quality, than is produced by solitary trees. This is supposed to be owing to the male flowers in one of the trees being more effective than those of the other.



living among the rocks, and frequenting the highest tops of the snowy peaks; very timid, and difficult to be taken. The length of the full grown animal scarcely ever exceeds three feet, and in its general aspect it resembles the deer; the head is not very unlike that of a hog; the eyes are black and full; and, projecting from the upper jaw, the teeth hang pointing downwards, over the lower jaw; the fleece is coarser than that of the stag, but very light and soft, and varying in colour, at different seasons of the year and different periods of life, chiefly from brown to nearly black, hoary underneath, and sometimes, but rarely, whitish. The tail is very short. Between the navel and the prepuce is an oval bag, flat on one side and convex on the other, about three inches long and two broad, projecting, with a very small orifice and beset with short hairs. This is the musk bag. It is empty in the young animal; but in the adult contains from ʒjss. to ʒij. of musk, in a liquid state. The animal often expresses part of the contents of the bag, when it becomes too full, by rubbing itself against stones; and the matter thus ejected is said to be a purer musk than that which is brought to this country. The bag is generally cut from the animal while it is yet alive, and an idea prevails that the animal must be caught alive in order to obtain the musk, which is said to be absorbed and lost if the deer be shot. As soon as the bag is cut away, a small hollow reed is inserted into it, that the musk may not suffer, which it would be apt to do from want of air; and the whole is tied around with a sinew of the animal.<sup>1</sup>

Musk is imported into England from China in caddies, which contain from twenty to sixty and one hundred ounces each; but an inferior kind is brought from Bengal and a still baser sort from Russia. The best is that which is in the natural follicle, or the *pod*, as it is denominated in mercantile language. This is a sack or bag, about the size of a pigeon's egg, of a brownish colour, lined with a very thin membrane, and covered externally with coarse hairs. The musk itself is in grains concreted together, dry, yet slightly unctuous and free from grittiness, when moistened and rubbed between the fingers, or chewed.

As musk is a very high-priced article, it is often adulterated. When this is the case, the bag, which should not have any appearance of having been opened, appears, if narrowly examined, slit or punctured in several places, through which sand, lead, and other heavy matters are inserted. The musk is sometimes nearly all abstracted, and a mixture of dried

---

<sup>1</sup> *Journ. of a Tour in the Himālā Mountains*, by J. B. Fraser, 4to. Lond. 1820. p. 352.



blood and asphaltum introduced into the bag; or both the bag and the musk are artificial, and only scented with real musk. The blood of the animal itself is often injected into the bag of musk, while both are warm, and they then unite. The first of these adulterations is easily detected. The presence of blood may be suspected, if the musk, when held over the flame of a candle on a thin spatula, emits, as it inflames, a fetid smoke: and asphaltum is discovered by its melting, and running before it inflames, if heated on a spatula; whereas real musk inflames without running, and is converted into charcoal.<sup>1</sup> The artificial bags are known by the inner membrane, which lines the real musk bags, being deficient.

*Qualities.* — The odour of musk is aromatic, but peculiar, extremely powerful, and durable<sup>2</sup>: the taste is bitterish and heavy; and the colour a deep brown, with a shade of red. Exposed to heat, it burns with a white flame, and leaves a light spongy charcoal. Trituration with potass developes ammonia. Boiling water dissolves it partially, alcohol better and sulphuric ether still more completely. The watery *infusion* has a yellowish-brown colour, a bitterish taste, and the strong odour of the musk; and reddens infusion of litmus. Solutions of oxymuriate of mercury and of sulphate of iron produce with it copious precipitates; as does also infusion of yellow cinchona bark. Solution of nitrate of silver throws down a whitish precipitate, which, on exposure to the light, changes to a livid blue; and nitrate of mercury produces a brownish precipitate. The *alcoholic tincture* is of a reddish-brown colour, transparent, but with scarcely any odour of the musk. Water renders it milky and gives out the strong musk odour; but with the other tests it presents the same results as the watery infusion. The *ethereal tincture* has a deep brown colour; and, when evaporated on the surface of water, deposits a brown, tenacious, nearly insipid resin, and renders the water milky. The resinous matter has the musk odour in perfection; while the substance which occasions the turbidness of the water possesses the properties of extractive. From these results musk appears to contain albumen, gelatin, muriate of ammonia, phosphate of soda, and an uncombined acid; but the greater part of it consists of a resin combined with a volatile oil, and a mucilaginous extractive matter.

*Medical properties and uses.* — Musk is stimulant and anti-

---

<sup>1</sup> The formation of ammonia, when rubbed with potass, has also been mentioned as a test of the presence of blood; but the fixed alkalies have developed ammonia in the best specimens of musk we have ever seen.

<sup>2</sup> “A clean cork, which stopped a phial in which there was musk, which it seemed never to have touched, in 1712, smelled of musk more than twenty years, after.” *Alston's Let. on Mat. Med.* vol. ii. p. 542.



spasmodic. Aëtius is the first writer who mentions it as a medicine; but it did not come into general use in this country till the beginning of the sixteenth century. It raises the pulse without much augmenting the heat of the body, and has a remarkable power of resolving spasm, and increasing the energy of the brain and nerves. Hence it is very efficaciously given in typhoid fevers, when low delirium, subsultus tendinum, and hiccough supervene; and in combination with ammonia to arrest the progress of gangrene. Its beneficial effects in all spasmodic diseases are well established; and Cullen says, he can vouch for its powers in retrocedent gout, which in many instances he had seen suddenly relieved by large doses of musk.<sup>1</sup> It checks the vomiting in cholera, at the same time that it allays the tormina of the intestines. In epilepsy I have seen more benefit derived from musk in combination with calomel than from any other remedy; and I am inclined to attribute much of the disappointment which others have experienced, either to the remedy not having been genuine, or to the smallness of the dose. To obtain the full benefit of musk in this disease, the dose must be much larger than that which is usually given; it should be repeated at shorter intervals, and its use longer continued. In an old confirmed case, in which three and four fits were experienced daily, musk, given to the extent of ʒss. four times a-day, reduced the number of fits to one in three months. Upon the whole, I agree with Cullen “that musk is one of the most powerful antispasmodics we are acquainted with,” and regret that the high price of the drug necessarily limits very much its employment.<sup>3</sup>

As a local remedy, musk is said to be useful in atonic deafness when inserted into the ear with cotton; and it is recommended in the form of enema in the convulsions of children arising from the irritation of dentition.

Musk is best given in substance, in the form of bolus. The dose may be from grs. vj. to ʒj., repeated at intervals of six or eight hours.

Officinal preparations. *Mistura Moschi*, L. *Tinctura Moschi*, D. MYRISTICA. *Spec. Plant. Willd.* iv. 869.

Cl. 22. Ord. 13. Diœcia Monadelphia. Nat. ord. Lauri, Juss. G. 1851. Male. *Calyx* bell-shaped, trifid. *Corolla* none. *Filament* columnar. *Anthers* six or ten united.

Female. *Calyx* bell-shaped, trifid, deciduous. *Corolla* none. *Style* none. *Stigma* two. *Drupe*, a nut involved in an arillus (*Mace*,) with one seed.<sup>4</sup>

<sup>1</sup> *Mat. Med.* ii. 381.

<sup>2</sup> *Mat. Med.* l. c. 380.

<sup>3</sup> A mixture of musk and cinnabar in arrack is used by the Tonquinese as a remedy in hydrophobia. *Phil. Trans.* xlv. 78. No benefit has been derived from its use in this disease in this country, although it has been fairly tried.

<sup>4</sup> In our description, we have followed Gærtner, who denominates the fruit “*Bacca monosperma*,” although, in our translation of the generic character by Willdenow, we have not altered the term “*Drupe*.”



*Species* 1. *M. moschata*. The nutmeg-tree. *Med. Bot.* 2d edit. 698. t. 238. *Rumphius. Herb. Amboin.* ii. lib. 11. c. 5. t. 4.

*Officinal.* MYRISTICÆ NUCLEI, *Lond.* MYRISTICÆ MOSCHATÆ NUCLEI, INVOLUCRUM vulgo MACIS, *Edin.* NUX MOSCHATA; OLEUM ESSENTIALE, OLEUM EXPRESSUM, INVOLUCRUM MACIS DICTUM, *Dub.* Nutmeg; Mace; Oil of Nutmeg, essential and expressed; Oil of Mace.

*Syn. of the Nutmeg.* Noix muscade (*F.*), Moskatnuse (*G.*), Nooten-moskat (*Dutch*), Muskad (*Dan.*), Muskot (*Swed.*), Nosce moscata (*I.*), Neuz moscada (*S.*), Noz moscada (*Port.*), Jaëphial (*H.*), Jattipullum (*Cyng.*), Jouzbewa (*Pers.*), Jouzalteil (*Arab.*), Woh pala (*Jav.*), Jadikai (*Tam.*), Jayaphula (*Beng.*), Jatiphalo (*San.*), Pela (*Malay*), — *Of the Mace.* Moshat blumen (*G.*), Macis (*I.*), Macias (*S.*), Jawatri (*H.*), Jatipatri (*San.*), Jadiputrie (*Tam.*), Bunga-buæ-pala (*Malay*), Kambangpala (*Jav.*), Wassawasel (*Cyng.*), Bezbaz (*Pers.*). *The oil.* Jadikai tylum (*Tam.*), Jatipullum tail (*Cyng.*).

The nutmeg-tree is a native of the Molucca islands. It has, however, been nearly extirpated from the greater number of them by the narrow policy of the Dutch, and is cultivated at Banda<sup>1</sup>, and, also, at Bencoolen, in the island of Sumatra, where a sufficient quantity is reared to supply with mace and nutmegs the whole of Europe. It rises to the height of thirty feet, producing many erect branches, which, as well as the trunk, exude, when wounded, a red glutinous juice, and are covered with a smooth ash-coloured exterior bark. The leaves, which stand alternately on short petioles, are elliptical, pointed, undulated, entire, and obliquely nerved; of a bright green colour on the upper surface and greyish underneath, with an aromatic taste. The flowers are present at the same time with the fruit, inodorous, small, supported on axillary peduncles; and male and female on the same, and on separate trees: the calyx in both is fleshy, smooth, and divided at the edge into three, moderately spreading segments. There are no corollas. The filaments in the male flowers are short, united into one columnar bundle, inserted into the receptacle, and bear each a linear anther, which surrounds the upper half of the filament. The germ in the female is superior, oval, and crowned with a style terminated by two stigmas. The fruit is an elliptico-spheroidal, one-celled, superior berry, marked with a shallow longitudinal groove on one side, fleshy, smooth, one-celled, and the size of a small peach; the flesh is thick, rather solid, and finally dries up to a coriaceous crust, which opens at one side, and displays the nutmeg in its shell covered with an arillus, which is the officinal *mace*, and by it fixed to the bottom of the cell. The Mace is a fleshy, coriaceous, saffron, or yellow coloured substance, divided deeply into many slips, which closely invest the shell of the nutmeg. The kernel, which is the proper nutmeg, is of a

<sup>1</sup> This term includes six smaller islands, Neyra, Lenteira, Pulo-Aya, Goenenga, Apia, Polerona, and Rossengenia; but the three first only bear nutmeg-trees.



roundish oval form, marked on the outside with many vermicular furrows, within of a fleshy-farinaceous substance; variegated whitish and bay, and having a cavity at the bottom for the embryo.<sup>1</sup>

The nutmeg-tree produces fruit at the age of seven years; its productiveness is at the height at fifteen; and it continues to bear for seventy or eighty years, in the Mollucas. It yields three crops annually: the first in April, the second in August, and the third, which is the best, in December, yet the fruit requires nine months to ripen it. When it is gathered, the outer coriaceous covering is first stripped off, and then the mace carefully separated, is flattened by the hands in single layers, and dried in the sun. The nutmegs in the shell are next exposed to heat, not exceeding 140° of Fah., and to smoke for three months. Much care is necessary in drying them; as they require to be turned every second or third day. The criterion of due preparation is their rattling in the shell. They are then broken, and the kernels thrown into a strong mixture of lime and water, at Banda; but at Bencoolen they are simply rubbed over with dry lime; after which they are cleaned and packed up in casks and chests, smoked, and covered within with a coating of lime and water.<sup>2</sup> This process is necessary for their preservation, and with the same intention the mace is sprinkled with salt water. There are several varieties of the tree; but that denominated the Queen Nutmeg, which bears a small round nut, is the best. They are imported in chests which contain each from 100 to 140lbs. weight; the mace comes in chests also of different sizes; the essential oil, which is obtained in Banda by the distillation of the nuts, is brought in bottles and the expressed oil in stone jars. Nutmegs are frequently punctured and boiled in order to obtain the essential oil, and the orifices afterwards closed with powdered sassafras. The fraud is detected by the lightness of the nutmeg. Both the nutmeg and the mace are perforated by the larva of *Dermestes Surinamensis*.<sup>3</sup>

*Qualities.* — The nutmeg has a fragrant, agreeable, spicy odour and a warm aromatic taste. It is easily cut with a knife, but not very pulverulent. When cut transversely and examined by the microscope, the dark coloured veins which run through its substance appear to consist of cellular matter filled with oil, which is the active matter of the nutmeg. Alcohol and ether extract completely the active qualities of nut-

---

<sup>1</sup> *Gærtner de Fructibus*, t. 41.

<sup>2</sup> *Report on the Cultivation of Spices at Bencoolen, &c.* by J. Lumsdaine, Esq. — *Edin. Phil. Journ.* vol. vii. p. 127.

<sup>3</sup> *Leeuwenhoek, Epist.* 92.



meg. When the ethereal tincture, which is limpid, and of a golden-yellow colour, is evaporated on water, a small portion of volatile essential oil unites with the water, whilst a white, opaque, granular, sebaceous substance, heavier than water, which has much the appearance of the expressed oil, is deposited. When alcohol is digested on this substance, it dissolves very little of it, but becomes yellow, and acquires the qualities of a spirituous solution of the essential oil. The undissolved substance, if washed in water, is nearly insipid, melts at a temperature of  $150^{\circ}$ , and, on cooling, concretes into a translucent brittle cake which has the properties of wax. The part of the nutmeg insoluble in ether is chiefly gum and starch. In distillation with water, nutmegs yield  $\frac{1}{32}$  part of their weight of essential, volatile oil, and by expression one-third of a sebaceous, fixed oil.<sup>1</sup> Hence, the components of the nutmeg seem to be starch, gum, volatile oil, wax, and a fixed, fat oil. The *volatile oil* possesses the odour and taste of the nutmeg in a concentrated degree, is of a pale straw colour, limpid, transparent, and lighter than water. The *expressed oil*, which is erroneously called *oil of mace*, when first drawn, is limpid and yellow, but on cooling acquires the consistence of spermaceti and somewhat of the appearance of Castile soap, being whitish, mottled with reddish brown. It is imported wrapped up in flag leaves. Its odour is agreeable and slightly aromatic: its taste fatty, pungent, and bitterish. It appears to be a vegetable cerate, or a triple compound of fixed oil, volatile oil, and wax. Besides the genuine expressed oil, there are two other sorts found in the shops; one, which is said to come from Holland, of a paler colour, and in flat square cakes; and another, which is an artificial composition of suet, palm oil, and spermaceti, scented with a little volatile oil of nutmeg. *Mace* resembles the nutmeg in its odour and taste, but is more pungent and bitter. It is in lacinated, flexible, thin pieces, unctuous to the feel, and of a deep reddish-yellow colour. Alcohol and ether extract its active principles; and when the ethereal tincture is evaporated on water, a thick, deep yellow-coloured, very pungent, and odorous oil is left on drops on the surface of the water, with some resin: and a small portion of extractive is deposited, but no waxy granular matter.

*Medical properties and uses.*<sup>2</sup> — As the medical properties of nutmeg and mace depend on the essential oil they contain, they agree in these circumstances; and both are stimulant, carminative, and, in large doses, narcotic. Mace is more

<sup>1</sup> *Neumann's Chemistry*, 404.

<sup>2</sup> Avicenna first noticed nutmegs as a medicine



generally used as a culinary spice; but the Nutmeg and its volatile oil are in frequent use to cover the disagreeable taste of other medicines; and are, sometimes, ordered in cases of languors, vomiting and diarrhœa, and in flatulent colic. On account of the narcotic property of the oil, the Nutmeg should be cautiously employed in apoplectic and paralytic habits. In India, its dangerous effects have been frequently felt<sup>1</sup>; and, in this country, instances have occurred in which the Nutmeg, taken in large quantity, produced drowsiness, great stupor and insensibility; and, on awakening, delirium which alternated with sleep for several hours.<sup>2</sup> The volatile oil is sometimes used as an external stimulant: — the expressed oil is seldom employed for any other purpose. The dose of the Nutmeg and the Mace is from grs. v. to ʒj; that of the volatile oil, ʒij, to ʒvj.

Official preparation. *Spiritus Myristicæ*, L. E. D.

MYROXYLON. *Spec. Plant. Willd.* ii. 546.

Cl. 10. Ord. 1. Decandria Monogynia. Nat. ord. Lomentaceæ, Linn. Leguminosæ, Juss.

G. 829. *Calyx* bell-shaped, five-toothed. *Petals* five, the upper one larger than the others. *Germen* longer than the corolla. *Legume* with one seed only at the point.

*Species* 1. *M. peruiferum*. Sweet-smelling Balsam-tree. *Hernandez Nova Plant. &c. Mexican. Hist. fol.* 51. *cum figura*.

*Official*. BALSAMUM PERUVIANUM, *Lond.* MYROXYLI PERU-IFERI BALSAMUM, *Edin.* Peruvian Balsam.

*Syn.* Baume de Perou (*F.*), Peruvianischer Balsam (*G.*), Peruviaanische Balsam (*Dutch*), Swart Perubalsam (*Swed.*), Balsamo del Peru (*I.*), Balsamo de Quinquica (*S.*).

The Peruvian-balsam tree is a native of the warmest provinces of the continent of South America<sup>3</sup>; growing in the mountains of Panatalmas, in the forests of Puzuzu, Munā, Cuchero, Paxaten, and many other places near the river Maranon, blossoming in August, September and October. It is a very beautiful tree, with a smooth, thick, straight trunk, covered with a grey, coarse, compact, heavy bark, which is interiorly of a straw colour and very resinous, as is every part of the tree. The branches extend almost horizontally. The leaves are alternate and abruptly pinnate. The leaflets in two pairs, nearly opposite, petiolate, ovato-lanceolate, with the apex lengthened and somewhat blunt and emarginate, entire, shining, veined and very smooth. The midrib, which runs through the whole length of the under surface of the leaf, is raised and pubescent; the common petiole is round and pubescent. The flowers are scattered, on axillary, erect

<sup>1</sup> Bontius de *Medicina Indorum*, 20.

<sup>2</sup> Cullen, *Mat. Med.* ii. 204.

<sup>3</sup> Mutis discovered it, and sent a branch of the tree to the younger Linnæus, about the year 1781.



racemes, longer than the leaves. The peduncles are roundish and pubescent; each supported by a small, erect, ovate, concave bracte, appearing to the naked eye like a tubercle, the pedicels are erect. The calyx is bell-shaped, dark green, divided into five, small, nearly equal teeth, but with one of them so far separated as to be found under the germen. The corolla consists of five, white petals; four narrow, equal, lanceolate, and larger than the calyx; the fifth reflex, broad and double the size of the others. The stamens are ten, inclining and inserted into the calyx; bearing elongated, sharp-pointed, sulcated anthers. The germen is oblong, pedicillated, inclining; the style short, subulate and crooked, crowned with a simple stigma. The seed vessel is a straw-coloured, club-shaped, somewhat curved, pendulous legume, globular near the top and terminated by the curved style. It contains, in a cell in the curved part, one seed only, which is crescent-shaped and projects from the cell. The substance of the leaves is full of translucent, linear points, like the leaf of the orange tree.

This tree is called *quinquino* by the natives, who use the bark as a perfume. The balsam, which is procured in a liquid state, from incisions made early in the spring, is collected in bottles, and is called *white liquid balsam*. What is found in the shops, is obtained by boiling the twigs in water. It is imported in jars, each containing from twenty to forty pounds weight. When the Indians collect the white balsam in calabashes, which is the case in Carthagena and in the mountains of Tolu, it condenses and hardens, and forms dry, white balsam, or the balsam of Tolu. Ruiz says there is no difference in these three balsams, excepting in name, colour and consistence. A mixture of resin and some volatile oil with benzoin is often sold for Peruvian balsam; and the fraud is not easily detected.

*Qualities.* — The Balsam which we receive has a fragrant, aromatic odour, much resembling that of benzoin, with a warm bitterish taste, leaving a slight sensation of burning in the throat after it is swallowed, with some degree of sweetness. It is viscid, of a deep, reddish-brown colour, being that which is obtained by boiling the twigs, and of the consistence of fluid honey. Water boiled on the balsam becomes acidulated, and deposits on cooling, crystals of benzoic acid. In distillation with water, a small portion of a volatile, limpid oil comes over, and benzoic acid sublimes in the neck of the retort. Its remaining matter is a resin. Ether, in small quantity, dissolves it readily and completely; alcohol also dissolves it, but the quantity of menstruum must be con-



siderable. Sulphuric acid converts it into artificial tannin and charcoal. Treated with nitric acid, some prussic acid is evolved, benzoic acid sublimes, and the residual matter is artificial tannin.<sup>1</sup> The alkalies and their carbonates form with it thick masses, which, on the addition of sulphuric acid, lets fall a resinous matter, and benzoic acid crystallizes. Hence Peruvian balsam appears to consist chiefly of resin, volatile oil, and benzoic acid: but according to Stoltze the oil is of a peculiar nature, differing from volatile oil. The result of his analysis is that 1000 parts of the balsam consist of 24 of brown, nearly insoluble resin, 207 of soluble resin, 690 of the above oil, 64 benzoic acid, and 6 of extractive matter.

*Medical properties and uses.* — Balsam of Peru is stimulant and tonic. It has been regarded as expectorant also, and recommended in catarrh and other pulmonary affections; but it is contra-indicated wherever any inflammatory action is present; and to its stimulant operation on the pulmonary exhalants we may ascribe its use in chronic asthma and old obstinate coughs.<sup>2</sup> In gleet, leucorrhœa, palsy, and chronic rheumatism, its tonic powers have proved beneficial; as well as in many other cases of debility. It may be given to the extent of fʒj for a dose. As a local stimulant it is employed externally with great advantage for cleansing and stimulating foul and indolent ulcers; and a mixture composed of ʒj of the balsam and ʒiij of ox gall, I have found extremely useful when dropped into the ear every day, after syringing with a solution of soap, in foetid discharges of the ear.

MYRRHA.<sup>3</sup> *Lond. Edin.* MYRRHA; GUMMI-RESINA. *Dub.* Myrrh, a gum-resin.

*Syn.* Myrrhe (*F.*), Myrrhen (*G.*), Mirra (*I.*), Mirra (*S.*), Murra (*Russ.*), Murr (*Arab.*), Ból (*H.*), Vóla (*San.*), Valatipolum (*Tam.*), Manisan lebah (*Malay*), Madu (*Jav.*).

The tree or plant which produces this gum-resin is a native of the eastern coast of Arabia Felix, and of Abyssinia, growing, according to Mr. Bruce's account, behind Azab, along the coast towards the straits of Babelmandel. It is undescribed by naturalists; and the conjectures of Mr. Bruce in favour of its being a *Mimosa* are by no means satisfactory.<sup>4</sup> The appearance of the best myrrh, as we receive it, affords reasons for supposing that it is an exudation from the plant. It is imported in chests, each containing from one to two hundred

<sup>1</sup> Hatchet. *Phil. Trans.* 1806. *Thomson's Chemistry*, iv. ed. v. 126.

<sup>2</sup> Sydenham gave it in phthisis.

<sup>3</sup> *Σμυρρα* Dioscoridis. The name *Myrrha*, used by Hippocrates, is derived from *μυρον*, an ointment. Professor Verey (*Journ. de Pharm.* 1820.) derives it from the Phœnician word *mor* or *mur*. Myrrha, the daughter of Cinyras, king of Phœnicia, was metamorphosed into a tree, for being criminally in love with her father.

<sup>4</sup> *Phil. Trans.* lxx. 413.



weight. The Abyssinian myrrh comes to us through the East Indies, while that produced in Arabia is brought by the way of Turkey.

*Qualities.* — Myrrh has a peculiar, rather fragrant odour, and bitter, aromatic taste. It softens in the mouth, adheres to the teeth when chewed, and is in small, irregularly shaped pieces, which can scarcely be called tears; they are translucent, of a reddish-yellow colour, brittle, breaking with a resinous fracture and easily pulverized. It does not melt when heated, and is not very inflammable. Its specific gravity is 1.360.<sup>1</sup> Such are the characters of good myrrh; but it is often opaque, mixed with many impurities, and either white or of a dark colour approaching nearly to black, with a disagreeable odour, in which case it should be rejected.

Myrrh is partially soluble in water, alcohol, and ether. In distillation with water, it yields an oil heavier than water. When it is triturated with very soft or distilled water, nearly the whole appears to be dissolved, forming an opaque, yellowish solution; but the greater part is deposited by rest, and not more than one-third of the gum-resin is actually dissolved. The alcoholic tincture is rendered milky and opaque when mixed with water, but no precipitate appears. Braconnot asserts, that 100 parts of myrrh consist of 23 of resin and 77 of gum<sup>2</sup>; but my experiments lead to a somewhat different conclusion, and accord more with those of Pelletier, who found the proportions to be, 34.68 of resin and 66.32 of gum. Ether, digested on powdered myrrh, dissolved three parts in eight, and the tincture, evaporated on water, deposited two grains and a half of very bitter resin and half a grain of extractive matter, which also tasted bitter. The part insoluble in the ether was nearly all soluble in water, and afforded a solution resembling that of acacia gum; but differed from it in being precipitated by solutions of muriate of mercury and of superacetate of lead. Myrrh triturated with crystallized alkalies is reduced to the form of a tenaceous fluid. When treated with nitric acid it yields oxalic acid. Hence myrrh seems to consist of resin, essential oil, extractive, and mucus, rather than gum.

*Medical properties and uses.* — Myrrh is tonic and expectorant. In moderate doses it stimulates the stomach, promoting the appetite and digestion; but, in large doses, increases the frequency of the pulse and augments the general heat of the body.<sup>3</sup> As a tonic, it is efficaciously given in

<sup>1</sup> Brisson.

<sup>2</sup> *Annales de Chimie*, lxxviii. 52.

<sup>3</sup> Cullen, *Mat. Med.* ii. 193.



cases of debility<sup>1</sup>, as, amenorrhœa, chlorosis and convalescencies; and in phthisis pulmonalis, when the inflammatory symptoms and hectic fever do not run high. Its use in phthisis has, indeed, been condemned by several physicians of great repute<sup>2</sup>; but, when there is an evident ulceration of the lungs without much hectic, and the patient's strength is considerably reduced by the quantity of the expectorated matter, the proper exhibition of myrrh is certainly productive of much benefit. In the first-mentioned diseases, it is advantageously combined with aloes, cinchona, or other bitters and chalybeates; and in phthisis, with nitre, digitalis, opium, camphor and the sulphate of iron or of zinc. Combined with oxide of zinc, it has been found extremely useful in the peculiar cough which sometimes accompanies pregnancy, and continues after abortion. As an expectorant, it is often employed in humoral asthma and chronic catarrh: and with the same view also has been given in phthisical affections; but as it cannot be employed with propriety in pulmonic cases, where there is much inflammatory action or hectic present, any advantage derived from its use in phthisis probably depends altogether on its tonic operation counteracting the exhaustion which is produced by a copious purulent expectoration. As a local stimulant, the alcoholic solution of myrrh diffused in water is used as a lotion in a spongy state of the gums, and for correcting the foetid discharge of vitiated ulcers, particularly when connected with caries of the bone; and as a gargle in cynanche maligna.

Myrrh is administered in substance, or in the form of watery infusion, or of tincture properly diluted. The watery infusion is much less stimulant than any of the other preparations. A watery extract is ordered in some foreign pharmacopœias, and preferred by many physicians, from an idea that it is less heating than the gum-resin; but it is equally bitter, and is perhaps not different from a diminished dose of the myrrh.

Officinal preparations. *Tinctura Myrrhæ*, L. E. D. *Tinctura Aloes et Myrrhæ*, E. *Tinct. Aloes Ætheria*, E. *Mistura Ferri comp.* L. *Pilulæ Aloes cum Myrrha*, L. E. D. *Pilulæ Ferri cum Myrrha*, L. *Pil. Galbani comp.* L. D. *Pil. Assafœtidæ comp.* E. *Pil. Rhei composita*, E.

MYRTUS. *Spec. Plant. Willd.* ii. 967.

Cl. 12. Ord. 1. Icosandria Monogynia. Nat. ord. Hesperideæ, Linn. Myrti, Juss.

<sup>1</sup> It was formerly used for the cure of quartans; ʒj, in a glass of warm Cretan wine, being given an hour before the paroxysm, and thrice repeated before the accession. Mathiolus, cap. 67, says "Hoc medicamento ego ipse curatus sum."

<sup>2</sup> Cullen. Fothergill.



G. 973. *Calyx* five-cleft, superior. *Petals* five. *Berry* two or three-celled, many-seeded.

*Species* 28. *Myrtus Pimenta*. Pimenta, or All-spice tree. *Med. Bot.* 2d edit. 541. t. 194.

*Officinal.* PIMENTÆ BACCÆ, *Lond.* MYRTÆ PIMENTÆ FRUCTUS, *Edin.* PIMENTO (*Piper Jamaicaense*); BACCÆ, *Dub.* Pimenta Berries. Jamaica Pepper.

*Syn.* Poivre de Jamaïque (*F.*), Nelkenpfeffer (*G.*), Jamaika pepper (*Dutch*), Krydd peppar (*Swed.*), Pimenti (*I.*), Pimienta (*S.*).

This tree is a native of South America where it is called Pumake (in the Maypure language), and of the West India islands. It grows in great plenty on the hilly parts, on the north side of the island of Jamaica; flowering in June, July and August, and soon afterwards ripening its fruit. It is a handsome tree, rising in height about thirty feet, straight, branching, and covered with a very smooth gray bark. The leaves, which are supported on footstalks at the ends of the twigs, are elliptical, pointed, of different sizes, but the largest are five inches long, and two broad in the middle, smooth, thin, entire, shining and of a deep-green colour; the flowers are produced in terminal branches, or rather trichotomous panicles: the calyx is four-cleft; the petals four, reflected, of a pale green colour, inclosing many longer, spreading filaments of the same colour, supporting pale-yellow, roundish anthers: the fruit is a spherical berry, crowned with the persistent calyx; when ripe, it is black, or dark purple, smooth, shining, and bilocular, with the seeds enveloped in a moist, green, pungent, aromatic pulp.<sup>1</sup>

The fruit, which is the part of this plant medicinally used, is gathered before it is ripe<sup>2</sup>, and exposed to the sun for many days, spread thin upon cloths. They require to be frequently turned and carefully preserved from the dews. By degrees, under this management, they become wrinkled, and change from green to a brown colour; after which, they are packed in bags and hogsheads for the European market. The more fragrant and smaller they are, the better are they accounted.<sup>3</sup>

*Qualities.*—Pimenta has an aromatic, agreeable odour, resembling that of a mixture of cinnamon, cloves, and nutmegs, with the warm pungent taste of the cloves; qualities which reside chiefly in the cortical part of the dried berry. Water, alcohol, and ether extract its virtues. The watery infusion is of a brown colour, and reddens litmus infusion. With solution of sulphate of iron it immediately strikes a deep-black colour, and slowly lets fall a precipitate. Nitrate of mercury

<sup>1</sup> Sloane, *Phil. Trans.* xvii. 462.

<sup>2</sup> When the berries ripen, they lose much of the aromatic warmth for which they are esteemed, and acquire a taste similar to that of juniper berries. <sup>3</sup> Sloane, l. c.



precipitates it of a yellowish-brown; superacetate of lead of a dirty-green; and nitrate of silver, of a deep reddish-brown colour. It is also precipitated by infusion of yellow cinchona bark. The sulphuric and muriatic acids redden it, and throw down pale, rose-coloured precipitates. The nitric acid forms no precipitate, but gives the infusion a yellow hue. The alcoholic tincture is rendered milky, and slowly precipitated by water: the ethereal, when evaporated on water, deposits drops of a greenish-yellow volatile oil, a pellicle of a pungent, nauseous tasted resin and some extractive. Hence Pimenta appears to contain a volatile oil, resin, extractive, tannin and gallic acid.

*Medical properties and uses.*—Pimenta is stimulant and tonic. It is useful as an adjunct to bitters in dyspepsia attended with much flatulence, and in arthritic and hysterical affections. The watery infusion of it sweetened with sugar, and with the addition of a little milk, is very readily taken by children; and is an excellent cordial in malignant measles, scarlatina, confluent small-pox, and the other exanthemata, when the fever assumes the typhoid type. But the principal use of Pimenta in medicine is to cover the disagreeable taste of other remedies, or to give them warmth. The dose of the berries is from gr. v. to ℥ij, in powder or swallowed in their entire state.

Officinal preparations. *Aqua Pimentæ*, L. E. D. *Oleum Pimentæ*, L. E. D. *Pilula Opiata*, E. *Syrupus Rhamni*, L.

NICOTIANA. *Spec. Plant. Willd.* i. 1014.

Cl. 5. Ord. 1. Pentandria Monogynia. *Nat. ord.* Luridæ, Linn. Solaneæ, Juss.

G. 379. Corolla funnel-shaped with the border plaited. Stamens inclined. Capsules two-valved, two-celled.

Sp. 1. N. *Tabacum*.<sup>1</sup> Tobacco. *Med. Bot.* 2d ed. 208. t. 77.

Officinal. TABACI FOLIA, *Lond.* NICOTIANÆ TABACI FOLIA, *Edin.* NICOTIANÆ FOLIA, *Dub.* Tobacco leaves.

<sup>1</sup> This plant was first discovered by the Spaniards in Yucatan in 1520, and was there called *petun* or *pete-ma*. Humboldt says it has been cultivated from time immemorial by the native people of the Oroonoko; and was smoked all over America at the time of the Spanish conquest. He found only two of the species cultivated in Europe, the *N. paniculata* and *N. glutinosa*, growing wild; but the *N. loxensis* and *andicola*, which he found on the Andes, 1850 toises of elevation, closely resemble the *tabacum* and *rustica*. It was transported to the West Indies and North America; and brought to Europe by Hernandez de Toledo, who came from Florida to Portugal in the beginning of the 16th century. The seeds were sent from Portugal to Catherine de Medicis by Jean Nicot, an agent of Francis II., after whom it received its generic name *Nicotiana*: the specific appellation being taken from *tabac*, the name of an instrument used by the natives of America in smoking the herb. The following are the names by which it is known in America: *yeth* in the Mexican or Azteek tongue; *sema* in Algonkiu; *oyugoua* in the Huron; in the Peruvian it is *sayri*; in Chiquito, *pâis*; in Vilela, *tusup*; Mbaja, *nalodagadi*; Moxo, *sabare*; Omagua, *potema*; Tumanac, *cavai*; Maypure, *jema*; and Cabre, *sema*. — Humboldt, *Person. Narr.* vol. v. p. 666.



*Syn.* Tabac (*F.*), Taback (*G. Dutch*), Tobak (*Dan. Swed.*), Tabak (*Pol.*), Tabacco (*I.*), Tobaco (*S. Port.*), Poghei elly (*Tam.*), Bujjer hany (*Arab*), Doorkole (*Cyng.*), Tam braco (Malay), Tambroco (*Jav.*), Tambácu (*H.*), Támracuta (*San.*).

Tobacco is an annual plant, a native of America, and partially cultivated in Europe; flowering in July and August. The root is large and fibrous, and sends up an erect, branching stem, about four feet in height, round, villous, slightly viscid and furnished with numerous, large, alternate, entire, pointed leaves, the lowermost of which are about two feet long and four inches broad; they are sessile, a little decurrent, with a strong midrib, and of a pale green colour on the upper surface and still paler underneath. The flowers are in large terminal panicles, with long, linear-pointed bractes at the base of each division: the calyx is bell-shaped, obscurely pentangular, villous, slightly viscid, and cleft into five, acute, erect segments: the corolla is very viscid, its tube twice the length of the calyx, of a pallid-greenish hue, and swelling into an oblong cup, which expands into five, pointed, plaited, pale-red or rose-coloured segments: the stamens are the length of the tube of the corolla, and support awl-shaped, compressed, oblong anthers: the style, which is the length of the corolla, and crowned with a capitate, slightly cleft stigma, rises from a conical germen, that changes to an ovate capsule containing many, reniform, small seeds, and opening at the apex.

Tobacco was at one period raised to a considerable extent in Yorkshire<sup>1</sup>; but the cultivation of it for the purposes of trade has been long prohibited; and this country, as well as the greater part of Europe, is chiefly supplied from Virginia, where the plant is cultivated in the greatest abundance. There are two varieties of this species, known by the name of Virginian tobacco, a broad and a narrow-leaved sort; but they do not differ in their medical properties. In Virginia, the plant is not allowed to attain its full height, but is topped whenever a certain number of leaves is thrown out. It is cut down in August, and the plants hung up in pairs, in sheds, to dry, after which the leaves are separated from the stem, bound up in bundles, and packed in the hogsheads in which they are exported.

*Qualities.* — The recent leaves possess very little odour or taste; but when dried their odour is strong, narcotic and somewhat foetid; their taste bitter and extremely acrid. When well cured, their colour is yellowish-green. They emit sparks in burning, and give out a suffocating smoke; and

---

<sup>1</sup> It was first cultivated in England in 1570, according to Lobel's account.



when distilled, yield an essential oil of a green colour, on which their medicinal properties are supposed to depend, and which is said to be a very virulent poison.<sup>1</sup> This oil is dissipated by the long coction of tobacco with water; yet, in distillation with ether, water, or alcohol, no oil comes over. By infusion, however, it yields its active principles to both these fluids. Its deflagration shows the presence of nitrate of potass; and Bouillon la Grange discovered muriate of potass in its inspissated juice.<sup>2</sup> According to Vauquelin, tobacco appears to contain albumen or gluten, supermuriate of lime, acetic acid, nitrate and muriate of potass, muriate of ammonia, a red matter soluble in alcohol and water, a green fecula, and a peculiar substance, on which the properties of the plant appear to depend, and which has been therefore named *nicotin*.<sup>3</sup>

*Medical properties and uses.*—Tobacco is narcotic, sedative, emetic, diuretic, cathartic, and errhine, whether it be taken into the stomach, or externally applied. The three first mentioned properties are sufficiently obvious, even from the effects which smoking or chewing it produce on persons unaccustomed to its use.<sup>4</sup> From Mr. Brodie's experiments, the infusion of Tobacco produces its effect on the heart, through the medium of the nerves. The symptoms are,

<sup>1</sup> The poisonous effects of this oil are very powerful. Mr. Barrow, speaking of the use which the Hottentots make of tobacco oil for destroying snakes, says, "A Hottentot applied some of it from the short end of his wooden tobacco-pipe to the mouth of a snake, while darting out his tongue. The effect was instantaneous as an electric shock: with a convulsive motion that was momentary, the snake half untwisted itself, and never stirred more; and the muscles were so contracted, that the whole animal felt hard and rigid, as if dried in the sun." *Travels in Africa*, p. 268.

<sup>2</sup> *Journal de Physique*, xxxix. 193.

<sup>3</sup> This substance is colourless, acrid, has the odour of tobacco, and like it occasions violent sneezing. It is volatile, poisonous, and produces colourless solutions with alcohol and water, from which it is thrown down by tincture of nutgalls. Vauquelin regards it as approaching the volatile oils in its properties. Vide *Ann. de Chimie*. tome lxxi. p. 139.

<sup>4</sup> The custom of smoking tobacco was introduced into England by Sir Walter Raleigh; and was at one time extremely prevalent, but is now confined chiefly to the lower class of the people. In some parts of Europe, however, it is still regarded as the greatest solace and pleasure of the luxurious. It is a curious fact, that in England an edict was published against its use, the reason of which was probably the apprehension thus stated by Camden—"Anglorum corpora in barbarorum naturam degenerasse, quum iidem ac barbari delectentur." *Annal. Eliz.* p. 143. Urban VIII. anathematized those who used it in churches; and in Constantinople, where its use is now so general, the custom was in the beginning of the 17th century thought so ridiculous and hurtful, that any Turk who was found smoking was conducted in ridicule through the streets with a pipe transfixed through his nose. Tobacco, which has been introduced into the Sandwich Islands by Europeans, "is now," says Kotzebue, (vide *Voyage of Discovery*) "so generally used, that young children smoke before they learn to walk, and grown up people have carried it to such an excess, that they have fallen down senseless, and often died in consequence."



very severe sickness, headach, extreme debility, cold sweats, and sometimes, even convulsions. The production of such a state of the habit, however, being useful for relieving violent spasmodic constriction, tobacco is advantageously employed in obstinate constipation, ileus, suppression of urine, and incarcerated hernia, when other remedies fail of affording relief. The smoke is either thrown into the rectum by means of a pair of bellows of a peculiar construction, or an infusion of the leaves is exhibited in the form of enema.<sup>1</sup> From its narcotic power also, the smoking or chewing Tobacco has been found useful in allaying the pain of toothach; and smoking it has been recommended, and in some instances found useful in shortening and rendering more supportable the paroxysm of spasmodic asthma. The infusion has been used as an emetic; but the practice cannot be recommended: and, notwithstanding the success of Dr. Fowler<sup>2</sup>, who employed it in dropsy and dysury, its general effects are too violent for internal exhibition, and it is not equal as a diuretic either to squill or foxglove, which are more manageable remedies. In dysury, however, as Dr. Pearson has observed, its antispasmodic properties are of advantage, and consequently its use in that complaint is less objectionable.<sup>3</sup> The external application of a strong infusion of Tobacco, or of a cataplasm of the moistened leaves themselves, is sometimes employed as a local stimulant in porrigo, scabies, and some other cutaneous eruptions; but, even in this mode of using it, Tobacco is apt to induce the same virulent effect as when it is internally administered in large doses.

But Tobacco is chiefly employed as a sternutatory, and is the basis of all the kinds of *snuff* generally used.<sup>4</sup> The powdered leaves, when snuffed up the nostrils of those unaccustomed to the use of snuff, excites vehement sneezing, and promotes a considerable discharge from the nostrils, answering all the purposes for which errhines are employed. As a luxury, snuff has been used upwards of two hundred years in Britain, and has been taken in great quantities without any perceptible bad consequence; although it has been asserted that its immoderate use weakens the sight, produces lethargy, and gives a tendency to apoplexy. After the use of it has become habitual, it cannot be relinquished without consi-

<sup>1</sup> The native doctors in India apply the leaves to the orifice of the anus. Vide *Ainslie's Mat. Med. of Hindostan*, 4to. p. 48.

<sup>2</sup> *Med. Reports on the Effects of Tobacco, &c.* <sup>3</sup> *Practical Synopsis, &c.* 228.

<sup>4</sup> In the manufacture of snuff, salt, urine, muriate of ammonia, and even powdered glass, are added to the tobacco. The difference of flavour depends, in some, on the species of *Nicotiana* employed, but chiefly on the preparation of the leaves, and these having undergone fermentation. *Macouba* derives its flavour from the leaves being fermented with an addition of the best cane juice.



derable risk, arising from the suspension of the artificial discharge it produces, as Dr. Cullen observed from his own experience.<sup>1</sup>

The London College has given a formula for an infusion proper to be used as an enema; as a diuretic, the infusion employed by Dr. Fowler is made with ℥i. of the dried leaves, and Oj. of boiling water, and given in doses of ℥℥. to ℥℥℥., twice a day.

Officinal preparations. *Infusum Tabaci*, L. *Vinum Nicotianæ Tabaci*, E.

OLEA. *Spec. Plant. Willd.* i. 44.

Cl. 2. Ord. 1. Diandria Monogynia. *Nat. ord.* Sepiariæ, Linn. Jasmineæ, Juss.

G. 36. *Corolla* four-cleft, with subovate segments. *Drupe* one-seeded.

*Species* 1. *O. Europæa*.<sup>2</sup> European Olive. *Med. Bot.* 2d edit. 280. t. 93. *Sibthorp Flora Græca*, t. 3.

*Officinal.* OLIVÆ OLEUM, *Lond.* OLEÆ EUROPÆÆ OLEUM FIXUM, *Edin.* OLEUM OLIVARUM, *Dub.* The oil of the olive.

*Syn.* Huile d'Olive (*F.*), Olivenhöl (*G.*), Olyfoly (*Dutch*), Bamolja (*Swed.*), Olio d'Ulivo (*I.*), Azeite (*S.*), Zeet (*A.*).

The Olive tree is a native of the south of Europe and the north of Africa, where it is named Zituna; but it is cultivated in France, Spain, and Italy. It has been raised in the open air in England, but its fruit, it is said, has never been ripened.<sup>3</sup> It grows upon the most rocky, calcareous soil, seldom exceeds twenty feet in height, and has a solid, upright, much-branched stem, covered with a grey bark: the leaves are evergreen, opposite, spreading, nearly sessile, stiffish, lanceolate, from two to three inches long, and half an inch broad in the middle, with the margin a little turned back; of a full-green colour, smooth and even on the upper surface, and pale on the under. The flowers are in opposite, axillary clusters, half the length of the leaves, on short flower-stalks, with small, concave, obtuse, hoary bractes: the calyx is deciduous, four-cleft, and regular: the corolla white, four-parted, regular, spreading, with ovate, obtuse, obscurely three-nerved segments; the stamens are shorter than the corolla, divaricated, supporting large pale-yellow, elliptical anthers; the stigma is bipartite on an erect style, rising from a roundish, superior germen, the fruit is a smooth oval plum or *drupe*, about three-fourths of an inch in length, and half an inch in diameter; of a deep-violet colour when ripe, whitish and fleshy within, bitter and nauseous, but replete with a bland oil<sup>4</sup>; and covering an osseous, oblong, pointed, rough nut.

<sup>1</sup> *Materia Medica*, ii. 437.

<sup>2</sup> Ελαια αγραια Dioscoridis.

<sup>3</sup> *Miller's Gardener's Dictionary*, ed. 1797, art. *Olea*. I have been informed, however, that it has ripened its fruit in Devonshire.

<sup>4</sup> The unripe fruit, when pickled in a strong solution of common salt, is a well known luxury of the table.



There are several varieties of the Olive tree, of which the variety  $\gamma$ , or *longifolia* of Willdenow, is most esteemed, as affording the best oil. The young plant bears at two years old, and at six years is in full bearing: but the best oil is procured from the fruit of trees, which have been grafted. The value of the tree continues to increase until it have passed its one hundredth year; after which it declines. The mode of obtaining the oil from the ripe fruit was known very early in Egypt; and it is chiefly for this purpose that the tree is now cultivated in Spain, Provence, and Italy. To procure the oil, the ripe fruit is gathered in November, and immediately bruised in a mill, the stones of which are set so wide as not to crush the nut. The pulp is then subjected to the press in bags made of rushes; and, by means of a gentle pressure, the best oil which is called virgin oil, flows first: a second sort is got by breaking the marc, moistening it with warm water and returning it to the press; and, lastly, a very inferior kind is obtained, either by boiling the magma, or by breaking, moistening and fermenting it in large cisterns, and again submitting it to the full force of the press. When the olive is not sufficiently ripe, the recent oil has a bitterish taste, and when too ripe it is fatty. After the oil is drawn, it deposits by standing a white, fibrous, albuminous matter; from this the clear oil is poured off, and a second deposition takes place; after which, if put into clean, glass flasks, there is no further alteration.<sup>1</sup>

The best oil is made in Provence, its excellence arising from the Olives being carefully cleaned and garbled; but what we receive in this country comes from Lucca and Florence. Sicily also furnishes some, but it has a resinous flavour.<sup>2</sup> Good oil has lately been brought from Samos. It is imported in jars, half-jars, and what are called half-chests, which are wooden packages containing flasks.

*Qualities.* — Pure olive oil is an insipid, inodorous, pale greenish-yellow coloured, viscid fluid; unctuous to the feel; inflammable, incapable of combining with water and nearly insoluble in alcohol. It is fixed in any temperature under 600°, suffering considerable expansion, but not evaporating; and congeals at 38° of Fahrenheit. It is the lightest of the fixed oils, its specific gravity being 0.9153. When kept for a great length time, or much exposed to the air, its components<sup>3</sup> are partially separated, the sebacic acid and water are

<sup>1</sup> A very old olive tree, near Gerecomio, yielded 240 English quarts of oil, in 1809. *Three Months near Rome*, by Maria Graham, p. 49.

<sup>2</sup> This flavour has been ascribed to the Sicilian olives being grown on dry hilly situations. *Galt's Letters from the Levant*, 8vo. p. 129. <sup>3</sup> Vide *Expressed Oils*.



formed, and the oil acquires a disagreeable smell and sharp taste, becomes thick, brown-coloured, and is then said to be rancid. The rancidity is hastened by heat and by the admixture of Poppy oil, with which it is often adulterated. The purity of Olive oil is readily discovered by adding to it solution of nitrate of mercury<sup>1</sup>, which solidifies Olive oil, but leaves the oil of poppy, linseed, or other grains liquid.

*Medical properties and uses.*—Olive oil is demulcent, relaxant, and laxative. It is used internally as a demulcent in catarrh and other pulmonary affections, diffused in water by means of mucilage; and is also given, internally, in large quantities, to mitigate the action of acrid substances, as some poisons, taken into the stomach; and in cases of worms. Externally applied, it is a very useful relaxant, and instead of stopping up the cutaneous exhalants, appears to promote the excretion of sweat; on which account it has been employed with great advantages in frictions in the commencement of plague. The body is ordered to be very briskly rubbed all over with a clean sponge dipped in warm olive oil: copious perspiration generally follows, and the operation must be repeated once a day until symptoms of recovery appear. Mr. Jackson relates, that the Coolies, who are employed in the oil stores at 'Tunis, smear themselves all over with oil, and are seldom afflicted with the plague when it rages in that city<sup>2</sup>; an effect which may be owing to the oil forming a coating to the skin, so that it cannot come directly in contact with the contagion. Frictions with it are useful in ascites.<sup>3</sup> It is, however, more generally used as a vehicle for more active substances, in the formation of embrocation: thus, it is an excellent solvent of opium, which can, through its means only, be used in frictions with any advantage.<sup>4</sup> It is also used as an injection in gonorrhœa and as an adjunct to glysters in dysentery and intestinal abrasions. It is extensively used in pharmacy, in the composition of ointments, cerates and plasters.

The dose of Olive oil is from fʒj. to fʒj., triturated with mucilage, or mixed with water by means of a few drops of solution of potass. In cases of poisons or of worms, as much may be given as the stomach can bear.

ONISCUS. *Syst. Nat. Gmelin.* v. 3009.

*Cl.* 5. *Ord.* 7. Insecta Aptera.

<sup>1</sup> This nitrate is prepared by dissolving, in the cold, 6 parts of mercury in  $7\frac{1}{2}$  of nitric acid, sp. grav. 1.36.

<sup>2</sup> *Reflections on the Commerce of the Mediterranean*, p. 64.

<sup>3</sup> Lord Bacon, speaking of Inunction, says, — “Ante omnia igitur usum olei vel olivarum vel amygdali dulcis, ad cutem ab extra unguendum, ad longævitatē conducere existimamus.” *Opera*, fol. 1665. p. 536.

<sup>4</sup> The nostrum called *Roche's Embrocation*, for hooping cough, consists of olive oil, with about half its quantity of the oil of cloves and oil of amber.



G. 272. *Jaw* truncated, toothed. *Lip* bifid. *Palpi* unequal. *Feelers* bristly. *Body* oval. *Feet* fourteen.

Sp. 14. *O. asellus*.<sup>1</sup> Slaters.

*Officinal*. MILLEPEDÆ; SPIRITUS VINI VAPORE ENECATÆ, *Dub*. Slaters killed by the vapour of spirit of wine.

*Syn*. Cloporte (*F.*), Kellerwürmer (*G.*).

These insects are found on roofs of houses, old walls and under stones; they are rather more than half an inch in length, whitish on the belly, with seven pairs of legs, each terminated by a sharp horny claw. The head is somewhat pyramidal, and furnished with two articulated feelers; and the whole of the animal on the upper part is guarded by a callous, brownish, livid-coloured, jointed armour, consisting of fourteen semicircular scales, within which the insect rolls itself like a ball when touched. Like some other insects it casts the skin, and carries the young in valvular follicles under the abdomen. Slaters are prepared by hanging them, inclosed in a canvas bag, in the steam of hot alcohol, till they are killed.

*Qualities*. — Prepared slaters have a fetid odour, and a sweetish nauseous taste; on analysis, they furnish an alkaliescent fluid and an inert oil.

*Medical properties and uses*. — These insects were formerly regarded as expectorant and diuretic, and used in humoral asthma, dropsy, jaundice, and a long list of diseases. The retention of them in the list of materia medica exhibits the remains of a barbarous practice, which the good sense of modern practitioners should altogether explode. Their value, as a medicine, was justly estimated by Professor Alston, when he observed, “Upon the whole there is reason to think *Millepedarum* ʒj. is good for nothing, and ʒss. not much worth.”<sup>2</sup>

OPIUM. See *Papaver*.

OPOPONAX. See *Pastinaca*.

ORIGANUM. *Spec. Plant. Willd.* iii. 132.

Cl. 14. Ord. 1. Didynamia Gymnospermia. *Nat. ord.* Verticillatæ, *Linn.* Labiatæ, *Juss.*

G. 1116. *Strobile* four-cornered, spiked, collecting the calyces. *Corolla* with the upper lip erect and flat, the under three-parted, with the segments equal.

*Species* 10. *O. vulgare*. Common Marjoram. *Med. Bot.* 2d edit. 344. t. 123. *Smith Flor. Brit.* 639. *Eng. Bot.* 1143.

*Species* 15. *O. Majorana*. Sweet Marjoram. *Med. Bot.* 2d edit. 345. t. 124.

#### 1. ORIGANUM VULGARE.<sup>3</sup>

*Officinal*. ORIGANUM, *Lond.* Common Marjoram leaves.

*Syn*. Oriang (*F.*), Dort, Wohlgemuth (*G.*), Mariolein (*Dutch*), Meiran (*Dan.*), Mejram (*Swed.*), Majeran (*Pol.*), Origano (*I.*), Origano Sylvestre (*S.*), Mange rona (*Port.*).

<sup>1</sup> Ὀνισκοί Dioscoridis.

<sup>2</sup> Σαμψυχον Dioscoridis.

<sup>3</sup> *Lectures on Mat. Med.* ii. 496.



This plant is indigenous and perennial, growing on dry, chalky and gravelly hills, flowering from July to September. The root is creeping and fibrous, sending up erect, branching, trichotomous, tetragonous stems, about eighteen inches in height, downy and of a purplish hue. The leaves are ovate, entire, somewhat hairy, ciliated, punctured, and of a deep yellowish-green colour. The flowers are in terminal panicles, of a pink-purple or rose colour, and furnished with ovate, sessile, brownish-red bractes. The calyx is tubular, toothed; the segments being nearly equal: the corolla is funnel-shaped, with the upper lip bifid and obtuse, and the under trifid, blunt, and spreading. The filaments are furnished with double anthers, and the style is filiform, with a bifid reflected stigma.

*Qualities*. — The odour is agreeable and aromatic, and the taste warm and pungent, much resembling thyme. In distillation with water it affords a very acrid, penetrating, volatile oil, on which its qualities depend.

*Medical properties and uses*. — Common Marjoram is regarded as tonic, stomachic, and emmenagogue. It was formerly used in debilities of the stomach; but is now neglected. The dose is from grs. x. to ℥j., in powder.

Officinal preparation. *Oleum Origani*, L.

## 2. ORIGANUM MAJORANA.<sup>1</sup>

*Officinal*. ORIGANI MARJORANÆ HERBA, *Edin.* MAJORANA; HERBA, *Dub.* Sweet Marjoram.

*Syn.* Marjolaine (*F.*), Majoran (*G.*), Maggiorana (*I.*), Origano (*S.*), Mirzunjoosh (*Arab.*), Marroo (*Tam.*).

This is an annual plant, a native of Portugal and Syria; but cultivated in our gardens for culinary and medicinal purposes, and flowering in July and August. The root is long, brown, and fibrous; the stems numerous, woody, branching, and rising a foot and a half in height. The leaves are downy, entire, ovate, petiolate, and of a pale-green colour. The flowers are small, white, appearing successively among the bractes, which are numerous, and form roundish, compact, terminal spikes. The calyx is tubular, five-toothed, with the teeth acute: the corolla funnel-shaped and bilabiate; the upper lip erect and roundish; the lower cut into three pointed segments.

It is cut for medicinal use when it begins to flower in July.

*Qualities*. — The odour is pleasant, and the taste moderately warm, bitterish and aromatic. Both alcohol and water extract the virtues of sweet Marjoram; and, in distillation with water, it yields a large portion of volatile oil, which, on being long kept, becomes solid.

<sup>1</sup> *Opovavos* Dioscoridis.



*Medical properties and uses.* — Sweet marjoram is tonic, and was formerly regarded as possessing errhine powers. It is scarcely ever used except as a culinary herb, or as an adjunct to cephalic snuffs, to which, however, it adds no efficacy.

**OSSA.** Bones. *Edin.*

*Syn.* Des Os (*F.*), Knochen (*G.*), Ossi (*I.*), Huesos (*S.*).

The bones of animals are composed of earthy salts, gelatin, albumen, and oil. According to the analysis of Fourcroy and Vauquelin, the components of Ox bones are, in 100 parts, 51 of animal matter, 37·7 of phosphate of lime, 10 of carbonate of lime, and 1·3 of phosphate of magnesia. Besides these, M. Hatchett detected sulphate of lime, and Berzelius some fluuate of lime in bones.

**OSTREA.** *Syst. Nat. Gmelin.* vi. 3315.

*Cl.* 6. *Ord.* 3. Vermes Testacea.

*G.* 313. *Animal Tethis.* Shell bivalve; the valves unequal, and somewhat eared. *Hinge* toothless, but furnished with an ovate hollow cavity, with lateral transverse furrows. *Vulva*, or anus, none.

*Species* 105. *O. edulus.* The common Oyster. *Pennant's British Zoology*, iv. 102. *t.* 62.

*Officinal.* TESTÆ, *Lond.* The shells.

*Syn.* Ecailles des Huitres (*F.*), Austerschaalen (*G.*), Conchiglia d' Ostrica (*I.*), Cascara (*S.*).

This well-known shell-fish inhabits the European and Indian oceans throughout; and is particularly plentiful on the British coasts, which were early famed for producing the best oysters to supply the stews of ancient Rome, in the most luxurious period of its history.<sup>1</sup> They are naturally attached to shelving rocks; but, for the facility of always obtaining them for the purposes of aliment, they are generally laid down near the shore. They are hermaphrodite, and throw out a spat in spring which gradually enlarges to a perfect oyster. The nature of the shell in some degree, and the taste and goodness of the fish, depend on the soil of the bed. They are tender and friable on a calcareous bottom, thick and solid on rocks, more glutinous on marle, and oily and luscious on a slimy bed. The green colour of those fed in pits on the coast of Holland has been supposed to be owing to copper; but it arises from a species of conferva which cover these stagnant pools.<sup>2</sup> The best oysters on the British shores are found at Purfleet, the worst near Liverpool. The oyster, when good, is very digestible and nutritious, particularly when eaten raw; and forms

<sup>1</sup> Sergius Ovata was the inventor of stews for oysters among the Romans. *Pliny*, lib. xiv. cap. 54.

<sup>2</sup> Beckman's Observations, *Phil. Mag.* vi. 97. In Scotland, oysters laid down to feed near the salt-works on the shore, attain a large size and a great richness of flavour: they are called *Pandoors*, and are much esteemed.



an excellent article of food for the phthisical, and convalescents. When they are sick, which is known by a black substance on the fringe or fin, or a very milky appearance of it, they are unwholesome. The shells only are officinal.

*Qualities.* — Oyster shells consist of alternate layers of carbonate of lime and an animal matter supposed to be coagulated albumen. When thrown into a fire, they emit a great deal of smoke, the animal matter is destroyed, the carbonic acid dissipated in the form of gas, and pure lime remains.

*Medical properties and uses.* — Oyster shells are antacid; but as, in their unburned state, they are less so than chalk, and, when burned, differ in nothing from lime, their retention in the list of materia medica is unnecessary.

Officinal preparation. *Testæ præparatæ*, L.

OVIS. *Syst. Nat. Gmelin.* i. 197.

Cl. 1. Ord. 5. Mammalia Pecora.

G. 31. Horns concave, rough, inclined outwards, and spirally twisted. *Cutting Teeth* eight in the lower jaw. *Tusks* none.

*Species* 1. *O. Aries*.<sup>1</sup> The Sheep. *Buffon Hist. Nat.* v. p. 1. t. 1, 2.

*Officinal.* SEVUM, *Lond.* ADEPS OVILLUS, *Edin.* SEVUM; OVILLUM, *Dub.* Mutton Suet.

*Syn.* Graisse de Mouton (*F.*), Hammeltalg (*G.*), Grasse duro (*I.*), Grassa (*S.*), Aatoo kalupoo (*Tam.*), Lemak (*Malay*), Elloomusstail (*Cyng.*), Addjavuppa (*Sans.*).

The sheep is too well known to require any description. It is an inhabitant of almost every climate, and delights in dry, saline, moderately elevated and warm pastures. It is the most innocent, simple, and timid of quadrupeds: scarcely ever living beyond fourteen years of age; yet liable to many diseases. There are several natural varieties of the sheep in the British islands, the largest of which is found in Lincolnshire, and the smallest in Zetland; and the number of these is much increasing by the cross breeds, which, for the improvement of the wool and flesh, are annually effected. Mutton is less dense than beef, very digestible and wholesome, and is at its greatest perfection when about five years old. It is very much improved by the castration of the animal, and is then called wether mutton. The broth made of it does not agree so well as light beef tea or veal tea, with delicate and weakened stomachs; but it forms an excellent emollient enema, in cases of ulceration or abrasion of the rectum and that state of the bowels of infants which occasions green stools and aphthæ.<sup>2</sup> The suet,

<sup>1</sup> Προβατον *Aristot. Hist. Animal.* v. cap. 11.

<sup>2</sup> The milk of the Ewe is seldom used either as aliment or medicine. It contains more cream and less whey than Cow's milk, but the butter yielded by it never acquires a proper consistence. It is made into cheese in Scotland, which is bitterish; and, when old, warm, biting, and resembling Parmesan cheese in flavour.



which is the officinal part of the animal, is chiefly obtained from about the kidneys and loins.

*Qualities.* — Suet is the most consistent of the real animal fats. It is white, has some degree of brittleness, is inodorous, and requires a temperature of 127° Fahrenheit to melt it. When distilled, it yields a large proportion of oleic and margaritic acids. In other respects, it agrees with the other animal fats. (See the qualities of fat under *Sus scrofa*).

*Medical properties and uses.* — Like the other fats, Suet is emollient. It is sometimes boiled in milk, in the proportion of ʒij of the suet to 0j of milk; and a cupful of the mixture may be administered in chronic diarrhœa, when there is much acrimony of the contents of the bowels; but its principal use is to give consistence to ointments and plasters.

Officinal preparation. *Sevum preparatum*, L.

OVUM. See *Phasianus*.

OXALIS.<sup>1</sup> *Spec. Plant. Willd.* ii. 772.

Cl. 10. Ord. 5. Decandria Pentagynia. Nat. ord. Gruinales, Linn. Gerania, Juss.

G. 918. Calyx five-leaved. Petals connected by claws. Stamens unequal, the five shorter exterior ones connected at the base. Capsules opening at the corners, five-cornered.

\* \* \* leaves ternate, scape one-flowered.

Species 25. *O. Acetosella*. Wood-sorrel. *Med. Bot.* 2d edit. 563. t. 201. Smith, *Flor. Brit.* 491. *Jacquin's Oxalis*, 114. t. 80. f. 1.

Officinal. ACETOSELLA, Lond. Wood-sorrel.

Syn. Oseille des Bucherons (F.), Sauerklee (G.), Zuurklaver (Dutch), Skovsyne (Dan.), Harsyra (Swed.), Szezawik (Polish), *Acetosa salvatica* (I.), *Oxalide acederilla* (S.), *Trevo azedo* (Portug.).

This is an indigenous perennial plant, found in woods, under hedges and other shaded places, and flowering in April and May; the root is horizontal, toothed, fleshy, and of a reddish colour. The leaves are all radical, ternate like the trefoil, and petiolate; with the leaflets, obcordate, very entire, hairy, of a yellowish-green colour, and purplish underneath. The scape or flower-stalk is furnished with two, scaly bractes, placed about an inch and a half beneath the flower, which is sub-nutant, delicate, and of a flesh-colour streaked with red. The calycine leaflets are oblong, oval, acute, ciliated, and purple at the tip. The corolla is bell-shaped; with the claws of the petals upright, and the borders obovate, rounded, and spreading: the filaments are somewhat connate at the base, and furnished with oblong, incumbent anthers; and the styles smooth, rising from an ovate germen. The capsule is membranous, and contains two seeds in every cell. Each seed is invested with a fleshy

<sup>1</sup> *Oxalis Dioecoridis*.



white aril, which is, at first, smooth and closed on every side, but at length, opening at the apex elastically, it rolls back and throws off the seed with considerable force.<sup>1</sup>

*Qualities*. — This plant is inodorous, and has a pleasant, acidulous taste. The expressed juice reddens vegetable blues; coagulates milk, and instantly precipitates lime from its solutions. Its active principle is superoxalate of potass, which is obtained crystallized from the expressed juice, and sold in the shops under the name of *Essential salt of lemons*.<sup>2</sup> The same salt may be formed by cautiously dropping a solution of potass into a saturated solution of the oxalic acid, obtained from sugar by the action of the nitric acid; the superoxalate precipitates as soon as the proper quantity of alkali is added.<sup>3</sup>

*Medical properties and uses*. — Wood-sorrel is refrigerant and antiseptic. Boiled with milk it forms a pleasant whey, which may prove a useful refrigerant in fevers, as may also the expressed juice, or the superoxalate obtained from it diluted with water: but although they are much extolled in inflammatory, bilious and putrid cases, by the continental physicians, yet their place is well and easily supplied by lemon juice, or the citric acid, dissolved in water. The recent herb eaten as a salad, may be serviceable in scorbutic affections.

PAPAVER. *Spec. Plant. Willd.* ii. 1144.

Cl. 13. Ord. 1. Polyandria Monogynia. Nat. ord. Rhœadææ, Linn. Papaveracæ, Juss.

G. 1015. Corolla four-petalled. Calyx two-leaved. Capsule one-celled, opening by pores under the persistent stigma.

\*\* with smooth capsules.

Sp. 5. *P. Rhoeas*. Corn or Red Poppy. *Med. Bot.* 2d edit. 387. t. 139. *Smith, Flora Brit.* 567. *Eng. Bot.* 645.

Sp. 7. *P. somniferum*. White Poppy. *Med. Bot.* 2d edit. 376. t. 138. *Smith, Flora Brit.* 568.

#### 1. PAPAVER RHOEAS.<sup>4</sup>

*Officinal*. RHOEADES PETALA, *Lond.* PAPAVER ERRATICUM; PETALA, *Dub.* Petals of the Red Poppy.

*Syn.* Coquelicot (*F.*), Die blumen der klapprose (*G.*), Klapperroos (*Dutch*), Klapperrose (*Dan.*), Kornros (*Swed.*), Papavero rosolaccio (*I.*), Adormidera sylvestre; Amapola (*S.*), Papoileira (*Port.*).

<sup>1</sup> *Gærtner de Fructibus*, ii. 152. t. 115. fig. 5.

<sup>2</sup> This salt is prepared on the continent by the following process: The juice is allowed to subside after being slightly heated, and then clarified by adding to it water, in which a small portion of fine clay is suspended. This clarified juice is next boiled till a pellicle forms on its surface, and put aside for a month to crystallize; the operation being repeated until the whole of the salt is obtained, when it is purified by a second crystallization. *Annales de Chimie*, xiv. 7. The essential salt of lemon of the shops is generally one half cream of tartar.

<sup>3</sup> *Crell's Annals*, (trans.) i. 107.

<sup>4</sup> *Portus Theophrasti et Dioscoridis*.



This species of the poppy is an indigenous annual, growing in the greatest abundance in corn-fields, and waste places, and flowering in June and July. Its geographic situation extends from 60° N. lat. towards the tropics; but it is not found in America. The stem rises about a foot in height, is branched, and every where furnished with stiffish, horizontally spreading hairs. The leaves are sessile, pinnatifid, sometimes doubly so, serrated or cut, and generally hairy. The flowers are solitary, on slender, hairy peduncles; the calyx consists of two ovate, rough, concave leaves, which fall before the petals expand: the petals are four, large, roundish, unequal, and spreading, of a full, bright-scarlet colour, and sometimes marked with a black spot at the base. The germen is ovate, smooth, with a convex, sessile, shield-like stigma, scalloped on the edge, having many purple-coloured rays; and becomes an urn-shaped capsule.<sup>1</sup>

The petals must be gathered when they begin to blow, as they very soon drop after they are fully expanded.

*Qualities.* — They have a faint, narcotic odour, and a mucilaginous, very slightly bitter taste. They yield their colouring matter to warm water; and on this account only are used, as they cannot be said to possess any anodyne properties. The capsules, however, of every species of poppy contain opium; and from the red, it has actually been procured for medicinal purposes, both by Boulduc<sup>2</sup> and Dr. Alston<sup>3</sup>; but the quantity is too small to make it an object of importance.

Official preparation. *Syrupus Rhoeades*. L. D.

## 2. PAPAVER SOMNIFERUM.<sup>4</sup>

*Official.* PAPAVERIS CAPSULÆ. OPIUM, *Lond.* PAPAVERIS SOMNIFERI CAPSULÆ. OPIUM, *Edin.* PAPAVER ALBUM; CAPSULÆ; OPIUM; SUCCUS CONCRETUS, *Dub.* Poppy capsules or heads; and Opium.

*Syn.* Capsules des pavots blancs; Opium (*F.*), die Köpse des Weissen Mohns, Mohnsaft (*G.*), Wittamnu Hewl, Turks Heulzap (*Dutch*), Valmuesaft (*Danish*), Wallmo Opion (*Swedish*, and *Russian*), Capi del Papavero; Oppio (*I.*), Adormideras; Opio (*S.*), Afeeoan (*Arab.*), Affion (*Turkish*), Afiun (*H.*), Jya-Pien (*Chinese.*), Abinie (*Tam.*), Ufyoon (*Malay.*), Sheerkhuskah (*Pers.*), Abim (*Cyng.*), Apium (*Jav.*), Caruppa (*Malab.*).

<sup>1</sup> This form of capsule easily distinguishes it from *Papaver dubium*, which has a long, narrow capsule, but in other respects closely resembles the Corn Poppy.

<sup>2</sup> *Mem. de Acad. de Paris*, 1712.

<sup>3</sup> *Alston's Mat. Med.*

<sup>4</sup> Μηκων ἡμερος Theophrasti et Dioscoridis. Homer notices the Somniferous Poppy under the name of μηκων, as a garden plant; and it is said to be nourishing, by Hippocrates: an expression which is explained by the fact that, at this day, in Persia, when the plants rise too thick in the fields, in which they are sown, those which are taken up, when they are young, are used as pot herbs. The seeds are not narcotic. The following are the names by which the Poppy is known in the greater part of Europe. Pavot (*F.*), Papavero (*I.*), Mohn, Magen (*G.*), Dormidera, Cascak (*S.*), Mák (Boh. and Hung.), Maczek (Polish), Maan (Flemish), Valmue (*Danish*).



The Somniferous or White Poppy is a native of Asia; and although it is found growing wild in the southern parts of Europe, and even in England, yet there is every reason for thinking, that its seed must have been carried to these parts. It was very early cultivated in Greece, perhaps, at first, solely for the sake of its seed, which was used as food. It is extensively cultivated in most of the states of Europe<sup>1</sup>, in the present age, not only on account of the opium, for which it is reared in Turkey, Persia, and India, but also on account of the capsules, and of the bland oil obtained from the seeds. It is an annual plant, flowering in June and July, in Europe; and in February, in India. The stem is glaucous coloured, smooth, erect, and round; rising to the height of five or six feet, when in a favourable situation. The leaves are large, simple, obtuse, lobed and crenated, and embracing the stem on which they are alternately placed. The flowers are large and terminal; the calyx is formed of two, smooth, ovate, bifid, concave leaves, that drop on the expanding of the

<sup>1</sup> In England, it has been cultivated for the purpose of obtaining opium. Mr. Ball, in 1796, received a premium from the Society for the Encouragement of Arts, for a specimen of British opium little inferior to the Oriental. *Transactions of the Society of Arts*, xiv. 260 to 270. Messrs. Cowley and Stains, in 1823, collected 196 lbs. of opium, which sold for 30s. 6d. per lb. from little more than 12 acres of land. But it has been most successfully cultivated by Mr. Young, who found that the best and most lucrative mode of cultivating the the poppy in this country is in wide drills, about 12 inches between the rows. The plants sown about the middle of April, are ready for bleeding about the middle of July. The subsoil should be sand, or at least pervious. The instrument used by Mr. Young for bleeding the capsules is a double bladed, convex-edged knife, covered with sealing wax, except so much of the edge as is necessary to make the incisions. These are made vertical, then obliquely; and the proper time for this operation is a week after the flowers fall. When these scarifications are finished, he slices off the capitulum or stigma, and collects the milky juice that exudes. The juice is collected on a common hair brush known to painters by the name of a sash tool, rounded a little at the point. This, when it is charged with juice, is scraped on the side of a tin flask, fastened to the breast of the gatherer.

Mr. Young supposes, that by sowing the poppies between early potatoes, the following may be the probable return—at least such was the result of his experiments, for one acre,

56 lbs. of opium, at 36s. per lb.	.	:	£100	16	0
36 boles of early potatoes, at 24s.	.	.	43	4	0
250 lbs. of oil, cold drawn, at 1s. 6d.	.	.	18	15	0
125 ——— warm, at 6d.	.	.	3	2	6
500 oil cakes, at 18s. per 100.	.	.	4	10	0
			£170	7	6
Expenses			60	—	—
Profit			£110	—	—

But I will suppose this to be over-rated one half. See *Edin. Philosophical Journ.* vol. i. p. 258—270. See, also, a valuable paper on this subject in the *Quarterly Journ. of Science*, vol. iv. p. 69.

In Bengal, the seeds are sown in quadrangular areas, and the intervals formed into channels for conveying water to each area.



petals ; which are four in number, large, roundish, entire, somewhat undulated and white ; occasionally of a silver-grey colour, and tinged with violet at the base. The filaments are very numerous, slender, shorter than the corolla, and support erect, compressed anthers ; and the germen, which is globular and smooth, is crowned with a many-rayed stigma. The capsule, which stands on a short pedicel, is globular when well grown, smooth, glaucous, from two to four inches in diameter, a little flattened at the top and bottom, and crowned with the persistent stigma, the segments of which stand erect, and have an elegant appearance. The seeds are small, white, or grey, reniform, and very numerous ; and escape, when ripe, through small openings under the points of the stigma.

All the parts of the Poppy contain a white, opaque, narcotic juice ; but it abounds more in the capsules : thence these are the only officinal parts of the plant, and for them chiefly is the plant cultivated in this country. They are gathered as they ripen ; and as this happens at different times, there are annually three or four gatherings. They are brought to market in bags, each containing about 3000 capsules, and sold to the druggists. <sup>1</sup>

The milky juice of the Poppy, in its more perfect state, is extracted by incisions made in the capsules, and inspissated ; and in this state forms the Opium of commerce. <sup>2</sup> The period for commencing this operation is when the petals fall and the capsule assumes a whitish hue. The mode of obtaining it, appears to have been nearly the same in the time of Dioscorides, as it is at this day. The plants, which should be six inches apart during their growth, are carefully watered and manured, the watering being more profuse as the period of flowering approaches, and until the capsules are half grown, when it is discontinued, and the collection of the Opium commences. At sun-set, longitudinal incisions are made upon each half-ripe capsule, with an instrument which has two sharp points. The incisions pass from below upwards,

<sup>1</sup> The London market is chiefly supplied from Mitcham in Surrey. The average price of each bag containing 5000 capsules, is about 4*l.* 10*s.* — *Stevenson's Survey*, 582.

<sup>2</sup> In tracing the origin of the name *Opium*, we find that the ancient inhabitants of India and of Egypt, and the Arabians, called the inspissated juice of the capsule of the Poppy, *affion* ; the Persians, *afiuun* ; the Moors, *affiun* ; and by the modern Turks, it is termed *affioni*. The Greeks named it *opion*, a word derived from *opos*, juice. Ὀπιον ἀπὸ τοῦ ὀποῦ, adding sometimes, *μεκωνος*, the juice of the Poppy, or ὀπος τῶν κωθειῶν, the juice of the capsule. Some suppose, that the *Nepenthes* of Homer (*Odyssey*, iv. 220. v.) was *Opium* ; but this opinion is completely disproved by Dr. Christen, in his excellent work entitled, *Opium Historice, chemice atque Pharmacologic Investigatum*. Vindobonæ. 8vo. 1820.



and do not penetrate to the internal cavity. The night dews favour the exudation of the juice, which is collected in the morning by old women and children, who scrape it from off the wounds with a small iron scoop, and deposit the whole in an earthen pot<sup>1</sup>, where it is worked by wooden spatulas in the sun-shine, until it attain a considerable degree of spissitude. It is then formed by the hand into cakes, which are laid in earthen basins to be further exsiccated, when it is covered over with poppy or tobacco leaves.<sup>2</sup> Such is the mode followed in India, and according to Kœmpfer's account nearly the same is practised in Persia<sup>3</sup>: and when the juice is drawn in a similar manner in this country, and inspissated, it has all the characters of pure Opium.

Opium is brought to this country in chests from Turkey and India. The *Turkey Opium* is in flat pieces, covered with leaves, and the reddish capsules of some species of *Rumex*, which is considered an indication of its goodness, as the inferior kinds of Opium have none of these capsules adhering to them. Turkey Opium generally contains about one-fourth part of impurities. *East Indian Opium* is in round masses, covered with the petals of the Poppy in successive layers, to the thickness nearly of one-fourth of an inch. Mr. Kerr relates, that at Bahar, it is frequently adulterated with cow-dung, the extract of the Poppy procured by boiling, and various other substances. In Malwa<sup>4</sup>, it is mixed with oil of Sesamum, which is often one-half of the mass: ashes, the dried leaves of the plant, and catechu are also used. It is also adulterated with the aqueous extract of the capsules, the extracts of *Chelidonium glaucum*, *Lactuca virosa*, and *Glycyrrhiza glabra*; and sometimes with gum arabic, tragacanth, aloes, and many other articles. Excellent opium has been lately produced in England as I have already noticed.

Opium is regarded as bad when it is either very soft, greasy, light, friable, of an intensely black colour, or mixed with many impurities. A weak or empyreumatic odour, a slightly bitter or acrid, or a sweetish taste, or the power of marking a brown or black continuous streak, when drawn across paper, are also symptoms of inferior opium.

*Qualities.* — 1. The dried capsule of the Poppy is inodorous, and nearly insipid, a slight degree of bitterness only being

<sup>1</sup> In some parts it is collected in a small brass pot, or a cocoa-nut shell, containing a little linseed oil.

<sup>2</sup> *Med. Observ. and Inquiries*, v. 317.

<sup>3</sup> According to Kœmpfer, the produce of the first incisions is of a pale yellow, and called *gobaar* in Persia; and is esteemed much superior in strength and goodness, in every respect, to the other collections.

<sup>4</sup> In Malwa about 350,000 pounds, avoirdupois weight, of opium, are annually produced, 210,000 of which are for exportation. *Malcolm's Mem. on Central India*, vol. i. p. 8.



perceptible, when it is long chewed. Water by coction extracts its virtues; and when the decoction is evaporated, an extract is obtained, with properties similar to opium, but much less powerful.

2. *Turkey Opium* has a peculiar, strong, heavy, narcotic odour, and a bitter taste, which is accompanied with a sensation of acrid heat, or biting on the tongue and lips, if it be well chewed: and if long kept in the mouth of a person unaccustomed to chew it, blistering is produced. Its colour, when good, is a reddish-brown, or fawn colour; its texture compact and uniform. Its specific gravity is 1.336. When soft, it is tenacious; but, when long exposed to the air, it becomes hard, breaks with an uniform, shining fracture is pulverulent, and affords a yellowish-brown powder: which is again aggregated by a heat so low as that of the hand. It is inflammable, [and partially soluble in water, vinegar, lemon juice, wine, alcohol, and ether. By long boiling in water under exposure to the air, its narcotic powers are impaired; yet nothing rises with water, when it is distilled with that fluid.<sup>1</sup> When carefully triturated with hot water, about five parts in twelve of the opium are dissolved and retained in solution, nearly six parts are simply suspended, and rather more than one part remains perfectly insoluble, of a viscid, plastic nature, somewhat resembling the gluten of wheat, but of a dark colour. Bucholz regarded this as caoutchouc; according to Proust it contains wax; and Gren supposed it to be analogous to gluten. By digesting alcohol on this substance, I found that it dissolved a small portion of it, acquired a reddish-yellow colour, and became milky when added to water. Sulphuric ether digested on it, broke it down, and dissolved a portion of it, forming a yellowish tincture, which when evaporated on water left resin, a bitter extractive, and some acicular crystals of a salt which Derosne, erroneously, supposed to be the narcotic principle. The insoluble part, after the action of the ether, was subjected to a set of comparative experiments with the gluten of wheat, when it afforded similar results with the majority of the tests employed. Hence this part of Turkey Opium appears to be a modification of gluten combined with resin, extractive, and peculiar salts.

3. *East Indian Opium* has a strong, empyreumatic smell, but not much of the peculiar, narcotic, heavy odour of the Turkey Opium. The taste is more bitter, and equally nauseous, but it has less acrimony. It agrees with the Turkey Opium in its other sensible qualities, except that its colour is blacker, and its texture less plastic, although it is as tenacious.

---

<sup>1</sup> Beaumé, however, asserts that the odorous part of the opium is an oil.



It is more friable, and when triturated with water, no insoluble, plastic residuum is left, but it is altogether taken up; eight parts in twelve being dissolved, and the remainder suspended in the fluid.

The aqueous solutions of both kinds of Opium are transparent when filtered, that of the East Indian having the deepest brown colour; both redden litmus paper; neither is decomposed by alcohol, but both are precipitated by the carbonates of potass and of soda, and by pure ammonia; precipitates are also formed by solutions of the muriate, and nitrate of mercury, the sub-acetate and acetate of lead, the nitrate of silver, and the sulphates of copper, of zinc, and of iron. They are also precipitated by infusion of galls; the precipitate, as Dr. Duncan justly observes, resembling more that produced by cinchonin, than that by gelatine.<sup>1</sup> The solution of acetate of barytes does not alter the solutions of Turkey Opium, but produces a copious precipitate with those of the East Indian; oxalic acid precipitates both, but the latter more copiously.

No article of the Materia Medica has occupied the attention of chemists so much as Opium. I shall detail the more important results of their labours: but I may preface this account by stating that from the experiments to which it has been submitted, the components of Opium appear to be gum, resin, bitter extractive, *two peculiar crystallizable salts*, *two new acids*, alum, and sulphates of lime and of potass, the latter of which appears to be very abundant in the East Indian Opium: the Turkish contains besides, a species of gluten, and a substance resembling caoutchouc.

According to Bucholz the proportion of *extractive*, in 100 parts of opium, is 36; of *gum* 30; of *resin* 9; gluten 11; substance like caoutchouc 5; sulphate of potass 2; and of sulphate of lime 1; the remainder consisting of an oily or balsamic matter and waste. Nothing more was known until lately: but, as the sedative power of Opium evidently could not depend on any of the above named principles, some other was to be looked for; and has been at length discovered. When Ether is used as a menstruum for Opium, and the resin and extractive which it takes up are separated by evaporating the tincture on the surface of water, the pellicle of resin deposited is nearly insipid, while the extractive dissolved in the water has an intensely bitter taste; from this fact, and the circumstance already mentioned of Opium becoming inert when boiled in water, we might venture to conclude that the sedative principle resides in the extractive. Derosne, in 1804,

---

<sup>1</sup> *Edinburgh New Dispensatory*, 5th edit. 552.



asserted that the activity of Opium depends on a peculiar salt. He evaporated a watery infusion of Opium to the consistence of syrup, and digested the gritty precipitate formed by this evaporation in hot alcohol: as the solution cooled, a salt formed, which by repeated solutions and crystallizations was obtained free from the resin, of a white colour, and in rectangular prisms with rhomboidal bases; these were inodorous, insipid, insoluble in cold water, but soluble in 400 parts of boiling water; soluble in 100 parts of cold, and 24 of boiling alcohol; soluble in hot ether and the volatile oils, but separating as these fluids cooled; and very soluble in all the acids. Given to dogs, it produced the effects of a strong dose of Opium; but these were readily relieved by vinegar. In repeating the experiments of Derosne, I obtained a much greater proportion of crystals of this peculiar salt from East Indian than from Turkey Opium, which I conceived to militate against his idea of its being the sedative principle, inasmuch as larger doses of that variety of Opium than of Turkey are required to produce its sedative effect on the system. I have had no opportunities of ascertaining the power of this salt; but some experiments by M. Orfila<sup>1</sup> shew that, although it exerts a deleterious effect on the animal economy, yet, that the symptoms differ from those produced by Opium; and, even from Derosne's account, it is not much more powerful as a sedative than Opium itself.<sup>2</sup> My scepticism on this subject was further confirmed by the discovery of M. Sertuerner of Eimbeck in Hanover. The first experiments of this chemist were made public about a year after those of Derosne: but they excited little attention until he published a second memoir in 1817. According to Sertuerner the salt of Derosne is not the sedative principle of Opium, that principle being, according to him, an alkaline salt, which he had obtained in a separate state. This salt has been named *morphia*. Robiquet confirmed the statements of Sertuerner regarding the existence of *morphia*.<sup>3</sup> To obtain morphia, he orders a concentrated solution of Opium to be boiled for a quarter of an hour, with a small quantity (184 grains to lib. j of Opium) of magnesia. A greyish precipitate forms, which is to be separated by filtration, washed on the filter with cold water, dried, and then digested for some time with weak spirit, in a moderate heat, in order to separate the colouring matter. The residue is now again to be separated by the filter, washed with a little cold alcohol, and then boiled

<sup>1</sup> *Nouveau Journ. de Med.* tom. x. p. 154.

<sup>2</sup> *Annales de Chimie*, lxxv. 270. Derosne concludes from the effects of nitric acid and caloric on this salt, that it is composed of oxygen, hydrogen, azote, and carbon. It does not redden vegetable blues. *Ibid.* p. 279.

<sup>3</sup> *Annales de Chimie et de Phys.* tome v. p. 276.



in a larger quantity of rectified alcohol: on filtering the solution whilst it is yet boiling, morphia, beautifully crystallized and almost free from colour, is deposited as it cools. By repeating the last part of the operation three or four times, with the residue of the previous boilings, the whole of the morphia is obtained. The best process, however, is the following lately proposed by M. Hottot. Macerate  $\text{℥vi}$  of opium in a sufficient quantity of cold water to exhaust the residuum, and evaporate the liquors to sp. gr. 1.012: when half cooled, add grs. xxivss. of ammonia, and allow the precipitate to fall, then decant, and again add 141 grains of ammonia, and leave the mixture to deposit the precipitate for 12 hours, after which filter, wash the residuum with cold water, and treat it with  $\text{f℥xv}$  of alcohol of sp. grs. 847, and grains 141 of animal charcoal, in a sand bath; and when the alcohol boils<sup>1</sup> filter. As it cools the Morphia will crystallize. The salt thus procured is colourless, bitter, inodorous, and crystallized, in regular parallelopipeds. It burns like vegetable matter, leaving carbon as a residue; restores, like the alkalies, the colour of reddened turnsole paper, browns turmeric paper, and readily combines with acids forming neutral salts.<sup>2</sup> It is nearly insoluble in water, and not very soluble in cold alcohol, or in ether; but it is readily soluble in the two latter fluids in the boiling state; the salt being again precipitated in crystals as the solutions cool. It is soluble also in oil, both fixed and volatile. As an alkali, it holds the next place to Ammonia, having less affinity for the acids than either that salt or Magnesia. Its ultimate components are, according to Dumas and Pelletier 72.02 of carbon, 5.57 of azote, 7.61 of hydrogen, and 14.34 oxygen in 100 parts.<sup>3</sup> Morphia being scarcely soluble in water or in the fluids of the stomach in its uncombined state, does not display in a striking manner its properties when exhibited alone; but these are very striking when it is combined with an acid, particularly the acetic, the citric, or the codeic; with the latter of which it exists naturally in Opium in a state of combination.

According to Sertuerner and Robiquet's experiments, Morphia is combined with a peculiar acid, which Sertuerner named *meconic*, in its natural state in Opium; but that Morphia is combined in Opium with an acid different from the

<sup>1</sup> *Journ. de Pharm.* 475.

<sup>2</sup> The nitric acid of commerce, when dropped on morphia, communicates to it a beautiful red colour. The following are the characters of the neutral compounds of morphia which may be medicinally employed. The *acetate of morphia* crystallizes in soft, silky prisms, which are very soluble. The *sulphate* in arborescent or branching crystals, soluble in two parts of water at 60°. The *carbonate* in short prismatic crystals, soluble in four parts of water at 60°.

<sup>3</sup> *Ann. de Chimie*, T. xxiv. p. 185.



meconic, has been well presumed to be demonstrated by the experiments of M. Robinet.<sup>1</sup> According to him the meconic acid is found to be united with soda in Opium, forming an acidulous meconate of soda, and the Morphia, united with a completely distinct acid, which he has named *Codeic*, from  $\chi\omicron\delta\eta$ , a poppy head, forming a *Codiate*. M. Robinet was led to this discovery by employing a solution of sea-salt instead of pure water to dissolve the opium. When such a solution is filtered and evaporated, at the moment the sea-salt begins to deposit itself, a brown oily substance forms upon the surface of the fluid, and is the Morphia in combination with the Codeic acid. The Codiate of morphia crystallizes in silky *needles*, which aggregate in the form of stars. It is more soluble in hot than in cold water, is soluble in alcohol, but insoluble in ether. It reddens concentrated nitric acid, and dissolves without effervescence in diluted sulphuric and hydrochloric acids. The only metallic salts which precipitate it are the muriate of platina and the subacetate of lead. It strikes a deep blue colour with the super salts of iron; but it is not precipitated red by the salts of titanium, which distinguishes it from the Gallate of Morphia.

M. Robinet separated the Codeic acid from the Morphia, by dissolving it in water, and precipitating by subacetate of lead, and decomposing the precipitate by sulphuretted hydrogen. The acid remains in the solution; but when this was filtered and evaporated, it crystallized. It combines with potass, barytes, ammonia, and magnesia, forming crystallizable salts; but its properties have not yet been fully examined. The sedative properties therefore, of opium, appear to depend on the Morphia it contains; which acts very powerfully on the animal economy: but the Codeic acid in its separated state exerts no peculiar effect on the animal system. Its principal chemical characteristics are the producing an intense blue colour with solution of iron oxidized *ad maximum*; whilst the meconic acid produces a colour with that salt. MM. Majendie and Robiquet have lately endeavoured to prove, that the salt obtained by Derosne, and which they have named *narcotine*, is that principle which produces the excitement experienced by those who take opium, before its sedative effects are felt.<sup>2</sup> The components of Narcotine are carbon 68.88, azote 7.21, hydrogen 5.91, oxygen 18.00 in 100 parts.<sup>3</sup> Robiquet has proposed a mode of preparing an extract free from narcotine, and yet containing Morphia. (See *Extractum Opii*.)

<sup>1</sup> *Journ. de Pharmacie. Aout. 1825. p. 371.*

<sup>2</sup> Narcotine is readily obtained by digesting ether in opium, and slowly evaporating the filtered ethereal tincture: the narcotine crystallizes.

<sup>3</sup> *Ann. de Chimie, T. xxiv. p. 186.*



In repeating Sertuerner's and Robiquet's experiments, I obtained from good *Turkey* Opium, nearly three times the quantity of morphia yielded by the same weight of *East India* Opium; that from the latter was also more coloured, and in smaller crystals. Although the utmost nicety of manipulation was not attended to in these experiments, yet as both specimens were treated exactly in the same manner, the experiments are sufficient to shew the comparative richness of these varieties of the drug in this salt; and the result is certainly in favour of the opinion that the sedative property of Opium depends on this alkaline salt; and accounts for the fact which has been stated above, that much larger doses of the *East Indian* than of *Turkey* Opium are required to produce its sedative effect on the system. Mr. Brande procured a larger quantity of Morphia from a carefully prepared sample of *English* Opium than from the same weight of *Turkey* Opium.<sup>1</sup> Practitioners and physiologists in this country have not yet made any decisive experiments with Morphia, to determine its effects on the animal economy; but this point has been investigated by MM. Orfila and Majendie.<sup>2</sup> The experiments of the former being made upon dogs, they can be regarded as important only inasmuch as they confirm the opinion, that opium owes its soporific powers to this salt, and displays the comparative activity of the different combinations of the salt. Of the saline compounds he found the acetate the most powerful; but a solution of Morphia in olive oil acts with still greater intensity, and with more than double the effect of the aqueous extract of Opium. The experiments of M. Majendie being made on the human subject, are more interesting. He found that a quarter of a grain of Acetate of Morphia produces the most beneficial effects that can be expected from an anodyne, allaying pain and procuring sleep without, in any degree, affecting the cerebral functions: and I have been able, from my own experience, to verify this fact. The Sulphate acts in a similar manner, but with less energy. No experiments have yet been made to determine the value of the *Codiate*; but as I am now repeating Robinet's experiments, I hope soon to be able to determine that point.

*Medical properties and uses.* — Poppy heads or capsules possess anodyne properties; they are chiefly employed, boiled in water, as fomentations to inflamed and ulcerated surfaces; and a syrup prepared with the inspissated decoction is used as an anodyne for children, and to allay the tickling cough in chronic catarrh, and phthisis.

---

<sup>1</sup> *Manuel*, p. 128.

<sup>2</sup> *Nouveau Journ. de Medicine*, tom. i. p. 1. 23.



*Opium* is more generally used than any other remedy, the average annual importation exceeding 40,000 lbs; and consumption in Great Britain alone being upwards of 16,500 lbs.<sup>1</sup> It appears to have been introduced into the *Materia Medica* before the time of Hippocrates; but was little employed as an internal remedy until 300 years after his period. It operates at first as a powerful and very diffusible stimulus, but its primary operation is followed by narcotic and sedative effects in a degree much greater than could be expected from the previous excitement it induces. It acts directly on the nervous system, and when taken into the stomach destroys irritability, and allays pain in the most distant parts of the body, independent of the circulation, and without inducing any change on the composition of the blood. As the principle, therefore, on which *Opium* acts is the same over all the body, the topical application of it is capable of producing similar effects, only in a diminished degree, to those resulting from it when it is taken into the stomach. The larger the dose is, the more quickly its primary action is extended over the whole habit; and as every part is excited nearly at the same moment of time, the general, consequent exhaustion must necessarily more rapidly follow, than when the dose is merely sufficient to induce a degree of excitement scarcely exceeding the powers of the system on which it operates. Hence either the stimulant or the sedative effects of *Opium* may be rendered obvious by the extent of the dose in which it is exhibited, and the early knowledge of this truth might have saved much of the keen controversy which this subject at one period occasioned.

In moderate doses *Opium* increases the fulness, the force, and the frequency of the pulse, augments the heat of the body<sup>2</sup>, quickens respiration, and invigorates both the corporeal and mental functions, exhilarating even to intoxication, effects which are referred to the narcotine it contains<sup>3</sup>: but by

---

<sup>1</sup> The quantity of *Opium* imported into England from Bengal and other places, from the year 1786 to 1801, amounted to 286,271 lbs, and the consumption to 247,619 lbs. *Morewood on Inebriating liquors*, p. 106.

<sup>2</sup> It is extraordinary that Dioscorides, Galen, Aurelianus, and many of the ancients, believed that it produced cold.

<sup>3</sup> The Turks call opium *afioni*; and in the *teriakihana*, or opium shops of Constantinople, they take it in graduated doses from ten grains to one hundred grains in a day. It is mixed with rich syrup and the inspissated juices of fruit to render it more palatable and less intoxicating; and is taken with a spoon, or made up into small lozenges stamped with the words, *Mash Allah*, literally, "The work of God." The Tartar couriers who travel great distances, and with astonishing rapidity, take nothing else to support them during their journeys. (*Dallaway's Constantinople*, 4to. 78.) There is, however, some reason for supposing that the *Mash Allah*, or *Maslach* of the Turks, contain other narcotics, as those of *kemp* and of *lolium*, as well as opium. The Chinese, the Bernese, and the Sumatrans, smoke opium as Europeans smoke tobacco.



degrees these effects are succeeded by languor, lassitude, and sleep; and in many instances headach, sickness, thirst, tremors and other symptoms of debility, such as follow the excessive use of ardent spirits, supervene. In very large doses the primary excitement is scarcely apparent, but the pulse seems to be at once diminished, drowsiness and stupor immediately come on, and are followed by delirium, sighing, deep and stertorous breathing, cold sweats, convulsions, apoplexy, and death. The appearances on dissection are those which indicate the previous existence of violent inflammation of the stomach and bowels; but notwithstanding the symptoms of apoplexy which an overdose, when it proves fatal, occasions, no particular appearance of an inflammatory state or fulness of the vessels of the brain are perceived.

Opium is efficaciously given in some diseases of debility, as, for instance, fevers of the typhoid kind, and intermittents; and combined with calomel to check the progress of gangrene. In typhus, when given in small doses frequently repeated, it is an useful assistant to wine and tonics in supporting the *vis vitæ*; and at the same time allaying irritation, and obtunding the susceptibility of those morbid impressions which occasion watchfulness, delirium, tremors, and subsultus tendinum. It is to this effect of it that Alibert and others ascribe its power, when moderately used, of rendering the human body less susceptible to different diseases.<sup>1</sup> Some caution, however, is required in its exhibition; for if the heat of the body be much above the natural standard, and the skin dry, Opium increases these symptoms, augments thirst, and occasions restlessness. But if moisture be coming on, Opium accelerates it, and tranquillity and sleep follow. Hence the propriety of Dr. Currie's advice, not to give the evening dose of Opium in these fevers till very late or about one or two o'clock in the morning, when the heat is subsiding; or first to lower the temperature, and excite sensible perspiration by the effusion of cold water, or tepid sponging.<sup>2</sup> It is hurtful, also, where there is any disposition to local inflammation, particularly of the chest; and where there is much determination to the head. Opium very materially assists the bark in curing intermittents, and prevents it from running off by the bowels. When given at the approach of the paroxysm, it sometimes checks its attack, or shortens and renders it milder, and abates the violence of the hot stage by determining to the surface, and inducing sleep.

In acute rheumatism Opium is given united with ipecacu-

<sup>1</sup> *Nouveaux Elémens de Thérapeutique*, &c. 4 edit. tome xi. p. 76.

<sup>2</sup> *Medical Reports on the Use of cold and warm Water*, i. 290.



anha or antimonials<sup>1</sup>, and nitre, and always relieves when it determines to the surface. In the other phlegmasiæ<sup>2</sup> however, it cannot with propriety be used in the early stages; but after the inflammatory action is subdued, it is useful in quieting cough, allaying pain, and procuring sleep.

In eruptive diseases, particularly small-pox, the liberal use of Opium is found to be highly beneficial, when convulsions precede the appearance of the eruption, or if the accompanying fever assume the typhoid type. In malignant scarlatina, pemphigus, and several others of the exanthemata, it is equally valuable; but its use is contra-indicated in this class of diseases when the fever is inflammatory.

In the hæmorrhagiæ it is useful when the discharge arises chiefly from an increased degree of irritability, and where the pulse, instead of being strong and full, is small, quick, and intermitting. Hence its efficacy in the floodings of weakened habits after abortions, and in phthisical hæmoptysis. It has been recommended also after blood-letting, in the hæmoptysis and hæmatemesis of the latter months of pregnancy.

Although opiates are hurtful at first, and check expectoration in catarrh, yet, when the cough remains obstinate, their good effects are undoubted; and in the contagious catarrh or influenza, an opiate at bed-time is requisite for quieting the cough in every stage of the disorder. In dysentery, also, the benefit to be derived from Opium depends very much on the bowels having been previously well cleared, in which case it allays the tormina and tenesmus; and the same remark applies to diarrhœa.

But the spasmodic and convulsive diseases are those in which Opium is most evidently useful. In tetanus, although it does not always succeed, even when given in the largest doses, yet, many cases have occurred in which the continued exhibition of large doses has overcome the spasm, and cured the disease; particularly when it has been judiciously combined with cathartics: often, however, very large quantities of the remedy have been taken without any sensible effect on the state of the habit, and without relieving the disorder; and the same is the case in hydrophobia, in which 180 grains of solid opium have been taken, in the space of twelve hours, without producing any apparent effect. It has been found beneficial in chorea; but, as in tetanus, it is necessary to precede its use by

<sup>1</sup> I know of no remedy which so effectually relieves the excruciating pain of acute rheumatism, which generally makes its attack at night, as the following combination: R. Submuriatis Hydrargyri, gr. j. Antimonii tartarizati gr.  $\frac{1}{4}$  Opii gr. jss. fiat pilula horâ decumbentis sumenda.

<sup>2</sup> Were it allowable in this work to criticise nosological arrangements, we might justly question the propriety of placing rheumatism among the phlegmasiæ.

R  
 and q  
 Pol Tan q  
 - Ghu q  
 pil h. s. 15.



strong cathartics, or at least to give it in combination with these.<sup>1</sup> In epilepsy it proves useful when given in combination with musk; and it has been recommended by highly respectable authority<sup>2</sup> in eclampsia; but its efficacy in this complaint is rather doubtful. In spasmodic asthma it shortens the paroxysms; abates the violence of the cough in pertussis, when given after the primary fever subsides; and is more especially useful in pyrosis and cholera than any other medicine. Solid opium, either alone or united with camphor, is the most effectual remedy for checking obstinate vomiting proceeding from a morbid irritability of the stomach. In colic and ileus it is given in combination with laxatives, and allays the spasm and pain; nor is it less efficacious in flatulent colic with hernia. As a remedy in lues venerea Opium is still relied on by some foreign practitioners, but the idea of its anti-venereal powers has been justly exploded in this country; and it is properly regarded only as an useful adjunct to mercury in this disease: "by diminishing the sensibility of the stomach and bowels, it prevents many of those inconveniences which this mineral is apt to excite in the primæ viæ, and allows it to be more easily introduced into the system."<sup>3</sup> In short, in all cases where the irritability is morbidly increased, and where it is of importance to lessen pain, and procure sleep, Opium is undoubtedly the most valuable article of the materia medica.

Opium is contra-indicated in all morbid states of the body where a strong inflammatory diathesis exists; in pulmonary affections, when the cough is dry and hard, and the expectoration difficult and scanty; and, if not hurtful, its use is at least doubtful in mania, in which it generally occasions restlessness instead of procuring sleep.

Externally used, Opium is almost as efficacious as when it is taken into the stomach, and produces its narcotic effects without affecting the head or producing nausea. It is applied in the form of frictions, either combined with oil, or with the Camphor liniment, or in the form of tincture: and thus applied, it may be used in all the diseases above enumerated. We have often seen its good effects in colic; and have also witnessed its singular efficacy in symptomatic trismus, when rubbed on the jaw, and applied to the scrobiculus cordis by means of pledgets soaked in the tincture. A piece of solid Opium stuffed into a carious tooth relieves the pain of tooth-ach; and introduced into the rectum, either in the solid form or dissolved in water as an enema, it affords relief in tenesmus, in painful affections of the prostate gland, and in spasmodic strictures.

<sup>1</sup> *Observations on the Administration and Utility of Purgative Medicines, &c.* 86.

<sup>2</sup> Denman. Bland.

<sup>3</sup> *Pearson's Observations, &c. on Articles used in the Cure of Lues Venerea*, p. 60.



A weak, watery solution of it, also, is an useful adjunct to injections in gonorrhœa, and to collyria in ophthalmia; and the vinous tincture dropped into the eye removes the suffusion which often remains in that disease after the inflammation has been subdued, and in ecchymosis; and restores the tone of the diseased organ. The aqueous solution also lessens the pain of open cancer, when cloths soaked in it are laid over the sore; and it is as useful in lacerated wounds of tendinous and aponeurotic parts.

Opium is generally exhibited either in substance as a pill, or under the form of tincture. It is necessary to avoid combining it with substances which decompose it; and therefore solutions of oxymuriate of mercury, acetate of lead, sulphates of zinc, iron, and copper; of the carbonates of alkalies, lime water, infusion of galls, and infusion of yellow cinchona bark, are incompatible in prescriptions with Opium. In combination, however, with vinegar, the vegetable acids, and oil, its narcotic power is much increased.<sup>1</sup> The combinations of Morphia, its Acetate and Citrate, may be used in the same diseases and with the same intention as Opium. The result of my own experience inclines me to regard the Acetate as well adapted for cases of Phthisis and inflammatory affection, where it is of importance to obtain the sedative effect of the remedy free from the exciting quality. As the *Codiate* is the

<sup>1</sup> The effects of vegetable acids in augmenting the efficacy of opium is explained by what has been said on the combinations of morphia. The greater power of that preparation of opium, which has been known, for upwards of a hundred years, under the name of "*Black Drop*," appears to depend on its containing an acetate of morphia. The following is the mode of preparing it, as published by Dr. Armstrong, (vide *Practical Illustrations of Typhus*,) from the papers of the late Edward Walton, of Sunderland, one of the near relations of Edward Tonstall, of Bishop's Auckland, by whom it was originally prepared. "Take half a pound of opium sliced; three pints of good verjuice; one and a half ounce of nutmegs; half an ounce of saffron. Boil them to a proper thickness, then add a quarter of a pound of sugar, and two spoonful of yeast. Set the whole in a warm place near the fire, six or eight weeks, then place it in the open air until it becomes a syrup; lastly, decant, filter, and bottle it up, adding a little sugar to each bottle." One drop of this preparation is calculated to be equal to three drops of the Tincture of Opium, of the London College. It evidently owes its efficacy to the *acetate of morphia*, which is formed by the verjuice decomposing the codiate of the opium. The acetate itself is a more elegant preparation; and produces its effects in doses  $\frac{1}{6}$  of a grain. There is also some reason for thinking that another preparation of opium, the *Liquor Opii Sedativus*, of Mr. Batley, of Fore-Street, London, which has been justly esteemed one of the best preparations of the drug hitherto discovered, owes its efficacy to the acetate of morphia. The mode of preparing it is as yet kept secret; but I know that the whole of the resinous part of the opium employed is separated and rejected; and I am inclined to believe that acetic acid is employed to separate the gummy part. Dr. Paris (*Pharmacologia*) states as an objection to this preparation, that it undergoes some important change on being kept. Justice obliges me to say, that my experience does not allow me to concur in this remark. I used the remedy before it was sold to the Profession, and gave it the name it bears; and although I have, since, constantly prescribed it, and kept the preparation in rather a warm situation, yet I have not observed the change of which Dr. Paris has spoken.



natural compound of Morphia in Opium, it is likely to prove still more useful.

The dose of opium should be regulated by the nature of the disease, and the peculiar intention for which it is ordered. The circumstance of the patient having been previously accustomed to its use must also regulate the extent of the dose; for in this case a dose, which to one unaccustomed to its use would prove fatal, may perhaps to another in the habit of taking it be scarcely sufficient to produce its sedative effects. A quarter of a grain, or even less, frequently repeated, is, in general, sufficient to keep up its stimulant effect; and from gr. j. to grs. ij. act as a sedative, and produce sleep; while in tetanus, hydrophobia, and some other diseases, fʒvss. of laudanum have been given in twenty-six hours, without occasioning any bad effects, or even producing sleep.<sup>1</sup>

The use of opium for the purpose of exhilarating the spirits has long been common in Turkey, Syria, and China<sup>2</sup>; and of late years it has been unfortunately adopted by many, particularly females, in this country. Russell<sup>3</sup> says that in Syria, when combined with spices and aromatics, he has known it taken to the amount of ʒiij. in twenty-four hours. Its habitual use cannot be too much reprobated. It impairs the digestive organs, consequently the vigour of the whole body, and destroys also gradually the mental energies. The effects of opium on those addicted to its use, says Russell, are at first obstinate costiveness, succeeded by diarrhœa and flatulence, with loss of appetite and a sottish appearance. The memories of those who take it soon fail, they become prematurely old, and then sink into the grave objects of scorn and pity.<sup>4</sup>

<sup>1</sup> *Currie's Medical Reports*, &c. i. 138.

<sup>2</sup> The inhabitants of these countries regard it also as an aphrodisiac. "Ad venerem enim ciere integræ nationes norunt, et in hunc usum adhibent: sic Japonenses, Chinenses, magis Indiæ, Persæ, Ægyptii et Turcæ aphrodisiacum opium, referentibus Pr. Alpino, Saar (*Itinerar. Ind. orient.*), Cleyer (*Eph. N. C.* 11. x. 35.) Foeminas turcicas opio viros incitare refert Jahn (*Mat. Med.* ii. 265.) Vide *Opium Hist. Chem. atque Pharm. invest. per. C. A. Christen*, 8vo. p. 53.

<sup>3</sup> *History of Aleppo*, i. 128.

<sup>4</sup> Mustapha Shatoor, an opium eater in Smyrna, took daily three drachms of crude opium. The visible effects at the time were the sparkling of his eyes, and great exhilaration of spirits. He found the desire of increasing his dose growing upon him. He seemed twenty years older than he really was; his complexion was very sallow, his legs small, his gums eaten away, and the teeth laid bare to the sockets. He could not rise without first swallowing half a drachm of opium. (*Phil. Trans.* xix. 289.) Some years ago, I was consulted by a lady who took a wine pint and a half of laudanum every week, and who, as she began to experience its bad effects on her constitution, was anxious to discontinue it, but was uncertain how to proceed. I recommended her to get a three-pint bottle of the drug, and to continue her usual dose; but, after taking each portion out of the bottle, always to replace it with water; so that, in the progress of time, the bottle would contain water only, and her propensity would be cured. She continued the plan for one week only, and having left my neighbourhood, I have had no opportunity of knowing the consequence of her return to the abuse of



When Opium has been taken in an overdose, the first thing to be done for counteracting its bad effect, is the exhibition of a powerful emetic; and for this purpose ℥j. of sulphate of zinc, or from grs. v. to grs. x. of sulphate of copper dissolved in water should be immediately swallowed, and the vomiting kept up for a considerable time, and urged by irritation of the fauces; but as these substances sometimes fail in producing vomiting, the following draught has been recommended: *R.* Ammoniae Subcarbonatis, ℥j; Pulv. Ipecacuanhæ, ʒss.; Aquæ Menthæ pip. fʒij.; Tinct. Capsici, fʒij.<sup>1</sup> This has certainly the advantage of rousing the system, at the same time that it evacuates the stomach. Large draughts of vinegar and water, or other acidulated fluids should be frequently taken, after the stomach has been evacuated, and the powers of the habit supported by brandy, coffee, and cordials. The sufferer should be kept awake, and, if possible, in continued gentle motion. Currie recommends the affusion of warm water at 106° or 108°<sup>2</sup>, for removing the drowsiness. The costiveness which supervenes, when the patient recovers, should be treated with aromatic purges, and the tone of the habit restored by stimulating tonics, and the shower-bath. When either simple Morphia or the Acetate of that salt is taken in an over-dose, say two or three grains, it operates as a virulent poison. The symptoms are head-ach, frightful reveries, vertigo, dimness of sight, subsultus, violent agitation, and obstinate vomiting. The pupil in the majority of cases is contracted. There is pain also in the epigastric region, and in the course of the intestines, but the breathing is not much affected. Pruritus of the skin is almost a constant symptom of poisoning by morphia.<sup>3</sup> M. J. L. Lassaigue has proposed the following method of detecting it when it has proved fatal. Digest the stomach or its contents, evaporating nearly to dryness in alcohol and precipitate by the subacetate of lead. Filter the solution, and pass through it a stream of hydrosulphuretted gas to precipitate any excess of the salt of lead; then evaporate the solution in vacuo, and add to it nitric acid. If morphia or its acetate be present, a yellow-orange colour, which changes to a blood-red, will be produced. *Bulletin des Sciences Médicales*, t. i. p. 147.

Officinal preparations. Of the poppy capsules — *Decoctum Papaveris*, L. *Extractum Papaveris*, L. E. *Syrupus Papaveris*, L. E. D. Of Opium — *Opium purificatum*, D. *Confectio Opii*,

---

**Opium.** Opium appears to have no detrimental effect on white ants, who eat it freely and in large quantities.

<sup>1</sup> See a paper by Mr. Sprague, *Medical Repository*, vol. xviii. p. 125.

<sup>2</sup> *Reports on Water*, i. 80.

<sup>3</sup> *Lond. Med. Repos.* vol. i. *New Ser.* p. 80.



L. *Elect. Opii*, E. *Elect. Catechu*, E. *Elect. Catechu comp.* D. *Extractum Opii*, L. E. D. *Pilulæ Opii*, E. *Pilulæ Saponis cum Opio*, L. *Pulvis opiatas*, E. *Pulvis Cornu usti cum Opio*, L. *Pulvis Cretæ comp. cum Opio*, L. *Pulvis Ipecacuanhæ comp.* L. E. D. *Tinctura Opii*, L. E. D. *Tinctura Camphoræ composita*, L. D. *Tinctura Opii ammoniata*, E. *Trochisci Glycyrrhizæ cum Opio*, E. *Vinum Opii*, L.

PASTINACA. *Spec. Plant. Willd.* i. 1465.

Cl. 5. Ord. 2. Pentandria Digynia. Nat. ord. Umbellatæ.

G. 558. Fruit elliptical, compressed, flat. Petals involute, entire.

Species 3. *P. Opoponax*.<sup>1</sup> Opoponax, or Rough Parsnip. Med.

Bot. 2d edit. 122. t. 47.

Officinal. OPOPONACIS GUMMI RESINA, Lond. Opoponax.

Syn. Opoponax (F.), Panax gummi (G.), Opoponace (I.), Opopanaco (S.), Gawsheer (Pers.), Iáwesheer (Arab.).

This species of Parsnip is a perennial plant, a native of the south of Europe, flowering in July. The root is as thick as the human arm, branched, of a yellow colour, and covered with a corky bark: the stem rises about five feet in height, the thickness of a man's finger, round, striated, scariose at the base, angular at the summit, and shining: the radical leaves are simple, cordate, and crenated; those of the stem, ternate and quinate, with the terminal leaflet cordate and very large: the whole are petiolate, with the petioles sheathing, and the leaflets hairy on the under surface; the umbelliferous branches are very smooth; first alternate, erect; then two, three, or four together, in a sort of whorl, two or three inches long, with one or two spathaceous leaflets towards the middle or at the top; the universal umbels have seven or eight rays, an inch long, of a yellowish green colour: both the involucre and involucels consist of from four to six very short leaflets, frequently permanent. The fruit is flat.

In the Levant, where this plant grows, the milky juice which exudes from incisions made in the roots, and dried in the sun, forms the opoponax of the shops. That which is obtained from plants grown in France, contains scarcely any resin. It is imported from Turkey and India in chests, and is sometimes in tears or drops, but more usually in irregular lumps.

Qualities. — Opoponax has a strong, disagreeable smell and a bitter, acrid taste. The masses are of a reddish-yellow colour, speckled with white on the outside, paler within, and frequently variegated with large white pieces. Its specific gravity is 1.622.<sup>2</sup> It appears to be a compound of gum 33.40, resin 42., starch 4.20, extractive 1.60, wax .30, malic acid 2.80, a trace of caoutchouc, essential oil 5.90, and woody fibre 9.80 in 100 parts.<sup>3</sup> When triturated with water, about one-half of it dissolves, forming an opaque, milky solution, which deposits,

<sup>1</sup> Ὀποπᾶναξ Dioscoridis.

<sup>2</sup> Brisson.

<sup>3</sup> Pelletier, *Ann. de Chim.* lxi. p. 90.



on standing, a portion of resinous matter, and becomes yellowish. Alcohol acts feebly on it; and in distillation, either with spirit or with water, the odour of the opoponax is very strongly communicated to the fluids, but scarcely any oil is obtained in a separate state.

*Medical properties and uses.* — This gum resin is regarded as antispasmodic and emmenagogue, and as such has been used in hysteria and chlorosis, but it is very seldom ordered. The dose may be from grs. x. to  $\bar{3}$  ss.

PETROLEUM. Vide *Bitumen*.

PHASIANUS. *Syst. Nat. Gmelin.* i. 737.

Cl. 2. Ord. 5. Aves, Gallinæ.

G. 101. Beak short, strong. Cheeks made smooth, with a naked skin. Feet spurred.

Species 1. *P. Gallus*. The Dunghill Fowl. *Willd. Ornith.* 154. t. 26.

Officinal. OVA, Lond. The Egg.

*Syn.* Œuf (*F.*), Ein, Ey (*G.*), Ovo (*I.*), Huevo (*S.*), Ey (*Dutch*), Aeg (*Dan.*), Agg (*Swed.*), Ovo (*Port.*), Jaizo (*Russ.*), Jaie (*Poln.*), Wegce (*Bohm.*), Muna (*Finnl.*), Moune (*Lappl.*), Jemurda (*Turk.*), Tochem (*Pers.*), Ménnik (*Greenl.*), Mannig (*Esquimaux*), Mootay (*Tam.*), Tookim (*Pers.*), Anda (*Sans.*).

The common Domestic Fowl is too well known to require any description. The country whence it originally came has not been correctly ascertained, although it is conjectured that it was brought from Persia by the Phœnicians, about 500 years before the birth of Christ.<sup>1</sup> As an article of food it is the least stimulating of animal substances; and the broth made of the young fowl or chicken is not only the best restorative diet for the convalescent, but is also a useful diluent in cholera, dysentery, and other disorders of the bowels. After they are a year old, their flesh becomes less and less digestible; but the Capon and Poulard retain their tenderness longer.

The Egg consists of two distinct fluid matters, the white and the yolk; the membranes which inclose these, and the shell.<sup>2</sup>

*Qualities.* — The *white* is inodorous and insipid, of a glary viscid nature, readily dissolving in water, coagulable by a heat of 165° Fahrenheit, and also by acids and alcohol. When coagulated it becomes sapid, and is no longer soluble, either in cold or hot water. From the experiments of Dr. Bostock, it appears to be composed of water 85.0, albumen 1.2, in 100 parts; and, besides, shows traces of uncoagulable matter 2.7, and salts 0.3, sulphuretted hydrogen gas, and benzoic acid. The *yolk* is also inodorous, but has a bland oily taste; and

<sup>1</sup> *British Zoology*, i. 280. There is a variety of the common fowl, named the *Dorking Fowl*, from being generally procured at Dorking, in Surrey, which has two toes behind instead of one. Another variety is found at Mozambique, and at Siam, which has the skin, bones, periosteum, and sometimes the flesh, quite black, and yet is esteemed good eating.

<sup>2</sup> Hens have been known to lay eggs when twenty years old. *Supplement to Latham*, 207.



when agitated with water forms a milky emulsion. When long boiled, it becomes a granular solid, and yields by expression a yellow, insipid, fixed oil. It consists of four constituents, water, oil, albumen, and gelatine; on the presence of the albumen depends the hardness of the boiled yolk. The *shell* consists of carbonate of lime, phosphate of lime and animal mucus. When it is burnt, the carbonic acid is dissipated, the animal cement destroyed, and pure lime, with phosphate of lime obtained. As long as the yolk remains suspended in the centre of the albumen, an egg is supposed to be good; but it spoils as soon as the yolk touches the shell. The preventing the admission of air through the pores of the shell, preserves eggs for a longer period than they otherwise could be kept good. This is effected by covering the egg with grease, or dipping it into lime water. A fresh or good egg appears semitransparent, when placed between the eye and the light: but when it is opaque or irregularly cloudy, it must be rejected.

*Medical properties and uses.* — The yolks of raw eggs are gently laxative, and have been thought serviceable in jaundice and other hepatic obstructions. Beaten up with sugar and wine, they are extremely nutritive, and are consequently useful in convalescencies, and other cases of debility. In pharmaceutical operations, the yolks are used for rendering oil and balsams miscible with water; and the whites for clarification.<sup>1</sup> The shells are antacid; but possess no advantages over chalk when unburned, or lime when they are burned.

PHYSETER. *Syst. Nat. Gmelin*. i. 227.

Cl. 1. Ord. 7. Mammalia Cete.

G. 39. *Teeth* in the lower jaw, but none in the upper. *Tube* in the head, or great front.

*Species* 2. *P. Macrocephalus*. *Spermaceti* Whale. *Willough.*

*Pisc. t. A. 1. f. 3. Phil. Trans. lx. 321. t. 9.*

*Officinal.* CETACEUM, *Lond.* SPERMACETI, *Edin.* SPERMA CETI; SEVUM, *Dub.* *Spermaceti.*

*Syn.* *Spermaceti*; Céline (*F.*), Wallrath (*G.*), Walschot (*Dutch*), Walraf (*Swed.*), *Spermaceti* (*I.*), *Espermaceti* (*S.*).

This species of whale inhabits chiefly the Southern Ocean, although occasionally seen in the European seas. It is a large fish, generally measuring about sixty feet in length, and thirty in circumference at the thickest part of the head, which is blunt, and about nine feet in height. It is of a blackish colour on the upper part of the body, and white on the belly. There are forty-six double teeth in the lower jaw, which is shorter than the upper; and in the head is a triangular, bony cavity, covered by the common integuments only,

<sup>1</sup> Owing to peculiar idiosyncrasy, the smallest portion of the white of egg cannot be eaten by some persons without occasioning pain, sickness, and an erythematic eruption on the skin.



and filled with an oily fluid, which, on the death of the fish, congeals into a spongy mass. The eyes are small: the pectoral fins near the angles of the mouth; and the tail forked.

The spongy, oily mass, is dug out from the cavity of the head, and the oil separated from it by dripping.<sup>1</sup> In this state it has a yellow unctuous appearance, and is brought to England in barrels. The following is the mode of purifying it in the great way. The mass is put into hair bags, and pressed between plates of iron, in a screw press, until it becomes hard and brittle. It is then broken in pieces and thrown into boiling water, where it melts, and the impurities, rising to the surface, are skimmed off. After being cooled, and separated from the water, it is put into fresh water in a large boiler, and a weak ley of the potass of commerce added to it by degrees. This part of the process is thrice repeated, after which the whole is poured into coolers, where the spermaceti concretes into a white semi-transparent mass, which, on being cut into small pieces, assumes the flaky aspect it has in the shops.<sup>2</sup>

*Qualities.* — Purified spermaceti is a white, crystallized, friable, semi-transparent, unctuous substance, nearly inodorous and insipid. Its specific gravity is 9.433. It melts at 112° Fahrenheit<sup>3</sup>, and at a higher temperature, 500°, it evaporates, very little altered; although by repeated distillations it is partly decomposed, and a brown, acid liquor obtained. Like the fixed oils, it leaves, when heated, on paper, a greasy stain, and can be diffused in water by means of the yolk of egg or mucilage. It is soluble in hot alcohol, ether, and oil of turpentine, but concretes again as the fluids cool; and is completely soluble in the fixed oils. When boiled with alcohol, it becomes fusible at 49°, more brilliant and less unctuous, less odorous, and more soluble in alcohol.<sup>4</sup> Of the acids, the sulphuric only acts on it, dissolving it, and forming a dark-coloured, thick, soapy solution, which has a faint smell of sulphur. The alkaline carbonates do not affect it, but it is partially dissolved in the pure alkalies; and with hot ammonia it forms an emulsion, which is not decomposed by cooling. Long exposure to hot air renders it rancid; but it may be again purified by being washed in a warm ley of potass.

*Medical properties and uses.* — Spermaceti is demulcent and emollient. It however possesses no advantages for internal use over the fixed, bland oils. It is used in dysentery and irritations of the alimentary canal, and in catarrh and phthisis: but in the latter cases it is less beneficial than the bland oils;

<sup>1</sup> An ordinary sized whale will yield upwards of twelve large barrels of crude spermaceti.

<sup>2</sup> *Monthly Magazine*, August 1809.

<sup>3</sup> Bostock, *Nichol. Journ.* iv. 134.

<sup>4</sup> Chevreuil, *Ann. de Chim. et de Phys.* tom. vii. p. 157.



for, as these are readily united with water by means of alkalies and mucilages, the compounds formed with them are more viscid, and better adapted for smearing the fauces. Several imaginary healing virtues were, formerly, supposed to belong to *Spermaceti*; on which account it was, and still is, often given to women in child-bed. It is, however, when combined with water by means of the yolk of egg, a pleasant vehicle for tincture of opium, when the after-pains are troublesome. It forms a part in the composition of several ointments.

The dose is from ʒss. to ʒjss. rubbed with sugar, or in the form of emulsion.

Official preparations. *Ceratum Simplex*, E. *Ceratum Cetacei*, L. *Unguentum Cetacei*, L. D.

PIMPINELLA. *Spec. Plant. Willd.* i. 1471.

Cl. 5. Ord. 2. Pentandria Digynia. Nat. ord. Umbellatæ.

G. 562. Fruit ovate-oblong. Petals inflected. Stigma nearly globular.

Species 8. *Pimpinella Anisum*.<sup>1</sup> Anise. *Med. Bot.* 2d edit. 135.t. 52.

Official. ANISI SEMINA, *Lond.* PIMPINELLÆ ANISI SEMINA, *Edin.* ANISUM; SEMINA, *Dub.* Anise seeds.

Syn. Graines d'Anis (*F.*), Anis (*G.*, *Dan.*, *Swed.*, *Russ.*), Anice (*I.*), Herbadoce (*Portug.*), Anis; Matalahuga (*S.*), Anys (*Dutch*), Anyz (*Poln.*, *Bohm.*), Anison (*Arab.*), Souf (*H.*), Seri nisü (*Japan.*), Rosianeh roomi (*Pers.*), Adis manis (*Javanese*), Somboo (*Tam.*), Jera manis (*Malay*).

This is an annual plant, a native of Egypt; but it is cultivated abundantly in Malta and Spain, and in our physical herb gardens<sup>2</sup>; flowering in July. It is a delicate plant, and rises about a foot only in height. The stem is striated, smooth, jointed, and branching: the lower leaves are roundish, lobed, and toothed; but the upper ones are divided into narrow, pinnated segments: the flowers are small and white, in flat, terminal umbels, without involucre; the seeds are oblong, swelling, striated, and of a greenish colour.

The Anise grown in this country ripens its seed sufficiently to be gathered about the middle of August. A greater quantity of seed, however, than is grown here, is annually imported from Malta and Spain. The Spanish is small, and generally preferred. The heaviest are to be preferred.

*Qualities*.—Anise seeds have an aromatic odour, which is increased when they are rubbed in the hands, and a sweetish, warm, grateful taste. Both alcohol and water extract their virtues; and in distillation with water they yield a yellowish volatile oil, which concretes at a temperature of 50° of Fahrenheit. An oil of a greenish colour also is obtained from Anise seeds by expression; it consists of a bland, fixed, inodorous oil, mixed with a large portion of the proper essential oil.

<sup>1</sup> Ἀννησον Dioscoridis, who says — “lactis ubertatem præstat, venerem stimulat.” l. 5. c. 65.

<sup>2</sup> A considerable quantity is cultivated at Mitcham in Surrey, chiefly for the use of the rectifiers of British spirits. *Stevenson's Survey*, 379.



*Medical properties and uses.*—These seeds are carminative; and are supposed to possess the power of promoting the secretion of milk. They are chiefly used in flatulencies, and in the tormina of infants. They are given in substance bruised, in doses of from grs. x to ʒij.

Officinal preparations. *Oleum Anisi*, L. E. D. *Spiritus Anisi*, L. PIMENTÆ BACCÆ. Vide *Myrtus*.

PINUS. *Spec. Plant. Willd.* iv. 494.

Cl. 21. Ord. 8. Monœcia Monadelphia. Nat. ord. Coniferæ.

G. 1711. Male. Calyx four-leaved. Corolla none. Stamens many. Anthers naked.

—— Female. Calyx strobiles, with a two-flowered scale. Corolla none. Pistil one. Nut with a membranous wing.

\* With double leaves.

Species 1. *P. sylvestris*. The Wild Pine, or Scotch Fir. *Med. Bot.* 2d edit. 1. t. 1. *Smith Flora Brit.* 1031. *Lambert, Description of the Genus Pinus*, i. t. 1. *Michaux, North American Sylva*, vol. iii. pl. 138.

\*\*\*\* With fascicled leaves.

Species 24. *P. Larix*. The Larch. *Med. Bot.* 2d edit. 7. t. 4. *Lambert*, 53. t. 35.

\*\*\*\*\* With solitary leaves, distinct at the base.

Species 27. *P. Balsamea*. Balm-of-Gilead Fir. *Lambert*, 48. t. 31. *Pursh*, ii. p. 639. *Abies balsamifera*. *Michaux*, iii. pl. 150.

Species 32. *P. Abies*. Norway Spruce Fir. *Med. Bot.* 2d edit. 4. t. 2. *Lambert*, 37. t. 25. *Abies picea*. *Michaux, North American Sylva*, pl. 146.

#### 1. PINUS SYLVESTRIS.

Officinal. α. TEREBINTHINA VULGARIS, *Lond. Edin.* TEREBIN'THINA VULGARIS; RESINA, *Dub.*

β. TEREBIN'THINÆ OLEUM, *Lond.* PINI OLEUM VOLATILE, *Edin.*

γ. RESINA FLAVA, *Lond.* RESINA PINI, *Edin.* RESINA ALBA, *Dub.* PIX NIGRA, *Lond.*

δ. PIX LIQUIDA, *Lond. Dub.* RESINA EMPYREUMATICA, VULGO PIX LIQUIDA, *Edin.*

Common Turpentine. Oil of Turpentine. Resin. Black Pitch. Tar.

*Syn.* α. Térébinthe (*F.*), Gemeiner Terabenthin (*G.*), Trementina (*I. & S.*), Kota (*Nepaulese*), Ratingeroo mie (*Arab.*), Zungbarie (*Pers.*), β. Huile essentielle de Térébinthe (*F.*), Terbenthinöhl (*G.*), Olio della Trementina (*I.*), Azeyte de Trementina (*S.*), γ. Resine blanche et flave (*F.*), Fichtenharz (*G.*), Ragia (*I.*), Resina (*S.*), δ. Gondron (*F.*), Theer (*G.*), Pece liquida (*I.*), Brea (*S.*).

The Wild Pine, or Scotch fir, so named from its growing wild on the Scotch mountains<sup>1</sup>, is common in most of the northern parts of Europe. It is a straight, abruptly-branched tree, rising, in a favourable soil, to the height of eighty-feet,

<sup>1</sup> When Cæsar asserted that the fir did not grow in Britain, he must have meant the *P. Abies*. The ancient name of the fir in Scotland was *Gius*, in Ireland *Giumhus*, and in Wales *Fynniduydh*. The following, also, are synonyma of this tree. *Die Kiefer* (*G.*), *Pynboom* (*Dutch*), *Furr* (*Dan.*), *Tall* (*Swed.*), *le Pin* (*F.*), *il Pino* (*I.*), *el Pino* (*S.*), *O. Pinheiro* (*Port.*), *Sosna* (*Russ.*), *Mandy* (*Finl.*), *Betze* (*Lapl.*) *Maats* (*Japan*), *Sum* (*Chinese*), *Bor.* (*Scalv.*). It prefers an arid, siliceous soil.



and four or five in diameter, covered with a rough, cracked, brownish-coloured bark, and always clothed with foliage. The leaves are short, linear, entire, pointed, concave on one side, and convex on the other, about two inches long, twisted, of a bright-green colour, and issuing in pairs from a white truncated sheath: the flowers are white, or rather of a yellowish tint: the *male catkin* is densely spiked, bracteated, elliptical, obtuse, with numerous scales crested on the upper side, but on the under bearing a sessile anther: the *female* is inferior, often ternate, or three together round the branches, peduncled, smooth, of a green colour, and changes into a small, nearly pointed, conical greyish strobile or cone, which appears tessellated and warty, and bears within each scale two small, black winged seeds.<sup>1</sup>

This tree is at its perfection when between seventy and eighty years old; but it yields turpentine at the age of forty.<sup>2</sup> Those trees which are most exposed to the sun, and have the thickest bark, afford it in the greatest abundance. The operations for procuring it commence in the month of May; the outer bark is stripped off for six inches, so as to expose the inner smooth bark, near the foot of the tree, and a wound made with a sharp tool three inches square, and an inch deep. The resinous juice soon begins to exude in transparent drops, which fall into a hole previously dug at the foot of the tree: fresh incisions are, successively, made till September, when the cold checks the further exudation. The warmer the weather is, the greater quantity of turpentine is obtained: and a healthy tree may thus yield from six to twelve pounds of turpentine annually, for a century of years. Part of the juice concretes in the wounds, and is called *Galipot* in Provence, and *Barras* in Guienne; but although it contains oil, yet it is not used for the purpose of procuring it. The proper turpentine is purified by being exposed to liquefy in the sun's rays, in barrels perforated in the bottom, through which it filters.

The *Oil of Turpentine* is obtained by distilling the resin with water in a common still, when the oil is found in the receiver swimming on the water, from which it is easily separated: the average proportion is 60lbs. of oil from 250lbs. of good turpentine. This process is carried on both abroad and at home; but the oil drawn in this country is always preferred.

*Common Resin*, or *Yellow Resin*, is the residue of the distillation of turpentine. It receives different appellations ac-

<sup>1</sup> Essence of spruce is a fluid extract prepared by coction of the twigs; which, when fermented with sugar or molasses and water, forms spruce beer.

<sup>2</sup> Great ravages are committed on this tree by *Bostrichus piniperda*, an insect which, by introducing itself into the cellular integument separates the bark from the wood, and thus destroys the life of the plant.



according to the mode in which the process is carried on. When the distillation is performed without addition, and continued to dryness, the residue is called *Common Resin*, or *Colophony*<sup>1</sup>, but when agitated with about one-eighth of fresh water while yet fluid, it is named *Yellow Resin*. A similar resin is made by melting and agitating the *Galipot* in water; and this is preferred in general to the former kind, on account of its greater ductility, which arises from its containing a portion of oil.

*Tar*<sup>2</sup> is the last officinal preparation from this species of Fir which we have to notice. The greater part of the tar imported into Britain is brought from the Baltic, i.e. Russia, Sweden, and Denmark; but a considerable quantity comes from the United States, where it is chiefly obtained from the *Pinus australis*, long-leaved pine of Michaux, and is still prepared in nearly the same method as described by Dioscorides to have been practised by the ancients. The branches of the trees are cut into billets, and piled up in large stacks which are covered with turf. Fire is then applied to the wood, and it is suffered to burn with a slow smothered flame: during which the tar is formed by the decomposition of the resinous juice, flows to the bottom, and runs out through a small channel cut for the purpose. The stacks are generally built on the slope of a hill, so that the tar is easily collected, and put into barrels; in which state it is brought to this country.

#### 1. PINUS LARIX.<sup>3</sup>

*Officinal.* α. RESINA LIQUIDA, *vulgo*, TEREBINTHINA VENETA, *Edin.* TEREBINTHINA VENETA; RESINA, *Dub.*

β. PINI OLEUM VOLATILE, *Edin.* Venice Turpentine; Oil of Turpentine.

*Syn.* Térébinthe de Venise (*F.*), Venetischer Terbenthin (*G.*), Trementina Veneta (*I.*).

There are two varieties of the Larch tree, one of them a native of America, the other of the south of Europe and Siberia. It rises erect to the height of fifty feet, sending off slender, spreading branches, which droop at their extremities. The leaves are deciduous, soft, and of a bright-green colour; in tufts, generally containing forty or more, springing from short, thick, corrugated sheaths, and spreading like a painter's brush; linear, somewhat obtuse, and entire: the male flowers, which appear in April, are in small, lateral, cylindrical catkins, with the apexes of the anthers inflated; the female are in erect, ovate catkins, twice as large as the male, and in some instances purple at the top: the strobiles or cones are about an inch long, obtuse, and purplish at the apex, with the scales

<sup>1</sup> The Colophonia of the ancients was a liquid resin, named from Κολοφών, a town of Ionia in Asia Minor, whence it was brought.

<sup>2</sup> Κωνον Græcorum.

<sup>3</sup> Πίτυς Theophrasti. Lärchenbaum (*G.*), Iorkenboom (*Dutch*), Lerketree (*Dan.*), Meleze (*F.*), Larice (*I. S.*), Lariceo (*Port.*), Listweniza (*Russ.*). The larch tree has been cultivated in England since 1629. *Hort. Kew.*



smooth on the surface, lacerated at the edges, and concealing under each two, winged seeds.

The Larch tree grows to very great perfection in the forests of Baye in Provence, where a very large proportion of the Venice turpentine of commerce is procured. It is obtained by boring a hole with an auger into the heart of the tree, at about two feet from the ground, and fitting into it a small pipe, through which the turpentine flows slowly into vessels placed for its reception. This process is begun in May, and continued till September; when the different quantities collected are put together, and purified by straining through cloths or hair sieves. No trees under twelve inches in diameter are tapped; but vigorous trees will yield, annually, seven or eight pounds for forty or fifty successive years, or during the term of their life.<sup>1</sup> Much of the Venice turpentine of the shops is brought from America; and is perhaps procured from a different species of fir.

The *Essential Oil* is separated from it by distillation in the same manner as from the common turpentine.

### 3. PINUS BALSAMEA.<sup>2</sup>

*Officinal.* TEREBINTHINA CANADENSIS, *Lond.* BALSAMUM CANADENSE,<sup>3</sup> *Dub.* Canada Turpentine. Canada Balsam.

*Syn.* Baume de Canada (*F.*), Kanadischer balsam (*G.*).

This tree is a native of the coldest regions of North America, flowering in May. It is a straight, elegant tree, rarely exceeding 40 feet in height, and 12 or 15 inches in diameter, covered with a smooth, whitish-grey bark. The leaves are in double rows like a comb, short, not exceeding 8 lines in length, and linear, but broader than the former two species, and less pointed; of a bright-green colour on the upper surface, marked with whitish lines underneath, and fragrant: the cones, which ripen in October, stand erect on the branches, are large, nearly cylindrical, of a beautiful, deep, glossy, purple colour, inclining to black; and exude a great quantity of transparent resin, which gives them a very rich, beautiful appearance.

The manner in which the Canada balsam, or fine turpentine, yielded by this tree is collected, is by incisions in the body of the tree from which it exudes. It exists in great

<sup>1</sup> Besides turpentine the Larch tree exudes a species of manna, which is named Briançon manna. It is in little white concrete drops, which adhere to the leaves, and tastes sweet like new honey; but it has the flavour of turpentine, which it contains. The inner part of the tree yields also a gum similar in its properties to acacia gum, of a reddish colour, with a slight resinous taste. In Russia it is officinal, and sold, as Pallas observes, under the improper name of Orenberg gum, being obtained from the Uralian Forest. *Flora Rossica*, i. p. 2, 3.

<sup>2</sup> *Balsamtanne* (*G.*), *Le beaumier de Gilead* (*F.*).

<sup>3</sup> The Dublin college have erred in retaining this name; for, as this substance does not afford benzoic acid, it cannot be ranked as a balsam.



quantity in vesicles between the wood and the bark. Canada balsam is brought to this country in casks, each containing about one hundred weight.

#### 4. PINUS ABIES.<sup>1</sup>

*Officinal.* α. ABIETIS RESINA, *Lond.* Resin of the spruce fir.

β. PIX ABIETINA, *Lond.* RESINA SPONTE CONCRETA, *vulgo.*

—PIX BURGUNDICA, *Edin.* PIX BURGUNDICA, *Dub.* Burgundy Pitch.

The Norway spruce Fir is a native of Europe, and of the moist parts of northern Asia; flowering in April. It is a lofty, noble tree, rising 150 feet in height; and from 3 to 5 feet in diameter; is straight, pyramidal, and covered with a reddish, scaly bark. The leaves are short, thickly set upon the branches, slightly carinated on both sides, of a dusky-green colour, shining on the upper surface, and often curved: the male catkins are ovate, purplish, and scattered in the axils of the leaves; the female are also purple, and generally terminal; the strobiles or cones are long, nearly cylindrical, greenish before they are ripe, but afterwards purple, and always pendent: the scales, which are arranged in eight spiral rows, have an oval shape terminating in a point, and become ragged at the edges.<sup>2</sup>

The *Resin* or *Thus* of the old London Pharmacopœia exudes spontaneously from the bark of the Norway Spruce Fir, and concretes as it exudes. It undergoes no preparation, but is brought to us in the form of tears or small masses, packed in casks, each containing from one to two hundred weight. The greater part comes from Germany, but a small quantity of a purer description comes from France.

*Burgundy Pitch* is obtained by making incisions through the bark so as to lay bare the wood. It concretes in the form of flakes at the incisions, which are detached by an iron instrument once a fortnight during the summer, and fresh incisions successively made. The flakes, after being detached, are put into large boilers with a sufficient quantity of water, melted, and then strained through coarse cloths under a press. The greatest quantity is collected in the neighbourhood of Neufchatel, whence it is brought to this country packed in casks. A fictitious sort is made in England, and found in the shops under the title of *Common Burgundy pitch*. It may be distinguished by its friability, want of viscidness, and unctuousness, and the odour which characterizes the genuine sort.

<sup>1</sup> Ἐλάτη Theophrasti. *Die Fichte* (G.), *Hartsboom* (Dutch), *Gran.* (Dan., Swed.), *La Pesse* (F.), *Picea* (I. S.), *Peuce* (Port.), *Jel* (Russ.). This species of Fir is cultivated in Britain, but it does not appear to have been introduced before 1739.

<sup>2</sup> Like the Scotch Fir, it is attacked by the *Bostrichus piniperda*.



*Qualities.* — **TURPENTINES.** Although these are produced from different species of the Pine tribe, and one sort from the *Pistacia Terebinthus*, yet all of them possess the same general and chemical properties. They have a peculiar, somewhat aromatic odour, and a warm, pungent, bitterish taste; are semifluid, tenacious, translucent, combine readily with fixed oils, and are inflammable, burning with a white flame and much smoke. Alcohol and ether dissolve them entirely, leaving the impurities; but water takes up only their flavour. When distilled with water a volatile oil comes over, and towards the end of the operation succinic acid rises<sup>1</sup>; resin remains in the retort: the turpentine therefore are compounds of these three substances. But each sort of turpentine has characteristic qualities which require to be noticed; 1. *Common Turpentine* has a strong, somewhat fragrant odour, and a bitter, disagreeable taste; its consistence is greater than that of honey; its colour is dirty yellow, and it is more opaque than the other sorts. 2. *Venice Turpentine* is more fluid, having the consistence of new honey, a yellowish colour, and is less unpleasant to the smell and taste than the common. 3. *Canadian Balsam* (or more correctly *Turpentine*) has a strong not disagreeable odour, and a bitterish taste; is transparent, whitish, and has the consistence of *Copaiva Balsam*. 4. *Chian or Cyprus Turpentine* (see *Pistacia*) is very fragrant, but almost insipid, nearly transparent, thick, tenacious, and of a whitish colour.

*Oil of Turpentine* has a strong, penetrating, peculiar odour, and a hot, pungent, bitterish taste. It is perfectly limpid and colourless: extremely light, volatile, and inflammable; and dissolves completely in six parts of sulphuric ether; but although hot alcohol readily dissolves it, yet it again separates in drops as the spirit cools, and is very sparingly soluble in the cold in the strongest alcohol: one hundred parts in volume, however, unite with 20 of alcohol, and form a homogeneous solution, which does not become turbid by water.<sup>2</sup> In all other respects it agrees with the other essential volatile oils. A stream of oxymuriatic gas passed through it converts it into a yellow resin.

*Tar* has a strong odour familiar to every person; a resinous, subacid, bitterish taste: and a coarse, thick consistence, with a deep brown colour, approaching to blackness, derived from the charring of the wood during its formation. It consists principally of empyreumatic oil, resin, and acetic acid; is partially soluble in water; and inspissated by boiling into pitch.

*Yellow and White Resin* are varieties of the same substance.

<sup>1</sup> *Annales de Chim.* xxi. 328.

<sup>2</sup> *Ibid.* xix. p. 279.



They are nearly inodorous when cold; but, when heated, emit a slight terebinthinate odour. Their taste is slightly acrid and bitterish; and their colour a dull whitish-yellow, or a greenish-yellow. The mass of resin is semipellucid, brittle, breaks with a true vitreous fracture, and adheres moderately to the fingers. Its specific gravity is 1.0742. It melts when heated, then inflames, and burns with a yellow flame giving out much smoke. It is insoluble in water, but entirely soluble in alcohol, ether, the fixed oils, and the alkalies. The acids also dissolve resin, and convert it into artificial tannin; with the exception of the acetic acid, which only dissolves it. When sulphuric acid is employed, charcoal, in the proportion of forty-three per cent. of the resin acted on, is produced.<sup>1</sup> The *Resin of the Norway Spruce* possesses nearly the same properties. It is in the form of solid, brittle tears, of a brownish-yellow colour on the outside, and internally white; and emits a very agreeable odour when burning.

*Burgundy Pitch* has a terebinthinate odour and taste, is brittle, opaque, and of a light yellow, or reddish-brown colour. It softens moderately in the heat of the hand, appears unctuous, and has a considerable degree of tenacity.

*Medical properties and uses.*—The *Turpentine*s and their *Essential Oil* are stimulant, cathartic, diuretic, and anthelmintic; and externally rubefacient. Of those which I have described, the Venice and Canada turpentine are more generally employed for internal purposes; the common turpentine proving offensive to most stomachs, and the Chian not being easily procured. The ancients were well acquainted with the medicinal properties of turpentine<sup>2</sup>; and, besides the diseases for which they are prescribed by the moderns, gave them liberally in coughs and all pulmonary affections. Turpentine seems to derive their virtues from the oil they contain. When swallowed, they produce a sensation of warmth in the stomach, at first increasing the quickness and force of the pulse, but afterwards diminishing it; and if the dose be large, some degree of nausea is excited, with slight vertigo; and soon, but not always, a copious discharge from the bowels; but if the dose be small, they act chiefly upon the kidneys. The cathartic operation of large doses of the oil, in particular, seems to counteract the determination to the kidneys, which smaller doses produce; for in doses of even fʒx and fʒxij, no other effect on the urinary organs is perceived than the violet smell of the urine.<sup>3</sup> The odour of violets is produced by the oil, even when it is not taken into the stomach,

<sup>1</sup> Hatchett, *Phil. Trans.* 1806.

<sup>2</sup> See *Dioscor.* lib. i. cap. 91. p. 50. *Aretæus passim.* *Alpinus de Med. Egypt.* lib. iv. &c.

<sup>3</sup> *Transactions of the London Medical Society*, i. part 1. 212. 227.



or rubbed upon the skin; for if a quantity of Oil of Turpentine be poured on a table in a room, this odour will be perceived in the urine of any one who remains in the room for half an hour, or even a shorter time. Turpentine is chiefly prescribed in gleet, leucorrhœa, mucous obstructions of the urinary passages, and calculous affections; but in the latter cases their stimulant operation on the kidneys requires that they be given with caution. The Oil is justly regarded as a useful remedy in lumbago, sciatica, and some other varieties of chronic rheumatism and neuralgia, particularly when combined with the Cinchona bark. Dr. Copland, in a valuable paper on terebinthinous remedies<sup>1</sup>, recommends the Oil strongly in the hæmorrhagiæ, particularly in atonic epistaxis and hæmoptysis. He also confirms Dr. Percival's statement of its efficacy in epilepsy; and extols its powers in infantile convulsions, arising from a disordered state of the alimentary canal and erysipelas. It exerts a powerful influence on the uterine organs; and, thence, is useful in chlorosis. Dr. Copland states some cases of ovarian dropsy, in which the effects of the Oil were such as to recommend its employment in incipient cases of this disease; and, also, in other dropsies, not even excepting hydrocephalus. For the expulsion of the tape-worm the power of the Oil of Turpentine is now generally known. It differs in its action from the other remedies which have been employed against tape-worm, by killing the worm before it throws it out; and thence is more permanently useful.<sup>2</sup> Neither wine nor spirits should be drunk during the use of the Oil; the usual quantity of food should be diminished: and the use of the Oil should be immediately discontinued, if an eruption resembling eczema appear on the skin; which is apt to arise from its employment in some habits. As local stimulants, turpentine, and the Oil of Turpentine have been efficaciously exhibited in the form of enema, in cases of colic, obstinate costiveness, and ascarides. The Oil is useful when dropped into the ear in deafness from defect of wax; and is an excellent addition to embrocations in acute rheumatism, bruises and paralysees of the extremities. As a discutient, it is applied to indolent tumours, and is a useful, primary application to burns.

Turpentine is given in doses of grs. x. to ʒj, either made into pills with powdered liquorice root, or diffused in water by means of almonds, mucilage, or yolk of egg and sugar. The dose of the Oil may be ℥x to ʒj, to produce its diuretic effect: but in doses of fʒj to fʒij, its effects are

<sup>1</sup> *Med. and Phys. Journ.* vol. xlv. p. 185. 206.

<sup>2</sup> In all the cases of the expulsion of tænia by oil of turpentine, the ejected worm has generally had a livid hue, without any appearance of animation.



more general on the system. In these doses, it may be combined with aromatics and spices, and rubbed up with mucilage or honey. Dr. Copland recommends the addition of the tincture of Capsicum, for correcting the nauseating and unpleasant effects which the Oil frequently produces on the stomach. For the expulsion of tænia, it is necessary to give from fʒss to fʒij of the oil, repeated every eight hours till the worm be ejected: and in these large doses it is more easily taken when exhibited uncombined, or when merely floated upon water with the addition of a drop or two of any aromatic oil. If it do not operate by stool in four or five hours after it has been taken, a dose of castor oil should be exhibited.

*Tar* is stimulant, diuretic, and sudorific; and externally detergent. In the solid state, made into pills, it has been beneficially used in Ichthyosis; but it is, more generally, employed in the form of Tar Water, *Aqua Picis liquidæ*. As an external application, it has been found beneficial in porrigo scutulata, foul ulcers, and some other cutaneous diseases.

The *Resins* and *Burgundy Pitch* are adapted for external use only; the former entering into the composition of some ointments and plasters; the latter being used as a rubefacient plaster. The pitch excites some degree of inflammation, sometimes a pimply eruption, and a serous exudation from the part over which it is applied, without raising the cuticle. It is used in cases of catarrh, pertussis, dyspnœa, and chronic rheumatism; and seems to be chiefly serviceable from the length of time it adheres to the skin, and by which its action can be continued.

Officinal preparations. Of turpentine — *Oleum Terebinthinæ*, D. Of the Oil — *Linimentum Terebinthinæ*, L. Of Yellow Resin — *Emplast. Resinæ*, L. E. D. *Ceratum Resinæ*, L. E. Of Burgundy Pitch — *Emplast. Picis. comp.*, L. E. Of Tar — *Aqua Picis liquidæ*, D. *Unguentum Picis liquidæ*, D.

PIPER. *Spec. Plant. Willd.* i. 159.

Cl. 2. Ord. 3. Diandria Trigynia. Nat. ord. Piperitæ, Linn. Urticæ, Juss.

G. 74. Calyx none. Corolla none. Berry one-seeded.

Sp. 1. *P. nigrum*. Black pepper. *Med. Bot.* 2d edit. 721. t. 246. Melago codi. *Rheede Hort. Malabar.* vii. 23. t. 12. Marsden, p. 105.

Sp. 12. *P. longum*. Long pepper. *Med. Bot.* 2d edit. 724. t. 247. Cattu-tirpali. *Rheede Hort. Malabar.* vii. 27. t. 14.

#### 1. PIPER NIGRUM.<sup>1</sup>

Officinal. PIPER NIGRI BACCÆ, *Lond.* PIPERIS NIGRI FRUCTUS, *Edin.* PIPER NIGRUM; BACCÆ, SEMEN, *Dub.* Black pepper.

Syn. Le poivrier commun (*F.*), Gemeine Pfeffer (*G.*), Gemeene Peper (*Dutch*), Swarteller Stark peppar (*Swed.*), Pepe nero (*I.*), Pimienta (*S.*), Mame

<sup>1</sup> Πεπερι Dioscoridis.



(*Japan*), Molago-Codi (*Malabar*), Fulful-filfil (*Arab.*), Mirch (*H.*), Maricha (*San.*), Mellaghoo (*Tam.*), Gummeris (*Cyng.*), Lada (*Malay*), Maricha (*Jav.*).

This species of Pepper is a native of the East Indies; and is very abundantly cultivated at Malacca, Java, and Sumatra, whence the whole of Europe is supplied. It is a climbing plant, the stem being round, smooth, jointed, and swelling towards each joint, woody, slender, branched, and from eight to twelve feet in length. The leaves are petiolate at the joints of the branches, ovate, entire, pointed, seven-nerved, and of a dark green colour. The flowers are diœcious, sessile, white, small, each supported by a scale: they cover, thickly, a cylindrical spadix; without any regular calyx or corolla. The filaments are two or more, flat, awl-shaped; and the anthers roundish. The germen is ovate, and crowned with a deeply divided stigma: the fruit is a globular berry of a red-brown colour.

In Sumatra, the Pepper vines are propagated by cuttings or suckers. In growing they are supported by props called *Chinkareens*, which are cuttings of *Morinda citrifolia*, or of an *Erythrina*, at the root of each of which two vines are planted.<sup>1</sup> The plants are three years old before they bear fruit, and they bear for eight years. The berries are four or five months in coming to maturity; are gathered as soon as any of them ripen; then spread upon mats to dry, and trodden to separate the fruit from the stalk; when they become black, and more or less shrivelled. The vines yield two crops yearly; the first in December, the second in July. White Pepper is the ripe and perfect berries, freed of their outer coat by means of a preparation of lime and mustard oil, called *Chinam*, applied before it is dried. The Pepper is now also cultivated to a considerable extent in India.

*Qualities.*—Black Pepper has an aromatic odour, and a hot, pungent taste. Its virtues are entirely extracted by ether and alcohol, and partially by water. The aqueous infusion is brown, and reddens vegetable blues; and the decoction of the ground Pepper forms a precipitate with infusion of galls, which dissolves again when the fluid is heated to 120°. When the alcoholic infusion is distilled, a green, resinous, oily matter is left, which appears to be the source of the odour and the taste of the Pepper. M. Pelletier<sup>2</sup> has lately ascertained, that when this green fatty matter is washed in warm water, and again dissolved in hot alcohol, it deposits, after some days, a number of small crystals, which, when purified, are

---

<sup>1</sup> Dr. Roxburgh began the cultivation of Black Pepper in the Circars in 1787. The prop trees he used were the Moochy wood tree, *Erythrina Corallodendron*. One thousand plants yield from 500lbs. to 1000lbs. of pepper.

<sup>2</sup> *Annales de Chim. et Phys.* xvi. p. 20.



insipid; and have the singular property of imparting a blood-red colour to strong sulphuric acid. This substance, which has no affinity with alkalies, M. Pelletier has named *Piperin*. He found in Pepper the following components. — Piperin; a very acrid, concrete oil, on which the acrimony of the pepper depends, a volatile balsamic oil, a gummy coloured matter, extractive, malic, and tartaric acids, starch, lignin, and earthy and alkaline salts. Ether digested on powdered pepper takes up three parts in ten; and when evaporated on water deposits an intensely hot, biting, yellowish, oily resin, with the odour of the pepper, and insipid extractive matter. Black Pepper, when purchased in the state of powder, is generally adulterated. It is often mixed with the powdered husks of mustard, which are openly sold by the makers of mustard for this purpose, under the title of P. D. (*Pepper dust*).

*Medical properties and uses.* — Black Pepper is stimulant and carminative. Its use, as a condiment, is well known; and although in general it is not hurtful, but rather useful to those who have a weak digestion, yet, even in small quantities, it proves injurious in inflammatory habits, and to those subject to piles.<sup>2</sup> As a medicine, Pepper is found sometimes serviceable in checking nausea and vomiting, and removing hiccough. It is also used as a stimulant in retrocedent gout, and in palsy. The watery infusion forms a useful gargle in relaxation of the uvula. The *Peperin* which it yields has been employed successfully, by Dr. Meli and other Italian physicians, in intermittents and other fevers, and is said to be more efficacious than the preparations of Quinia.

The dose of Black Pepper may be from grs. x to ʒj.

Officinal preparation. *Unguentum Piperis Nigri*, D. *Emplastrum Meloes vesicat. comp.*, E.

## 2. PIPER LONGUM.

*Officinal.* PIPERIS LONGI FRUCTUS, *Lond. Edin.* PIPER LONGUM; FRUCTUS, *Dub.* Long pepper.

*Syn.* Poivre longue (*F.*), Langer Pfeffer (*G.*), Lange peper (*Dutch*), Long Peppar (*Swed.*), Pepe lungo (*I.*), Pimienta larga (*S.*), Pimenta longa (*Port.*),

<sup>1</sup> Dr. Meli has proposed the following method of procuring pure Piperin. Digest lbj. of Black Pepper powdered, in ʒjss. of alcohol at 36°, in a gentle heat. Then raise it to ebullition, after which leave it to cool, and having poured off the fluid, repeat the operation with a fresh quantity of alcohol. Pour upon the mixed tinctures, ʒj. of distilled water and fʒjss. of hydrochloric acid. A fatty precipitate will fall down, which should be separated by filtration. The crystals which will form on the sides of the vessels and on the filter are pure Piperin. Adding water till a precipitate ceases to be produced, a fresh quantity is obtained. *Ann. Univ. di Med. t.* 27.

<sup>2</sup> It nevertheless is an ingredient in a celebrated nostrum for the cure of piles, which is sold under the name of *Ward's paste*. This consists of equal parts (lbj.) of black pepper and elecampane; of fennel-seeds, lbjss., and of honey and sugar, equal parts (lbj.) beaten together and well mixed in a mortar. The dose is the size of a nutmeg three times a day.



Darfilfel (*Arab.*), Pipel (*H.*), Tipiliē (*Tam.*), Pippāli (*San.*), Filfil drāz (*Pers.*), Tabee (*Malay*), Chabijawa (*Jav.*).

This plant is a perennial, a native of Malabar and Bengal. The stems are round, smooth, branched, slender, and scandent: the leaves are commonly cordate, pointed, nerved and of a deep green colour: the flowers are small, in dense, short, terminal spikes, nearly cylindrical. In other circumstances, the fructification agrees with the former species. The fruit consists of very small one-seeded berries or grains imbedded in a pulpy matter, forming what Mirbel terms “*serose*.”

The fruit is hottest in its immature state, and is, therefore, gathered while green and dried in the heat of the sun. It is imported in the entire spikes, which are about one inch and a half long, and indented on the surface.

*Qualities.* — Long Pepper has a weak, aromatic odour, an intensely fiery, pungent taste, and a dark grey colour. Its constituents appear to be similar to those of Black Pepper. Ether digested on powdered long pepper takes up two parts and a half in ten parts, and when evaporated on water deposits a resin less hot than that of Black Pepper, but more permanent, and a smaller proportion of extractive. According to an analysis of Dulong, the constituents of Long Pepper are Piperin, a concrete fatty matter, in which the pungency and acrimony resides, a small portion of volatile oil, extractive, a coloured gummy matter, starch, a large quantity of bassorine, a malate, and some other saline substances.<sup>1</sup>

#### PIPER CUBEBA.

*Officinal.* CUBEBA, *Lond.* CUBEBS.

*Syn.* Cubebes (*F.*), Kubeben (*G.*), Koebar (*Swed.*), Koebeben (*Dutch*), Cubebi (*I.*), Cubebas (*S.*), Cobibas (*Port.*), Cubab chinie (*Hind.*), Komuchus (*Batavian*), Val Mellaghoo (*Tam.*), Duncke mirchie (*Duk.*), Koomoonkoos (*Malay*), Salavamirrialoc (*Til.*), Kebabeh (*Arab.*), Walgummeris (*Cyngal.*), Sirgandha marichum (*Sans.*), Kumukus (*Javan.*).

The plant which yields this spice is a native of Java, Nepaul, Batavia, Guinea, and the Isle of France. The younger plants differ from the older: their branches are long, creeping, and rooting; their leaves cordato-lanceolate, attenuated, about one and a half inch in length, and supported on petioles the length of the leaf. In the older plants the branches are flexuose and tetragonous; the leaves are less than an inch in length, unequal at the base, and supported on channelled footstalks half an inch in length. The fruit is a berry growing in clusters.<sup>2</sup>

Cubebs are brought to this country packed in cases. The best are about the size of a small pea, round, plump and heavy. They have a short stalk attached to each, which appears to terminate in raised veins on the surface of the berry.

<sup>1</sup> *Journ. de Pharm.* Fevrier 1825.

<sup>2</sup> *Medical Repository*, Dec. 1820, p. 523.



*Qualities.* — Cubebs when chewed have a pungent, aromatic, slightly bitter taste; which, however, leaves a sensation of coolness on the palate resembling that which is produced by peppermint. Their odour is fragrant and agreeable. According to Vauquelin, Cubebs contain a volatile, almost concrete oil, a resin resembling that of copaiva, another coloured resin in small quantity, extractive, and some saline substances.<sup>1</sup> The colour of the powder is darker than that of the berries. It should be kept in well corked bottles, and should even be dispensed in stoppered phials, as it quickly loses its active part when it is kept in paper. It is often adulterated with pimenta, which may be detected by the odour.

*Medical properties and uses.* — Cubebs are diuretic and slightly purgative. In gonorrhœa and gleets they have been long used by oriental practitioners: and the Arabs employ them in seasoning food.<sup>2</sup> They have been found beneficial in this country in gonorrhœa, in which they moderate the inflammation and consequently the discharge; this effect is soon apparent, and unless it be so the remedy is seldom useful.

The following statement of the results of fifty cases treated by Mr. Broughton, gives some idea of the time in which Cubebs produce their effects; and of their value as a remedy in gonorrhœa:

“Patients cured in from two to seven days	-	10
- - - eight to fourteen	-	17
- - - fifteen to twenty-one	-	18
- - - twenty-two to thirty	-	1
- - - in fifty-five days	-	1
Patients in whom no sensible effects were produced		3

Total 50”<sup>3</sup>

Mr. Jeffrey says, that, even when they failed, he found the symptoms afterwards yield readily to Copaiva; an observation which is confirmed by our own experience, and that of Mr. Broughton. They have also been lately given with advantage in inflammation of the mucous membrane of the intestines.

The powder, which is the best form in which Cubebs can be administered, sometimes nauseates: it acts as a diuretic in large doses, imparts an odour to the urine; and gives a cool sensation to the rectum, in passing the fæces. It sometimes, also, produces a quick pulse, a burning heat in the palms of the hands and the soles of the feet, flushing of the face, headaches, and frequently a slight degree of giddiness. The dose of the powder is from ʒi. to ʒjss. four times in the day. Of

<sup>1</sup> *Pract. Observ. on the Use of Cubebs in Gonorrhœa.* Lond. 1821.

<sup>2</sup> Martyn's Gardener's Dict. Art. Piper.

<sup>3</sup> *Medico Chirurg. Trans.* vol. xii.



The tincture, made by digesting ℥iij. of the bruised pepper in in Oj. of rectified spirit of wine, 3j. may be given in a glass of water three times a day.

PISTACIA. *Spec. Plant. Willd.* iv. 752.

Cl. 22. Ord. 5. Diœcia Pentandria. Nat. ord. Amentaceæ, Linn. Terebintaceæ, Juss.

G. 1782. Male. Calyx five-cleft. Corolla none. Female. Calyx three-cleft. Corolla none. Styles three. Drupe one-seeded.

Species 4. *P. Terebinthus*. Chian Turpentine-tree. *Med. Bot.* 2d edit. 29. t. 12. *Du Hamel Arbres*, ii. t. 87.

Species 6. *P. Lentiscus*. Mastich-tree. *Med. Bot.* 2d edit. 26. t. 11. *Du Hamel Arbres*, ii. t. 136.

#### 1. PISTACIA TEREBINTHUS.<sup>1</sup>

Officinal. TEREBINTHINA CHIA, *Lond.* Chian Turpentine.

*Syn.* Térébinthe de Chio (F.), Zyprischer Terebinthin (G.).

The tree which yields the Chian Turpentine is a native of Barbary and the south of Europe. It is cultivated in the islands of Chios and Cyprus, and also bears the severity of our climate; where, however, it is cultivated only as an ornamental tree, flowering in June and July. It is low in stature, sending off many spreading branches, and is covered with a smooth bark: the leaves are pinnate, composed of three pair of lanceolate, ovate, veined, entire leaflets, with a terminal one: the male and female flowers are on different trees. The *male* are in amentum, with the calyx divided into five small, ovate segments, the filaments four or five in number, very short, and supporting large, brown, erect, quadrangular anthers. The *female* are placed on a common peduncle in alternate order: consisting of a calyx of three small squamous segments, and an ovate germen crowned with two or three styles, with reflected clubbed stigmas. The fruit is subovate, reddish, smooth, and gibbous towards the top on one side.

The Turpentine is gathered chiefly in Chios, by making incisions in the bark of the trunk of the tree, in the month of July. It is allowed to flow upon stones placed at the bottom of the tree, and after being condensed by the cold of the night, is scraped off the following morning before sun-rise. It is then reliquefied by the heat of the sun, and strained to free it from any extraneous matter; and in this state is imported into this country in casks. On account of its high price, Chian Turpentine is often adulterated with common turpentine.

*Qualities.*—Chian Turpentine has a fragrant odour, a moderately warm taste, devoid of acrimony or bitterness; and a white or very pale yellow colour: it has the consistence of thick honey, is clear, transparent, and tenacious; and in its

<sup>1</sup> Τέρρινθος Dioscoridis. *Der Terpentibaum* (G.), *Terpentintrae* (Dan.), *Le Terebinthe* (F.), *Il Terebinto* (I.), *Cornicabra* (S. Port.), *Skipidarnoe derevo* (Russ.).



other qualities, as well as its medicinal properties, resembles the other turpentine. See *Pinus*.

## 2. PISTACIA LENTISCUS.<sup>1</sup>

*Officinal.* MASTICHE, *Lond.* PISTACHIE LENTISCI RESINA, *Edin.* Mastic.

*Syn.* Mastic (*F.*), Mastix (*G.*), Mastice (*I.*), Almastiga, Almaciga (*S.*), Almaceda da India (*Port.*), Roomie Mustike (*Tam.*), Sakes (*Turk.*), Arah (*Arab.*), Kinneh (*Pers.*).

The Lentisck, or Mastich tree, is a native of the Levant, particularly the island of Chios. It flowers in May, and ripens its fruit in August.<sup>2</sup> It is a low tree, seldom exceeding twelve feet in height, and eight inches in thickness; is covered with a smooth brown bark; and towards the top sends off numerous branches: the leaves are abruptly pinnate; consisting of five or six opposite pairs of narrow, ovate leaflets, of a full lucid-green colour on the upper, and a pale hue on the under side; they are sessile on the common footstalk, which has a narrow, foliaceous membrane or wing on each side, running from one pair of leaflets to the other. The male and female flowers are on distinct trees, and resemble those of the former species: the fruit is a drupe, containing an ovate, smooth nut, of a brownish colour when it is ripe.

Mastic is most abundantly obtained in the island of Chios. Transverse incisions are made in the trunks and branches of the Lentisck trees, from the 15th to the 20th of July, from which the Mastic slowly exudes, some dropping on the ground, which is made smooth and hard as a pavement for the purpose of receiving it; and some remaining fixed on the trees, and hardening so as to require, for its detachment, the aid of a sharp iron chisel. In both instances it concretes into a yellowish, semi-transparent substance. It is not gathered until August, when fresh incisions are made, and a second gathering takes place about the middle of September: no more incisions are made after this period of the year, but the gathering is continued, twice a week, until the 19th of November.

The low trailing Lentiscks yield the finest Mastic, and in the greatest quantity. Chios exports annually about 1508 cwts. part of which is brought to this country packed in chests.<sup>3</sup> That which is in the form of brittle grains is the best.

<sup>1</sup> *Exivos* Dioscoridis. *Der Mastixbaum* (*G.*), *Mastikboom* (*Dutch*), *Mastix-træ* (*Dan.*), *Lentisque* (*F.*), *Lentisco* (*J. S. Port.*), *Xihudia* (*Turk.*).

<sup>2</sup> It appears to have been cultivated in Britain so early as 1664. But it never attains here any degree of perfection.

<sup>3</sup> *Olivier's Travels* (translation) ii. 90. Olivier says, a soft mastic having all the quantities of mastic, except in its consistence, which is that of turpentine, is procured by engrafting the lentisck on the Chian turpentine tree.



*Qualities.* — Mastic is almost inodorous, unless when rubbed or heated, when it exhales an agreeable fragrant odour. It is nearly insipid; and when chewed, at first crumbles, feeling gritty between the teeth, but by degrees becomes soft and white. When it is heated it melts. Ether dissolves it entirely; but in alcohol about one-fifth remains undissolved, which has, when moist, the characters of caoutchouc<sup>1</sup>, but becomes brittle when dried; and therefore appears to be a peculiar, vegetable principle. It appears to amount to nearly a fifth of the Mastich.<sup>2</sup> In some respects this principle resembles the pure resins, being brittle, semitransparent, fusible, insoluble in water, and soluble in ether; but it differs in being insoluble in alcohol. Mastich, when distilled with either water or alcohol, gives over very little with these liquids<sup>3</sup>; and this is, perhaps, an essential oil.

*Medical properties and uses.* — Mastic has generally been regarded as astringent and diuretic, and is ordered for the same diseases as turpentine: but its virtues, if it has any, are very trifling. The wood and leaves of the Lentisk were used by the ancients in fluor albus and ulcerations of the uterus; and the Turkish and Armenian women use the resin as a masticatory for cleaning the teeth, and giving an agreeable smell to the breath. It is employed to fill the cavities of carious teeth, for which purpose it is well adapted, from its property of softening in the mouth, and imparting little taste.

PIX ABIETINA. Vide *Pinus Abies*.

PIX LIQUIDA. Vide *Pinus sylvestris*.

PIX NIGRA. Vide *Pinus sylvestris*.

PLUMBUM, *Edin.* Lead.

*Syn.* Plomb (*F.*), Blei (*G.*), Lood (*Dutch*), Blye (*Dan.*), Bly (*Swed.*), Olow (*Polish*), Piombo (*I.*), Plomo (*S.*), Chumbo (*Port.*), Swinez (*Russ.*), Soorb (*Pers.*), Anuk (*Arab.*), Sisa (*H.*), Eeeum (*Tam.*), Sisaca (*San.*), Sheesh (*Duk.*), Temaetain (*Malay*), Ak-kil-lě-look (*Esquimaux*).

This is a metal of a blueish grey colour, occurring in great abundance in most countries of both hemispheres of the globe, in primitive transition, and floetz formations. It is found, A. in its metallic state:

i. Sulphuretted.

Sp. 1. *Galena*.

Var. a. Common.

b. Compact.

2. *Blue lead ore*.

a. and combined with }  
antimony.

3. *Antimonial sulphuret*.

<sup>1</sup> *Crell's Annals*, 1794, ii. 185. *Thomson's Chemistry*, 4th edit. vol. v. 93.

<sup>2</sup> See Mr. Mathew's experiments, *Nicholson's Journ.* vol. x. 247.

<sup>3</sup> Hoffman (*Observ. Phys. Chim. Select.* p. 68.), however, states, that by rubbing the mastich in a mortar, with its weight of carbonate of potass, and then distilling with alcohol, the liquid which comes over has both the smell and the taste of mastich.



## B. united with oxygen :

## ii. Oxides.

1. *Yellow oxide*.
2. *Native minium*.

## iii. combined with carbonic acid. }

1. *Carbonate of lead*.
2. *Earthy lead ore*.  
Var. *a*. Indurated.  
    *b*. Friable.
3. *Black lead ore*.
4. *Murio-carbonate of lead*.
5. *Phosphate of lead*.  
Var. *a*. Brown lead ore.  
    *b*. Green lead ore.
6. *Arsenia-phosphate*.
7. *Chromate of lead*.  
    *lead of Siberia*.
8. *Sulphate of lead*.
9. *Molybdate of lead*.
10. *Arseniate of lead*.

*b*. — with muriatic acid.*c*. — with phosphoric acid.*d*. — with chromic acid.*e*. — with sulphuric acid.*f*. — with molybdenic acid.*g*. — with arsenic acid.

Galena is the ore from which metallic Lead is commonly procured. When brought up from the mine, the ore is broken in pieces, and the impurities, which are mostly iron pyrites, quartz, calcareous spar, and clay, are separated by picking and washing : it is then exposed to a strong heat in a common reverberatory furnace till the sulphur is all separated, after which the metal is brought into a state of fusion ; and some spadefuls of lime being thrown in, the scoriæ, which are thus rendered solid on the surface of the melted metal, are raked to the side of the furnace, while the Lead is run out into moulds through an aperture near the bottom ; and in this state it is called *Pig-lead*. It frequently contains silver, which is separated by oxidizing the lead into Litharge, and freeing the silver from what remains by cupellation.

*Qualities*. — Pure metallic Lead is of a light blueish colour, and, immediately after being melted or cut, has a very considerable degree of lustre, which it quickly loses on exposure to the air. It is nearly insipid ; and emits, when rubbed, a peculiar unpleasant odour. It stains the fingers and paper of a blueish colour, and has a specific gravity of 11.352, which is somewhat diminished after it is well hammered.<sup>1</sup> It is the softest and least elastic of the solid metals ; and although its ductility be trifling, yet it is very malleable, and may be reduced into thin leaves and drawn into wire : it melts at a temperature of 594<sup>2</sup>, and at a greater heat is volatilized. Its susceptibility of oxidizement is very considerable, and is the cause of its diminished lustre when exposed to the air. Ac-

<sup>1</sup> Muschenbroeck.<sup>2</sup> Irvine, *Chemical Essays*, 55.



cording to Dr. Thomson's experiments, it is capable of uniting with four doses of oxygen, and forming four distinct oxides, 100 parts of each of which contain the following proportions of lead and oxygen.<sup>1</sup>

Yellow ( <i>protoxide</i> )	contains of lead	91·5.	—oxygen	8·5.
Yellow ( <i>duetoxide</i> )	. . . . .	90·5	. . . . .	9·5.
Red ( <i>tritoxide</i> )	. . . . .	88·	. . . . .	12·.
Brown ( <i>peroxide</i> )	. . . . .	80·	. . . . .	20·. <sup>2</sup>

*Medical properties and uses.*—Lead has no action on the animal system in its pure metallic state; but, when oxidized, or in combination with acids, it produces very deleterious effects. Hence, metallic Lead taken into the stomach may prove a poison, from its meeting with acids in the primæ viæ; and liquors which are apt to become in any degree acidulous, if kept in leaden cisterns, may, from the same cause, be productive of much danger to those who drink them. I know an instance of the officers of an East Indiaman having been nearly poisoned from drinking water which was kept in a leaden cistern, and which, by the constant agitation of it, from the rolling of the ship, had oxidized the Lead: and there have been instances also of plumbers being poisoned by the volatilized particles of lead, which in great part form into a grey oxide. But the greater number of cases of poisoning from this metal, are produced by the preparation of it we are about to describe; and, therefore, the mode in which Lead acts on the animal system will be more properly noticed under it than in this place.

### 1. SUBCARBONATE OF LEAD.

*Officinal.* PLUM'BI SUBCARB'ONAS, *Lond.* CARBONAS PLUMBI, *vulgo, Cerussa, Edin.* CERUSSA. *Subacetas Plumbi, Dub.* Carbonate of lead. White oxide of lead. Ceruse.<sup>3</sup>

*Syn.* Plomb carbonaté (*F.*), Bleiweiss (*G.*), Loot wit (*Swed.*), Cerussa (*I.*), Blanco de Plomo (*S.*), Asfeedās (*Arab.*), Suffida (*H.*), Vūllay (*Tam.*).

This substance, which is known in commercial language by the name of *White-lead*, appears, from the analysis of it, by several of the most expert chemists, to be a Carbonate of Lead.<sup>4</sup> It is prepared in the large way in the following manner:—Sheets of Lead, about two feet long, five inches broad, and a quarter of an inch thick, cast in a mould, and not afterwards flattened, are rolled up into loose coils and placed in

<sup>1</sup> *Thomson's System of Chemistry*, 4th edit. i. 274—277.

<sup>2</sup> This oxide was first discovered by Proust. Mr. Murray thinks there is reason for believing it to be a subsalt. *System of Chemistry*, iii. 266.

<sup>3</sup> Σαρδύξ Dioscoridis.

<sup>4</sup> The various appellations given to this substance by the different colleges, arise from the indeterminate ideas which prevail of its composition,



earthen pots, which are wider at the mouth than at the bottom. Each pot is capable of holding six pints of fluid, but into it as much vinegar only is poured as will rise so high as not to wet the lead, which rests on a ledge half-way down. In some manufactories, however, the pots are made to contain about a pint of vinegar only; and the Lead, instead of being coiled up, is in form of a simple plate, which is laid over the mouth of the pot. The vinegar and Lead being arranged, the pots are buried in fresh stable litter, where they remain for about two months; during which time the vapours of the vinegar, elevated by the heat of the dung, oxidize the surface of the Lead, converting it into the yellow oxide, which combines with the carbonic acid gas evolved from the fermenting materials of the bed. The Carbonate (for it is not a *subcarbonate*) appears as a white, scaly, brittle matter, on the surface of the lead, and is separated “by spreading the coils upon a perforated wooden floor covered with water, and drawing them to and fro by rakes, which detaches the carbonate, and causes it to sink through the water and the holes of the floor to the bottom of a vessel placed below.”<sup>1</sup> In some places, this operation is performed by merely scraping off the Carbonate with a knife. It is afterwards ground in mills fitted for the purpose. It was formerly ground dry, and the workmen suffered very severely; but it is now ground in water, and the Carbonate afterwards dried in earthen pans placed in stoves, heated by means of flues. The beauty of the Carbonate depends on the purity of the Lead.

*Qualities.* — Carbonate of lead is inodorous, and nearly insipid; in the form of a heavy white powder, insoluble in water, but soluble in pure potass. When exposed upon charcoal to the action of the blowpipe, a button of metallic Lead is produced. Its constituents, according to Berzelius, are yellow oxide of lead 83·5 and carbonate acid 16·5 in 100 parts.<sup>2</sup> It is often adulterated with chalk, which may be discovered by pouring distilled vinegar on the suspected Carbonate, and then adding oxalic acid, or oxalate of ammonia to the solution. The formation of a precipitate proves the presence of chalk.

*Medical properties and uses.* — This preparation of Lead is a very powerful astringent. It is used externally only, being sprinkled on inflamed and excoriated parts. Some writers, however, regard even its external employment dangerous, but we have ordered it a thousand times, without witnessing any bad effects to follow. It enters into the composition of some ointments.

It is from this preparation that most of the cases of poison-

<sup>1</sup> *Aikin's Dictionary*.

<sup>2</sup> *Annales de Chim.* lxxvii. p. 85.



ing from the internal use of lead occur. This often happens to painters, and to those employed in grinding White Lead, from the want of cleanliness in not washing their hands before eating, by which some of the White Lead is introduced into the stomach with their food; and also from the criminal custom of putting White Lead, as well as Sugar of Lead, into acid wines for the purpose of sweetening them, and into hollands to deprive the spirit of the colour which it acquires when long kept in the wood. The symptoms which preparations of Lead produce are obstinate costiveness, pain in the stomach, and vomiting; the pulse becomes small and hard; the respiration laborious; and tremors ending in paralysis of the extremities, or death, ensue, when its operation is not counteracted by medicine. The exhibition of cathartics, particularly castor oil, combined with opium or henbane, plentiful dilutions with mucilaginous liquids, the warm bath, and injecting mutton broth per anum, are the best antidotes.

When the presence of any Salt of Lead is suspected in a dry substance, it may be discovered by reducing it to a metallic state with the blowpipe upon charcoal; and in a liquid, by dropping into it a watery solution of sulphuretted hydrogen gas<sup>1</sup>, when it is made obvious by a dark brown precipitate, which is insoluble in tartaric acid, the Salt of Lead being formed into an insoluble hydrosulphuret.

Officinal preparations. *Plumbi Superacetæ*, L. E. D. *Unguentum Cerussæ*, D.

## 2. SEMI-VITRIFIED OXIDE OF LEAD.

*Officinal*. PLUMBI OXYDUM SEMI-VITREUM, *Lond. Edin.* LITHARGYRUM, *Dub.* Litharge.

*Syn.* Litharge (*F.*), Bleiglätte (*G.*), Loot glans (*Dutch*), Silver-glitt (*Swed.*), Piombo semi-vitreo (*I.*), Almastago (*S.*), Marudar Singhie (*Tam.*), Moordar singh (*Pers. Duk. Hind.*).

This Oxide is prepared by the simple action of heat and air upon Lead. It is generally obtained during the calcination of Lead, when separating the silver with which this metal is often combined. The Lead is placed in a wind furnace, on a large cupel, or hollow dish made of ashes, and kept at a red heat with the blast of a large pair of bellows directed upon its surface; a scaly, yellowish white, glistening oxide is soon produced and successively formed by raking it off and exposing new surfaces till the whole of the Lead is thus converted into litharge. The varying of the circumstances of the process varies the colour of the oxide; some kinds of it from having

---

<sup>1</sup> To prepare this solution, put into a phial a paste made of iron filings and sulphur; then after some time add to it a small portion of sulphuric acid, and receive the gas which is produced, through a bent tube connected with the phial, into a flask filled with distilled water, and inverted in a basin or pneumatic trough full of water.



a silvery gloss are denominated Litharge of silver; and others, from the colour being a reddish yellow, litharge of gold.

*Qualities.* — Litharge is inodorous and insipid: it is in flakes with a vitreous lustre, dissolves in many of the acids; and, according to the experiments of Dr. Thomson, 100 parts of it contain 86·9 of lead and 9·1 of oxygen, forming 96 of yellow oxide and 4 of carbonic acid; which last, however, does not appear to be so essential as to constitute litharge a sub-carbonate of lead. For an account of the action of this oxide on fixed oils, see *Plasters*.

*Medical properties and uses.* — Litharge, like the other preparations of lead, is a powerful astringent. The ancients were acquainted with it. It is never given internally; and is used only for pharmaceutical purposes. Litharge is sometimes added to wines which are sour. It may be detected, either by passing sulphuretted hydrogen gas through the suspected wine; or evaporating this to the consistence of syrup, and then reducing the Lead with charcoal, in a crucible. In all cases of poisoning by Salts or Oxides of Lead the best antidote is a solution of magnesiæ sulphas, with the addition of some sulphuric acid; which uniting with the deleterious Salt of Lead forms an inert sulphate of lead, which is carried out of the bowels by the sulphate of magnesia.

Officinal preparation. *Emplastrum Plumbi*, L. E. D.

### 3. RED OXIDE OF LEAD.<sup>1</sup>

*Officinal.* OXYDUM PLUMBI RUBRUM, *Edin.* Red oxide of Lead.

*Syn.* Minium (*F.*), Mönnige; Mennig (*G.*), Minio (*I.*), Vermillon (*S.*), Isrenj (*Arab.*), Sindur (*H.*), Segāpoo Sindoorum (*Tam.*), Sindura (*San.*), Temamera (*Malay*).

This preparation is Lead in the highest state of oxidizement. It is prepared in a reverberatory furnace, vaulted like a baker's oven, and having two internal walls rising from the floor of the surface, but not reaching to the roof. The coals are placed between these internal walls and the wall of the furnace, by which means the flame is drawn over the top, and reflected from the roof down upon the surface of a quantity of Lead placed on the floor. The metal soon melts, and is altogether converted into a yellow oxide, or *massicot*, by successively raking off the pellicles which form on its surface; this is then ground in a mill, and washed, to separate any metallic Lead, by which it becomes of an uniform yellow colour, and, after being replaced in the furnace, is exposed to the flame while it is constantly stirred for about forty-eight hours, when it is converted into Red Oxide of Lead.<sup>2</sup> By this process, 20 cwts. of Lead produce on an average 22 cwts. of Red Lead, notwith-

<sup>1</sup> Ψιμυθιον Dioscoridis.

<sup>2</sup> Watson's *Chemical Essays*, iii. 338. *Aikin's Dictionary*.



withstanding a portion is necessarily volatilized. To save the previous calcination, litharge is sometimes employed.

*Qualities.* — Red Oxide of Lead is inodorous and insipid; in the form of a very heavy, scaly powder, its specific gravity being  $8.940^1$ , and of an intense red or scarlet colour, verging into orange. When heated to redness it gives out oxygen gas, and runs into a dark-brown, hard glass. According to Dr. Thomson's experiments, 100 parts of it contain 88 of lead, and 12 of oxygen.

*Medical properties and uses.* — Red Lead may be applied to the same uses as litharge, but is now rarely or never used. Its chief use is in the arts, as a pigment.

POLYGALA. *Spec. Plant. Willd.* iii. 871.

Cl. 17. Ord. 3. Diadelphia Octandria. *Nat. ord.* Lomentaceæ, *Linn.* Pediculares, *Juss.*

G. 1313. *Calyx* five-leaved, with two of the leaflets wing-shaped, and coloured. *Legume* obcordate, two-celled.

\*\*\* *Beardless; herbaceous, with a simple stem.*

*Species* 67. *P. Senega*. Seneka root. *Med. Bot.* 2d edit. 452. t. 162. *Amæn. Acad.* iii. 124. *Pursh.* iii. p. 464.

*Officinal.* SENEGÆ RADIX, *Lond.* POLYGALÆ SENEGÆ RADIX, *Edin.* SENEKA; RADIX, *Dub.* Seneka root.

*Syn.* Polygale de Virginie (*F.*), Senegawurzel (*G.*), Poligala Virginiana (*L.*).

This plant is a perennial native of North America, flowering in June.<sup>2</sup> The root is woody, branched, contorted, about half an inch thick, and covered with ash-coloured bark: it sends up several stems a foot in height, erect, slender, round, smooth, and of a dark-reddish colour. The leaves are petio- late, alternate, lanceolate, acute, and pale green: the flowers are in loose, terminal spikes, small, white and papilionaceous, with the calyx divided into three narrow, persistent segments: the fruit is an inversely cordate capsule, containing several small seeds.

The root is brought from Virginia in bales, each containing from two to four hundred weight.

*Qualities.* — Seneka root is inodorous: the taste is at first sweetish and nauseous, but after being chewed for less than a minute becomes pungent and hot, producing a very peculiar tingling sensation in the fauces. These qualities reside in the bark; which, on the dried root, is white within and covered with a brownish-grey, corrugated, transversely cracked cuticle: the central part is white, but woody and inert: alcohol extracts the whole of its active matter, which is precipitated from the tincture by the addition of water; and the ethereal tincture deposits a pellicle of resin, but no extractive. From six ounces

<sup>1</sup> Muschenbroeck.

<sup>2</sup> It was first cultivated in England by Mr. F. Miller, in 1759.



of the root, Peschier separated 100 grains of a peculiar alkaline principle, which he has named *Polygalina*, united with a new acid, which he has denominated the *Polygalinie*; and this salt he supposes is the active principle of seneka root.<sup>1</sup> Hot water extracts its virtues partially only; but in a sufficient degree to exert its influence on the animal system.

*Medical properties and uses.* — This root is a stimulating expectorant and diuretic; and in large doses emetic and cathartic: it increases absorption, and consequently augments the natural excretions, particularly that of urine; and frequently occasions a copious ptyalism. It was introduced to the notice of physicians by Dr. Tennant, who, having discovered that it was the antidote employed by the Senagaro Indians against the bite of the rattle-snake, and reasoning from the effects of the poison, and of the remedy in removing these, was induced to try it in pneumonic affections, and found it useful. On account of its stimulant properties, however, it can be employed in these complaints, only after the resolution of the inflammation by bleeding and evacuations. It proves more directly useful in humoral asthma, chronic catarrh and some kinds of dropsy. The extract of it combined with carbonate of ammonia has been found by Dr. Brandreth, of Liverpool, to be efficacious in some cases of lethargy; and in America the decoction given in divided doses, at short intervals, till it vomit or purge, has been employed with seeming success in croup<sup>2</sup>: it has also been used as a stimulant gargle, in the same disease.

It may be administered either in the form of powder or decoction, combined with aromatics, opium, or camphor, which check its nauseating qualities. Madeira wine, where it can be ordered, may be used to cover the taste of the powder. The dose in substance is from grs. x. to ʒj., repeated every three or four hours.

Officinal preparation. *Decoctum Senegæ*, L. E.

POLYGONUM. *Spec. Plant. Willd.* ii. 440.

Cl. 8. Ord. 3. Octandria Trigynia. *Nat. ord.* Holoraceæ, Linn. Polygoneæ, Juss.

G. 3. *Corolla* five-parted, calycine. *Seed* one, angular.

\* \* *Bistorts*, with a single spike.

*Species* 3. *P. Bistorta*.<sup>3</sup> Great Bistort or Snakeweed. *Med. Bot.* 2d edit. 668. t. 232. *Smith Flora Brit.* 427. *Eng. Bot.* t. 509.

*Officinal.* BISTORTA, *Lond.* POLYGANI BISTORTI RADIX, *Edin.* BISTORTA; RADIX, *Dub.* Bistort root.

<sup>1</sup> *Pharmacopœia Batava*, &c. editone De Joanne Frid: Neimann. Lipsiæ, 1824.

<sup>2</sup> *London Medical Review and Magazine*, iii. 426.

<sup>3</sup> *Bistorta*, quasi bis torta, twice twisted, *Alston, Mat. Med.* vol. i. 399.



*Syn.* Bistorte (*F.*), Natter-Wurzel (*G.*), Natter-Wortel (*Dutch*), Bistortat (*I. S. Port.*), Slangeort (*Dan.*), Sertecinaja trawa (*Russ.*).

This plant grows in many parts of Europe, Siberia, and Japan, and is indigenous to Great Britain: found generally in moist meadows, flowering in May and June.<sup>1</sup> The root is perennial, woody, and tortuous: the stem rises nearly two feet in height, is foliaceous, jointed, swelling at the joints, solid, smooth, and bending a little near the top: the leaves are ovate, those next the root cordato-lanceolate; the whole are entire, waved at the edge, veined, of a fine green colour on the upper surface, and glaucous below; the radical ones are on long winged footstalks, those of the stem almost sessile, amplexicaule, and sheathing. The flowers are small, of a pale rose colour, collected into a close, oblong, terminal spike an inch and a half long; the single flowers standing on short, white flower-stalks, which rise in pairs from membranous, withering, floral leaves. The corolla is divided into five obtuse segments, with nectareous glands at the base: the filaments are longer than the corolla, tapering, and supporting purple anthers; and the germen is triangular, of a red colour, crowned with three long styles, with small round stigmas. The seeds are three-sided, of a dark-brown colour, and shining.

*Qualities.* — The dried root is inodorous, and has a very austere taste. Water extracts its virtues. The decoction strikes a deep black with oxysulphate of iron, and precipitates gelatin. It contains a large proportion of tannin, some gallic acid, and much starch.

*Medical properties and uses.* — The root of Bistort is astringent and tonic. It is employed in hæmorrhagies, obstinate fluxes, leucorrhœa and all diseases in which simple astringents are indicated. It has also been given in intermittents, combined with gentian, or a coruscalamus. Externally, a strong decoction of it is a useful lotion for spongy gums and ill conditioned ulcers. But it is almost discarded from modern practice.<sup>2</sup>

The dose of the powdered root is from grs. xv. to ʒj., twice or thrice a day.

PORRI RADIX. Vide *Allium Porrum*.

POTASSÆ NITRAS, *Lond.* NITRAS POTASSÆ, *Edin.* NITRUM, *Dub.* Nitrate of Potass. Nitre.

<sup>1</sup> It is, however, not confined to low situations, being found on the Carpathian Alps, vegetating under *Pinus magnus*, at an elevation of 4476 feet. Vide *Wahlenberg's Flora Carpatorum*.

<sup>2</sup> In Iceland the recent root of Bistort is eaten raw, or converted into bread. It may, therefore, be reasonably enquired, what effect can it have as a medicine, when prescribed in the small doses usually ordered?



*Syn.* Nitrate de Potasse (*F.*), Salpetersaures kali (*G.*), Salpeter (*Dutch, Dan. Swed.*), Saletra (*Polish*), Nitro (*I. S. Port.*), Salitre (*S.*), Selitra (*Russ.*), Pottle Ooppoo (*Tam.*), Shora (*Pers.*), Sandaiva (*Malay*), Ubkir (*Arab.*), Wedie looneo (*Cyng.*), Shora (*H.*), Yavac Shora (*San.*).

This salt is well known in commerce under the name of Saltpetre or Nitre. It may be regarded both as a natural and artificial production, being found effloresced on the surface of the soil in Hungary<sup>1</sup> and some parts of Europe<sup>2</sup>, South America, Africa<sup>3</sup>, and very abundantly in India<sup>4</sup>, whence this country is chiefly supplied; while in some countries, as in Germany and France, it is artificially produced. Nitre is prepared by art by the same means as nature employs, the artificial composts being imitations only of the natural soils where it is most abundantly formed; by giving, therefore, an account of the former mode, both will be better understood. Glauber first suggested the formation of what are termed nitre beds. In France they consist of a compost of putrefying animal and vegetable matters, such as blood, offal, excrementitious matters, and decaying leaves, with street-sweepings, old mortar, chalk, and other calcareous matter; which are mixed in casual proportions, and lightly spread in long beds, covered with roofs to protect them from the weather. These are turned up occasionally, frequently moistened with putrid water, or urine; and at the end of two years, or less, are supposed to be fit to yield the nitre by lixiviation. The theory of this process, which is not yet completely elucidated, was not at all understood till the experiments of Thouvenel and the discovery of the composition of nitric acid by Mr. Cavendish removed much of the obscurity in which it was involved: the following is the explanation. The spontaneous decomposition of the animal and vegetable matter evolves azote, oxygen, hydrogen, and carbon, which reuniting by the operation of new affinities, new compounds are formed, and among these nitric acid by the union of the azote or nitrogen from the animal substances with the oxygen from the vegetable matter. The acid thus formed is attracted partly by the calcareous earth of the beds, and partly by a portion of potass, either contained in them ready formed, or, as some have supposed, formed during the process. The presence of

<sup>1</sup> *Beudant's Travels in Hungary.*

<sup>2</sup> A great repository of native Nitre in Europe, is the Pulo of Molfelta, in the province of Puglia, in the kingdom of Naples. It is a deep cavity, formed by the falling in of several caverns. The Abbé Fortis first drew public attention to this place, at which time it was lined with a crust of Nitre an inch thick, which on being scraped off was successively renewed in a few days.

<sup>3</sup> Near the city of Tlemsan, in the kingdom of Algiers, six ounces of Nitre is extracted by simple lixiviation, from one quintal of the common mould.—*Shaw's Travels*, 228.

<sup>4</sup> The presidency of Calcutta exports annually upwards of 8000 tons of Nitre.



animal matter, although it aid the formation of Nitre, yet, is not essential; for Dr. J. Davy found a rich impregnation of Nitre in a Nitre cave near Mensoora, in the district of Doombera, in Ceylon, in a decomposing rock consisting of calcspar, felspar, quartz, mica, and talc, in a humid state, exposed to the air, and perfectly free from any animal matter.<sup>1</sup> The presence of a certain degree of heat and humidity, of atmospheric air, of lime and some alkaline mineral, is absolutely necessary; for, besides fixing the nitric acid when formed, the affinities lime exerts to oxygen and azote favour very much their combination, and consequently the formation of the acid.

The compost, when ready to be lixiviated, is first mixed with wood ashes, or with pulverized, impure potass, to decompose the nitrate of lime. It is, then, put into a cask furnished with a cock at the bottom, and an inner, false, perforated bottom: a quantity of river water is now poured over it, and after some hours the cock is turned, and the liquor drained off, which is used instead of water for a second portion of earth; and this is successively repeated, till it is supposed to be sufficiently impregnated with the soluble matter of the compost. The lixivium, which contains chiefly Nitrate of Potass, and the muriates of potass and of soda, is now boiled and clarified with bullocks' blood or a solution of glue; and the boiling continued, the muriates as they form being withdrawn by perforated ladles, till the liquor is so concentrated, that a few drops poured on cold iron immediately crystallize. It is then, when nearly cold, poured into separate crystallizing dishes, in which after some days the salt is found deposited in a confused mass of opaque, dirty white, imperfect crystals, which, after being broken to pieces and drained, are known under the name of rough or crude nitre.

Nitre is brought from Bengal in an impure state, but crystallized, put up in bags, each containing two Bazar mounds, or 164lbs. weight.<sup>2</sup> The crystallized state of this impure Nitre arises from the lixivium of the soil having been slowly evaporated in shaded, shallow pits. To purify crude Nitre, it is repeatedly washed with cold water, which dissolves the deliquescent muriates: and then is boiled with half its weight of water, until a pellicle forms on the surface; after which the solution is poured into leaden coolers, and stirred till it is quite cold, by which means the salt is deposited in acicular crystals.<sup>3</sup>

<sup>1</sup> *Davy's Account of the interior of Ceylon*, 4to. p. 32. Lond. 1821.

<sup>2</sup> Each Bengal ship of 800 tons generally brings home in a period of war about 5000 bags of nitre.

<sup>3</sup> Nitre was unknown to the ancients; and Beckman thinks that it was not dis-



*Qualities.* — Pure Nitrate of Potass is inodorous; and has a bitterish, sharp taste, occasioning a sensation of cold both in the mouth and stomach. It is generally in white, pellucid, brittle, hexahedral prisms, terminated by two-sided summits, the specific gravity of which is 1.933. These crystals are soluble in seven parts of water at 60°, producing cold during their solution; but boiling water takes up its own weight of them. They are perfectly insoluble in strong alcohol. They are permanent in the air; melt when exposed to a moderate heat, and when cast into moulds, form *Sal Prunelle*. In a strong heat oxygen gas is disengaged at first, and afterwards azotic gas; and, in a continued intense heat, the acid is completely expelled and decomposed, leaving behind pure potass. Nitre when mixed with inflammable substances detonates in a strong heat; and if charcoal be used, a pure subcarbonate of potass remains behind. It is likewise decomposed by the sulphuric acid, alum, sulphate of magnesia, and the sulphates of zinc, copper, and iron, when aided by heat, and in the cold by barytes. According to the analysis of Berard, 100 parts of Nitrate of Potass contain 51.36 nitric acid, 48.64 potass.<sup>1</sup>

Nitrate of Potass sometimes contains muriate and sulphate of soda, and sulphate of potass. The muriate is discovered by nitrate of silver, throwing down a precipitate, every 100 grains of which denotes  $42\frac{1}{2}$  of muriate of soda. The sulphates are detected by nitrate of barytes.<sup>2</sup>

*Medical properties and uses.* — Nitrate of Potass is refrigerant and diuretic; and, when externally applied in solution, cooling and detergent. If taken in repeated small doses, it abates heat and thirst in diseases of increased excitement, diminishes the force and frequency of the arterial action, and increases the secretion of urine, in which, and in the alvine dejections, the salt may be detected by chemical tests. It is efficaciously given in all inflammatory cases, active hæmorrhages, and in herpetic eruptions. Although diuretic, yet it is of little use in dropsies, and is contra-indicated in typhus and hectic fever: in the latter of which, as Dr. Percival has justly observed, it lowers the pulse at first, but afterwards raises it higher than before. A small portion of it allowed to dissolve slowly in the mouth, often removes incipient inflammatory sore throat; and thence its utility in gargles in that complaint.

---

covered till the 13th century. The term *sal petrôsum* is first mentioned in the work of Albertus Magnus "*de Mirabilibus mundi*," in a prescription of Marcus Græcus, for making the Greek fire: but it is probable, as Beckman conjectures, that this salt was known long before this period in India, where he believes gunpowder also was invented, and brought by the Saracens from Africa to Europe. *History of Inventions*, vol. iv.

<sup>1</sup> *Ann. de Chim.* t. lxxi. p. 69.

<sup>2</sup> *Henry's Elements of Experimental Chemistry*, 7th ed. vol. ii. p. 464.



It is most advantageously given dissolved in mucilaginous fluids, as almond emulsion, in moderate doses not exceeding grs. xv. frequently repeated. In large doses it excites nausea, a lively sensation of cold at the epigastric region, and colic: and in very large doses, from ʒiv. to ʒj. for instance, which have sometimes been given by mistake for sulphate of soda, it occasions vomiting, hypercatharsis, bloody stools, convulsions, and sometimes death. Opium and aromatics are the best antidotes.

Officinal preparations. *Acidum nitricum*, L. E. D. *Trochisci Nitratis Potassæ*, E.

POTASSÆ SUPERTARTRAS, *Lond.* Vide *Supertartras Potassæ*.

POTA'SSA IMPU'RA. *Carbonas Potassæ impura*, *Lond.* SUB-CARBONAS POTASSÆ IMPURUS, *Edin.* CINERES CLAVELLATI. *Kali impurum*<sup>1</sup>, *Dub.* Impure Potass. Impure Sub-Carbonate of Potass. Potashes. Pearlashes.

*Syn.* Carbonate alalinule de Potasse (*F.*), Koloensuares Kali; Pottasche (*G.*), Potasch (*Dutch*), Potaske (*Dan.*), Potaska (*Swed.*), Potassa del Commercio (*I.*), Cenizas claveladas (*S.*), Mara Ooppoo (*Tam.*), Hindu loonoo (*Cyng.*), Char Lawanum (*San.*), Jarka nemack (*Duk.*).

This substance consists chiefly of Subcarbonate of Potass, mixed with some other salts. It is known in commerce by the name of Potash; and is brought to us principally from the Baltic and America. The manipulation of the process by which it is prepared differs in different countries; but the general features of it are every where the same. The dried stems and branches of plants are set fire to, and reduced to ashes; which are lixiviated by pouring over them, in proper vessels, hot or cold water, so as to dissolve the alkaline matter they contain. The impregnated solution, drawn off from the ashes, is then boiled to dryness in iron boilers, and leaves behind a solid saline mass, coloured brown by a small portion of vegetable inflammable matter, and which generally becomes moist. This is the *Potash* of commerce. After the colouring matter is destroyed, and a portion of the water dissipated by calcination in a reverberatory furnace, it assumes a spongy texture, with a blueish or greenish colour, and is then denominated *Pearl-ashes*.

Those vegetables only which grow at a distance from salt water, are employed to obtain this product. Herbaceous plants yield the largest proportion, and shrubs more than trees. Kirwan remarks, that although Fumitory produces

<sup>1</sup> Of these three appellations that of the Dublin College is the least exceptionable, because it does not convey an erroneous idea of the nature of the substance, which is the case with the others. As synonyma, those of the London College are at complete variance.



more of this salt than any other plant, and, next to it Wormwood, yet, that 1000 parts of the ashes of Wormwood yield more Potass than the same quantity of the ashes of Fumitory, in the proportions of 748 and 360. It has been said, that it was lately obtained in great abundance from the herbaceous part of the Potatoe, cut down just as the fruit is beginning to form : 40,000 lbs. of the dried stems, it has been stated, will yield 2200 lbs. of Impure Potass ; but the trials in this country have not confirmed these statements.<sup>1</sup> It is generally supposed, that at least the greater part of the Potass is contained ready formed in the vegetables ; but this is somewhat doubtful, and perhaps in living plants the base only of Potass exists as an element, and is oxidized so as to form the alkali during the combustion. Such is the conjecture of Dr. Murray<sup>2</sup>; and the same may take place during the spontaneous decomposition of plants where much water is present, for Potass can be obtained by the evaporation of dunghill water.<sup>3</sup>

The Pearl-ash of commerce is still a very compound mass, containing, besides the subcarbonate of potass, sand with which it is often adulterated to a great extent, sulphate of potass, muriate of potass, oxide of iron, and oxide of manganese ; to the last of which, according to Scheele, it owes its blueish or greenish colour. Different parcels of Pearl-ash must undoubtedly contain different quantities of Potass ; and thence no accurate standard of the proportion of the ingredients can be fixed. The following table drawn up by Vauquelin shows the comparative value of samples from different countries, examined by him. The quantity of each was 1152 parts.<sup>4</sup>

Kinds of Potass.	Real Potass.	Sulphate of Potass.	Muriate of Potass.	Insoluble Residue.	Carbonic Acid and Water.
Russian potass	772	65	5	56	254 = 1152
American do.	857	154	20	2	119 = 1152
Pearl-ash	754	80	4	6	308 = 1152
Potass of Treves	720	165	44	24	199 = 1152
Dantzic potass	603	152	14	79	304 = 1152
Potass of Vosges	444	148	510	34	304 = 1152

<sup>1</sup> *Phil. Mag.* vol. i. p. 340.

<sup>2</sup> *Murray's Chemistry*, 2d ed. ii. 193.

<sup>3</sup> See Birch's Experiments, *Phil. Trans.* for 1780, 345.

<sup>4</sup> *Annales de Chimie*, xl. 284.



The proportion of real alkali in any quantity of Pearl-ash may be ascertained in the following manner. Pulverize 500 grains of the Pearl-ash, and digest in successive portions of hot water as long as any thing is dissolved; which is known by adding a drop of the solution of oxymuriate of mercury, to a little of the lixivium. Mix the solutions, and drop in some diluted sulphuric acid (previously prepared by mixing one part of concentrated acid with thrice its bulk of water) from a phial containing a known quantity of it, till litmus paper indicate the slightest possible excess of acid. Next heat this mixture to expel the carbonic acid; and on trying it again with the litmus paper, if it show any excess of alkali, add a few drops more of acid. Ascertain now by weighing the phial of acid how much acid has been expended in saturating the alkali, and for every hundred parts of real acid set down 121.2 of Potass.<sup>1</sup>

The Pearl-ash of commerce is not sufficiently pure for medicinal use: and therefore it is used only for pharmaceutical purposes.

Officinal preparation. *Potassæ Subcarbonas*, L. E. D.

POTASSÆ SULPHAS, *Lond.* Sulphate of Potass.

This salt, which had formerly a place among the preparations only of the British Pharmacopœias, is now placed in the list of materia medica of the London College; the greater part of the salt found in the shops being made on a large scale. It is prepared from the residue of the distillation of nitrous acid from nitre and sulphate of iron. The residue of this process is a mixture of Sulphate of Potass and red oxide of iron, from which the Sulphate is separated by boiling water.

*Qualities: Medical properties and uses.* — See Part iii. page 606, 7.

PRUNUS. *Spec. Plant. Willd.* ii. 984.

Cl. 12. Ord. 1. Icosandria Monogynia. Nat. ord. Pomaceæ, Linn. Rosaceæ, Juss.

G. 982. Calyx five-parted, inferior. Petals five. Nut of the drupe with prominent sutures.

Species 29. *P. domestica*.<sup>2</sup> Common Plum-tree. *Med. Bot.* 2d edit. 520. t. 187.

Officinal. PRUNA, *Lond.* PRUNI DOMESTICÆ FRUCTUS, *Edin.*

PRUNUS GALLICA; FRUCTUS, *Dub.* Prunes.

*Syn.* Prunes (*F.*), Pflaumen (*G.*), Priumen (*Dutch*), Pruno domestico, o Prugna (*I.*), Ciruelas pasas (*S.*), Sliwnik (*Russ.*), Erik (*Turk.*).

<sup>1</sup> The value of the diluted acid must be previously ascertained, by adding to 100 grains of it muriate of barytes as long as any precipitate falls. This forms sulphate of barytes, which, when washed and dried at a low red heat, contains 33.3 per cent. of sulphuric acid; by which the proportion of real acid in the diluted acid may be known. *Aikin's Dictionary*, i. 263.

<sup>2</sup> ΚΥΚΚΥΜΗΛΕΑ Dioscoridis, *Barkuk* (Arab.), Muei Xu (*Chin.*).



The tree which yields this fruit is a native of Asia and Greece, although it is now completely naturalized to our climate, growing wild in coppices, and flowering in April and May. It rises about fifteen feet in height, with a moderately spreading head. The leaves are pale green, standing on short petioles, which have two glands near the base of the leaf: they are serrated, smooth, and when young convoluted and pubescent underneath: the flowers are large, on short solitary peduncles, with an erect calyx, and obovate white petals: the fruit is a superior, berried, oval drupe, swelling a little more on one side, and three-grooved; of a bluish violet colour on the outside, internally consisting of a yellow, fleshy, sweet pulp; and containing a smooth, almond-shaped nut.<sup>1</sup>

The dried fruit, which only is officinal, is imported from the continent in chests; and that which is brought from France is regarded as the best. The recent fruit, when perfectly ripe, is pleasant to the palate, and sufficiently wholesome; but when eaten too freely it occasions flatulence, griping, and diarrhœa, more readily than any other fruit.

*Qualities.*—Prunes are nearly inodorous, but have an agreeable, sweet, subacid taste. They contain chiefly mucus, saccharine matter, and malic acid.

*Medical properties and uses.*—Dried Plums or Prunes are gently laxative, and form a pleasant addition to purgative electuaries and decoctions. Simply boiled, they may be beneficially given to children who are habitually costive; and in fevers.

Officinal preparation. *Confectio Sennæ*, L. E. D.

PTEROCARPUS.<sup>2</sup> *Spec. Plant. Willd.* iii. 904.

*Cl.* 17. *Ord.* 4. Diadelphia Decandria. *Nat. ord.* Papilionaceæ. *G.* 1318. *Calyx* five-toothed. *Legume* falcated, leafy, varicose, surrounded with a wing, not gaping. *Seeds* solitary.

*Sp.* 6. *P. santalinus*. Red Saunders tree. *Med. Bot.* 2d edit. 430. *t.* 156. *Willdenow Spec. Plant.* iii. 906.

*Sp. nova*, *P. erinacea*. *Encycl. Method. Lam. Illust. Gen.* tab. 602. fig. 4.

#### 1. PTEROCARPUS SANTALINUS.

*Officinal.* PTEROCARPI LIGNUM, *Lond.* PTEROCARPI SANTALINI LIGNUM, *Edin.* SANTALUM RUBRUM. LIGNUM, *Dub.* Red Saunders wood.

*Syn.* Santale rouge (*F.*), Rothes Sandalholz (*G.*), Root Zandelhout (*Dutch*), Sandalo Rosso (*I.*), Sandolo rubio (*S.*), Ract Chandan (*H.*), Segapoo Shandandum (*Tam.*) Sundul Ashmer (*Arab.*), Sundul Soorkh (*Pers.*), Ruct Handoon (*Cyng.*), Ruckta Chunduna (*Beng.*), Racta Chandana (*San.*), Hoam pe mo (*Chin.*).

<sup>1</sup> Martyn, in his edition of Miller's Gardener's Dictionary, enumerates sixty varieties of the plum. The French prunes are the same as those which were formerly brought from Damascus.

<sup>2</sup> From πτερον, a wing, and καρπος, fruit.



This tree is a native of the mountains of India, particularly the rocky parts in the Onore district<sup>1</sup>, and of Ceylon. It is a lofty tree, with alternate branches, and a bark resembling that of the common alder. The leaves are petiolate and ternate, each simple leaf being ovate, blunt, entire, retuse, veined, smooth on the upper surface, and hoary beneath; the flowers are in axillary spikes, without bractes: the calyx is brown: the corolla papilionaceous, consisting of an erect, obcordate *vexillum*, turned back at the edges, denticulate, curled, and waved, and of a yellow colour, with red veins: yellow, spreading, denticulate *wings* waved at the edges; and an oblong *keel* a little inflated and curled at the tip: the filaments are yellow, and support globular white anthers; the germen is oblong, compressed, hirsute, with a curved style, and an obtuse stigma; the pod is pedicelled, compressed, smooth, keeled along the lower edge, and contains one round, compressed seed.

This tree, which yields the true officinal Red Saunders, was first detected by Kœnig in India. The wood is brought home in billets, which are very heavy and sink in water.

*Qualities*.—Red Saunders Wood has an aromatic odour, and is nearly insipid. It is extremely hard, of a fine grain, and a bright, garnet-red colour, which deepens on exposure to the air. It yields its colouring matter, which appears to be of a resinous nature, to ether and alcohol, but not to water.<sup>2</sup> The alcoholic tincture is red, but becomes yellow when largely diluted with water. Volatile oil of lavender also extracts its colouring matter; yet, it is scarcely affected by oil of turpentine, which acquires a pale yellow tinge only, even when assisted by heat. Neumann first noticed this fact<sup>3</sup>; and it has been suggested that the camphor contained in the oil of lavender may give it the above property; but camphoreted oil of turpentine has no more effect than the simple oil. I find that by shaking oil of turpentine, which has been digested over Red Saunders with a little alcohol, the slight tinge of colour it received is instantly taken up by the spirit, and the oil settles as a colourless substratum.

Red saunders has no medicinal properties, and is used only as a colouring matter.

\* PTEROCARPUS ERINACEA.

*Officinal*. KINO, Lond.

*Syn*. Gomme de Kino (F.), Kinoharz (G.), Chino (I.), Toomble hœan (Tam.).

<sup>1</sup> When transplanted to low situations and a richer soil, the tree degenerates; and is in all respects less esteemed. *Forbes's Oriental Mem.* 4to. vol. i. p. 808.

<sup>2</sup> Yet Willdenow, who received the description of the tree and its wood from Kœnig, says, "attritu humido pulchre rubrum tingens." The yielding no colouring matter to water affords an easy mode of distinguishing Red Saunders from Brazil wood, which was first pointed out by Dr. Lewis. *Thomson's Chem.* v. 208.

<sup>3</sup> *Neumann's Chem.* 337.



Although the Edinburgh college has inserted Kino as the inspissated juice of the *Eucalyptus resinifera* in the list of materia medica of its Pharmacopœia, and the Dublin College has considered it as the product of the *Butea frondosa*; yet we believe that the plant which yields the best kino is an African tree; and from a specimen sent home by Mungo Park in his last journey, which is in the possession of Sir Joseph Banks, it is a *Pterocarpus*, and that species which is described under the specific name *Erinacea*, in the Encyclopédie Méthodique. It is a native of Senegal; and is distinguished from the other species of the genus by long yellow spines on the fruit. The leaves are pinnated; composed of obtuse, oval leaflets, larger at the base, petiolate, entire, thin and smooth above; pubescent, and of a reddish hue below, where they are marked with fine parallel, alternate, oblique ribs or nerves, a little arched. The flowers are on peduncles. The calyx is bell-shaped, truncated, slightly toothed and pubescent. The fruit is a compressed, orbicular, pubescent pod, bulging on both sides in the middle, where it is covered with white hairs and long, numerous, yellow skins. It contains one seed only. It is nevertheless true that Kino, such as was brought from Botany Bay about twenty years ago, is the production of the above species of *Eucalyptus*, the Brown-Gum tree of that country<sup>1</sup>; but it differs in several of its qualities from the Kino described by Dr. Fothergill, who introduced this remedy into practice.<sup>2</sup> We are informed, that none of it has been brought to this country since the above period. Another sort is said to come from Jamaica, and is stated by Dr. Duncan, junior, to be the extract of the *Cocoloba uvifera*, or sea-side grape<sup>3</sup>; while Dr. Murray says, “he has been informed that it is the extract of the wood of mahogany.”<sup>4</sup> The Dublin College indicated the *Butea frondosa* on the authority of Dr. Roxburgh; but the red juice which this plant yields has been examined by Dr. Duncan, and found to differ very consi-

---

<sup>1</sup> This plant belongs to the first order of the twelfth class of the Linnean system. It is a lofty tree, exceeding an English oak in size; and bearing yellowish flowers in umbellated clusters. The calyx is hemispherical, perfectly entire in the margin, and afterwards becomes the capsule; on the top, just within the margin, stands a pointed calyptra, of the same colour as the calyx, and as long. This calyptra, which is the essential mark of the genus, is analogous to the corolla in other plants, but neither splits nor divides; on removing it a great number of red stamens appear, standing in a conical mass, very resinous, aromatic, and bearing small red anthers. In the centre is a simple style terminated by a blunt stigma, and rising from a transversely cut, trilocular germen. The quantity of juice obtained from incisions made into the wood of the trunk amounts, sometimes, to sixty gallons from one tree. See *White's Voyage*, 231.

<sup>2</sup> *Medical Obs. and Inquiries by a Society of Physicians in London*, i. 238. 243.

<sup>3</sup> *Edinburgh New Dispensatory*, 5th edit. 292.

<sup>4</sup> *System of Mat. Med. and Pharmacy*, ii. 304.



derably from Kino, although it may be used as a substitute for it. The Kino now found in the shops comes from India, and is the extract of the *Nauclea Gambir*. The branches and twigs are bruised and boiled in water. The decoction is then evaporated until it acquire the consistence of an extract which is Kino. It is imported in chests containing from one to two cwt., and on the inside of the lid of each chest is a paper, inscribed with the name of John Brown, the month and year of its exportation: and stating that it is the produce of Amboyna.

*Qualities.* — 1. *Kino* which was given to me as a specimen of true *African Kino*, is inodorous, and insipid when first taken into the mouth; but after some time it imparts a slight degree of roughness, with a scarcely perceptible sweetness, to the palate: feels gritty between the teeth when chewed, and does not colour the saliva. It is in very small, irregularly shaped, shining, deep ruby-brown coloured fragments, and intermixed with small twigs and minute bits of wood, which are white in the inside. It is pulverulent, affording a dark chocolate or reddish-brown powder. Water at 60° dissolves the larger moiety of it, and gives a brick-red, rather turbid infusion, which does not become clear after standing twenty-four hours. Alcohol dissolves nearly two-thirds of it, the tincture having a very deep brown colour: what remains undissolved is nearly colourless. Ether takes up about one-third; and the tincture, which is of a beautiful claret colour, when evaporated on the surface of water, leaves a pellicle of brittle brown resin; while a sweetish, red-coloured extractive matter remains dissolved in the water.

2. *Botany Bay Kino* is inodorous; tastes bitterish, and more austere than the African; it is in larger fragments, equally brittle, breaking with a glassy fracture; of a chocolate hue, and affording a brown-coloured powder, but it is not uniform in appearance, some of the fragments being of a lighter hue. Water at 60° dissolves nearly the same quantity as of the former variety, and the infusion is brown and transparent. Alcohol dissolves rather more than two-thirds of its weight, but the tincture is not so deep-coloured as that of the former variety. Ether takes up one-twentieth; a pale brownish-straw colour only is imparted to it; and when evaporated on water, the resinous pellicle is scarcely perceptible, and very little extractive is deposited.

3. The *Kino* said to have been brought from Jamaica, but of which none is now to be procured, is in bitterness and roughness nearly equal to the last variety, but these qualities are accompanied with a slight degree of acidity. It is in brittle fragments, of an almost black colour, having a shining,



resinous fracture, in which appear small air-bubbles. The powder is of a reddish-brown colour. With alcohol and ether it affords results very similar to those of the first variety. Water dissolves a greater portion of it than of the other two kinds, and forms an infusion intermediate in colour and transparency; approaching in colour to the first, and in clearness to the second variety.

4. *East India* or *Amboyna Kino*, is inodorous, very rough, and slightly bitter when first taken into the mouth; but it afterwards impresses a degree of sweetness on the palate. It is in small, uniform, deep brown, shining, brittle fragments, which appear like portions of a dried extract broken down; being perfectly uniform in their appearance. It is easily pulverized, affording a powder of a lighter brown colour than the fragments. Water dissolves two-thirds of it, forming a deep brown clear solution; whilst the portion that remains undissolved is long suspended, if mixed with a fresh portion of water.<sup>1</sup> Alcohol dissolves the greater part of this variety, forming a deep claret-coloured tincture, which is not rendered turbid on the addition of water. Ether takes up a portion of it, and forms a yellowish-red tincture, which, when evaporated on water, leaves no resinous pellicle.

All the varieties, according to Vauquelin, contain a species of tannin; but no gallic acid. They dissolve in solutions of pure potass and of ammonia, and no precipitation takes place on the addition of water. Some chemical change, however, is effected; and the astringent property of the kino is completely destroyed, a fact which ought to be kept in remembrance in prescribing this remedy.

The following tables show the result of some experiments with several chemical re-agents on the watery infusions of these three varieties of Kino.<sup>2</sup> They point out the distinctive features of the four varieties I have enumerated; but they have no pretensions towards advancing the knowledge of the chemical properties of Kino.

<sup>1</sup> In India, an aqueous solution of Kino is employed for dying the colour termed *Nankcen* on cotton and other clothes. Virey, *Bullet. de Pharm.* tom. iv. p. 364.

<sup>2</sup> The specimens subjected to these experiments, I have reason to think, were perfectly genuine. The African kino was brought home twenty years ago.



TABLE I. Precipitates formed in the Aqueous Solution of Kino, by Gelatine and Solutions of some Metallic Salts.

Variety of Kino.	Solution of Isinglass.	Solution of Oxy-sulphate of Iron.	Solution of Nitrate of Silver.	Solution of Oxy-muriate of Mercury.	Solution of Acetate of Lead.
1st.	copious, slowly formed, of a brick-red colour.	copious, quickly formed, of a dirty olive-black.	copious, slowly formed, of a deep reddish brown.	not very copious, slowly formed, reddish.	copious, flocculent, quickly formed, brown.
2d.	copious, almost instantly formed, of a pink colour.	very slowly formed, of a deep brownish black.	copious, quickly formed, of an olive-black.	copious, quickly formed, yellowish pink.	copious, flocculent, quickly formed, lilac.
3d.	scanty, slowly formed, of a pinkish colour.	copious, quickly formed, of a blue-black.	copious, quickly formed, reddish brown.	scarcely altered.	copious, flocculent, quickly formed, brownish-lilac.
4th.	the same as No. 1.	copious, and dirty olive-black.	copious, and quickly formed, reddish-brown.	quickly formed, reddish.	the same as No. 1.

TABLE II. Precipitates formed by Solution of Potass and Acids.

Variety of Kino.	Potass.	Sulphuric Acid.	Nitric Acid.	Muriatic Acid.
1st.	none, but renders it clear, and of a deep brown colour.	copious, pale brown.	scanty, slowly formed, reddish-yellow.	scanty, slowly formed, yellowish-brown.
2d.	flocculent, purplish.	copious, deeper brown.	copious, quickly formed, yellowish-brown.	scanty, more quickly formed, pale red-brown.
3d.	flocculent, brownish purple.	very copious, very deep brown.	copious, brown.	scanty, quickly formed, a beautiful red.
4th.	the same as No. 1.	copious, pale brown.	copious, quickly formed, brown.	quickly formed, yellowish-brown.



From these experiments there appears to be a considerable difference between three of the four varieties of Kino known in commerce, but the first and the fourth appear to be nearly the same. The most remarkable differences are, the small portion of resin which that from Botany Bay and Amboyna contain; the blue colour of the precipitate of the Jamaica variety by the oxy-sulphate of iron; and the effect of the solution of potass in rendering that from Africa transparent, while it precipitates the second and the third varieties. The predominant principles, in all the varieties, are tannin and extractive matter; and the portion of resin, in the first and third varieties, enables either to take up their colouring matter and some extractive, whilst the second variety is scarcely affected by it. Dr. Duncan<sup>1</sup> and Vauquelin<sup>2</sup> observed, that, although heat increases the solvent power of water over Kino, yet, that a substance insoluble either in water or in alcohol always remains. Vauquelin also found that the solutions form a precipitate with tartarized antimony and the salts of iron.

The best menstruum is diluted alcohol.

*Medical properties and uses.* — Kino is a powerful astringent. Like catechu, it is employed in obstinate chronic diarrhœas, lentergy, uterine and intestinal hæmorrhages, and fluor albus; but as it is supposed to be less certain in its qualities than catechu, it is less used. It has also been used for the cure of intermittents; and is said to increase the power of Cinchona. Externally, it has been applied as a styptic, and to give tone to, and diminish the ichorous discharge of, flabby ill-conditioned ulcers. The alkalies, as already stated, destroy its astringent qualities.

It may be exhibited internally in substance, or in the form of watery infusion, or of tincture. The dose in substance is from grs. x. to ʒss.; that of the infusion fʒj. ss. and of the tincture fʒj. In ordering the infusion or the tincture, it is necessary to recollect that solutions of isinglass, sulphate of iron, nitrate of silver, muriate of mercury, acetate of lead, tartarized antimony, the alkalies, and the strong acids, are incompatible in prescriptions with Kino.

Officinal preparations. *Tinctura Kino*, L. E. D.

PULEGIUM. Vide *Mentha Pulegium*.

PUNICA. *Spec. Plant. Willd.* ii. 981.

Cl. 12. Ord. 1. Icosandria Monogynia. Nat. ord. Pomaceæ, Linn. Myrtæ, Juss.

<sup>1</sup> *Nicholson's Journal*, vi. 234.

<sup>2</sup> *Annales de Chimie*, xlv. 521. Vauquelin states generally, that the salts of iron precipitate Kino green; but Dr. Duncan justly observes, that by the red sulphate it is precipitated black: the sulphate only precipitates it green.



G. 980. *Calyx* five-cleft, superior. *Petals* five. *Pome* many-celled, many-seeded.

Sp. 1. *P. Granatum*.<sup>1</sup> Pomegranate tree. *Med. Bot.* 2d edit. 531. t. 190.

*Officinal.* GRANATI CORTEX, *Lond.* GRANATUM; FLORES, PERICARPII CORTEX, *Dub.* Pomegranate bark, and flowers.

*Syn.* Le Grenadier (*F.*), Granatass felschale (*G.*), Granaat-boom (*Dutch*), Drzewo Granatowe (*Pol.*), Pomo Granato (*I.*), Romã (*Port.*), Granado (*S.*), Granatnik (*Russ.*), Rānā (*Arab.*), Anár (*H.*), Darim (*San.*), Delima (*Javanese*), Madalum pallum (*Tam.*), Dalema (*Malay*), Deloomghedie (*Cyng.*).

The Pomegranate tree is a native of the south of Europe, Asia, and Barbary; but in the West Indies, where it was introduced from Europe, the fruit is larger and better flavoured than in its native climates.<sup>2</sup> In its proper soil, which is a cretaceous one, it rises twenty feet in height, sending out branches from the whole length of the stem, some of which bear thorns. The leaves are opposite, about three inches long, half an inch broad in the middle, pointed at each end, entire, and of a light, lucid-green colour: the flowers which are terminal, and sessile, are three or four together: the calyx is thick, fleshy, of a fine red colour, and divided into five pointed segments: the petals are wrinkled, and of a scarlet colour: the fruit, which is cellular, according to Gærtner<sup>3</sup>, is a pulpy, many-seeded berry, the size of an orange, crowned with the calyx, which is sharply toothed, globular, a little flattened, and covered with a thick, coriaceous rind.

The red succulent pulp, which is not officinal, is pleasantly acid<sup>4</sup>, resembling that of the orange: it is cooling, and useful for quenching thirst, and gently aperient.

*Qualities.*—The *flowers*, which are named *Balaustines*, are inodorous, and taste bitterish and astringent. The *bark of the fruit* has the same sensible qualities. Water extracts the virtues of both; and the solutions strike a deep bluish-black with sulphate of iron.

*Medical properties and uses.*—Both the parts we have described are astringent. They are given in the form of decoction in chronic and colliquative diarrhœa, and the protracted stage of dysentery. They are supposed to prove beneficial also in checking the violent sweating which accompanies hectic fever; but the chief use of the decoction is an injection in leucorrhœa, or as a gargle in sore throats, after the local inflammation is moderated. Dr. Buchanan has stated, that the

<sup>1</sup> *Poa Dioscoridis*, *Han Xe lieu* (Chin.).

<sup>2</sup> It stands our winters, and even bears fruit, which, however, has not the proper flavour.

<sup>3</sup> *De Fructibus*, i. 183. t. 38. f. 1.

<sup>4</sup> Russel says there are three varieties of this fruit in Syria; one sweet, another very acid, and a third partaking of the qualities of both blended. *Nat. Hist. of Aleppo*, ii. 85.



bark of the *root* of the plant has been long used by the natives of Hindostan, and according to M. Deslandz, by the negroes of St. Domingo, for expelling tape worm; and its utility for this purpose has been fully confirmed by the experiments of Mr. Breton<sup>1</sup>, who gave it in the form of powder, in doses of ℥j., and of decoction, prepared by boiling, ℥ij. of the bark in Oj. ss. of water, and reduced to f℥ix., of which, when cold, a glassful was given every half hour, until four doses were taken. The worm was generally voided alive, a few minutes after the last dose. Celsus says, it was used by the ancients for a similar purpose.<sup>2</sup> The bark and flowers are given in the form of powder, in doses of a scruple increased to a drachm; or, of a decoction made with ℥iv. of the bark and f℥vj. of water, six fluid drachms may be given every three hours.

**PYRETHRI RADIX.** Vide *Anthemis Pyrethrum*.

**PYRUS.** *Spec. Plant. Willd.* ii. 1012.

*Cl.* 12. *Ord.* 5. Icosandria Pentagynia. *Nat. ord.* Pomaceæ, *Linn.* Rosaceæ, *Juss.*

*G.* 992. *Calyx* five-cleft. *Petals* five. *Pome* inferior, five-celled, many-seeded.

*Species* 17. *P. Cydonia.* The Quince tree. *Med. Bot.* 2d edit. 505. *t.* 182.

*Officinal.* CYDONIÆ SEMINA, *Lond.* Quince seeds.

*Syn.* Semen de Coignassier (*F.*), Quittenkörner (*G.*), Kwee (*Dutch*), Quaede (*Dan.*), Quitten (*Swed.*), Pigwa (*Pol.*), Semi de Cotogno (*I.*), Simiente de Membrillo (*S.*), Marmelo (*Port.*), Hubusnfirjul (*Arab.*), Belcheeybeej (*H.*), Behdanæ (*Tam.*), Bedana (*Pers.*).

The Quince tree was originally brought from Cydon<sup>3</sup> in Crete by the Greeks: but, it has been found growing wild in Germany and on the rocky shores of the Danube, and is cultivated to great perfection in England, and many other parts of Europe; flowering in May. It is a low, crooked tree, with many spreading branches, and covered with a brown bark. The leaves are ovate, very entire, about 2½ inches long, and 1½ inch broad, of a dusky green colour on the upper surface, paler and downy beneath: the flowers are large and solitary: the calyx, which is of the length of the corolla, is spreading, serrated, persistent, and villous: the petals rose-coloured or white, concave, roundish, and inserted into the calyx: the filaments are awl-shaped, purplish, and support yellow anthers: the fruit is, according to Gärtner, a berry.<sup>4</sup>

<sup>1</sup> Vide *Medico-Chirurg. Trans.* vol. ii. p. 501.

<sup>2</sup> Vide *Celsus de Medicina*, lib. iv. § xvii.

<sup>3</sup> Whence its Greek name *μηλεα Κυδωνια* (Theophrasti) is derived. It is supposed to be the apple of the Hesperides.

<sup>4</sup> *De Fructibus*, ii. 45. *t.* 87.



Its magnitude and shape are those of a moderate-sized pear.<sup>1</sup> It is of a yellow colour, downy, umbilicated; and, when ripe, has a pleasant odour and a very austere, acidulous taste<sup>2</sup>: each of its cells contains from eight to fourteen ovate, angled, reddish-brown, coriaceous seeds, placed erect in pairs.

*Qualities*.—The seeds are inodorous, and nearly insipid, having a slight bitterness only when long chewed. The inner coat contains a very considerable quantity of mucus, which can be extracted by hot water: but is not quite pure mucus, being mixed with fecula and the other soluble parts of the seeds. It contains also gallic and malic acid. For its qualities and medicinal use, see *Decoctum Cydoniæ* among the Preparations.

QUASSIA.<sup>3</sup> *Spec. Plant. Willd.* ii. 567.

*Cl.* 10. *Ord.* 1. Decandria Monogynia. *Nat. ord.* Gruinales, *Linn.* Magnoliæ, *Juss.*

*G.* 849. *Calyx* five-leaved. *Petals* five. *Nectary* five-leaved.

*Drupes* five, distant, bivalve, inserted into a fleshy receptacle.

*Species* 2. *Q. Simaruba*. *Simaruba Quassia*. *Med. Bot.* 2d edit. 569. t. 203. *Trans. of the Royal Society of Edin.* ii. 73—81.

*Species* 3. *Q. excelsa*. Lofty Quassia. *Trans. of the Royal Society of Edin.* iii. 205—210. t. 6.

#### 1. QUASSIA SIMARUBA.

*Officinal.* SIMARUBÆ CORTEX, *Lond.* QUASSIÆ SIMARUBÆ CORTEX, *Edin.* SIMARUBA; CORTEX, LIGNUM, *Dub.* *Simaruba Bark and Wood.*

*Syn.* Ecorce de Simarouba (*F.*), Simarubenrinde (*G.*), Corteccia de Simaruba (*I.*).

The *Simaruba Quassia*, or mountain Damson, as it is called in Jamaica, is a native of South America, Carolina, and the West India islands, growing in sandy places. It is a tall tree with alternate branches, and a smooth, grey bark, maculated with yellow spots. The leaves are pinnate, consisting of from two to nine leaflets placed alternately on short petioles, elliptical, acute, smooth, and of a deep-green colour above, and whitish beneath: the flowers are male and female on the same axillary panicles: the calyx in both is monophyllous and five-toothed: the petals lanceolate, yellowish, white, and inserted into the calyx: the nectary in the male is a small scale affixed to the inner part of the base of each filament; and the same

<sup>1</sup> The Quince-tree varies in the form of its fruit, which is sometimes globular, sometimes oblong, but more generally pyriform; and, also, in the magnitude of its leaves.

<sup>2</sup> Although the fruit of the quince is not good in its raw state, yet it affords an elegant sweetmeat, called Quince Marmalade, *Mira Cydoniarum*: and from the expressed juice an excellent and wholesome wine is prepared.

<sup>3</sup> Named after Quassia, a negro slave, who discovered to Rolander the wood of the *Quassia excelsa*, which he had employed with success as a secret remedy in the malignant endemic fevers of Surinam. The genus is placed in the class *Decandria*, because the *Quassia amara*, from which the generic description is taken, has hermaphrodite flowers.



in the female, except that the scales are placed in a regular circle; the filaments in the male are the length of the corolla; and in the female are five connate germens, with five striated styles and spreading stigmas; the fruit, according to Gærtner, consists of five smooth, ovate, black, one-celled berries, on a common receptacle: and opening spontaneously when ripe.<sup>1</sup>

The officinal part of this tree is the bark of the root; and although the wood is designated by the Dublin College, yet it is quite inert. The bark is imported in long pieces, a few inches in breadth, and folded lengthwise. It comes generally from Jamaica packed in bales.

*Qualities.* — Simaruba bark is inodorous, and has a bitter, but not disagreeable taste. The pieces are of a very fibrous texture, rough, scaly, warty, and of a full-yellow colour in the inside when fresh. Alcohol, Wine, and also Water take up all its active matter by simple maceration, at a temperature of 60° Fahrenheit, better than at a boiling heat. The infusion is stronger in taste than the decoction, which grows turbid and of a reddish brown colour as it cools. The infusion is not affected by sulphate of iron, and scarcely by muriate of tin. M. Morin and Rouen have analysed Simaruba and found in it resin, a volatile oil having the odour of Benzoin, malic and gallic acids, an ammoniacal salt, acetate of potass, malate and oxalate of lime, oxide of iron, alumen and a peculiar bitter principle which they have named *quassine*.<sup>2</sup>

*Medical properties and uses.* — This bark is tonic; and has been employed with advantage in intermittent fever, obstinate diarrhœa, dysentery and dyspeptic affections. It was first introduced at Paris in 1713 as a powerful remedy in dysentery; but its effects in this disease were previously known to the natives of Guiana, whence it was brought to France. Simaruba bark, however, was little known in this country till Dr. Wright's paper on it appeared in the Edinburgh Transactions. It cannot with propriety be used in the commencement of dysentery; but after the fever has abated, when the tenesmus continues with a weak, sinking state of the pulse, it allays this symptom and griping, promotes the secretion of urine, determines to the surface, and restores the tone of the intestines. It has also been highly commended as a remedy in fluor albus; but notwithstanding the high character which it acquired, Simaruba is not much employed by the British practitioner. It may be combined with aromatics and opium.

---

<sup>1</sup> Owing to this circumstance, Gærtner allows that the pericarp is something between a berry and a capsule. It is certainly not a drupe, although mentioned as such in the generic character by Willdenow.

<sup>2</sup> *Journal de Pharmacie.* Février, 1822.



The dose in substance is from ℥j. to ʒss.; but it is more frequently and commodiously given in the form of infusion.

Officinal preparation. *Infusum Sinarubæ*, L.

## 2. QUASSIA EXCELSA.

*Officinal*. QUASSIÆ LIGNUM<sup>1</sup>, *Lond. Edin. Dub.* The wood of Quassia.

*Syn.* Bois du Quassie (*F.*), Quassienholz (*G.*), Legno della Quassia (*I.*), Leno de Quassia (*S.*), Pao de Quassia (*Port.*).

This species of Quassia grows in the natural woods of Surinam, Jamaica, and the Caribbean islands, where it is called the bitter ash; and flowers in October and November. It is a beautiful tall tree, rising sometimes one hundred feet in height, with a straight, smooth, tapering trunk, often ten feet in circumference near the base; and covered with a smooth grey bark. The leaves are pinnate, consisting of from five to eight opposite pairs of leaflets, with a terminal leaflet: they are oblong and pointed; the ribs reddish; and the young leaves are covered with a fine brown down: the flowers are in clusters from the lower part of the last shoot before the leaves; they are small, of a yellowish-green colour, with a very small calyx: the male flowers are nearly similar to the hermaphrodite, except that they have the rudiments only of a style; the fruit is a small black drupe, round, the size of a pea, and attached in threes, sideways, to a round fleshy receptacle. It is ripe in December, and is not bitter.<sup>2</sup> The wood is sent to this country in billets, and is reduced to chips, or rasped by the druggists.<sup>3</sup>

*Qualities*. — Quassia wood is inodorous, and has an intensely bitter taste; it is of a pale yellow colour. Alcohol and water take up its bitterness, and when evaporated to dryness, leave a brownish-yellow, somewhat transparent, brittle extract, which has been regarded as a vegetable constituent *sui generis*, and named *quassine*, or the bitter principle.<sup>4</sup> I am inclined to believe that this principle, although not itself of a resinous nature, is connected with resin, as ether takes it up, and the tincture, when evaporated on water, which becomes intensely bitter, leaves an insoluble pellicle that has the character of a resin. The infusion is rendered muddy by nitrate of silver, a soft, flaky, yellow precipitate being formed; and acetate of

<sup>1</sup> The officinal Quassia was long erroneously supposed to be the wood of the Quassia amara, which is a very rare tree, and excels all the others in bitterness.

<sup>2</sup> *Edin. Phil. Trans.* iii. 207.

<sup>3</sup> It is asserted, that of late years the brewers have used Quassia wood instead of hops. Beer made with it certainly does not keep, but soon becomes muddy, flat, has a mawkish taste, and runs into the acetous fermentation. Mr. Brand says, an infusion of Quassia sweetened with brown sugar is an effectual poison for flies. *Manuel*, p. 146.

<sup>4</sup> *Thomson's Chemistry*, 4th edit. v. 32.



lead occasions a copious white precipitate: thence, these salts are incompatible in formulæ with it: but it is not affected by tartarized antimony, sulphate of iron or gelatin.

*Medical properties and uses.* — Quassia is tonic. It has been found efficacious in dyspepsia and nervous irritability; intermittent and bilious remittent fevers, chlorosis, diarrhœa; and, when combined with cretaceous powder and ginger, in atonic gout. It does not sensibly quicken the circulation nor augment the animal heat. We have given it, combined with nitric acid, with evident benefit in typhus, gout, and also in fluor albus. Infusion is the best form of administering Quassia; the raspings, for it cannot be properly pulverized, being too bulky: but it may, nevertheless, be given in substance in doses of from grs. x. to ʒj. three or four times a day.

Officinal preparations. *Infusum Quassiae*, L. *Tinctura Quassiae excelsæ*, E. D.

QUERCUS. *Spec. Plant. Willd.* iv. 423.

Cl. 21. Ord. 6. Monœcia Polyandria. Nat. ord. Amentaceæ. G. 1692. Male. Calyx commonly five-cleft. Corolla none. Stamens five to ten.

————— Female. Calyx one-leafed, entire, rough. Corolla none. Styles two to five. Nut coriaceous, surrounded at the base by the persistent calyx.

\* \* With toothed leaves.

Species 33. *Q. infectoria*. Dyers' Oak. *Olivier's Travels* (translation), ii. 41. t. 14, 15.

\* \* \* \* \* With sinuated leaves and beardless lobes.

Species 65. *Q. pedunculata*. Common White Oak. *Med. Bot.* 2d edit. 23. t. 10. *Michaux, North American Sylva*, vol. i. pl. 2. (*Q. Robur*.) *Smith, Flor. Brit.* 1026. *Michaux*, b. c.

#### 1. QUERCUS INFECTORIA. (*Quercus Cerris*, Edin.)

Officinal. GALLÆ, Lond. Edin. GALLÆ, Dub. The Gall.

Syn. Noix de Galles (*F.*), Galläpfel (*G.*), Galnoot (*Dutch*), Galdaebel (*Dan.*), Gallaple (*Swed.*), Galla (*I.*), Agalla de Levante (*S.*), Galha (*Port.*), Galdæble (*Dan.*), Maju P'hal (*H. and San.*), Makakai (*Tam.*), Afis (*Arab.*), Mazu (*Pers.*).

The London and Dublin Colleges have not named any particular species of oak as furnishing the Gall; the Edinburgh College has particularized the *Cerris*: but although it, as well as most of the other species of *quercus*, may produce galls, yet it is not the species from which the galls of commerce are obtained. Olivier has distinctly pointed out, from his personal knowledge, the species above named as the real tree: and as we know no reason for doubting his veracity, we shall copy his description of it.

The *Quercus infectoria* is scattered throughout all Asia Minor, from the Bosphorus as far as Syria, and from the coasts of the Archipelago as far as the frontiers of Persia. It has a crooked stem, seldom exceeds six feet in height, and more



frequently assumes the character of a shrub than that of a tree. The leaves, which are deciduous in autumn, are on short petioles, smooth, of a bright-green colour on both sides, and obtusely toothed: the acorn is elongated, smooth, two or three times longer than the cup, which is sessile, in a slight degree downy, and scaly: the gall comes at the shoots of the young boughs, and acquires from four to twelve lines in diameter: the insect which produces it is the *Cynips quercusfolii* of Linnæus, (*Diplolepis gallæ tinctoriæ* of Geoffroy,) a small hymenopterous insect or fly, with a fawn-coloured body, dark antennæ, and the upper part of the abdomen of a shining brown. The insect punctures the tender shoot with its sting, which is spiral, and deposits its egg in the puncture. This occasions a morbid irritation in the vessels of the part; the gall rises in a few hours and attains its full size in a day or two, before the larva is hatched: the egg grows with the gall; and it is the irritation which it keeps up, not as has been supposed by the maggot feeding on the juices of the plant, that the morbid excitement is maintained in the vessels of the part, sufficient for the production of this kind of vegetable wen.

Galls are gathered before the larva within them changes to a fly, and eats its way out; for, when this has happened, the galls become lighter, and contain less of the astringent principle. The first galls that are picked are named *yerli* by the natives, and are known in trade by the terms *black* or *blue galls*, and *green galls*. Those which are gathered afterwards, from the circumstance of their being pierced, are of an inferior quality, and are denominated *white galls*. The best galls are those of Aleppo, Smyrna, Magnesia, Karahisser, Diarbekir, and the interior of Natolia. Those which are brought to this country, come chiefly from Aleppo in bags and cases.

*Qualities.*—Galls are inodorous, and have a bitter, very astringent taste. They are nearly round, of different magnitudes, from the size of a pea to that of a hazel-nut; smooth on the surface, yet studded with tuberosities; and, when good, of a blackish-blue, or deep olive colour: a white or a red hue indicates an inferior quality.<sup>1</sup> They are heavy, brittle, break with a flinty fracture, and display a compact, striated texture. The whole of their soluble matter is taken up by about forty times their weight of boiling water, and what remains is tasteless. Alcohol, digested on powdered galls, takes up seven

---

<sup>1</sup> This is the character of the galls from which the insect has escaped, and which are also of a brighter colour. Another species of gall, produced by another species of the insect, is also, Olivier says, found on the same oak. It is spongy, very light, of a brown-red colour, covered with a resinous coat, and furnished with a circular row of tubercles placed nearly towards the most protuberant part. Their astringency is very inferior, and they are used only to adulterate the better sort.



parts in ten, and ether five parts. The watery infusion reddens tincture of litmus, and forms precipitates with solution of isinglass, the infusions of Cinchona bark, Cusparia bark, and Columba root; but not with infusion of Quassia. Sulphuric acid throws down a yellowish curdy, muriatic acid, and a flaky, whitish precipitate; while nitric acid changes the colour only of the infusion, first to a deep orange, and afterwards to a paler orange-yellow. The solution of ammonia occasions no precipitate, but deepens the colour: the carbonate, however, produces a precipitate. Carbonate of potass throws down a yellowish, flaky precipitate, and extricates ammonia; and lime-water a copious, deep-green precipitate. Precipitates also are formed with solutions of the following metallic salts: with acetate and superacetate of lead, greyish; with tartarized antimony, yellowish; with sulphate of copper, brown; with sulphate of iron, blueish-black; with sulphate of zinc, reddish-black, but very slowly formed; with nitrate of silver, deep olive; and with nitrate of mercury, bright yellow. The muriate of mercury renders the infusion milky and opaque, but no precipitate is formed. The alcoholic tincture reddens litmus, and is affected by the same re-agents as the watery infusion. The ethereal tincture, when evaporated on water, leaves on the side of the glass an opaque pellicle, and on the surface of the water small drops of an oily, resinous-like matter, while the substratum of water becomes charged with tannin and gallic acid. The pellicle and resinous-like matter is plastic, tenacious, resembling bird-lime treated with ether; and when subjected to heat, melts, swells, burns, and leaves a dense, black charcoal. These experiments show results which cannot altogether depend on the presence of tannin, gallic acid, extractive, or mucilage, which are supposed to be the constituents of galls. In Sir H. Davy's experiments, 500 grains of Aleppo galls yielded to pure water, by lixiviation, 185 grains of solid matter, of which 130 were tannin, 31 gallic acid and extractive, 12 mucilage and matter rendered insoluble by the evaporation, and 12 saline and earthy matter. From different experiments, the proportion of extractive, however, if any, is very small: none appears in the evaporation of the ethereal tincture; and Dr. Bostock's experiments render the existence of mucilage very doubtful. From the experiments of Professor Branchi, it appears that galls also yield, by distillation with water, a concrete, volatile oil<sup>1</sup>; and M. Braconnot has discovered in them a new acid, which he has rather affectedly termed *ellagic*, from the word *galle* reversed!<sup>2</sup> It is an insipid, inodorous, white

---

*Phil. Mag.* vol. .p. 401.

<sup>2</sup> *Ann. de Chim. et Phys.* ix. p. 187.



powder, with a slight tinge of red, and insoluble in boiling water. When mixed with nitric acid, and very gently heated, the mixture acquires a reddish tint, gradually passing to a deep blood-red. Hence we may conclude, that the constituents of galls, besides tannin and gallic acid, are the above oil and ellagic acid.

*Medical properties and uses.* — Galls are the most powerful of the vegetable astringents. They are seldom used as an internal remedy, although, in combination with bitters or aromatics, they have been given in obstinate diarrhœas, passive intestinal hæmorrhages, and intermittents. They are frequently ordered in the form of gargles and injections; and an ointment formed of galls in fine powder, with eight parts of simple ointment, and a small proportion of powdered opium, is a useful application to blind piles. For internal exhibition, the dose of galls may be from grs. x. to ℥j., which may be given twice or thrice a day.

Officinal preparation. *Tinctura Gallarum*, E. D.

## 2. QUERCUS PEDUNCULATA.<sup>1</sup>

*Officinal.* QUERCUS CORTEX, *Lond.* QUERCUS ROBORIS CORTEX, *Edin. Dub.* Oak Bark.

*Syn.* Ecorce de la Chene commune (*F.*), Eichenrinde (*G.*), Cortecchia della Quercia (*I.*).

This species of Oak is indigenous: but it is found also all over Europe, in the north of Asia, and the northern extremity of Africa. It is a well-known beautiful tree, often rising to the height of 90 or 100 feet, and attaining a great degree of thickness in the trunk, which is covered with a rough, brown bark. The leaves are alternate, supported on short petioles, ovate-oblong, deeply sinuated, forming obtuse lobes: deep green, smooth, and shining on the upper surface; paler, and nearly glaucous underneath: the flowers are in axillary catkins; the male lax, pendulous, many-flowered, and yellow; the female longer, peduncled, and only three-flowered: the calyx of the male flower is membranous, bell-shaped, often five-cleft; while that of the female is coriaceous, scaly, downy, and becoming hemispherical, entire and woody: the stamens are ten, longer than the calyx: the germen is ovate, crowned with a short cylindrical style and three stigmas: the fruit is an elliptical, coriaceous, smooth nut, fixed in the calyx as in a shallow cup, but at length dropping from it. The peduncles of the fruit, which are from two to three inches long, distinguish it from the *Q. Robur*, the fruit of which is sessile. It ripens its acorns in October.

<sup>1</sup> Δρυσ Græcorum, Eiche (*G.*), Eik (*Dutch*), Eeg (*Dan.*), Ek (*Swed.*), Darach (*Galic.*), Le Chene (*F.*), Quercia (*I.*), Roble (*S. Port.*), Dub (*Russ.*), Mesche (*Turk.*), Baalut (*Arab.*), Tamma (*Finnl.*), Pelut (*Pers.*).



Almost every part of the oak is astringent, but the bark only is officinal; and, as its epidermis is perfectly inert, it is taken for medicinal purposes from the smaller branches, the epidermis of which is still thin, and scarcely cracked. The bark cut in spring is preferable to that which is cut in winter, as it contains four times the quantity of the astringent principle or tannin.<sup>1</sup>

*Qualities.* — Oak bark is inodorous, has a rough astringent taste, and yields its virtues to both alcohol and water. The watery infusion is affected by all those tests which indicate the presence of gallic acid, tannin, and extractive (see *Decoctum Quercus*). Sir H. Davy<sup>2</sup> found that 3j. of the inner cortical part of young oak bark affords, by lixiviation, 111 grains of solid matter, of which 77 are tannin; the cellular integument, or middle-coloured part, yields grs. 43 only of solid matter, of which 19 are tannin; and the epidermis furnished scarcely any quantity either of tannin or of extractive. The quantity of tannin, however, varies according to the size and age of the trees, and the season at which they are barked. Vauquelin discovered that the infusion of Oak bark does not precipitate tartarized antimony, or the infusion of Santa Fe Cinchona, which resembles the officinal Red Cinchona, although both of these are precipitated by infusion of Galls. I find, however, that infusion of Oak bark forms a precipitate with infusion of yellow Cinchona bark.

*Medical properties and uses.* — Oak bark is tonic and astringent. It has been given, united with bitters and aromatics, with seeming advantage, in intermittents; but it is in every respect inferior to Cinchona, and cannot be depended on. It is, however, useful in obstinate diarrhœa and alvine hæmorrhages; and it is strongly recommended in the malignant coryza (*snuffles*) of infants, when, in spite of keeping the bowels regular, and the use of cordials, the child becomes weak and pallid.<sup>3</sup> Its principal use is as a local astringent. (See *Decoctum Quercus*.)

The dose in substance may be from grs. xv. to grs. xxx.; but it is so difficultly pulverized that it is seldom given in this form.

Officinal preparation. *Decoctum Quercus*, L.

RHAMNUS. *Spec. Plant. Willd.* i. 1092.

Cl. 5. Ord. 1. Pentandria Monogynia. Nat. ord. Dumosæ, Linn. Rhamni, Juss.

G. 405. Calyx tubular. Corolla, scales defending the stamens inserted into the calyx. Berry.

<sup>1</sup> Biggin. *Phil. Trans.* 1799.

<sup>2</sup> *Phil. Trans.* 1803.

<sup>3</sup> Underwood, *Diseases of Children*, 4th ed. i. 45.



\* *Thorny*.

*Species* 1. *R. catharticus*.<sup>1</sup> Purging Buckthorn. *Med. Bot.* 2d edit. 594. t. 210.

*Officinal*. RHAMNI BACCÆ, *Lond.* RHAMNI CATHARTICI SUCCUS, *Edin.* RHAMNUS CATHARTICUS; BACCÆ, *Dub.* Buckthorn Berries.

*Syn.* Nerprun (*F.*), Kreutz beeren (*G.*), Bacche del spino Cervino (*I.*).

This is an indigenous shrub, growing in woods and hedges near brooks; flowering in May and June, and ripening its fruit in October. It rises with a strong, rigid, woody, stem, sending off alternate, round branches, which terminate in a spine. The leaves are in fascicles, on footstalks, ovate, serrated, nerved; and the younger are downy: the flowers come from the same buds as the leaves; they are peduncled, of a greenish-yellow colour, four-cleft; and frequently, but not always, they are male and female upon different plants: the anthers are round, on short filaments which rise from the base of a small convex scale: the germen is ovate, with a slender style and four-cleft stigma: the fruit is a small, round, black, four-seeded berry, about the size of a pea, compressed on one side.<sup>2</sup>

These berries are said to be often mixed with those of the black-berried alder and of the dog-berry tree: but as the Buckthorn berry has four seeds, while the others have only two and one, it can be easily distinguished.

*Qualities*.—The odour of these berries is faint and unpleasant, the taste bitterish, acrid, and nauseous. They are very succulent, and yield by expression a deep green juice, or a purple juice if they be gathered late in the autumn.

*Medical properties and uses*.—The berries, and their expressed juice, are briskly cathartic; but their operation is accompanied with thirst and severe griping, which is not altogether mitigated by the most plentiful dilution. They were formerly much used as a hydragogue purgative, but are now very seldom prescribed. The dose of the recent berries is ʒj; of the dried ʒj, and that of the expressed juice fʒj.

*Officinal preparation*. *Syrupus Rhamni*, L. E.

RHEUM.<sup>3</sup> *Spec. Plant. Willd.* ii. 488.

*Cl.* 9. *Ord.* 3. Enneandria Trigynia. *Nat. ord.* Holoraceæ, *Linn.* Polygoniæ, *Juss.*

*G.* 803. *Calyx* none. *Corolla* six-cleft, persistent. *Seed* one, three-sided.

<sup>1</sup> *Purgierdorn* (*G.*), *Purgerende wegedoorn* (*Dutch*), *Korsbaerton* (*Dan.*), *Ge-tappel* (*Swed.*), *Ramno cartico* (*S.*), *Escambrociro* (*Port.*), *Pridoroschnaja igolka* (*Russ.*), *Szaklak krzewia* (*Pol.*).

<sup>2</sup> The pigment called sap green is the inspissated juice of this berry.

<sup>3</sup> *Ῥῆον* *Dioscoridis*. The Rhubarb of the Greeks was the root of *Rheum Rhaponticum*.



*Sp. 2. R. undulatum.* Wave-leaved Rhubarb. *Amœn. Acad.* iii. 212. *t. 4.*

*Sp. 3. R. palmatum.* Palmatum Rhubarb. *Med. Bot. 2d edit.* 662. *t. 231. Phil. Trans.* iv. 292. *t. 12, 13.*

### 1. RHEUM UNDULATUM.

*Officinal.* RHEUM UNDULATUM; RADIX, *Dub.* The root of Waved-leaved Rhubarb.

*Syn.* Rhubarb (*F.*), Rhabarber (*G., Dutch, Swed.*), Rabarbaro (*I.*), Ruibarbo (*S. Port.*), Variatookalung (*Tam.*), Rāwend (*Arab.*), Révand Chîni (*H.*), Ta Hoam (*Chinese*).

This species of Rheum was supposed by Boerhaave to be the true Chinese Rhubarb; and, as it is not unlikely that foreign Rhubarb is taken from several species, that which we receive by way of Canton, which certainly differs, more than simply in the drying, from that which comes through Russia, may be the produce of this plant; the Dublin College, therefore, is right in giving it a place in the list of materia medica. It is a native of China and Siberia, but grows well in this country. The root divides into a number of thick fibres, which run deep, and are extremely yellow within: the leaves, which appear early in the spring, are supported on moderately thick footstalks, channelled on their under side, and plain on their upper: the leaves are long, running to a point much waved on their edges, a little hairy on the upper surface, and very strongly veined on the under: the flower-stem is of a pale brownish colour, rising about four feet high, and dividing into several loose panicles, or bunches of white flowers, which appear in May, and are succeeded by triangular seeds that ripen early in the season.

### 2. RHEUM PALMATUM.

*Officinal.* RHEI RADIX, *Lond. Edin. Dub.* Rhubarb root.

*Syn.* Aechte Rhabarber (*G., Dutch, Dan., Swed.*), Ruibarbo (*S., Port.*), Hai houg (*Chinese*).

This species, like the former, is a native of China and Tartary; and arrives at considerable perfection, when cultivated in this country. The root is perennial, thick, oval, branched, externally brown, and internally of a deep-yellow colour: the stem, which rises eight or ten feet in height, is erect, round, hollow jointed, very slightly furrowed, and maculated with small, oblong, purple streaks: the lower leaves stand upon long, smooth petioles<sup>1</sup>; are numerous, large, divided into five segments, which are deeply sinuated, toothed, and strongly ribbed, the petiole being divided at its apex, into the five midribs of the segments; of a deep-green colour, rough above,

---

<sup>1</sup> These make a much better tart than the leaf stalks of *R. undulatum*, the species generally employed by the pastry cook.



and pale and villous below: those of the stem spring from the joints, are also petiolate, and gradually lessen in size towards the top of the stem: there is a sheathing stipule or *ochrea*, at the base of each stem leaf. The flowers spring from the axilla of the base in numerous panicked clusters: they appear in May: the corolla is divided into six obtuse segments, of a greenish-white colour, tinted with light pinkish-purple:<sup>1</sup> the filaments are nine, slender, the length of the corolla, and furnished with oblong, double anthers: the style is short, with three, reflected, capitate stigmas: and the germen is a triangular seed, inclosed in a capsule with three membranous, reddish margins or alæ.

This plant has been generally believed to be the species which yields the foreign Rhubarb; and under this belief, a very excellent and correct description of it was given by Dr. Hope, professor of Botany at Edinburgh, in the Philosophical Transactions for 1765. He had raised it from seed sent to him by Dr. Mounsey from Petersburg two years before, and found that the root possessed all the medicinal qualities of the best foreign Rhubarb. Since that period, many laudable attempts have been made to introduce the cultivation of Rhubarb into this country, in sufficient quantity to supply our domestic consumption of this valuable drug: but although many individuals have reared large quantities, and some of it extremely good, yet so powerful is prejudice, that very little of it can be sold, and the efforts, therefore, of the cultivators have of late very much relaxed.<sup>2</sup> It is still, however, uncertain which of the species yields the foreign Rhubarb; nor is it of very great importance, as the roots of the two species above described, and another, the *R. compactum*, accord so very closely in their medicinal powers, that any of them may be used with equal certainty of success.

Three varieties of Rhubarb are known in the shops, named from the places whence we receive them; *Russian* Rhubarb, *Turkey* Rhubarb, and *East Indian* or *Chinese* Rhubarb. The two first resemble one another in every respect, appearing to be the root of the same species of plant, prepared in the same mode; and although the East Indian is seemingly the root of a different species, yet we are informed by Dr. Rehman<sup>3</sup>, that it is the same, only prepared with less care.

---

<sup>1</sup> Qy. Whether is the floral envelope calyx or corolla?—It is white at first — becomes red and succulent, and is persistent.

<sup>2</sup> For an excellent account of those different trials, and some very judicious observations on the mode of cultivating rhubarb, see *Miller's Dictionary*, edited by Dr. Martyn, article *Rheum*.

<sup>3</sup> The best Treatise on the Commerce of Rhubarb, and from which much of the information contained in this article has been taken, is from the pen of Dr. Rehman. Vide *Mem. de la Société Imperiale des Nat. de Moscow*, 1809. t. ii. p. 126.



All the Rhubarb of commerce, known under the names Turkey or Russian, grows on the declivities of the chain of mountains in Tartary, which stretches from the Chinese town Sini to the lake Kokonor, near Thibet. The soil is light and sandy: and the Bucharrians assert that the best grows in the shade, on the southern side of the mountains. Rhubarb, however, is also cultivated in China, in the province of Shen-see, where it is called *Hai-houng*. In Tartary the roots are taken up twice a year, in spring and in autumn<sup>1</sup>, and after being cleansed, and decorticated, and the smaller branches cut off, the body of the root is divided transversely into pieces of a moderate size, which are placed on tables, and turned three or four times a day, during five or six days. A hole is then bored through each piece, by which it is hung up to dry, exposed to the air and wind, but sheltered from the sun. In about two months, the roots have lost seven parts in eight of their weight<sup>2</sup>, and are fit for the market. In China the roots are not dug up till winter<sup>3</sup>; and the cultivators, after cleaning, scraping off the bark, and cutting them, dry the slices by frequently turning them on stone slabs heated by a fire underneath; after which the drying is completed by hanging them up in the air, exposed to the greatest heat of the sun.<sup>4</sup> As soon as the Rhubarb has been dried where it is grown, it is conveyed to Si-ning, where it is again cleaned and aired, and after being cut into smaller pieces and sorted, a large hole is drilled through that intended for the Russian market, in virtue of the contract made with the Russian government, for the examination of the heart of the pieces. It is then packed up in Camel's-hair sacks, and conveyed to Mac-ma-tchin, where it is examined previously to its being transported to Kiachta. The whole of the trade in Rhubarb in China, is carried on by one Bucharrian family, which has enjoyed the monopoly since 1772; and it is even by the agents of this family, that it is sold to the English at Canton. This Bucharrian family resides at Sin-ning Fu, a town on the frontiers of Thibet, about 3000 versts from Kiachta, the town on the Russian frontier, where the Rhubarb is purchased on the account of the Russian government. Part of the Tartarian Rhubarb is carried to Turkey through Natolia; but the greater part is conveyed

---

<sup>1</sup> *Bell's Travels.*

<sup>2</sup> *Bath Papers*, iv. 175.

<sup>3</sup> *Bath Papers*, ii. 249.

<sup>4</sup> It is in the process of drying the roots that the British cultivators of Rhubarb are supposed to fail. Baumé proposes to steep the roots in water, to deprive them of their gummy matter, before drying them; then to lay them upon twigs in the open air for twelve hours, and lastly to place them in a stove heated to 120°, till they are dried. When sufficiently dry, the wrinkles must be rasped out, and the pieces shaken together in a barrel, turned on an axis, for half an hour, which covers them with a fine yellow powder formed by their attrition.



by the Bucharrians to Kiachta, where it is examined by a Russian apothecary. The best pieces only are selected and sent to Petersburg. The pieces are roundish, perforated with a large hole, of a yellow or reddish colour on the outside, somewhat soft and friable, and, when broken, exhibit many diverging streaks of a beautiful, bright-red colour. Agreeably to the contract with Russia, all the Rhubarb which is rejected must be burnt: and even that which is approved must undergo another cleaning before it is finally packed up for St. Petersburg.<sup>1</sup> The Chinese Rhubarb, at least what we receive under that appellation, is conveyed to Canton, and there purchased by the East India Company's agents, who purchase all qualities; whence it is brought to this country by sea. It is in oblong, sometimes flat pieces, seldom perforated; considerably heavier, more compact, and less friable than the former kind; of a brownish-yellow colour on the outside; and, when broken, the fracture is hackly, appears of a dull colour, and variegated with yellow, pink, and white. Both kinds are brought to this country in cases and chests.<sup>2</sup>

*Qualities.*— Good *Russian* or *Turkey* Rhubarb has a peculiar, somewhat aromatic odour, and a bitter, slightly astringent, subacid taste; feels gritty between the teeth when chewed, and tinges the saliva of a bright-yellow colour. It breaks with a rough, hackly fracture, is easily pulverized, and affords a powder of a bright buff-yellow colour. It should not be porous, but rather compact and heavy. Water at 212° takes up 24 parts in 60, forming an infusion of a brown colour, nearly clear, and reddening litmus paper. Alcohol extracts 2·7 from 10 parts, and gives a tincture of a rich, golden colour, which reddens tincture of litmus, is not altered in its transparency by the addition of water, and strikes a blackish-olive hue with solution of sulphate of iron, but no immediate precipitate falls. Sulphuric Ether takes up 1·5 in 10 parts of this Rhubarb; the tincture is of a golden yellow hue, and when evaporated on water, leaves a thin pellicle of yellow resin and abundance of extractive dissolved in the water, combined, however, with tannin. *East Indian* or *Chinese* Rhubarb has a stronger odour, and is more nauseous to the taste than the *Turkey*; breaks with a more compact and smoother fracture; and affords a powder of a redder shade. Water takes up 30 parts in 60; the infusion is not so deep-coloured as that of *Russian* Rhubarb, is more turbid,

---

<sup>1</sup> At this examination, each piece is struck with a small mallet, to detach from it any impurities, or decayed parts.

<sup>2</sup> Rhubarb sometimes appears wormeaten; a circumstance which arises from its being eaten, in its dry state, by a small beetle, *Sinondendrum pusillum*. Kirby and Spence's *Entom.* vol. i. p. 252.



and reddens also litmus paper. Alcohol extracts 4 parts in 10; the tincture is of a much deeper colour, and brownish; gives a deeper red to litmus tincture; is rendered slightly turbid by the addition of water; and strikes a green, not blackish olive with sulphate of iron, which it also quickly and copiously precipitates. Ether takes up 2 parts in 10; the tincture is deeper coloured, and when evaporated on water affords the same results as the former kind, except that the compound of tannin and extractive is more soluble.

The infusion of Chinese Rhubarb is more copiously precipitated by solution of isinglass than that of the Russian. Infusion of yellow Cinchona bark throws down a copious, greenish precipitate from infusion of Russian Rhubarb, and a less copious, but more dense, bright-yellow precipitate from that of Chinese Rhubarb.

The following Tables show the effects of re-agents on the aqueous infusions of the two varieties of Rhubarb.

When the residue, after the action of water, is digested in muriatic acid, and solution of ammonia added in excess, the liquid becomes of a deep purple colour, then somewhat milky, and deposits oxalate of lime. What remains consists of woody matter, a small portion of alumen and silex. Of the specimens which I examined, one drachm of Russian rhubarb yielded twenty-six grains of the oxalate, while the same weight of East Indian yielded only eighteen grains.

From the results of the above experiments, rhubarb appears to contain a large portion of *extractive matter*, a very small portion of *resin*, *mucus*, *tannin*, or rather a principle analogous to it, *gallic acid*, a *colouring matter*, much *oxalate of lime*, and minute proportions of *alumen* and *silex*.<sup>1</sup>

---

<sup>1</sup> According to some experiments published by Mr. John Henderson, in the *Annals of Philosophy*, rhubarb is supposed to contain also a peculiar acid, to which he has given the name of *Rheumic*; but M. De Lassaigues has proved, that this is the oxalic acid; which agrees with the result of our analysis. It is remarkable, that Mr. Brande, in an analysis of rhubarb, published in the *Quarterly Journal of Science*, vol. x., does not notice either oxalic acid, or any oxalate as being contained in this root, although they have been found and are mentioned by every other analyst who has examined rhubarb. But it is still more extraordinary, that Mr. Brande has asserted, that “no chemical investigation into the nature of rhubarb, if we except a few experiments upon it, given in Neumann’s Chemistry,” had been made prior to his own; while every one acquainted with pharmaceutical chemistry knows that (not to mention our analysis,) it has been examined by Scheele, Bayen, Deluval, Vauquelin, M. Clarion, De Lassaigues, M. Henry, &c.; all of whom mention oxalic acid and oxalate of lime among its components.



TABLE I. Precipitates formed by Acids, Alkalies, and Neutral Salts.

Variety of Rhubarb.	Sulphuric Acid.	Nitric Acid.	Muriatic Acid.	Oxymuriatic Acid.	Solution of Potass.	Solution of Subcarbonate of Potass.	Lime water.	Muriate of Barytes.	Silicated Potass.
Russian.	copious, greenish-yellow.	scanty, flocculent, pale yellow.	scanty, very slowly formed. yellow.	slowly formed, pale olive.	none, but strikes a deep lake colour.	none, but strikes reddish-brown.	scanty, slowly formed, brown.	scanty, olive-green.	none, but strikes a deep brown.
Chinese.	more copious, brownish-yellow.	less scanty, pale yellow.	scanty, quickly formed, brownish-yellow.	slowly formed, orange-yellow.	none, a deeper lake.	none, but renders it turbid, and deep reddish-brown.	copious, quickly formed, brown.	less scanty, orange-yellow.	none, but strikes a deep brown.

TABLE II. Precipitates formed by Solutions of Metallic Salts.

Variety of Rhubarb.	Solution of Oxysulphate of Iron.	Solution of Nitrate of Silver.	Solution of Nitrate of Mercury.	Solution of Nitrate of Lead.	Solution of Muriate of Mercury.	Solution of Acetate of Lead.	Solution of Tartarized Antimony.
Russian.	copious, nearly black.	scanty, pale greenish-yellow.	copious olive-yellow.	scanty, slowly formed, yellow.	scanty, slowly formed, pale olive.	scanty, greenish-yellow.	scanty, slowly formed, whitish.
Chinese.	copious, deep olive green.	copious, orange-yellow.	copious, heavy, bright yellow.	scanty, slowly formed, deeper yellow.	copious, quickly formed, heavy yellow.	copious, yellow.	scanty, still more slowly formed.



These tables show, that the two varieties differ from each other in several respects. The Russian contains more tannin, oxalate of lime, and resin; the Chinese more extractive and gallic acid. Besides these components, M. Henry found a *fixed oil* soluble in ether and alcohol, an amylaceous matter, supermalate of lime, sulphate of lime, and a salt with a base of potass.<sup>1</sup> But the purgative principle is still unascertained, although it appears to be combined with the extractive, and hence is soluble in water.<sup>2</sup>

*Medical properties and uses.* — Rhubarb is stomachic and astringent or purgative, according to the extent of the dose in which it is administered. With a view to the first-mentioned properties, it is usefully given in dyspepsia, hypochondriasis, and in a weakened, relaxed state of the bowels, combined with ginger, nutmeg, soda, or bitters.

As a purgative it operates mildly, and may be given to the youngest infants. Its operation is quickened by the addition of neutral salts and calomel, the purgative powers of which it also reciprocally augments; so that a compound formed of small portions of rhubarb and a neutral salt or calomel, acts with more certainty and quicker than large doses of either separately taken. Rhubarb is particularly adapted for the majority of cases of diarrhœa, as it evacuates any acrid matter that may be offending the bowels, before it acts as an astringent. Externally it has been applied by friction to produce its purgative effects<sup>3</sup>, and its powder is sometimes sprinkled over ulcers to assist their granulation and healing. It colours the urine after it is taken; and may be detected by the aid of an alkali. It disappears after an hour or two, but reappears, owing to a second absorption from the colon.<sup>4</sup>

The Chinese use it medicinally; but they chiefly employ it to colour a spirituous liquor.

Rhubarb is given in a variety of forms (see Preparations,) but its purgative properties are most powerful in substance. From  $\mathfrak{zj}$  to  $\mathfrak{zss}$  of the powdered root opens the bowels freely; and from grs. vi to grs. x may be given for a dose, when its stomachic properties only are required.

Official preparations. *Infusum Rhei*, L. E. *Vinum Rhei*, E. *Tinctura Rhei*, L. E. D. *Tinctura Rhei composita*, L. *Tinctura Rhei et Aloes*, E. *Tinctura Rhei et Gentianæ*, E. *Pilulæ Rhei compositæ*, E.

<sup>1</sup> *Journal de Pharm.* tom. iv.

<sup>2</sup> M. Navi of Milan, imagines he has discovered a new alkali in Rhubarb. See Account of the Experiments in *Brande's Journ.* N<sup>o</sup> 32. p. 172.

<sup>3</sup> *Nouveaux Elem. de Therap. par Alibert*, tome ii. p. 247.

<sup>4</sup> Sir E. Home.



**RHODODENDRON.** *Spec. Plant. Willd.* ii. 603.

*Cl.* 10. *Ord.* 1. Decandria Monogynia. *Nat. ord.* Bicornes, *Linn.* Rhododendra, *Juss.*

*G.* 867. *Calyx* five-parted. *Corolla* nearly funnel-shaped. *Stamens* declined. *Capsule* five-celled.

*Species* 7. *R. chrysanthum.* Golden-flowered Rhododendron.

*Med. Bot.* 2d edit. 299. *t.* 103. *Pall. Ross.* i. *p.* 44. *t.* 30.

*Officinal.* RHODODENDRI CHRYSANTHI FOLIA, *Edin.* The leaves of Rhododendron.

*Syn.* Rosage (*F.*), Gelber Alpbalsam (*G.*), Rhododendro Aureo (*I.*), Sabina (*Russ.*), Kaschkara (*Koibal*), Schei (*Tartar.*).

This beautiful shrub is a native of the mountainous parts of Siberia, flowering in June and July. It rises a foot in height, and sends off spreading branches which are covered with a brown bark. The leaves are terminal, oblong, ovate, and attenuated to the petiole; few, smooth, stiff, with the margin entire and bent in; the upper surface reticulated, rugged, and of a deep-green colour; and the under ferruginous or glaucous. The flowers are large, yellow, and terminate the branches, forming umbels: the calyx is persistent; the corolla monopetalous, inclining, and irregularly divided into five spreading segments: the filaments are slender, nearly as long as the corolla, supporting oval anthers; and the germen is pentagonal, bearing a long, slender style, crowned with a five-lobed stigma. The capsule is ovate, somewhat angular; and contains many small gray irregular seeds, like sawdust.

*Qualities.* — The leaves of this plant are inodorous, and have an austere, astringent, bitterish taste. Water extracts their virtues either by infusion or decoction.

*Medical properties and uses.* — Yellow rhododendron leaves are stimulant, narcotic, and diaphoretic. When taken, they first increase the arterial action and the heat of the body, producing diaphoresis; and these effects, according to Dr. Home's observations, are followed by a proportional diminution of excitement, the pulse, in one case, having been reduced thirty-eight beats. In large doses they produce nausea, vomiting, purging, delirium, and all the violent symptoms of intoxication. Both the plant and its effects were first described by<sup>1</sup> Gmelin and Steller, in 1747, as a Siberian remedy for rheumatism; but it was not much noticed till after 1779, when Koelpin strongly recommended it in that disease, and also in Gout and Lues venerea. Besides the effects we have already mentioned, it is said to excite a creeping sensation in the pained parts, which after a few hours subsides, and at the same time the pain is relieved. It has not been much used in this country, but,

<sup>1</sup> *Flora Sibirica*, iv. 121.



from the result of some trials of it in Scotland, it has obtained a place in the Edinburgh Pharmacopœia.

It is given in the form of decoction, made by boiling ʒiv of the leaves in fʒx of water in a close vessel, over a slow fire for twelve hours. The dose of the strained liquor is from fʒj to fʒij given twice a day, and gradually increased.

RHUS.<sup>1</sup> *Spec. Plant. Willd.* i. 1479.

Cl. 5. Ord. 2. Pentandria Digynia. Nat. ord. Dumosæ, Linn. Terebintaceæ, Juss.

G. 566. Calyx five-parted. Petals five. Berry with one seed.

\*\* With ternate leaves.

Species 17. R. *Toxicodendron*. Poison oak. Sumach. Kalm's *Travels*, ii. 318.

Officinal. TOXICODENDRI FOLIA, *Lond.* RHOIS TOXICODENDRI FOLIA, *Edin.* Sumach leaves.

Syn. *Toxicodendron*; Herbe à la Puce (*F.*), Giftsumach (*G.*), Vergiftboom (*Dutch*), Rus Tossicodendro (*I.*).

This shrub is a native of North America. It seldom exceeds three feet in height; the root sending up many stems, which divide into slender woody branches, and are covered with a brownish bark. The leaves are placed alternately, supported on long petioles; and are composed of three oval leaflets, about three inches long, and one inch and a half broad, angularly indented, hoary on the under surface, and of a deep-shining, green colour on the upper: the two lateral leaflets are nearly sessile, and the terminal one is considerably the largest of the three: the male flowers which are on a distinct plant from the female, spring from the sides of the stalks in close short spikes, and are of an herbaceous colour; the female, which are larger, are produced in loose panicles, and embosom a roundish germen, supporting three very short styles: the fruit is a striated berry.

The stems, if cut or broken, exude a milky juice, which was supposed to inflame the skin wherever it touched; but this is not the case, although the plant exudes a deleterious vapour.<sup>2</sup> The juice becomes black when it is exposed for a short time to the action of atmospheric air.<sup>3</sup>

*Qualities.* — The leaves of *Toxicodendron* are inodorous, and have a mawkish, subacid taste. Their virtues are completely extracted by water, and partially by alcohol. The

<sup>1</sup> Pous Dioscoridis.

<sup>2</sup> Vans Mons has proved that the acrimonious matter of the plant is exhaled during the night, combined with carbonated hydrogen gas. He collected a jar full of the gas, into which his brother, who was very susceptible of the poison of this species of rhus, plunging his arm, the skin was quickly inflamed and blistered. *Actes de la Soc. de Med. de Bruxelles.*

<sup>3</sup> This juice forms an indelible black stain on linen cloth; and is used in Japan, where the shrub is a native, as a varnish. *Phil. Trans.* xlix. 158.



aqueous infusion reddens litmus paper; precipitates the solution of sulphate of iron black; that of nitrate of silver brown; and throws down a precipitate with gelatine. Hence it contains gum, resin, gallic acid, and tannin; but a narcotic principle is also present, on which its effects principally depend.

*Medical properties and uses.* — The leaves are stimulant and narcotic. In the hands of Dr. Alderson of Hull, who introduced them as a remedy, they proved successful in several cases of paralysis; but we believe their efficacy in this disease has not been confirmed by the observations of other physicians. They excite a sense of heat and pricking, and irregular twitchings in the affected limbs. We believe some advantage has been found from their use in herpetic eruptions. They have also been found useful in the form of tincture, in cramp of the stomach.

The dose of the powdered leaves may be gr. ss, given twice or thrice a day, and gradually increased to grs. iv, in the form of a bolus.

RICINUS. *Spec. Plant. Willd.* iv. 564.

Cl. 21. Ord. 8. Monœcia Monadelphia. Nat. ord. Tricoccæ, Linn. Euphorbiæ, Juss.

G. 1720. Male. Calyx five-parted. Corolla none. Stamens numerous.

—— Female. Calyx three-parted. Corolla none. Styles three, bifid. Capsule three-celled. Seed one.

\* With palmated leaves.

Sp. 1. *R. communis*.<sup>1</sup> Common Ricinus or Palma Christi. *Med. Bot.* 2d edit. 625. t. 221. *Rheede Hort. Malab.* ii. 57. t. 32.

*Officinal.* RICINI SEMINA ET OLEUM, *Lond.* RICINI COMMUNIS SEMINA; ET OLEUM FIXUM, *Edin.* RICINUS; OLEUM E SEMINIBUS EXPRESSUM, *Dub.* The seeds, and the fixed oil of the seeds of the Ricinus. Castor oil.

*Syn.* Le Noix et l'Huile du Ricin (*F.*), Rizinuskorner; Rizinusöhl (*G.*), I semi e l'olio di Ricino (*I.*), Palmoel; Ricinsoel (*S.*), Erundatylum (*San.*), Duhnul Kirwa (*Arab.*), Sittāmoonāka unnay (*Tam.*), Endooroo tail (*Cyng.*), Rowgen Bedangur (*Pers.*), Miniak jarak (*Malay*), Linga jarak (*Jav.*).

This species of Ricinus is an annual plant, a native of Greece<sup>2</sup>, the East and West Indies, South America, and Africa.<sup>3</sup> It is of very quick growth, and sometimes attains to the height of sixteen feet.<sup>4</sup> The stem is round, thick, jointed, furrowed, glaucous in the lower part, but of a purplish-red

<sup>1</sup> KIKI Dioscoridis.

<sup>2</sup> At Pamisus, in the Morea, where it grows in great abundance, it is called *Agra staphylia*, or wild vine, from the resemblance of its leaves to those of the vine. *Gell's Journ. in the Morea*, p. 193.

<sup>3</sup> It was first cultivated in Britain in 1562.

<sup>4</sup> It has been asserted that this plant is in some places perennial, becoming a large tree. Willdenow, however, says, "Planta semper annua, nunquam fruticosa vel arborea, nec in calidissimis terræ plagis lignescit." *Spec. Plant.* iv. 564.



colour towards the top: the leaves are petiolate and subpeltate, large, deeply divided into seven pointed serrated lobes, and of a blueish-green colour: the calyx of the *male* flower is composed of five oval, pointed, purplish segments, inclosing many long stamens united at the base; the *female* is at the upper part of the spike, and is composed of a three-cleft, reddish calyx: the styles are three, slender, and forked at the apex: the capsule is a trilocular nut, covered with rough spines, and bursting elastically to expel the seeds, which are generally three, and of an oblong flat figure, and white colour.

The Oil, which is more frequently used than the Seeds, is obtained from the seeds both by coction and expression. The former method was generally used till lately; and was performed by tying up the seeds, previously decorticated and bruised, in a bag, which was suspended in boiling water till all the Oil was extracted and rose to the surface, when it was skimmed off. This mode of preparation is still preferred by many of the West Indian practitioners; but as the oil is apt to get soon rancid when thus prepared, it is now obtained both at home and abroad, by subjecting the seeds to the press, in the same manner as almonds to procure almond oil. The Oil obtained is equal to one-fourth of the weight of the seeds employed. The acrid principle is contained in the cotyledons; and not in the embryo, nor in the testa. It is of a volatile nature. It is often adulterated with olive oil, linseed oil, and poppy oil, which, however, may be readily detected by adding an equal quantity of *alcohol* sp. gr. 820 to any given quantity of the suspected oil; if it be pure, a uniform solution will take place, which will not happen should it be adulterated: and the same will be the case, if a weaker spirit be employed, with the addition of camphor.

*Qualities.*—Good expressed Castor Oil is nearly inodorous and insipid: but the best leaves a slight sensation of acrimony in the throat after it is swallowed. It is thick, viscid, transparent, and colourless, or of a very pale-straw colour; that which is obtained by coction has a brownish hue; and both kinds, when they become rancid, thicken, deepen in colour to reddish-brown, and acquire a hot, very nauseous taste. It has all the chemical characters of the other expressed oils, except that it is heavier, and is very soluble in alcohol, and also in sulphuric ether. Mr. Brande says that it sometimes undergoes such a change in the bowels that it passes off “in round and indurated nodules.”<sup>1</sup> But such a circumstance has never come within our knowledge.

<sup>1</sup> *Manuel*, p. 150.



*Medical properties and uses.*—The Seeds are drastic cathartics, but are scarcely ever ordered. The Oil is mildly purgative, operating very quickly<sup>1</sup>, and with so little irritation, as to render it peculiarly fitted for cases in which the stimulating purgatives would prove hurtful; as in ileus, colica pictonum, (in which it may be advantageously joined with henbane,) calculous affections, piles, and after surgical operations in which the abdominal viscera are concerned. In dysentery, in which this Oil is particularly indicated, the stomach will seldom retain it; but when this is the case it may be efficaciously exhibited per anum. It is also an excellent purgative for infants, even of the tenderest age, and for women in child-bed.

The dose is from fʒiv to fʒjss, either floated on a little water, and covered with a small quantity of any ardent spirit; or diffused in a cup of coffee; or in cinnamon or mint water, by means of mucilage of acacia gum, or the yolk of egg. The addition of some aromatic tincture is generally necessary to make this oil remain on the stomach.

ROSA. *Spec. Plant. Willd.* ii. 1063.

Cl. 12. Ord. 5. Icosandria Pologynia. Nat. ord. Senticosæ, Linn. Rosaceæ, Juss.

G. 997. *Petals* five. *Calyx* pitcher-shaped, five-cleft, fleshy, contracted at the neck. *Seeds* numerous, hispid, affixed to the inner side of the calyx.

\* \* *With ovate germens.*

*Species* 15. *R. centifolia.* Hundred-leaved Rose. *Med. Bot.* 2d edit. 495. t. 178.

*Species* 16. *R. gallica.* Red Rose. *Med. Bot.* 2d edit. 497. t. 179.

*Species* 31. *R. canina.* Dog Rose, or Hep Tree. *Med. Bot.* 2d edit. 493. t. 177. *Smith, Flora Brit.* ii. 540. *Gærtner,* i. 347. t. 73.

### 1. ROSA CENTIFOLIA.<sup>2</sup>

*Officinal.* ROSÆ CENTIFOLIÆ PETALA, *Lond. Edin.* ROSA DAMASCENA; PETALA, *Dub.* The petals of the Hundred-leaved Rose.

*Syn.* Petales de Rose à cent feuilles (*F.*), Blumen der Blassen Rose (*G.*), Rosa de Alexandria (*S.*), Rozen (*Dutch*), Rosa (*Russ.*), Wurd (*Arab.*), Gooläbupoo (*Tam.*), Tu Miuhoa (*Chin.*), Gul (*Pers.*), Mawar (*Malay*), Sewooandamull (*Cyng.*), Goolat (*H.*).

Although this species of Rose be commonly cultivated in almost every garden in Europe, yet the place whence it was originally brought is still undetermined, notwithstanding the assertion of Herodotus that it grows wild in Macedon; and

<sup>1</sup> The bark of the root of the tree is a powerful purgative.

<sup>2</sup> Ρωδωνία Theophrasti.



that of Loureiro that it is a native of China. In the Dublin Pharmacopœia it is improperly confounded with the Damask Rose, which is altogether a different species.<sup>1</sup> The bush rises with prickly stems about three feet in height. The leaves consist of two or three pair of leaflets, with a terminal one, attached on very short petioles to a rough common footstalk: the leaflets are oval, broad, smooth, of a deep green colour on the upper surface, hairy on the under, and serrated, with purple edges. The flowers, which appear in June, are large, supported on peduncles beset with brown, bristly hairs. The segments of the calyx are semipinnate; the petals large, of a beautiful, pale-red colour, fragrant, and very numerous.

The varieties of this species of rose are very many; but for medicinal purposes they may be indiscriminately used. The petals only are employed.

*Qualities.*—Their odour is extremely fragrant, and their taste sweetish, subacidulous, and at last very slightly bitter. In distillation with water, a small portion of a butyraceous oil is obtained; and the water is strongly impregnated with the odour of the Rose.

*Medical properties and uses.*—The petals of this species of Rose are slightly laxative<sup>2</sup>; and as such are ordered, combined with sugar, in the form of a syrup, as an adjunct to oil and other purgatives in infantile diseases: but they are chiefly employed for the distillation of Rose-water.

Officinal preparations. *Aqua Rosæ*, L. E. D. *Syrupus Rosæ*, L. E.

## 2. ROSA GALLICA.

*Officinal.* ROSÆ GALLICÆ PETALA, *Lond. Edin.* ROSÆ RUBRÆ PETALA, *Dub.* The petals of the Red Rose.

*Syn.* Fleurs des Roses rouges (*F.*), Essig-rosen (*G.*), Fransche rozen (*Dutch*), Aettikerosor (*Swed.*), Rosa domestica (*I.*), Rosa rubra à Castillara (*S.*), Rosa Vermelha (*Port.*).

This species is a native of the south of Europe, cultivated in our gardens, and flowering in June and July. The stalks rise about three feet in height, are erect, and almost destitute of prickles. The foliage resembles that of the *centifolia*; but the leaflets are not so large, scarcely tomentose below, and subacute. The flowers also are less doubled; the petals large, widely spread open, of a deep, rich crimson colour, fragrant, and displaying an abundance of yellow anthers, on thread-like filaments; with the papillary stigmas of numerous, connected, villose styles, rising from the germen. The petals of the unblown buds are the parts medicinally used. They

<sup>1</sup> See *Hort. Kew. Willdenow*, &c. Gerarde cultivated this species in 1596.

<sup>2</sup> Illustres mulier mihi asserebat, ex rosarum olfactu se fœcum alvinarum odorem percipere. *Plenck, Icones*, &c. cent. iv.



are cultivated in abundance in the neighbourhood of London for medicinal purposes.

*Qualities*.—The odour of this Rose is less fragrant than that of the former species, but it is improved by drying: the taste is pleasantly bitter and austere. Water at 212° extracts both its odour and taste; and the infusion strikes a black with sulphate of iron; and also forms a precipitate of a dark colour with sulphate of zinc.

*Medical properties and uses*.—The Red Rose is astringent and tonic. It forms an elegant and useful vehicle for the exhibition of mineral acids, nitrate of potass, and other neutral salts, in hæmorrhages, and many other diseases. (See the Preparations of it.)

Official preparations. *Confectio Rosæ*, L. E. D. *Infusum Rosæ*, L. E. D. *Mel Rosæ*, L. D. *Syrupus Rosæ*, E.

### 3. ROSA CANINA.<sup>1</sup>

*Official*. ROSÆ CANINÆ PULPA, *Lond*. ROSÆ CANINA FRUCTUS, *Edin*. The pulp of the Dog-rose fruit, or Hep.

*Syn*. Le fruit d'Eglantier de Chien (*F.*), Die frucht der wilden rose, Hagebutter (*G.*), Le polpa di fruti di Rosa canina (*I.*).

This species is a common but beautiful ornament of our hedge-rows, flowering in June, and exhaling a very fragrant perfume. It rises to the height of eight or ten feet, has a smooth stem; with two alternate, compressed, conical-hooked, bright red, internodial prickles; and elongated branches spreading from the upright. The leaves are pinnate, composed of seven ovate, pointed, inodorous leaflets, naked and smooth on both sides, but the upper shining, and of a deeper green than the under. The petioles are pubescent, prickly, and glandular. The floral peduncles generally form a kind of corymb; but are sometimes solitary and smooth. The calyx is pubescent within, on the margin: the petals are inversely cordate, generally five, of a pale flesh colour, often white, and odorous. The fruit is an ovate, fleshy, smooth, red-berried calyx, with the apex sometimes open, sometimes shut, containing about thirty long angular seeds, embedded among white silky bristles.

*Qualities*.—The fruit is inodorous, and has a pleasant, sweet, acidulous taste, depending on uncombined citric acid and sugar, which it contains.

*Medical properties and uses*.—The pulpy part of heps is cooling, but possesses no direct medicinal properties. It is used only for the formation of the confection.<sup>2</sup>

<sup>1</sup> Κυνοςβατον Dioscoridis. Handsrose (*G.*), Hondсроос (*Dutch*), Hybentorn (*Dan.*), Niupon (*Swed.*), Rosa Silvestre (*S.*), Rosa brava (*Port.*), Schippownich (*Russ.*), Shora polna (*Pol.*), Gul (*Tatar.*), Foo son (*Japan*).

<sup>2</sup> The confection is sometimes brought to table as a sweetmeat on the continent: and the leaves of the plant are drank by the Tartars and the Russians in Siberia



Official preparation. *Confectio Rosæ caninæ*, L.

ROSMARINUS. *Spec. Plant. Willd.* i. 126.

Cl. 2. Ord. 1. Diandria Monogynia. Nat. ord. Verticillatæ, Linn. Labiatæ, Juss.

G. 62. Corolla unequal, with the upper lip two-parted. Filaments long, curved, simple, with a tooth.

Species 1. *R. officinalis*.<sup>1</sup> Official Rosemary. *Med. Bot.* 2d edit. 329. *Sibthorp Flora Græca*, t. 14.

Official. ROSMARINI CACUMINA, Lond. ROSMARINI OFFICINALISCA CUMINA, Edin. ROSMARINA HERBA, Dub. Rosemary tops.

Syn. Romarin (*F.*), Rosmarin (*G.*, *Dutch*, *Dan.*, *Swed.*), Romarino (*I.*), Romero (*S.*), Rosmaninho; Alcerim (*Port.*), Klil (*Arab.*), Yong tsao (*Chin.*).

This plant is a native of the south of Europe, Greece, and Barbary, but has been long cultivated in Britain, where it flowers in April and May. It is an evergreen, erect, very branching shrub, rising about four feet in height; the branches thickly covered with leaves, and the smaller ones four-cornered and downy. The leaves are opposite, almost sessile, more than an inch in length, and one-sixth of an inch broad, linear, obtuse, entire, with the margin turned back; of a dark-green colour, smooth, and shining on the upper side; woolly, veined, and of a silvery hue on the under. The flowers, which are placed on little axillary, opposite, leafy branches, are axillary and terminal, erect, on short stalks. The calyx is bell-shaped, bilabiate, villous; the upper lip entire, the under cloven into two pointed segments: the corolla is ringent, downy, pale blue, variegated with purple and white; the tube longer than the calyx, a little compressed, the upper lip erect and bifid, the lower cleft into three segments, the middle one larger, concave, and notched. The stamens are longer than the upper lip, arched, furnished with a tooth above the base, and supporting an oblong, blue, terminal anther. The style is the length of the stamens, thread-like, arched, and terminated by a simple sharp stigma. The seeds are four, oblong, and lodged in the bottom of the calyx.

*Qualities*.—Both the leaves and flowers have a grateful aromatic odour, and a bitterish, warm, pungent taste, depending on an essential oil, which appears to be combined with camphor: Vide *Ol. rosmarini*. Alcohol extracts its virtues completely, but they are only partially given out to water. By distillation with water, its essential oil is obtained. The leaves afford the greatest quantity: the flowers the smallest.

*Medical properties and uses*.—Rosemary is stimulant, and,

instead of tea. The Russians of the Volga prepare a spirit from the flowers. *Lin. Trans.* vol. xii. part i. p. 227.

<sup>1</sup> *Διςανωτισ* Dioscoridis.



according to some, emmenagogue.<sup>1</sup> It has been given in the form of infusion in nervous head-ach, hysteria, and chlorosis, but it is now scarcely ever prescribed, unless as an odorous adjunct to sternutatory powders. The dose in substance may be from grs. x. to ℥ij.; and from ʒj. to ʒiss. in infusion.

Officinal preparations. *Oleum Rosmarini*, L. E. D. *Spiritus Rosmarini*, L. E. D.

RUBIA. *Spec. Plant. Willd.* i. 603.

Cl. 1. Ord. 1. Tetrandria Monogynia. Nat. ord. Stellatæ, Linn. Rubiaceæ, Juss.

G. 187. Corolla of one petal, bell-shaped. Berries two, one-seeded.

Species 1. *R. tinctorum*.<sup>2</sup> Dyers' Madder. *Med. Bot.* 2d edit. 173. t. 67.

Officinal. RUBIÆ RADIX, *Lond. Dub.* RUBIÆ TINCTORUM RADIX, *Edin.* The root of Dyers' Madder.

Syn. Garance (*F.*), Krappwurzels; Færberrothe (*G.*), Mee (*Dutch*), Krap (*Dan.*, *Swed.*), Radiche di Robbia o Garanza (*I.*), Rudia (*S.*), Ruida (*Port.*), Munjith (*H.*), Manjittittie (*Tam.*), Marzana (*Pol.*), Mariona (*Russ.*), Kermesa Buja (*Turk.*).

This plant is a perennial, with annual stems. It is a native of the south of Europe, the Levant, and Africa, and has been cultivated to a very great extent for upwards of 300 years in Zealand: it flowers in June.<sup>3</sup> The root is composed of many long, thick, succulent fibres, about the thickness of a man's finger, united at the top in a head, from which go off many side-roots, extending under the surface of the ground, and throwing up shoots, by which the plant may be propagated. The stems are quadrangular, jointed, procumbent, and furnished with rough, short, hooked points, by which they are supported on the neighbouring plants. The leaves, which are in whorls of four or five, are elliptical, pointed, rough, and ciliated, about three inches long, nearly one broad in the middle, and having the midribs armed with the same kind of spines as on the stems. The branches bearing the flowers spring from the joints of the stems. The flowers are small, terminal, with a campanulate, yellow corolla, cut into four oval segments: the filaments short, supporting simple, erect anthers; and the germen is inferior and double, crowned with a slender style bearing two globular stigmas, and becoming two, round, black berries.

<sup>1</sup> *Bergius Mat. Med. a Regno Veget.* p. 21.

<sup>2</sup> Ερεθόδαρον Dioscoridis.

<sup>3</sup> As madder is an article of great national importance as a dye-stuff, many attempts have been made to cultivate it in this country, but without success, the Dutch madder being both better and cheaper than ours. That it can be grown to great perfection in this country is certain, and the effort to introduce its culture should not be dropped. The best comes from Zealand; to which Britain alone is said to have paid 200,000*l.* annually for madder. *Bancroft on Permanent Colours*, 2d edit. v. ii. p. 222.



Madder root is dug up for use in the third summer of its growth. It is then dried gradually in a stove built in the form of a tower, containing several floors: and from the uppermost it is progressively removed to the lowest; after which it is thrashed to remove the cuticle, and then dried completely in a kiln. When perfectly dried it is pounded, and finally packed in barrels for the market. There are three descriptions of this powder. The first pounding separates and reduces to a powder the fibrillæ and the skins of the larger roots only, which is sold at a low price under the name of *mull*; a second pounding separates one-third of the remaining parts of the large roots, and is sold under the name of *gem-cens*; and a third pounding forms into a powder the pure bright residue of the roots, which is the best, and is simply called *crop madder*.<sup>1</sup>

*Qualities*. — Madder has an unpleasant but not strong odour, and a bitter, slightly austere taste. It attracts the moisture of a damp atmosphere, and is injured by it. To water, alcohol, and volatile oils, at a temperature of 60°, it imparts a red colour; but to water at 212°, the colour imparted has a deep tinge of brown. Its principal constituent is extractive, which is precipitated by solution of alum brownish red: by the alkaline carbonates and lime-water, blood-red or lake; and by acetate of lead, brown<sup>2</sup>. The taste and odour of the madder are imparted to water and alcohol by infusion.

*Medical properties and uses*. — Madder is usually regarded as emmenagogue, and was formerly much relied on in chlorosis, and scanty and difficult menstruation. It has also been recommended in jaundice, and the atrophy of infants: but its efficacy in any disease is extremely problematical. Its colouring matter, however, is carried into the circulation, tinges the urine a blood-red colour, and is deposited in the bones.<sup>3</sup>

The dose of madder may be from grs. xv. to ℥j., united with sulphate of potass, and given three or four times a day.

RUMEX. *Spec. Plant. Willd.* ii. 249.

Cl. 6. Ord. 2. Hexandria Digynia. *Nat. ord.* Holoraceæ, *Linn.* Polygoneæ. *Juss.*

G. 699. *Calyx* three-leaved. *Petals* three, converging. *Seed* one, three-sided.

\*\* *Hermaphrodites*: with naked valves, or not marked with a grain.

*Species* 18. *R. aquaticus*. Great Water-dock. *Smith, Flora Brit.* 394. *Med. Bot.* 2d edit. t. 299.

<sup>1</sup> *Bancroft*, l. c.

<sup>2</sup> *Annales de Chimie*, iv. 104.

<sup>3</sup> Vide *Phil. Trans.* xxxix. 287—299. The leaves of the plant are said to tinge the milk of cows reddish, when eaten by them.



\*\*\* *With declinous flowers.*

*Species* 31. *R. acetosa*. Common Sorrel. *Med. Bot.* 2d edit. t. 230. *Smith, Flora Brit.* 396.

### 1. RUMEX AQUATICUS.<sup>1</sup>

*Officinal.*——; *RADIX, Dub.* The root of Water-dock.

*Syn.* La Patience aquatique (*F.*), Wasser Ampfer (*G.*), Waterpatich (*Dutch*), Rabaca major (*Port.*), Wodanoi Schawel (*Russ.*), Kong-ō-lik (*Esquimaux*).

Water-dock is an indigenous perennial plant, growing in ditches and on the banks of rivers; flowering in July and August. The root is thick. The stem rises about five feet in height, straight, furrowed, and smooth. The leaves are almost glaucous, lanceolate, and pointed, and the lower ones obcordate at the base. The flowers are in approximate whorls. They are nodding, on capillary pedicles, thickened at the apex. The valves are large, ovate, veined, entire, sometimes a little toothed, and all marked with a small linear, often obscure grain: the seed is large.

*Qualities.*—The root is nearly inodorous, and has a very austere taste. It yields its virtues to water.

*Medical properties and uses.*—Water-dock root is powerfully astringent. It was formerly celebrated under the name *Herba Britannica*, as a remedy for scurvy, and some cutaneous affections. It is now scarcely ever employed, although it undoubtedly possesses considerable powers in scurvy. Linnæus, in a letter to Dr. Lind, describing the scurvy of the Laplanders, asserts, that it is the only remedy which proved efficacious in that disease when “the ulcers are healed, and the patient is attacked with asthma.”<sup>2</sup> I have ascertained that a decoction of one ounce of the sliced root of the common dock, *Rumex patientia*, in a pint of water, is extremely efficacious in obstinate ichthyosis. In a full dose, about two ounces of the decoction, it purges freely; but at the same time improves the tone of the stomach.

### 2. RUMEX ACETOSA.

*Officinal.* ACETOSÆ FOLIA, *Lond.* RUMICIS ACETOSÆ FOLIA, *Edin.* Common Sorrel leaves.

*Syn.* Oseille ordinaire (*F.*), Sauer Ampfer (*G.*), Veldznuring (*Belg.*), Suramper (*Dan.*), Angsyra (*Swed.*), Szsaw (*Polish*), Acetosa (*I.*), Azedera (*S.*), Azedas (*Portug.*).

This is an indigenous perennial plant, common in pastures, and flowering in June. The stem is round, striated, and leafy, and rises from one to two feet in height. The leaves are oblong, ovate, and arrow-shaped; the radical ones petiolate and obtuse; and those of the stem sessile, amplexicaule, pointed, and a little rolled back. The flowers are dioecious, in

<sup>1</sup> Βρεταννική ή Βερτονική Dioscoridis.

<sup>2</sup> Correspondence of Linnæus, vol. ii. p. 476.



branched panicles, and arranged in half whorls: the calyx and corolla small: the stamens very short, bearing large, yellow anthers, and the styles short, with large, crimson, bearded stigmas. The valves are ovate, entire, and graniferous.

*Qualities*. — Sorrel leaves are inodorous, and have a grateful, austere, acidulous taste, depending on the presence of superoxalate of potass, which they contain.

*Medical properties and uses*. — These leaves are refrigerant and diuretic. Their expressed juice diluted with water, or a decoction of them in whey, affords a useful drink in cases of inflammatory fever; and eating them in large quantities daily as a salad, may prove serviceable in some cutaneous affections. In France, the plant is cultivated for the use of the table; and Dr. Clarke mentions that the natives of *Wermeland*, on the confines of Sweden, make it into bread in seasons of scarcity, and that it is not unsalutary.<sup>1</sup>

RUTA. *Spec. Plant. Willd.* ii. 542.

Cl. 10. Ord. 1. Decandria Monogynia. *Nat. ord.* Multisiliquæ, *Linn.* Rutaceæ, *Juss.*

G. 827. Calyx five-parted. Petals concave. Receptacle surrounded by ten melliferous points. Capsule lobed.

Species 1. *R. graveolens*.<sup>2</sup> Common Rue. *Med. Bot.* 2d edit. 437. t. 174.

*Officinal*. RUTÆ FOLIA, *Lond. Dub.* RUTÆ GRAVEOLENTIS HERBA, *Edin.* The leaves and herbaceous part of Rue.

*Syn.* Ruë sauvage (*F.*), Raute, Garten raute (*G.*), Ruite (*Dutch*), Rude (*Dan.*), Winruta (*Swed.*), Ruta (*I.*), Ruta de derpesado (*S.*), Ruda (*Port.*), Ruda (*Russ.*), Arooda (*Tam., Cyng.*), Sadsah (*Malay*), Inghoo (*Jav.*), Sendib (*Arab.*), Saturi (*H.*), Inghoo (*Javanese*).

Rue is an evergreen perennial, a native of the south of Europe, but much cultivated in our gardens, flowering in June and September. It rises to the height of two or three feet, shrubby and branching, with the lower part of the stems ligneous, and covered with a rough, striated, grey bark; but the upper branches are smooth, and of a pale-green colour. The leaves are doubly pinnate; the pinnæ distant; and the leaflets obovate, sessile, decurrent, and very obscurely crenate, with the terminal one generally notched; the surface punctured, the texture rather thick, and the colour blueish-green or glaucous. The flowers are produced in terminal branched corymbs on subdividing peduncles. The flower which opens first has a five-parted calyx, and a five-petalled corolla; but the others have the calyx four-parted only, and a four-petalled corolla. The petals are concave, wrinkled at the edge, of a pale, greenish-yellow colour and very much spread: the stamens are awl-shaped, the length of the petals, and bearing

<sup>1</sup> *Travels, &c.* Part iii. p. 90. 4to. Lond. 1825.

<sup>2</sup> Ρυτη-Πηλανον Dioscoridis.



small, yellow, quadrangular anthers.<sup>1</sup> The germen is large, oval, punctured, deep green, with crucial furrows, and crowned with a short style and simple stigma: and the seeds are angular, rough, and blackish.

*Qualities*. — Rue leaves have a powerful unpleasant odour, and a hot, bitter, nauseous taste. In the recent state, the leaves possess so much acrimony as to inflame and blister the skin; but much of this is dissipated in drying. In distillation with water, they yield a pungent volatile oil, on which their virtues chiefly depend: consequently, decoction is a bad form of preparation of Rue.

*Medical properties and uses*. — Rue is stimulant, and antispasmodic, and is supposed to possess emmenagogue powers. It was in high estimation so early as the time of Hippocrates, who frequently ordered it in female complaints.<sup>2</sup> In modern practice it is chiefly used in hysteria and flatulent colic. I have found a strong infusion of it, exhibited per anum, of great service in relieving the convulsions of infants, arising from flatulence and other intestinal irritations. It may however inflame the mucous coat of the intestines, and therefore should be used with caution. The dose of the powdered leaves is from grs. xv. to ℥ij., given twice or three times a day.

Official preparations. *Oleum Rutæ*, D. *Extractum Rutæ graveolentis*, E. D.

SABINÆ FOLIA. Vide *Juniperus*.

SACCHARUM. *Spec. Plant. Willd.* i. 122.

Cl. 3. Ord. 2. Triandria Digynia. *Nat. ord.* Gramina.

G. 122. *Calyx* two-valved, involucred, with a long lanugo. *Corolla* two-valved.

*Species* 4. *S. officinarum*.<sup>3</sup> Common Sugar-Cane. *Sloane's Jamaica*, i. 101. t. 66. *Phil. Trans.* lxi. 207—278. t. 3.

*Official*. SACCHARUM. SACCHARUM PURIFICATUM, *Lond.* SACCHARUM, a. *non purificatum*; b. *purificatum*; c. *syrupus empyreumaticus*, *Edin.* SACCHARUM PURIFICATUM; SACCHARUM RUBRUM, EJUSDEMQUE SYRUPUS, (Molasses,) *Dub.* Unrefined Sugar. Refined Sugar. Molasses.

*Syn.* Sucre; Sucre-pur (*F.*), Zucker; Weisser Zucker (*G.*), Suiker (*Dutch*), Sukker (*Dan.*), Socker (*Swed.*); Zucchero brutto: Zucchero in pane; Melassa (*I.*), Azucar; Atriaca (*S.*), Assucar (*Port.*), Shukhir (*Arab.*), Chenee (*H.*), Sakkari (*San.*), Sakkarei (*Tam.*), Shukker (*Pers.*), Soola (*Malay*).

<sup>1</sup> These stamens display in a striking manner the spontaneous motions which take place in some plants. They are very stiff, and cannot be disturbed from the posture in which they happen to be; but nevertheless they rise, by a spontaneous movement, one or two at a time, and lean over the stigma till the pollen be shed, when they fall back again, and give place to others.

<sup>2</sup> *De Morbis Mulier.*

<sup>3</sup> Cannamelle (*F.*), Zuckerrohr (*G.*), Suikerriet (*Dutch*), Cannamele (*I.*), Cana de azucar (*S.*), Quasab (*Arab.*), Can che (*Chin.*).



The Common Sugar-cane is a native of both the East and West Indies. It is cultivated in Persia, and very abundantly in the West Indies. There are two varieties of the officinal sugar-canes; the Creole and the Otaheite. They are distinguished chiefly by the colour of the leaves; those of the former being much deeper than those of the latter. The root is jointed, and sends up several jointed stems, which rise in general to the height of eight or ten feet. A leaf springs from each joint, and the base of it embraces the stem to the next joint above its insertion, before it expands. From this point each leaf is about three or four feet long, and comparatively narrow, like a blade of grass; with the midrib broad, and prominent on the under side, and the edges thin and sharply toothed. The flowers are in terminal panicles, two or three feet in length, and composed of subdivided spikes, with long flexuose down or lanugo, which incloses the flowers and hides them from the sight. The seed is oblong-pointed, and ripens in the valves of the flowers.

Although the sugar-cane is undoubtedly a native of the American continent and its islands, yet the culture of it, and the art of making sugar, were carried from Spain to the Canary Islands<sup>1</sup>, and thence extended, about the end of the fifteenth century, to the West Indies and the Brazils; the former of which supplies the greater part of the consumption of Europe<sup>2</sup>, a small proportion only being brought from the East Indies. The quantity of sugar yielded by the plant is varied by climate: thus the Otaheitean cane contains one third more crystallizable matter than the Cane of any other place.<sup>3</sup>

In the West Indies, the plant is propagated by cuttings of the stalk taken from near its top, and laid horizontally in the ground. It requires a rich, permeable soil; and from fourteen to seventeen months before it ripens; good land will furnish five crops of shoots without transplanting. The canes are cut for the purpose of making sugar, between the sixth and thirteenth month of their growth, when the stems have acquired from seven to ten feet in height, a proportionable size, and

---

<sup>1</sup> At one time, sugar was the staple commodity of Maderia, although there is now one sugar-mill only on the island; but the sugar is uncommonly fine, and has an agreeable odour, not unlike that of violets. Sprengel says, the first notice of the sugar-cane is found in the Itinerary of Abusaidi, in which it stated that it grows at Siraf; and Abulfed says, it grows spontaneously at Almansura, in India. Ebu Alvan first described the mode of collecting and preparing the juice.

<sup>2</sup> The average importation into England and Scotland between 1787 and 1790, amounted annually to 1,952,262 cwt. *Mosley's Hist. of Sugar*, p. 154.

<sup>3</sup> The Batavian which is less productive, is undoubtedly a distinct species. The foliage is purple, and very broad.



the cuticle appears smooth, dry, and brittle. This generally happens in the months of February, March, and April. As soon as they are cut, the canes are stripped of their leaves, and immediately crushed between iron rollers, to express the juice, which is received into large leaden vessels, called *receivers*, whence it is conveyed into a large copper vessel, named the *clarifier*, where it is mixed with lime, in the proportion of one pint to 100 gallons of juice, and heated to the temperature of  $140^{\circ}$ .<sup>1</sup> A thick scum soon forms on the top, from under which the clear liquor is drawn off by a cock into a large copper boiler, where it is boiled till the bulk of the liquor is very considerably diminished. The boiling is successively repeated in four other coppers progressively smaller; and from the last, which is called the *teache*, it is conveyed into shallow wooden coolers, where it grains, and the concreted mass separates from the uncrystallizable matter or molasses. This mass is then put into empty hogsheads, having holes in the bottom, through each of which the stalk of a plantain-leaf is thrust; and when the molasses has drained off, the process is finished. In this state the sugar is brought home, under the name of *raw*, or *muscovado* sugar. In Europe, however, sugar undergoes another process for its purification. It is coarsely ground, dissolved in lime-water, and clarified with bullocks' blood; then boiled down to a proper consistency, the impurities being skimmed off as they rise, and poured into conical earthen vessels, where it is allowed to grain. The point of the cone is perforated; and the base covered with moist clay, the moisture of which percolates the sugar, and runs off through the perforated apex, which is placed undermost, carrying with it any uncrystallized impure syrup. In this state it is called *Loaf* sugar; and requires a second purification before it is considered as completely *Refined* sugar.

*Qualities*.—*Raw* or *Muscovado* sugar is inodorous, and sweet to the taste. It is in concreted masses, consisting of small, dry, sparkling, irregular crystals of a yellowish colour. *Refined* sugar is also inodorous, and sweet to the taste. Its colour is pure white; and the mass or loaf in which it is concreted should be hard, extremely brittle, pulverulent, and persistent in the air. One hundred parts of sugar in its ordinary state contains, according to Berzelius, 5·3 of water: and it requires its own weight only of water at  $48^{\circ}$  for its solution. When united at a higher temperature with a smaller quantity of water, it remains dissolved, forming syrup. Four

---

<sup>1</sup> The lime extricates carbonic acid from the juice, and forms with the *herbaceous* or *feculent* matter an insoluble compound, which rises to the surface, and forms the scum.



parts of boiling alcohol dissolve one part of sugar; but by rest a moiety of the sugar again separates in crystals. Oils also readily combine with it, and the mixture is miscible with water. Lime and the fixed alkalies unite with sugar, and form compounds, without any sweetness of taste. The concentrated strong acids dissolve and decompose sugar, but the weaker simply dissolve it; and the alkaline and earthy hydrosulphurets, sulphurets and phosphurets, decompose it, and resolve it into a substance resembling gum.<sup>1</sup> “When sugar is boiled,” says Vogel, “with peroxide of mercury and acetate of copper, these salts are converted into protoxides: oxy-muriate of mercury is converted into calomel; and sulphate of copper and nitrate of mercury are reduced to a metallic state. But sugar does not decompose the salts of iron, zinc, tin, and manganese.” Its ultimate constituents, according to the experiments of Lavoisier, are, sixty-four oxygen, twenty-eight carbon, and eight hydrogen, in 100 parts<sup>2</sup>: but according to Thenard and Gay Lussac, the proportions are oxygen 50.63, carbon 42.47, hydrogen 6.90, with which the last analysis of Berzelius nearly coincides.

*Molasses* has a peculiar odour, and a sweet empyreumatic teste. It is of a brown or black colour, thick, and viscid; and is constituted chiefly of the uncrystallizable part of the juice of the sugar cane, which Proust has denominated liquid sugar. It is more soluble in alcohol than sugar.

*Medical properties and uses.*—Raw sugar and molasses are laxative; and refined sugar externally applied is escharotic. All the kinds are extremely nutrient, and more generally used as articles of diet than for medicinal purposes; except it be to cover the taste of nauseous drugs. Sugar, however, is said to be a preventive of worms; and to prove useful in scurvy; but it is hurtful to those of bilious, hypochondriacal, and dyspeptic habit. Milk boiled with fine sugar will keep good for a considerable time. But perhaps the most important use of sugar is as an antidote to the poison of verdegriis. It requires to be given in large quantities, both in the solid form and in solution of water. It appears to act chemically on the poison, and also by increasing the peristaltic action of the bowels.<sup>3</sup>

Officinal preparations. *Syrupi omnes*, L. E. D. *Trochisci omnes*, E. *Confectiones omnes*, L.

**SAGAPENUM.** *Lond. Edin. Dub.* *Sagapenum*.

*Syn.* *Sagapenum* (F.), *Sagapengummi* (G.), *Sagapeno* (I.), *Sugbeenuj* (Arab.), *Kundel* (H., Sans.).

<sup>1</sup> *Rollo on Diabetes*, 452. *Thomson's Chemistry*, 4th ed. vol. iv. 660.

<sup>2</sup> A sugar in every respect resembling common sugar is obtained from the maple.

<sup>3</sup> Vide *Traité des Poisons*, &c. Par M. P. Orfila, tom. i. p. 281.



This gum-resin, which is brought to this country from Smyrna, Aleppo, and Alexandria, is the concrete juice of an unknown Persian plant. Dioscorides mentioned it as the juice of a ferula growing in Media<sup>1</sup>; and nothing more is known of its source at this day, although Willdenow supposes it to be the *Ferula Persica*.<sup>2</sup>

*Qualities*. — Sagapenum has an alliaceous odour, and a hot, acrid, bitterish taste, not unlike that of assafoetida, only weaker. It is in agglutinated drops or masses, of an olive, or brownish-yellow colour, slightly translucent, and breaking with a horny fracture. It softens and is tenacious between the fingers; melts at a low heat, and burns with a crackling noise and white flame, giving out abundance of smoke, and leaving behind a light, spongy charcoal. Water and strong alcohol dissolve it partially; but it is almost completely soluble in proof spirit. In distillation with water it yields a little volatile oil; and impregnates the water strongly with its flavour. Its constituents, according to Pelletier, are 31.94 of gum, 54.26 of resin, 0.40 acidulous malate of lime, 11.80 of volatile oil, and 0.60 of peculiar matter<sup>3</sup>, on which its virtues probably depend.

*Medical properties and uses*. — This Gum-resin is antispasmodic, and emmenagogue; and externally discutient. It is sometimes employed in hysteria, chlorosis, and other cases in which assafoetida has been found serviceable; but it is much inferior in its powers. It is usually given in substance, in doses of from grs. x. to 3ss. made into pills.

SALIX. *Spec. Plant. Willd.* iv. 703.

Cl. 22. Ord. 2. Diœcia Diandria. *Nat. ord.* Amentaceæ.

G. 1756. Male. *Amentum* cylindrical. *Calyx* a scale. *Corolla* none. *Gland of the base* nectariferous.

———— Female. *Amentum* cylindrical. *Calyx* a scale. *Corolla* none. *Style* bifid. *Capsule* one-celled, two-valved. *Seeds* downy.

\* *With smooth serrated leaves*.

*Species* 10. *S. fragilis*. Crack Willow. *Smith, Flora Brit.* 1051. *Med. Bot.* 2d edit. 13. t. 8. *Hoffman Sal.* ii. 9. t. 31.

\*\*\* *With villose leaves*.

*Species* 33. *S. alba*. White Willow. *Smith Flor. Brit.* 1071. *Hoffman Sal.* i. 41. t. 7, 8.

*Species* 101. *S. caprea*. Great round-leaved Sallow. *Smith, Flora Brit.* 1067. *Hoffman Sal.* i. 25. t. 3. f. 1.

#### 1. SALIX FRAGILIS.

*Officinal*. — CORTEX, *Dub.* Bark of the Crack Willow.

<sup>1</sup> *Dioscoridis*, lib. 3. c. 95. (Σαγαπηνον.)

<sup>2</sup> This plant was fully described by Dr. Hope, as the plant which yields the assafoetida, which, however, is the produce of another species. See *Phil. Trans.* lxxv. 36. t. 3, 4.

<sup>3</sup> *Journ. de Pharm.* tom. 3.



*Syn.* Ecorce de Saule (*F.*), Weidenrinde (*G.*), Corteccia di Salcio (*I.*), Cor-teza de sauce (*S.*).

This species of Willow is indigenous, growing upon the banks of rivers, and flowering in April and May. It grows to a considerable height, sending off upright branches; which are covered with an even brownish-yellow bark, and are very fragile at the base. The leaves are petiolate, from three to five inches in length, lanceolate, pointed, obtusely serrated, inflected and glandular; smooth on both surfaces, shining on the upper; and, in the younger ones, ciliated at the apex. There are sometimes no stipules; but, when present, they are rounded and obscurely toothed. The male catkin is pale, cylindrical, rather lax, with ovate, downy scales. The nectary is composed of two yellow glandular scales, the larger between the stamen and the receptacle, and the smaller between the stamen and the scale. The stamens are two, filiform, and smooth. The female catkin resembles the male; with the germen egg-shaped, supporting two bifid, erect stigmas. The capsule is ovate, and contains many small seeds. The bark requires to be dried in an oven moderately heated.

*Qualities.*—The dried bark is inodorous, bitter, and austere.

## 2. *SALIX ALBA*.<sup>1</sup>

*Officinal.* *SALIX CORTEX*, *Dub.* Willow Bark.

*Syn.* Saule (*F.*), Weide (*G.*), Wilg (*Dutch*), Piil (*Dan.*), Pihl (*Swed.*), Koro Wierzbowa (*Pol.*), Salice (*I.*), Sauce (*S.*), Salgueiro (*Port.*).

The White Willow is indigenous, growing in woods and moist places, and flowering in April and May. It is a large tree, with a cracked bark; and furnished with many round, spreading branches; the younger of which are silky. The leaves are alternate, on short petioles, lanceolate, pointed, acutely and regularly serrated, with the lower serratures remote and glandular: pubescent on both sides, and silky beneath: the younger ones are altogether silvery and convoluted. There are no stipules. The catkins are terminal, cylindrical, elongated, slender and many-flowered, with elliptical, lanceolate, brown, pubescent scales. The stamens are yellow, and a little longer than the scales: the style is short; and the stigmas bipartite and thick. The capsules are nearly sessile, ovate, brownish, and smooth.

The bark of this species is easily separated all the summer. It has been used for tanning leather; and the inner part of it affords the miserable inhabitants of Kamschatka a substitute for bread.

*Qualities.*—The same as those of the former species.

<sup>1</sup> Ἰτέα λευκή Theophrasti.



3. SALIX CAPREA.<sup>1</sup>

*Officinal.* SALICIS CORTEX, *Lond.* Willow Bark.

This species of Willow is indigenous, very common in woods, flowering in April. It is a middling-sized tree, with the branches round, even, shining, and brownish; and the shoots pubescent. The leaves are alternate, petiolate, and varying in shape, being sometimes elliptical or roundish, pointed, large, undulated, waved, or serrated; smooth and dark green on the upper surface, and densely tomentose and veined on the under. The stipules are crescent-shaped or roundish, recurved, waved, and tomentose. The petioles are linear, and densely villose. The catkins appear before the leaves, are ovate, thick, many-flowered, with obovate, very hairy scales. The stamens are yellow: the stigmas nearly sessile, undivided, but at last occasionally cleft. The capsules are pedicelled, ovate, bellied at the base, and downy.

*Qualities.* — The bark of this species, like that of the two former, is inodorous, bitterish and astringent.

The bark of the White Willow only has been chemically examined; but as the other two species agree with it in their sensible qualities, it is probable that they agree also in other respects. Water extracts its virtues, and affords a decoction of a reddish colour, which is precipitated by a solution of isinglass, the carbonates of potass and of ammonia, and by lime-water, which throws down a precipitate, at first blue, and afterwards buff-coloured: sulphate of iron, also, produces a dark green precipitate. The watery extract is reddish, brittle, has a bitter taste, and does not deliquesce. Digested in alcohol this bark affords a greenish-yellow tincture, which water renders turbid. When evaporated, the extract is of a bright-yellow colour, bitter, melts at a moderate heat and emits an aromatic odour.<sup>2</sup> The constituents, therefore, of White Willow bark, and probably of the two other species also, are tannin, bitter resin, extractive, and gluten.

*Medical properties and uses.* — These barks are tonic and astringent. They have been given as a substitute for the Cinchona bark; and in some cases intermittents and remittents have yielded to their use.<sup>3</sup> They have also been efficaciously administered in cases of debility, dyspepsia, and pulmonary

<sup>1</sup> *Siler Virgilii*.

<sup>2</sup> *Ann. de Chimie*, liv. 290. *Thomson's Chemistry*, 4th ed. vol. v. p. 221.

<sup>3</sup> The bark of the white Willow was first used by the Rev. Edmund Stone, of Chipping Norton, Oxfordshire. He gave it successfully in doses of one drachm of the powder every hour between the paroxysms, in tertians, and added one-fifth of Peruvian bark to augment its power, in obstinate quartans. *Phil. Trans.* iii. 195.



hemorrhagies; and have apparently been more serviceable in phthisis and hectic fever than the Cinchona. They may be given either in substance, or in the form of decoction. Of the powdered bark from ʒss. to ʒj. may be given for a dose, combined with aromatics, myrrh, or the Cinchona bark, as circumstances direct.

**SALVIA.** *Spec. Plant. Willd.* i. 127.

*Cl.* 2. *Ord.* 1. Diandria Monogynia. *Nat. ord.* Verticillatæ, *Linn.* Labiatæ, *Juss.*

*G.* 63. *Corolla* unequal. *Filaments* affixed transversely to a pedicel.

*Species* 7. *S. officinalis*.<sup>1</sup> Garden Sage. *Med. Bot.* 2d edit. 352. *t.* 127.

*Officinal.* SALVIÆ OFFICINALIS FOLIA, *Edin.* SALVIA, *Dub.*  
The leaves of Sage.

*Syn.* Sauge (*F.*), Salbei (*G.*), Salei (*Dutch*), Salvie (*Dan.*), Salvia (*Swed.*), Szalwia (*Pol.*), Salvia (*I.*), Salvia (*S.*), Salva (*Port.*), Schalweja (*Russ.*), Sayseley (*Tam.*), Saulbēy (*Pers.*).

The common or officinal Sage is a perennial plant, a native of the south of Europe, cultivated abundantly in our gardens, flowering in June. It rises about two feet in height, with a quadrangular, shrubby, branching stem. The younger branches are whitish, and downy. The leaves, which stand in pairs on footstalks, are ovate-lanceolate, wrinkled, crenate, and sometimes tinged reddish or purple. The flowers are on long terminal spikes, in six-flowered distant whorls, accompanied with ovate, acute, deciduous bractes. The calyx is striated, purplish on the upper part, and notched into three acute teeth above, and two below: the corolla is tubular and bilabiate, of a beautiful blue, variegated with purple and white; the upper lip obtuse, notched, and concave, the under three-lobed, the lateral lobes bent backwards. The filaments are affixed transversely at their middle to short pedicels, on a moveable axis, and are curved threads bearing a gland on the lower end, and on the upper a yellow, oblong anther: the style is long, curved, purple with a bifid stigma, and rising from the centre of four naked seeds in the bottom of the calyx.

There are many varieties of common Sage, but their properties are the same. It is cut when in flower, and hung up in a shady place to dry.

*Qualities.* — The odour of Sage is fragrant, and the taste warm, bitterish, and aromatic: qualities depending on an essential oil, which can be obtained separate in distillation with water. Sulphate of iron strikes a deep black colour with the infusion.

<sup>1</sup> Ελαιωφάκον Dioscoridis.



*Medical properties and uses.* — Sage is stimulant, carminative and slightly astringent. The estimation in which it was held by the ancients is sufficiently well known; but it does not support the character it formerly acquired; and “*salvia salvatrix naturæ conciliatrix*”<sup>1</sup> is very little regarded by the modern practitioner. Infusions of the leaves, if strained before too much of the bitter is extracted, prove very grateful to the stomach, when nausea is troublesome in febrile complaints; and when drunk cold they are said to check hectic perspiration<sup>2</sup>, and those which frequently attend convalescencies. The infusion either alone, or mixed with honey and vinegar, is a well known gargle in cases of sore throat, and relaxation of the uvula. The dose of the pulverized leaves is from grs. xv. ʒss.; or, of an infusion made with ʒj. of the dried leaves and Oj. of boiling water, fʒij. may be taken every three or four hours.

SAMBUCUS. *Spec. Plant. Willd.* i. 1494.

Cl. 5. Ord. 3. Pentandria Trigynia. *Nat. ord.* Dumosæ, *Linn.* Caprifoliæ, *Juss.*

G. 569. *Calyx* five-parted. *Corolla* five-cleft. *Berry* three-seeded.

*Species* 3. *S. nigra*.<sup>3</sup> Common Elder. *Med. Bot.* 2d edit. 596. *Smith, Flora Brit.* 336. *Eng. Bot.* 476.

*Officinal.* SAMBUCI FLORES, *Lond.* SAMBUCI NIGRÆ FLORES; BACCÆ CORTEX, *Edin.* SAMBUCUS; CORTEX INTERIOR, FLORES, BACCÆ, *Dub.* The flowers, berries, and inner bark of Common Elder.

*Syn.* Sureau ordinaire (*F.*), Fliederblumen (*G.*), Coortecchia; bacche, e fiori di Sambuco (*I.*), Sabuco (*S.*), Uktee (*Arab.*).

The Common Elder is a very abundant, indigenous, middle-sized shrubby tree, growing commonly in hedges; flowering in June, and ripening its berries in September. It is much branched near the top, and covered with a roughish, gray bark. The wood is white, hard, and has a large, spongy pith. The leaves are pinnated, composed of five oval, pointed, serrated leaflets, nearly equal at their base. The flowers are in terminal cymes, consisting of five, principal branches, and many small ones, with some of the flowers sessile. They are cream-coloured: with the calyx superior, and permanent, and the corolla monopetalous, rotate, and somewhat convex. The berries are globular, and when ripe of a purplish-black colour.

*Qualities.* — The flowers have a peculiar, faint, sickly odour, and bitterish taste, which are imparted to water by infusion, and also by distillation, during which a small portion of buty-

<sup>1</sup> *Schola Salernitana*, c. 38. p. 406.

<sup>2</sup> *Van Swieten's Comment.* ii. 370.

<sup>3</sup> *Ἀκτὴ* Dioscoridis. The leaves laid in the subterraneous passages of moles are said to drive them away.



raceous oil is separated. The *berries* are inodorous, have a sweetish taste; and yield on expression a fine purple juice<sup>1</sup>, which contains saccharine matter, jelly, and the malic acid. The *inner bark* is inodorous, and has a faint, sweetish taste, which is succeeded by a slight bitterness, and a very permanent acrimony. Both water and alcohol extract their virtues.

*Medical properties and uses.* — The flowers and berries are diaphoretic and aperient. The berries were formerly much used in febrile diseases, rheumatism, gout, and eruptive diseases, but they are now scarcely ever ordered. The flowers are used chiefly in fomentations and cooling ointments; and to afford their odour to water in distillation. The bark is a hydragogue purgative, and in large doses proves emetic at the same time. It is said to prove efficacious in dropsy; and in smaller doses to be a useful aperient and deobstruent in various chronic affections. The dose of the bark is from grs. x. to ʒss., given in wine: or ʒj. may be boiled in Oij. of milk or of water down to Oj, and the fourth part taken for a dose.

*Officinal preparations.* *Succus spissatus Sambuci Nigræ*, E. D. *Unguentum Sambuci*, L. D.

#### SAPO. Soap.<sup>2</sup>

Soap is a compound of margaritic and oleic acids<sup>3</sup> with an alkaline, or an earthy, or an oxidized metalline base. The first kind is that which is employed in medicine, and has been longest known, having been invented by the Gauls at a period antecedent to historical record. Alkaline soap is of two kinds; one made with soda, and oil either animal or vegetable, or tallow, and called *Hard Soap*; the other made with potass and similar oily matters, and called *Soft Soap*. For medical purposes it is essential that both kinds be made from the purest materials; and therefore the soap made in countries which produce olive oil, as the south of France, Italy, Tripoli, and Spain, is preferable to the soap of this country, which is generally manufactured from grease, tallow, and other kinds of fat.

##### 1. HARD SOAP.

*Officinal.* SAPO DURUS, *Lond.* SAPO DURUS; *ex soda confectus*, *Edin.* SAPO; DURUS HISPANICUS, *Dub.* Hard Soap. Spanish Soap.

<sup>1</sup> M. A. Chevalier has ascertained that paper stained with this juice is as delicate a test of the presence of alkalies and acids, as litmus paper. *Journ. de Pharm.* Avril 1820.

<sup>2</sup> The name is derived, according to Beckman, from the old German word *Sepe*. — *History of Inventions*, iii. 239.

<sup>3</sup> *Chevrue*, whose experiments have elucidated the nature of soap more than those of any other chemist, has ascertained that fixed oils and tallow consist of two substances, one solid, which he has named *steräin*, and the other fluid, which he has named *clain*. These are altered by salifiable bases, and converted in the above named acids, vide *Ann. de Chimie et de Phys.*



*Syn.* Savon blanc (*F.*), Spanische Seife (*G.*), Spaansche zeep (*Dutch*), Silkestwal (*Dan.*), Sapone duro (*I.*), Xabon (*S.*), Nāt Sowcārum (*Tam.*), Sāboon (*Duh.*).

Hard Soap is manufactured in Spain in the following manner: To five parts of barilla, coarsely ground to powder, one part of quicklime rendered fluid with a small portion of water, is added; and after some time the clear liquor, which is a strong solution of caustic soda, is drawn off, and called the *first ley*; with the residue more water is then mixed, and drawn off after some time, and called the *second ley*; and a *third ley* is procured by another portion of water treated in a similar manner. This last ley is then mixed with a quantity of olive oil equal in weight to the barilla employed, and the mixture boiled in an iron vessel, the second ley and a portion of the first being added in a gradual manner during the boiling. The boiling mixture is constantly stirred with a wooden pole, and when it becomes tolerably thick, a small portion of common salt is added, and the boiling continued for half an hour. The fire is then damped, and after some hours the clear liquor, which has separated, is drawn off, and the half-made soap again boiled with a little fresh water and the residue of the first ley. After the separation of the fluid of this boiling, it is again heated with a little water, and then poured into wooden vessels called frames, where it cools, and in a few days acquires a sufficient degree of hardness. Three parts of oil and three parts of soda produce five parts of firm Soap.<sup>1</sup> Castile Soap is made in the same manner, except that the marbled appearance which it presents is produced by the addition of sulphate of iron to a part of the alkaline ley, after the Soap is fully boiled, which gives the blue colour; and the stirring in red oxide of iron, when the Soap is almost made, gives the red colour.

*Qualities.*—Well made Hard Soap, fit for medical use, has very little odour, and a nauseous, alkalescent taste; is white, and of a firm consistence; does not feel greasy, and is devoid of any saline efflorescence on the surface. With water it forms a milky, opaque solution; and with alcohol a nearly transparent, somewhat gelatinous solution.<sup>2</sup> It is decomposed by all the acids, and acidulous salts; by alum, the muriate and the sulphate of lime, and sulphate of magnesia; thence hard water, which contains sulphate of lime, does not properly dissolve Soap. Nitrate of silver; ammoniated copper; tincture of muriated iron; ammoniated iron; acetate, submuriate and oxymuriate of mercury; superacetate of lead; tartarized iron;

<sup>1</sup> *Annales de Chimie*, xix. 253.

<sup>2</sup> The alcoholic solution of soap is a convenient test for discovering earthy salts in mineral waters.



tartarized antimony; sulphates of zinc, copper, and iron; and all astringent vegetable solutions also decompose it. According to the experiments of Darcet, Lelievre, and Pelletier, 100 parts of newly made soap consists of 60·94 oil, 8·56 alkali, and 30·50 water: but part of the water is lost by keeping, and the soap becomes lighter.

## 2. SOFT SOAP.

*Officinal.* SAPO MOLLIS, *Lond. Edin.* Soft Soap.

*Syn.* Savon Moû (*F.*), Sapone Molle (*I.*).

This Soap is prepared in the same manner as the former: a caustic ley of potass, however, being used instead of the soda ley. It was this variety of Soap which was originally made by the Gauls and Germans, who employed wood ashes to afford their ley; and these are still used in many places.

*Qualities.* — Soft Soap differs from Hard Soap chiefly in its consistence, which is never greater than that of hog's lard.

*Medical properties and uses.* — Soap is regarded as purgative and lithontriptic; externally applied it is stimulant and detergent. For internal use the Hard Soap only is employed. It is occasionally ordered in habitual costiveness, and in jaundice, combined with rhubarb, or some bitter extract; but its power as a purgative is very limited, and it cannot act in any other way in relieving jaundice. It is more useful in calculous habits, in which, however, its action is altogether confined to the stomach; for as Soap is decomposed by the weakest acids, its alkaline base corrects the acidity so prevalent in the stomachs of calculous patients, and thus, at least, assists in checking the increase of the disease. Soap is also beneficial in decomposing some metallic poisons when taken into the stomach; and, as it is the antidote which can most readily be procured, should always be early resorted to. It is necessary, in this latter case, to give it in solution; of which, a tea-cupful should be drank at short intervals, till the effects expected from it be produced. In other cases it is preferable to give it in substance. As an external remedy, Soap is efficaciously used in frictions to sprains and bruises; and we have seen much benefit derived from rubbing the tumid bellies of children labouring under mesenteric fever, with a strong lather of Soap every morning and evening. The dose internally is from grs. iij. to ʒss. made into pills.

*Officinal preparations.* *Pilulæ Saponis cum Opio*, L. *Pil. Scillæ comp.*, L. *Pil. Aloet.*, E. *Pil. Aloes et Assafœtidæ*, E. *Pil. Aloes cum Zinzibere*, D. *Pil. Colocynth comp.* D. *Emplastrum Saponis*, L. E. *Ceratum Saponis*, L. *Linimentum Saponis comp.*, L. *Linimentum Saponis cum Opio*, E.

SARSAPARIL/LÆ RA'DIX. Vide *Smilax*.

SAS'SAFRAS LIG'NUM ET RA'DIX. Vide *Laurus*.



SCAMMONIÆ GUMMI RESINA. Vide *Convolvulus*.

SCILLA. *Spec. Plant. Willd.* ii. 125.

Cl. 6. Ord. 1. Hexandria Monogynia. Nat. Ord. Coronariæ, Linn. Asphodeli, Juss.

G. 640. Corolla six-petalled, spreading, deciduous. Filaments threadlike.

Species 1. *S. Maritima*.<sup>1</sup> Officinal Squill. *Med. Bot.* 2d edit. 745. t. 255.

Officinal. SCILLÆ RADIX, *Lond. Dub.* SCILLÆ MARITIMÆ RADIX<sup>2</sup>, *Edin.* Squill root (bulb).

Syn. Scille (*F.*), Meerzwiebel (*G.*), Zeeajuin (*Dutch*), Skille (*Dan.*), Siölok (*Swed.*), Scilla (*I.*), Cebolla abarruna (*S.*), Albarra (*Port.*), Alvazraä (*Port.*).

This species of Squill is a native of Spain, Sicily, Syria<sup>3</sup>, and Barbary, flowering in April and May. The bulb is large, sometimes nearly the size of the human head, of a pear shape, and formed of fleshy scales, attenuated at both edges, and closely applied one over the other. The roots are fibrous attached to a radical plate at the bottom of the bulb. The stem is round, smooth, and succulent, rising about three feet in height, from the centre of several radical, sword-shaped, straight, pointed, long leaves of a deep green colour. The flowers are produced in a long close spike upon purplish peduncles, with a linear, twisted, deciduous bracte at the base of each peduncle. The corolla consists of six, white, ovate, spreading petals, with a reddish mark in the middle of each: the filaments are shorter than the corolla, tapering, and furnished with oblong, transversely-placed anthers: the germen is roundish; the style and stigma are simple; and the capsule is oblong, smooth, three-celled, and contains many roundish seeds.

There are two varieties of the officinal Squill, one with a white bulb, and the other with a reddish bulb; but both are indiscriminately used, and do not differ in their virtues. The bulbs are brought from the Levant generally in bulk. They are preserved fresh in sand; but as they are apt to spoil, it is preferable to keep them in the dried state. (See *Preparations*.)

*Qualities.* — The Squill bulb is inodorous; its taste is bitter, nauseous, and acrid; and when much handled it inflames and ulcerates the skin. The expressed juice slightly reddens litmus paper. The acrimony on which its virtue depends is partially dissipated by drying and long keeping, and completely destroyed by any heat above 212°: it is extracted by water,

<sup>1</sup> Σκίλλα Dioscoridis. The trivial name *maritima* has been objected to, as it does not generally grow on the sea coast.

<sup>2</sup> All the colleges have erred in designating the root as the officinal part of the squill. The bulb is the part employed.

<sup>3</sup> The soil at Navarino, which is remarkable for the production of an infinite quantity of squills, “is of a red colour.”—*Gell's Journ. in the Morea*, p. 21.



alcohol, and vinegar. The expressed juice, when diluted with water, filtered, and boiled, does not yield flakes of albumen as has been stated.<sup>1</sup> Nitrate of mercury and superacetate of lead separate from the juice white, curdy precipitates. Gelatin throws down a copious precipitate: and in a less degree the same effect is produced by lime water and the alkaline carbonates. Infusion of galls forms in it pale brownish flakes; sulphate of iron throws down a copious green precipitate: lime evolves ammonia. When the insoluble part of dried Squill is digested in muriatic acid, filtered, and ammonia added in excess, a copious precipitate is thrown down, which is citrate of lime. Ether, digested on dried Squill, acquires a pale green hue, and, when evaporated on the surface of water, a thin pellicle of very bitter, resinous matter is deposited; while the water acquires an intensely bitter taste, and yields copious precipitates, with solutions of acetate of lead and nitrate of silver. From these imperfect experiments, Squills appear to contain extractive, a small portion of resin, mucus, carbonate of ammonia, the bitter principle, and citrate of lime. Vogel, from a careful analysis of Squill, gives the following as its principles: Gum 6 parts. Bitter principle (*scillitina*<sup>2</sup>) 35. Tannin 24. Citrate of lime 0. Saccharine matter 6. Woody fibre 30<sup>3</sup>, in 100 parts of the dried bulb.

*Medical properties and uses.*—Squill in small doses is expectorant and diuretic; in larger doses, emetic and purgative. Its medicinal powers were very early known, and it still retains its character as a remedy of great efficacy when judiciously exhibited. Although it operates powerfully as an expectorant, yet from its stimulating properties it cannot be given with propriety in pulmonary inflammations, until the fever and inflammatory action be previously greatly subdued by bleeding, and other evacuants; after which, by promoting a more copious excretion from the mucous follicles, it rapidly unloads the chest, and relieves the congestion and difficulty of breathing. It is more useful when combined with nitrate of potass, tartarized antimony, or ipecacuanha; and in asthma and dyspnoea without fever, Squill combined with ammoniacum is perhaps the best remedy we can employ. In dropsies, conjoined with a mercurial and opium, the efficacy of Squill is well ascertained.

<sup>1</sup> But when the expressed juice is boiled, till one half is dissipated, a white precipitate is thrown down, which when washed with alcohol, appears to be citrate of lime. *Annales de Chimie*, vol. lxxxiii. p. 149.

<sup>2</sup> *Scillitina* is white, transparent; breaks with a resinous fracture; and is pulverulent; but it attracts moisture rapidly from the atmosphere until it becomes fluid. It has an intensely bitter taste, with a slight degree of sweetness; and is very soluble in water, and in alcohol.

<sup>3</sup> *Annales de Chimie*, vol. lxxxiii. p. 158.



Its diuretic powers are much increased by this combination; perhaps depending on the absorbents being powerfully excited by the mercury, while the Squill determines to the kidneys. Cullen recommends <sup>1</sup> the oxymuriate of mercury as the best adjunct; but I have seen every purpose answered by calomel. Squill is a very uncertain emetic, a very small dose producing the most cruel vomiting in some persons, while in others, the largest doses do not even excite nausea; where, however, it readily and moderately induces vomiting, it proves more useful in whooping-cough and croup, than any other emetic. To produce its expectorant and diuretic effects most effectually, Squill must be given in substance; but to excite vomiting, its infusion in vinegar, or the oxymel, is more usually employed. Of the dried Squill gr. j. in the form of a pill, may be given at first for a dose, morning and evening, or every six hours; gradually increasing the dose to grs. v. or grs. vj., or until some degree of nausea is induced, and its expectorant or diuretic operation is obtained. In an over-dose it vomits, purges, produces strangury, bloody urine and stools, severe gripings, convulsions, and cold sweats.

Official preparations. *Acetum Scillæ*, L. E. D. *Oxymel Scillæ*, L. D. *Pilulæ Scillæ comp.*, L. E. D. *Pulvis Scillæ*, E. D. *Syrupus Scillæ maritimæ*, E. *Tinctura Scillæ*, L. D.

SCROPHULARIA. *Spec. Plant. Willd.* iii. 269

Cl. 14. Ord. 2. Didynamia Angiospermia. Nat. ord. Personatæ, Linn. Scrophulariæ, Juss.

G. 1152. Cal. five-cleft. Corolla subglobular, resupine. Capsule two-celled.

Species 2. *S. nodosa*. Knobby-rooted Figwort. Smith, *Flora Brit.* 663. Eng. Bot. 1544.

Official. SCROPHULARIA; HERBA, Dub. Figwort herb.

Syn. La Scrophulaire aquatique (F.), Die Wasser-Braunwurz (G.), Waterspeenkruid (Dutch), Scrofularia aquatica (L.), Escrofularia aguatica (S.), Escrofularia dos rios (Port.).

This is an indigenous, perennial plant, growing in woods and about hedges, flowering in July. The root is tuberous, and knobbed or granulated. The stem rises three feet in height, is erect, simple, sharply quadrangular, smooth, and leafy. The leaves are opposite, petiolate, cordate, pointed, unequally serrated, veined, smooth, cut away at the base to the two small lateral veins, and as if three-nerved. The flowers are in terminal bunches, erect, with the peduncles opposite, dichotomous, and bracteolated. The flowers are slightly drooping of a dark blood-red colour. The capsules are ovate oblong.

Qualities. — The recent leaves have a rank, fetid odour, re-

<sup>1</sup> *Materia Medica*, ii. 558.



sembling that of elder leaves, and a bitterish, disagreeable taste; but, both these qualities are nearly lost by drying. They yield their virtues to water; and the infusion precipitates sulphate of iron brown.

*Medical properties and uses.* — Figwort is supposed to possess diuretic and sedative properties. It has been used in scrophula, whence its name; and is recommended as a fomentation to piles, malignant tumours, spreading ulcers, and cutaneous eruptions; but is very little known in practice.

SENEGÆ RADIX. Vide *Polygala*.

SENNÆ FOLIA. Vide *Cassia*.

SERPENTARIÆ RADIX. Vide *Aristolochia*.

SEVUM. Vide *Ovis*.

SIMAROUBÆ CORTEX. Vide *Quassia*.

SINAPIS.<sup>1</sup> *Spec. Plant. Willd.* iii. 534.

*Cl.* 15. *Ord.* 2. Tetradynamia Siliquosa. *Nat. ord.* Siliquosæ, *Linn.* Cruciferae, *Juss.*

*G.* 1246. *Cal.* spreading. *Cor.* claws erect. *Gland* between the shorter stamens and pistil, and the longer stamens and calyx.

*Species* 4. *S. alba*. White Mustard. *Smith, Flora Brit.* 721.

——— 5. *S. nigra*. Common Mustard. *Med. Bot. t.* 151. *Smith, Flora Brit.* 722.

#### 1. SINAPIS ALBA.

*Officinal.* SINAPIS ALBÆ SEMINA, *Edin.* SINAPI; SEMINA, *Dub.*  
White Mustard-seed.

*Syn.* Moutarde (*F.*) Senfsamen (*G.*), Mosterd (*Dutch*), Senep (*Dan.*), Senape bianca (*I.*), Grano de Mostaza (*S.*), Mostardeira (*Port.*), Gortschiza (*Russ.*), Kabar (*Arab.*), Sinap (*Swed.*), Gerzykare (*Pol.*).

This species of Mustard is an indigenous annual plant, growing in the fields and by road sides; but it is also much cultivated for salads in which the young plant is used. It flowers in June. The root is spindle-shaped, and the stem round, strong, branched, and rising nearly two feet in height. The lower leaves are deeply pinnatifid; the upper sublyrate, and the whole rough with strong hairs on both sides, toothed, and of a pale green colour. The flowers are in racemes, with striated peduncles; the leaflets of the calyx are linear, and green; the petals yellow, with the claws narrow, and the border obovate and entire. The pods are spreading, on almost horizontal peduncles, hispid, roundish, ribbed, swelling where the seeds are placed, and furnished with a very long, ensiform, keeled, greenish rough beak. The seeds are large, for the size of the pod, globular, and of a light yellow, or pale yellowish-brown colour.

<sup>1</sup> Σινηπι Dioscoridis.



2. SINAPIS NIGRA.<sup>1</sup>

*Officinal.* SINAPIS SEMINA, *Lond.* Mustard-seeds.

*Syn.* Moutarde noir (*F.*), Schwarzer Senfe (*G.*), Senape (*I.*), Mostaza nigra (*S.*).

Common Mustard is an indigenous annual; and although very plentiful in its wild state, yet it is cultivated for domestic and medicinal purposes. It flowers in June. The root is small. The stem, which rises three or four feet in height, is very much branched, and spreading. The leaves are petiolate, variously lobed and toothed; those nearest to the root being rugged, while those of the stem are smooth; and the uppermost narrow, quite entire, and hanging down, which distinguishes it at first sight from its congeners. The flowers are yellow, and small. The calyx is coloured. The pods erect, parallel, to the stem, short, quadrangular, frequently smooth, many-seeded, and furnished with a short quadrangular beak. The seeds are small, globular, and of a deep brown colour.

Although the seeds of these two species of mustard differ in their botanical characters, yet they agree in other respects, the common being only rather more pungent; and they may be indiscriminately employed. Reduced to a fine powder, they form the common condiment every day used at our tables.

*Qualities.* — These seeds, in the entire state, are nearly inodorous, but when bruised they have a pungent, penetrating odour. Their taste is bitterish, acrid, and biting. Unbruised mustard seeds, when macerated in boiling water, yield only an insipid mucilage, which, like that of linseed, resides in the skin: but, when bruised, water takes up all their active matter, although it is scarcely imparted to alcohol. They contain a fixed bland oil, which is purgative in large doses. In distillation with water, mustard-seeds yield a very acrid, volatile oil, on which their virtues are supposed to depend. It is united in the seed with fecula or starch; its force appears to be obtunded by the soft, insipid, fixed oil, which can be separated by pressure, and the cake left after the expression is considerably more pungent and acrid than the unpressed seeds. It is not dissipated by drying, nor by keeping the seeds, and is rendered considerably more active by the addition of vinegar. When the seeds are triturated with lime and a few drops of water, ammonia is plentifully evolved. Hence their constituents appear to be starch, mucus, a bland fixed oil, an acrid volatile oil, and an ammoniacal salt.

*Medical properties and uses.* — Mustard-seeds are stimulant, emetic, diuretic, and rubefacient. Swallowed whole, they

<sup>1</sup> Νάπυ Hippocratis.



have been found useful in dyspepsia, chlorosis, and the torpid state of the intestines which accompanies paralysis. The bruised seeds, or the powder, to the extent of a large teaspoonful mixed with water, form an excellent emetic in paralytic, epileptic, and some apoplectic cases, often operating quickly and fully when other emetics fail. In small doses, they are found to promote considerably the secretion of urine, and consequently prove beneficial in dropsies. In these affections, however, perhaps the best mode of exhibiting Mustard is in the form of whey, which is made by boiling 3 iv. of the bruised seeds in Oj. of milk, and straining to separate the curd. A fourth part of this quantity may be taken for a dose three times a day. But Mustard is more frequently employed as an external remedy. The flour rubbed on the skin, or applied in the form of a cataplasm, made with crumbs of bread and vinegar, soon excites a sense of pain, considerable inflammation, and sometimes vesication. The volatile oil is a violent rubefacient. In these forms it has been found serviceable in paralysis; and when applied to the soles of the feet, in the delirium of typhus, and in comatose affections.

Officinal preparations. *Cataplasma Sinapis*, L. D. *Emplastrum Meloes comp.*, E.

SIUM. *Spec. Plant. Willd.* i. 1431.

Cl. 5. Ord. 2. Pentandria Digynia. *Nat. ord.* Umbellatæ.

G. 544. *Fruit* subovate, striated. *Invol.* many-leaved. *Pet.* cordate.

*Species* 4. *S. nodiflorum*, Procumbent Water Parsnip. *Med. Bot.* 2d edit. 139. *Smith, Flora Brit.* 313. *Eng. Bot.* 639.

*Officinal.* SIUM; HERBA, *Dub.* Water Parsnip herb.

*Syn.* Berle; Ache d'eau (*F.*), Wasser-Partinake (*G.*), Knoophloemige watereppe (*Dutch*), Sio; Gorgolestro (*I.*), Sio; Sion (*S.*).

This is an indigenous perennial plant, common in ditches and brooks, flowering in July and August. The root is creeping. The stem procumbent or floating, branched, round, somewhat striated and leafy. The leaves are unequally pinnate, consisting of from five to nine sessile, ovate, serrated leaflets. The umbels are solitary, nearly sessile, and opposite to each leaf. The umbellets are from five to seven, divaricated, and composed of twelve flowers. The involucre is of one piece, and often there is none. The involucl is ovate, concave, and the length of the pedicels. The calyx is scarcely visible, and the fruit ovate.

This plant has been omitted by the London College in the last edition of its Pharmacopœia, and we are inclined to believe that its pretensions even to the character of an antiscorbutic require further confirmation.

SMILAX.<sup>1</sup> *Spec. Plant. Willd.* iv. 774.

<sup>1</sup> Σμίλαξ Dioscoridis.



Cl. 22. Ord. 6. Diœcia Hexandria. Nat. ord. Sarmenaceæ,  
Linn. Asparagi, Juss.

G. 1800. Male. Calyx six-leaved. Corolla none.

———— Female. Cal. six-leaved. Cor. none. Styles three.

Berry three-celled. Seeds two.

\* Stem prickly, angular.

Species 9. *S. Sarsaparilla*.<sup>1</sup> Sarsaparilla. Med. Bot. 2d edit.  
161.

Officinal. SARSAPARILLÆ RADIX, Lond. Dub. SMILACIS SARSAPARILLÆ RADIX, Edin. Sarsaparilla root.

Syn. Racine de Salsepareille (F.), Sarsaparille (G.), Sarzaparille (Dutch), Radiche della Salsapariglia (I.), Zarzaparilla (S.), Salsaparilha (Port.), Juapecanha (Brasil.).

This plant is a perennial, a native of South America and Virginia, flowering in July and August. The root is divided into pedicels, which are somewhat thicker than a goose-quill, straight, externally brown, internally white, and three or four feet in length. The stems are shrubby, long, slender, scandent, and beset with spines: the leaves alternate, ovate, pointed, and petiolate, with long tendrils at the base. The flowers stand three or four together upon a common peduncle. The calyx of the *male* flower is bell-shaped, with the segments oblong, spreading, and reflected at their points; the filaments are six, simple, and bearing oblong anthers: the calyx of the *female* flower is also bell-shaped: the germen ovate, supporting three, minute styles, with oblong, reflexed, hairy stigmas; and the fruit a round, three-celled berry, containing two, globular seeds. But several species of *Smilax* are gathered under the name of *Sarsaparilla*. The best *Sarsaparilla* grows wild on the borders of a lake on the north of the Cerra Unturan, not far from Esmeralda. It is celebrated all over South America by the name of *Zarza del Rio Negro*.<sup>2</sup>

The dried root is imported from the Spanish West Indies, packed in bales. It is known in the London market, by the names Lisbon, Honduras, and Vera Cruz; but, of late years, it has been also brought from Jamaica. Humboldt states that nearly 5000 quintals are annually exported from Vera Cruz.<sup>3</sup> The Lisbon root, which is the produce of Brazil, has a reddish or dark brown cuticle; is internally farinaceous, and more free from fibre than the other kinds: the Honduras has a dirty brown, and sometimes whitish cuticle; is more

<sup>1</sup> Bauhin derives the name from *Zarsa*, which, he says, is the Spanish for red; and *parilla*, a little vine. The latter part of the derivation is correct; but we are inclined to think the first part must be referred to *Zarza*, a brier or bush; hence, *Zarzaparilla* would imply a *bushy little vine*.

<sup>2</sup> It is purposely smoked in drying it. *Humboldt's Person. Nar.* vol. v. p. 378. trans.

<sup>3</sup> *Polit. Essay*, vol. 2. p. 442. Clusius asserts, that Europe received the first *Sarsaparilla* from Iucaton and the island of Puna, opposite Guayaquil.



fibrous, and has more ligneous matter than the Lisbon and the Vera Cruz. Is in long slender twigs, covered with a wrinkled brown cuticle, and has a small woody heart. The Jamaica differs from the others, in having a deep red cuticle of a close texture; and the red colour partially diffused through the ligneous part.

*Qualities.*—This root is inodorous, and has a mucilaginous very slightly bitter taste. The whole of the efficacious part of the plant resides in the bark, the ligneous part being tasteless, inert, woody fibre. The root when macerated communicates to cold as well as to boiling water, and partially to alcohol and ether, any active matter it possesses. The watery infusion has a brown colour, reddens litmus paper, and yields a precipitate with infusion of galls, which is again dissolved when the infusion is heated. It is precipitated also by lime-water, and solution of nitrate of mercury, and superacetate of lead, but is not affected by sulphate of iron, or any other of the metallic oxides. The alcoholic tincture has a yellowish red hue, is rendered turbid by the addition of water, and yields an extract slightly bitter and pungent. Ether takes up two parts in ten of the powdered root; and the tincture, which has a golden yellow colour, when evaporated on water, leaves a small portion of reddish yellow insipid resin, and a larger of yellowish extractive dissolved in the water. The root brought from Jamaica yields rather more than two parts for one of extract yielded by the other kinds, and is, therefore, the best for medicinal use. M. Galileo-Pallota has separated from it an alkaloid substance, in which he supposes the virtue of the rest resides, and has named it *Parillina*. This salt is white, pulverulent, light, permanent in the air, austere, slightly astringent, nauseous, and having a peculiar odour. It is insoluble in cold, but very soluble in hot water. Slightly soluble in cold alcohol, very soluble in hot. It reddens turmeric paper, fuses at  $212^{\circ}$ , and is decomposed at a higher temperature: forms a sulphate with diluted sulphuric acid, but is decomposed by the strong acid. It forms neutral salts with the other acids.<sup>1</sup>

*Medical properties and uses.*—Sarsaparilla is demulcent, and said to be diuretic. It was brought to Europe about the year 1530, and introduced as a medicine of great efficacy in the cure of lues venerea; but it fell into disrepute and was little used, till it was again brought into esteem by Dr. William Hunter and Sir William Fordyce, about the middle of the last century; not however as a remedy fitted to cure

---

<sup>1</sup> *Journal of Science*. xix. p. 164.



syphilis ; but of much efficacy in rendering a mercurial course more certain, and after the use of mercury.<sup>2</sup> Experience, however, has not verified the encomiums bestowed on it; and the extensive observations of Mr. Pearson have fixed the degree of benefit which is to be expected from this root in syphilitic complaints. “The contagious matter, and the mineral specific, may,” he observes, “jointly produce, in certain habits of body, a new series of symptoms, which, strictly speaking, are not venereal<sup>3</sup>; which cannot be cured by mercury, and which are sometimes more to be dreaded than the simple and natural effects of the venereal virus. Some of the most formidable of these appearances may be removed by Sarsaparilla, the venereal virus still remaining in the system; and, when the force of the poison has been completely subdued by mercury, the same vegetable is also capable of freeing the patient from what may be called the sequelæ of a mercurial course.”<sup>4</sup> Sarsaparilla is, also, recommended in scrofula, elephantiasis or cutaneous affections resembling it, and chronic rheumatism; but its efficacy is doubtful. From experiment made upon himself, M. Pallota considers that Parilline is a powerful debilitating medicine, diminishing the vital energy in proportion to the dose.

The dose of the powdered root is from ʒj. to ʒj., given three or four times a day. Of the Parilline from 2 to 13 grains may be given for a dose.

Officinal preparations. *Decoctum Sarsaparillæ*, L. E. D. *Decoctum Sarsaparillæ compositum*, D. *Extractum Sarsaparillæ*, L.

SODÆ MURIAS, *Lond.* MURIAS SODÆ, *Edin.* SAL COMMUNE. *Murias Sodæ*, *Dub.* Muriate of Soda. Common Salt.

*Syn.* Muriate de Soude (*F.*), Salzaures Natrum (*G.*), Zout (*Belg.*), Salt (*Dan. Swed.*), Sal commune (*I.*), Sal (*S.*), Méllh (*Arab.*), Poppoo (*Tam.*), Loon (*H.*), Nemuck (*Pers.*), Locnoo (*Cyng.*), Lavana (*Sans.*), Garam (*Malay*), Uyah (*Jav.*), Tarrëöke (*Esquimaux*).

This salt is one of the most abundant productions of nature, being found in almost every country of every quarter of the globe; either existing in mineral springs<sup>5</sup> or

<sup>1</sup> The celebrated Mutis, in a letter to the younger Linnæus, says, “scarcely any *Lues Venerea* resists my method of administering a drink of this medicine.” (*Correspondence of Linnæus*, vol. ii. p. 549.) But we must recollect that syphilitic complaints are as benign as they are common, both among the whites and the mixed casts, in South America; and as they yield to this remedy, the quantity of sarsaparilla employed in the Spanish colonies is very considerable. See *Humboldt's Person. Nar.* vol. v. p. 379.

<sup>2</sup> *Medical Observations and Enquiries*, vol. i.

<sup>3</sup> The symptoms alluded to, are nocturnal pains in the limbs, painful enlargements of the knee and elbow joints, membranous nodes, and cutaneous ulcerations, arising after a full course of mercury.

<sup>4</sup> *Pearson on Remedies for Lues Venerea*, 24.

<sup>5</sup> The salt spring of Luneburg yields 75,600 gallons of brine in twenty-four hours, of which  $\frac{1}{4}$ th is saline matter, making the annual produce 55,000,000 lbs. of salt. *Kirwan's Geo. Essays*, 392.



lakes<sup>1</sup>; spread in strata under the surface of the ground<sup>2</sup>, or rising from it into mountains<sup>3</sup>; and to its presence also the ocean owes its saltness.<sup>4</sup> In all these situations, however, it is generally mixed with earths and other matters; and, therefore, must undergo several processes to bring it to the degree of purity in which it occurs as an article of commerce.

In Cheshire, where the greater part of the salt used in this country is made, the brine is pumped up from very deep wells, and evaporated in wrought iron pans, which are generally about twenty or thirty feet long and broad, and nine or twelve inches deep; strongly set upon masonry over a large furnace, from which flues proceed under every part of the pan. They are protected from the weather by light pyramidal roofs of boards, sufficiently open, however, to admit of the escape of the steam from the boiling brine. When the brine attains the temperature of 100° Fahrenheit, it grows turbid, and carbonate of lime and of iron are separated. These are partly taken off by skimming, but much of the mass falls to the bottom, and cannot be removed until the first deposition of crystallized salt gives it a sufficient body to enable the workmen to rake it out. After this is carefully done, the evaporation is continued at a boiling heat, when the salt gradually forms, and falls to the bottom of the pan in beautifully white, delicate crystals, which are fished out, as they collect, with wooden vessels, and poured into large wooden, hollow cones, having a hole at the apex, which is undermost. When the salt has sufficiently drained, the cones filled with it are taken to a large room, heated by stoves, where they remain until thoroughly dry.<sup>5</sup> In warm climates, the sea water is evaporated in shallow ponds by the heat of the sun; and in this mode, what is denominated bay salt, is made; but, in colder countries, the evaporation is carried on by artificial heat in a way similar to the Cheshire process. The crystals of the salt procured by these means are more perfect

<sup>1</sup> These lakes are generally dry in the summer, being formed by the small streams from the hills settling in the valleys, and dissolving the salt of the soil. There is a lake or valley of this description eighteen miles from Aleppo, called in Arabic *Subkhet al Jibool*, or Valley of Salt; in which the salt is found, in the summer, crystallized from half an inch to two inches thick. *Russel's Aleppo*, 2d edit. i. 55.

<sup>2</sup> The stratum of rock-salt in Cheshire is 50 feet thick. The salt mine of Wiliska, near Cracow, in Poland, is 6691 feet long, 1115 feet broad, and 743 feet deep. *Coxe's Travels*, i. 197.

<sup>3</sup> Near Cordova, in Spain, is a mountain of common salt 500 feet high, and nearly three miles in circumference.

<sup>4</sup> The average quantity of salt contained in sea water varies in different latitudes. Between 10° and 20° south, it amounts to rather more than one twenty-fourth; between 18° and 34° north, it is rather less than one twenty-fourth; and at the equator it is nearly one twenty-fifth. *Thomson's Chemistry*, 4th edit. vol. iv. 141.

<sup>5</sup> *Aikin's Dictionary of Chemistry*.



and purer, the more slowly the evaporation is conducted. The following table, drawn up by Dr. Henry, shows the components in the different varieties of salt used in this country :

100 Parts, by Weight, of the following Salts.		Insoluble Matter.	Muriate of Lime.	Muriate of Magnesia.	Total earthy Muriates.	Sulphate of Lime.	Sulphate of Magnesia.	Total Sulphates.	Total Muriates.	Pure Muriate of Soda.
For Bay Salt.	St. Ube's - - -	9	trace	3	3	23½	4½	28	40	960
	St. Martin's - -	12	do.	3½	3½	19	6	25	40½	959½
	Olevon - - -	10	do.	2	2	19½	4½	23¾	35¾	964¼
Sea-water from Brit. Salt	Scotch, (common,)	4	—	28	28	15	17½	32½	64½	935½
	Scotch, (Sunday,)	1	—	11½	11½	12	4½	16½	29	971
	Lymington, (com.)	2	—	11	11	15	35	50	63	937
	Lymington, (cut,)	1	—	5	5	1	5	6	12	988
Cheshire Salt.	Crushed Rock - -	10	0· <sup>1</sup> / <sub>6</sub>	0· <sup>3</sup> / <sub>6</sub>	0¼	6½	—	6½	16¾	983¼
	Fishery - - - -	1	0· <sup>3</sup> / <sub>4</sub>	0· <sup>3</sup> / <sub>4</sub>	1	11¼	—	11¼	13¼	986¾
	Common - - - -	1	0· <sup>3</sup> / <sub>4</sub>	0· <sup>3</sup> / <sub>4</sub>	1	14½	—	14½	16½	983½
	Stoved - - - -	1	0· <sup>3</sup> / <sub>4</sub>	0· <sup>3</sup> / <sub>4</sub>	1	15½	—	15½	17½	982½

The common salt of commerce, however, still contains small portions of muriate of magnesia, muriate of lime, and sulphate of lime. To separate these, dissolve the salt in four times its weight of pure water, and drop into the filtered solution, first muriate of barytes, and then carbonate of soda, as long as any precipitate falls. Filter and evaporate the clear fluid slowly till the salt crystallizes, which is pure muriate of soda.<sup>1</sup>

*Qualities.* — Pure Muriate of Soda is inodorous; its taste is strictly salt; and, when pure, it is perfectly devoid of any degree of bitterness. It is in regular cubes, which are not affected by exposure to the atmosphere. When it deliquesces, it contains muriate of magnesia. Its crystals decrepitate when heated<sup>2</sup>, and melt in a red heat, losing about two per cent. of their weight; and, in a still greater heat, the salt is volatilized undecomposed in white fumes. Its specific gravity is 2·126. It is equally soluble in cold and in hot water, rather more than two and a half parts of either being required to

<sup>1</sup> *Thomson's Chemistry*, 4th edit. ii. 377.

<sup>2</sup> The term Muriate of Soda is not chemically correct when applied to the dry salt; for in this state it is a compound of chlorine and soda, or a chloride, and becomes a muriate only when water is added; which, being decomposed, forms muriatic acid by giving up its hydrogen to the chlorine, and soda by transferring its oxygen to the sodium or metallic base of that alkali. Dry, pure sea salt, therefore, is a chloride of sodium, the chlorine being in the proportion of 59·305 to 40·695 of the sodium, according to the analysis of Berzelius.



dissolve one of salt.<sup>1</sup> It consists, according to Berzelius, of 45·74 of acid, and 54·26 of soda, in 100 parts.<sup>2</sup> It is decomposed by sulphuric acid and nitric acid.

*Medical properties and uses.* — This salt is tonic and anthelmintic in moderate doses; purgative in larger, and externally stimulant. In the ordinary mode of using it, the tonic power of salt operates in assisting the process of digestion; and, consequently, taken more freely, it proves useful in dyspepsia, and in correcting the weakened state of intestines, which favours the propagation of worms. In large doses it is said to check vomiting of blood, and may be used as a purgative; although it is seldom employed.<sup>3</sup> As a local stimulant its solution in tepid water, in the proportion of  $\bar{3}$ ss. or  $\bar{3}$ j. to Oj. of water, forms the common domestic enema. It is used also as a fomentation to sprained and bruised parts; and, dissolved in a large proportion of water, it forms the best stimulant general bath, whether used cold, or in a tepid, or in a hot state. To act as a tonic, the dose of muriate of soda may be from grs. x. to  $\bar{3}$ j; but to operate by stool from  $\bar{3}$ ss. to  $\bar{3}$ j. is necessary. It should be largely diluted.

Officinal preparation. *Murias Sodæ siccatum*, E. D.

SO'DÆ SUB-BO'RAS, *Lond.* BORAS SODÆ, *Edin.*  
BORAX. *Sub-boras Sodæ*, *Dub.* Sub-borate of Soda.<sup>4</sup>

*Syn.* Borate alcalinule de Soude (*F.*), Borax (*G.*), Borace (*I.*), Borrax (*S.*), Tineal (*Portuguese*), Booruk (*Arab.*), Tancána (*San.*), Sohaga (*H.*), Val-ligarum (*Tam.*), Patteric (*Malay.*), Tunkav (*Pers.*), Piger (*Jav.*).

This is the purified state of a natural salt found in Persia and Thibet. In the latter country it is formed in the bed of a lake situated among the mountains, fifteen days' journey from Tisoolumboo, which is twenty miles in circumference, and supplied only by springs from the bottom.<sup>5</sup> The borax is dug in large masses from the edges and shallows of the lake; yet the quantity is not diminished, the cavities being gradually filled by a fresh deposition of the salt. In this state it is named *tincal*, and is brought home packed in chests, in masses of adhering crystals, of a grey-yellowish, or greenish-white colour, intermixed with sand and other impurities, and

<sup>1</sup> The following appears to be the solubility of chloride of sodium:—

100 parts of water at 13·89° Centigrade, dissolve	35·81 parts of the chloride.
16·90	35·88
59·93	37·14
109·73	40·38. <i>Ure, Chem. Dict.</i>

<sup>2</sup> Other chemists have obtained different results. Berard procured from 100 parts, 43 of acid, 57 alkali; Dr. Marcet, acid 46, alkali 54; Wenzel, acid 46, alkali 54.

<sup>3</sup> The purgative property of sea water does not altogether depend on this salt, as it contains a large proportion of muriate of magnesia, which is purgative.

<sup>4</sup> The term *borax* is a corruption of the Arabic word *buruk*, which signifies *brilliant*. Vide Asiatic Researches, 8vo. vol. iii. p. 255.

<sup>5</sup> Saunders, *Phil. Trans.*, vol. lxxix, 97.



covered with a greasy artificial production to prevent it from efflorescing. The purification of tincal was first discovered by the Venetians; and afterwards long carried on by the Dutch, who kept the process secret. It is now practised in England; and although the method has not yet been made public, yet, Pelletier has ascertained, that by destroying the unctuous matter by calcination, the salt may be obtained pure by solution and crystallization.<sup>1</sup>

*Qualities.* — Purified Sub-borate of Soda is inodorous, and has a styptic, cool, alkalescent taste. It is of a white colour, and usually in irregular, crystalline masses, approaching to the form of hexangular prisms, terminated by triangular pyramids. It effloresces slowly and very slightly in the air; dissolves in twelve times its weight of water at 60° of Fah., and in six times its weight of boiling water, the solution changing the vegetable blues to green. In a moderate heat it undergoes the watery infusion, loses four-tenths of its weight, and becomes a dry, white, spongy mass, without undergoing any decomposition. According to Bergman, 100 parts consist of 34 acid, 17 soda, and 49 water; the acid, however, is not sufficient to saturate the alkali, and consequently the salt is a sub-borate. It is decomposed by the majority of the acids; by potash, and the sulphates, muriates, phosphates, and fluates of the earths, and of ammonia.

*Medical properties and uses.* — This salt is refrigerant and detergent. It is not given internally; and its chief employment is in aphthous affections of the mouth, and excessive salivation. It is applied either in the form of powder mixed with sugar, or dissolved in water, and united with honey as a lotion.

Officinal preparation. *Mel Boracis*, L. E.

**SO'DÆ SUL'PHAS.** *Lond. Edin.* Sulphate of Soda.

*Syn.* Sulphate de Soude (*F.*), Krystallisirtes Schwefelsaures Natrum, Glaubersalz (*G.*), Sale Mirabile di Glaubero (*I.*), C'hara Nún (*H.*).

This salt is found native in combination with oxide of iron, and muriate and carbonate of soda, and sometimes effloresced on the surface of the soil in the neighbourhood of salt lakes, in Hungary and North America<sup>2</sup>; and very often forming part of the contents of mineral saline springs, as those of Cheltenham and of Carlsbad. But the greater part of it used in this country is artificially prepared, and chiefly in the large way,

<sup>1</sup> *Mémoires de Chimie*, vol. i. 82.

<sup>2</sup> To the north of Carlton House, on the river Skatchawan, lat. 53° 20' is a small lake, on the shores of which, in the summer time, it effloresces in the form of a fine powder, to the depth of two or three inches. *Franklin's Journey to the Polar Sea*, 4to. p. 506. *Appendix.* Captain Hall also relates that the valley of Copiapo, on the coast of Chili, is covered with a layer of Sulphate of Soda, several inches thick. The salt looks like snow upon the ground. *Journ. in South America*, vol. ii. p. 22.



during the manufacture of sal ammoniac from sulphate of ammonia and common salt. On this account the London College has inserted it in the list of materia medica; but as a formula is also given for its preparation, we shall defer the consideration of its qualities and uses till it comes under our notice among the Preparations.

**SODA IMPURA.** *Carbonas Sodæ impura*, Lond. **SUB-CARBONAS SODÆ IMPURUS.** *Barilla*, Edin. **BARILLA**, *Soda impura*, Dub. Impure Soda. Carbonate of Soda. Barilla.<sup>1</sup>

*Syn.* Soude (*F.*), Kohlensaures Natrum (*G.*), Charuni (*Malab.*), Soda (*I.*), Barilla (*S.*), Sejjimitti (*H.*), Sorjica (*San.*), Karum (*Tam.*), Loogzout (*Dutch*), Soda (*Russian*).

Carbonate of Soda is found native in Hungary, Syria, Egypt, Arabia, Thibet, India, China, Siberia, and South America<sup>2</sup>, on the surface of the earth, and on the margins of some lakes which become dry in summer. A large quantity is annually collected from the natron lakes of Hungary, in the vicinity of Debretzin, and from those of Egypt, situated in the valley Bahr-bela-ma, near the Delta. The Trona lake, which lies between Wadies Shiati and Ghrurbi, yields annually between 400 and 500 camel loads, each equal to about 4 cwt. It is obtained by a man wading into the lake and breaking it off in sheets, which are afterwards packed in the setose basis of the palm leaves.<sup>3</sup> It is named Szekso in Hungary, and *trona* by the natives of Egypt: but very little of it finds its way to Britain; and the greater part of that which is employed, in this country at least, is of vegetable origin, being prepared from the ashes of some species of *Algæ*, but more abundantly from those of the *Salsola Soda*, a plant which is cultivated on the shores of the Mediterranean, by the Spaniards, expressly for the purpose of yielding this salt. This plant is cultivated in salt marshes, chiefly in the vicinity of Alicant; and near Carthagena two hundred thousand quintals of Barilla are gathered annually. In September, when the seed is ripe, it is pulled up by the roots: after which it is dried, and in October burnt in simple furnaces, the heat of which is just sufficient to cause the ashes

<sup>1</sup> Νίτρον of the ancients. This term was converted in the middle ages into *natrum* or *natron*; while the real word Νίτρον was improperly applied to Nitre: but the ancients were unacquainted with Nitre, and the term *natrum* is never found in the works of the Greeks or the Romans, and not even in those of good writers of the middle ages. Beckman proves, that Nitre was not known, even in Europe, till the invention of gunpowder. *Hist. of Inventions*, iv. p. 537. It is remarkable that both the London and Dublin Colleges have fallen into the same error, in naming this salt *impure soda*; the composition of the salt being well ascertained.

<sup>2</sup> Native Carbonate of soda is found in considerable quantity in a small lake, near the Indian village, Lagunillas, situated to the south east of Merida, in the republic of Colombia. It is termed *Urao* by the natives. It contains one-fourth more soda than the Trona of Egypt; and more carbonic acid than the artificial carbonate. *Annales de Chimie*, tom. xxix. p. 110.

<sup>3</sup> Letter from Dr. Oudney to Professor Jameson. *Edin. Phil. Journ.* vol. xi. p. 384.



to enter into a state of semifusion, and concrete into compact, cellular masses, which form the *Barilla* of commerce. The *Barilla* of Ustica in Sicily is esteemed the finest in Europe: and its excellence is supposed to depend on the plant being burnt before it is thoroughly dry.<sup>1</sup> That which is obtained in this country by burning the sea-wrack (chiefly the *Fucus vesiculosus* and *serratus*) is denominated *kelp*; and is the worst description of this salt.<sup>2</sup> Sardinia also furnishes some *Barilla*, but inferior in quality to the Spanish. In Sicily *Barilla* is procured at *Catania*, *Trapani*, *Marsala*, *Terranora*, and *Girgenti*. Orkney furnishes annually about three thousand tons of kelp.<sup>3</sup>

Vauquelin has proved that the salt exists ready formed in *Salsola soda*, and is only set free by the burning of the plant.<sup>4</sup> He obtained from 500 parts of the plant, 100 parts of ashes, besides oil, ammonia, and prussic acid; but the plant also contains iodine. Five hundred grains of the ashes afforded 113 of muriate of soda, 68 dry subcarbonate of soda, 204 insoluble subcarbonate of magnesia, 100 of sand and oxide of iron, and 23 of water<sup>5</sup>; an analysis which may be regarded as exhibiting the general components of *Barilla*.

There are several varieties of *Barilla* brought from Spain; that which is known by the name of *Sweet Barilla* is the most esteemed.

*Qualities.* — Good *Barilla* is in hard, dry, spongy, sonorous masses of a greyish blue colour, and becoming covered over with a saline efflorescence when exposed to the air. It should not emit any unpleasant odour on solution; and when applied to the tongue should impress a sharp, alkaline taste.

*Use.* — Impure Subcarbonate of Soda is employed only for yielding the pure Subcarbonate.<sup>6</sup>

---

<sup>1</sup> *Smyth's Sicily and its Islands*, 4to. p. 15.

<sup>2</sup> The inhabitants of the Canary Isles extract Carbonate of Soda from the ashes of the *Mesembryanthemum crystallinum*, or ice plant, which yield one-third of their weight of the salt. *Phil. Mag.* vi. 187.

<sup>3</sup> The sea-wrack is cut from the rocks in April and May chiefly, dried in the air, and then burnt in a kiln, in which they are stirred with an iron rake into a fluid state; on cooling, the ashes condense into a dark blue or whitish mass, very hard and solid. The plants about three years old yield the largest quantity of kelp. The best kelp has an acrid caustic taste, a sulphurous odour; is compact, and of a dark blue-greenish colour. It yields about one-twentieth part of its weight of soda.

<sup>4</sup> *Annales de Chimie*, xiii. 65.

<sup>5</sup> *Annales de Chimie*, xviii. 76.

<sup>6</sup> A tolerably pure Subcarbonate of Soda is obtained in France from sea salt, muriate of soda. To a solution of one part of salt, three parts of finely pulverized litharge is added, and rather more than half a part of chalk. These are agitated well together, and then set apart: a double decomposition gradually takes place, muriate of lead and muriate of lime are formed, while the soda, uniting with the carbonic acid set free from the chalk, crystallizes, and is easily separated. In Britain it is obtained in considerable quantity by decomposing sulphate of soda either



Officinal preparation. *Sodæ Subcarbonas*, L. E. D.

**SOLANUM.** *Spec. Plant. Willd.* i. 1025.

*Cl.* 5. *Ord.* 1. Pentandria Monogynia. *Nat. ord.* Luridæ, Linn. Solanææ, Juss.

*G.* 383. *Corolla* wheel-shaped. *Anthers* slightly coalescing, opening by two pores at the apex. *Berry* two-celled.

\* *unarmed*.

*Species* 15. *S. Dulcamara*.<sup>1</sup> Woody Nightshade, or Bitter Sweet. *Med. Bot.* 2d edit. 240. t. 85. *Smith, Flora Brit.* 256. *Eng. Bot.* 565.

*Officinal.* DULCAMARÆ CAULES, *Lond.* SOLANI DULCAMARÆ CAULES, *Edin.* DULCAMARA; STIPITES AUTUMNO COLLECTI, *Dub.* The Stalks of Bitter Sweet.

*Syn.* Douce-amere (*F.*), Bittersusstangel (*G.*), Bitterzoet (*Dutch*), Bittersöde (*Dan.*), Quesved (*Swed.*), Psinki wodne (*Pol.*), Dulcamara (*I.*), Amara-dulcis (*S.*), Doçamarga (*Port.*), Solotucha Elsterbeeren (*Russ.*).

This species of *Solanum* is an indigenous shrub, growing in hedges and shaded spots; and flowering in June and July. The root is ligneous: the stem woody, roundish, branched, and climbing sometimes to the height of six or eight feet: the leaves are alternate, on footstalks, entire, smooth, soft, about two inches long and one broad, and of a dull-green colour; the lowermost cordate and undivided, and the uppermost halberd-shaped; the flowers are in elegant clusters opposite to the leaves, or terminal, drooping, and having the semblance, but not the structure, of a true cyme; each consisting of a small purplish calyx with blunt segments; a corolla of five, reflected, equally divided, pointed, bright violet-coloured segments, with two green dots at the base, and a longitudinal deeper-purple vein through the centre of each segment; and large, erect, almost sessile, lemon-yellow anthers: the berries, which ripen in September and October, are oval, scarlet, very juicy, bitter, and poisonous.<sup>2</sup>

The extreme twigs are the parts employed. They should be gathered in autumn, as at that season they are more powerful; depending perhaps on their being less succulent, and containing more of the peculiar secretion on which the virtues of the plant depend. The soil in which the plant grows also affects its medicinal powers. That grown on a high and dry situation is the best.

*Qualities.* — The fresh twigs have a peculiar odour, but

by means of subcarbonate of potass, or acetate of lime, or litharge; or by decomposing the sulphuric acid of the sulphate by charcoal. The latter method is practised in the West of Scotland, and affords a very pure Subcarbonate of Soda. "About 500 cwt. of Sulphate of Soda and 100 cwt. of charcoal are ground together, and the mixture exposed in a reverberatory furnace until it becomes pasty. It is then transferred into large casks, and lixiviated. The ley is afterwards evaporated and crystallized." *Duncan's New Edinburgh Dispensatory*, 216.

<sup>1</sup> *Citocattia* of the monastic age.

<sup>2</sup> They excite violent vomiting and purging.



the dried are inodorous. They have a slightly bitter taste, followed by a sweetness not unlike that of liquorice root, with a slight degree of acrimony. Boiling water extracts all their active matter. Scheele found that they contain citric acid. Their medicinal properties depend on an alkaline principle which has been termed *Solanine* or *Solania*.

*Medical properties and uses.*—Bitter Sweet is diuretic and narcotic. It has been found useful in humoral asthma, dropsy, chronic rheumatism, and in lepra vulgaris and alphas, scabies, and pityriasis. Willan<sup>1</sup> remarks, that it is not applicable for the cure of lepra nigricans: we can assert that it is not of the least use in acute rheumatism; and we believe of as little in fluor albus and suppression of the menses, in which it has been strongly recommended. When given in too large doses at first, it occasions nausea, vomiting, syncope, violent palpitation and convulsive twitchings in the eyelids, the lips and the hands. It therefore requires to be begun with small doses; and even when it is more cautiously administered, if these symptoms occur, the dose must be lessened, and some aromatic conjoined. The usual form under which it is used is that of watery infusion or decoction; but it may be also given in substance pulverized. The dose of the powder may be from grs. x. to ʒj. taken in a cupful of milk.

Official preparation. *Decoctum Dulcamaræ*, L.

SOLIDAGO. *Spec. Plant. Willd.* iii. 2053.

Cl. 19. Ord. 2. Syngenesia Superflua. *Nat. ord.* Compositæ Discoideæ, *Linn.* Corymbiferae, *Juss.*

G. 1292. Receptacle naked. Pappus simple. Corollets of the ray about five. Calyx scales imbricate, closed.

\*\*\* with erect racemes.

Species 35. *S. Virgaurea*. Common Golden Rod. *Smith, Flora Brit.* iii. 890. *Eng. Bot. t.* 301. *Purch.* ii. p. 542.

Officinal. VIRGA AUREA; FLORES, FOLIA, *Dub.* Flowers and leaves of Common Golden Rod.

This is an indigenous perennial plant, found in woods, copses, and among furze upon heathy ground; flowering from July to September. The root consists of long, simple fibres: the stem, which varies in height, from ten inches to three feet, is curved below, then erect, leafy, very slightly zigzag, angular, striated, rough, and a little downy. The leaves are deep-green on the upper surface, and pale beneath, on winged petioles, harsh and clothed with rigid down; the radical leaves are obovate and serrated: the stem leaves narrower, more entire, and alternate. The flowers are yellow, in terminal and axillary erect clusters, forming a dense panicle, with lanceolate, downy bractes; the scales of the calyx are membranous, with a downy

<sup>1</sup> *Description and Treatment of Cutaneous Diseases*, 147.



border; the rays of the corolla from five to nine or ten, oblong, elliptical, and toothed at the apex. The seeds are pubescent, and the pappus rough.

*Qualities*. — The odour is slightly aromatic, and the taste subastringent and somewhat aromatic. Boiling water extracts the active matter of the leaves; the infusion changes, slightly, to green the colour syrup of violets, and precipitates sulphate of iron black.

*Medical properties and uses*. — Golden rod is astringent, and has been regarded as lithontriptic. It may be of some use in a weak state of the viscera; and thence, like other tonics, may prove beneficial in calculous habits; but its effects are not sufficient to entitle it to much notice. The dose of the powdered leaves and flowers may be from grs. x. to ʒj. or more.

SPARTIUM.<sup>1</sup> *Spec. Plant. Willd.* iii. 926.

*Cl.* 17. *Ord.* 4. Diadelphia Decandria. *Nat. ord.* Papilionaceæ. *G.* 1332. *Stigma* longitudinal, villous above. *Filaments* adhering to the germen. *Calyx* produced downwards.

\* \* with ternate leaves.

*Species*. *S. scoparium*. Common Broom. *Med. Bot.* 2d edit. 413. *t.* 150. *Smith, Flora Brit.* iii. 753.

*Officinal*. SPARTII CACUMINA, *Lond.* SPARTII SCOPARII SUMMITATES, *Edin.* GENISTA; SEMINA, CACUMINA, *Dub.* The tops and seed of Broom.

*Syn.* Genet à balais (*F.*), Pfriemenkraut (*G.*), Bezembrem (*Dutch*), Gyfvel (*Dan.*), Pingsblomma (*Swed.*), Ginestra (*I.*), Esparto (*S.*), Giesta (*Port.*).

This is an indigenous shrub, growing on dry common pastures flowering in May and June. It usually rises from four to six feet in height, and sends off numerous, straight, angled, green, smooth, leafy branches: the leaves are ternate, small, and smooth; the upper ones, however, are frequently simple. The flowers are papilionaceous, axillary, solitary, peduncled, nodding, large, and showy; of a golden colour, sometimes tawny on the outside, and occasionally altogether of a lemon hue; the calyx is nearly bell-shaped, bilabiate, gaping, even and purplish, with a five-toothed lip; the stamens are all united into a tube at the base, and bear oblong saffron-coloured anthers; the germen is villous; the style bent almost to a circle; and the legume compressed, brown, ciliated and containing several compressed, shining seeds.

*Qualities*. — The tops, when bruised, have a disagreeable odour, and a nauseous bitter taste. Both water and alcohol extract their active matter.

*Medical properties and uses*. — Broom tops are diuretic and cathartic: the seeds are said to be emetic. The effects of this plant have been very long known to the common people;

<sup>1</sup> Σπαρτίον Dioscoridis,



and both Mead and Cullen found them useful in dropsy. The usual mode of exhibiting them is in the form of decoction, made by boiling ℥j. of the green tops in a pint of water down to half a pint. Speaking of this decoction, of which two table-spoonfuls were given every hour till it operated by stool, Cullen says<sup>1</sup>: “It seldom fails to operate both by stool and urine, and by repeated exhibition, every day, or every second day, some dropsies have been cured<sup>2</sup>,” Sydenham used the ashes, which contain an alkaline salt.<sup>3</sup>

Official preparation. *Extractum Cacuminum Genestæ*, D.

SPIGELIA. *Spec. Plant. Willd.* i. 824.<sup>4</sup>

Cl. 5. Ord. 1. Pentandria Monogynia. Nat. ord. Stellatæ, Linn. Gentianæ, Juss.

G. 308. Corolla funnel-shaped. Capsules twin, two-celled, many-seeded.

Species 2. *S. marilandica*. Perennial Worm-Grass. *Med. Bot.* 2d edit. 178. t. 69. *Edin. Phil. Trans.* iii. 151. t. 1.

Officinal. SPIGELLÆ RADIX, Lond. Dub. Edin. Worm-grass root.

Syn. Spigelia de Maryland Brinailler (*F*), *Spigelia* (*I.*).

This is a perennial plant, a native of the warmer parts of North America; flowering in July and August.<sup>5</sup> The stems are annual, simple, erect, rough, quadrangular and rigid; and about seven or eight inches in height. The leaves are opposite, sessile, ovate-lanceolate, quite entire, smooth and spreading. The flowers are in a solitary spike, with small opposite bractes; the calyx consists of five awl-shaped, persistent leaflets; the corolla is of a bright-red colour on the outside, and deep-orange within, pentangular above, gibbous at the throat, widening at the base; with the border five-parted; the segments being lanceolate and revolute; the stamens are five, shorter than the corolla, supporting sagitate, converging anthers; the germen is superior, with a round style, jointed below, with the upper part deciduous; the seeds are angular and rugged.

*Qualities.* — *Spigelia* root has a bitter taste, which is imparted to boiling water.

*Medical properties and uses.* — This root is purgative and anthelmintic. Its anthelmintic virtues were discovered by the Cherokee Indians, to whom it is known under the name of *unsteetla*; and many opportunities of proving its efficacy in worm cases have occurred both in America and this country. When in the recent state, and given in small doses, it occasionally produces giddiness, dimness of sight, and even con-

<sup>1</sup> *Mat. Med.* ii. 534.

<sup>2</sup> *Ibid.*

<sup>3</sup> *Tract. de Hydrope*, Opera, 466.

<sup>4</sup> The genus was named after Adrian Spigelius, a celebrated Professor of Anatomy at Padua.

<sup>5</sup> It was first cultivated in England in 1694, by Bobart.



vulsions: effects which are attributed to a narcotic principle it possesses, but which its powerful cathartic property prevents from acting, when the dose is large. It is usual to administer an emetic previous to the use of it; and to aid its purgative operation by the addition of two or three grains of calomel, or eight or ten of rhubarb. It has been found most powerful in expelling lumbrici; and its vinous infusion is said to have been found useful in intermittents. Dr. Barton recommends it in the protracted remitting fever of infants, which often lays the foundation of hydrocephalus. Spigelia root may be administered either in substance or in the form of aqueous infusion. The dose of the pulverized root is from grs. x. to ʒj, given every night and morning until the worms are expelled.

**SPIRITUS RECTIFICATUS**, *Lond.* **ALCOHOL FORTIUS**, *Edin.* **SPIRITUS VINOSUS RECTIFICATUS**, *Dub.* Rectified Spirit. Alcohol.

*Syn.* Eau de Vie Rectifié (*F.*), Rectifizirter Weingeist (*G.*), Acquavite rettificata (*I.*), Agua ardiente (*S.*).

This is Alcohol nearly in the highest state of concentration in which it can be easily prepared in the large way for the purposes of trade. The London and Edinburgh Colleges state its specific gravity to be to that of water as 835 to 1000; while the Dublin College states it as 840. The Edinburgh College names this spirit *Alcohol*; but as directions are given both by the London and the Dublin Colleges for the preparation of a still stronger spirit, the name *Alcohol*, in their Pharmacopœias, is judiciously retained for the stronger spirit, while that of *Rectified Spirit* is applied to the present preparation.

All substances which have undergone the vinous fermentation, and in which it is not completely over, contain alcohol ready formed, but combined with colouring matter, extractive, and other principles, and are capable of affording it by distillation. The first distillation of wines and fermented liquors afford ardent spirits, such as brandy, rum, arrack, whisky and gin.<sup>1</sup> These are all mixtures of alcohol, water, and a little oil or resin, which give them their characteristic flavour and colour, and the quantity and nature of which constitute

---

<sup>1</sup> We have no historical record of the period when the distillation of spirit was invented: the Greeks and Romans were ignorant of ardent spirits: but the use of the still was well known in the time of Geber, who lived in the seventh century, who describes very accurately the process of distillation by the alembics *per descensorium et filtrum*, in his work, entitled *Liber Investigationis Magisterii*. The first spirits known in Europe was made from grapes, and sold as a medicine both in Italy and Spain, under the name of *Alcohol*. The Genoese afterwards prepared it from grain; and sold it in small bottles at a very high price, under the name *Aqua Vitæ*. Vide Morewood's *Essay on Inebriating Liquors*.



the sole differences in ardent spirits. It is from the re-distillation or rectification of these that rectified spirit is produced.

Table of the different kinds of Spirits.

Names.	Materials from which they are distilled.	Countries producing them.
Agua ardiente ...	Pulque, the fermented juice of the Agave	Mexico.
Arrack .....	Coarse palm sugar, named jaggery, fermented with the bark of the <i>Mimosa leucophlea</i> : also from rice and the fermented juice of the Palm.....	India.
Var: Mahwah Arrack .....	Flowers of the Madhuca tree, <i>Bassia butryacea</i> .....	India.
Tuba .....	Palm wine .....	Phillipine Islands.
Araka .....	Koumis, fermented Mare's milk .....	Tartary.
Araki .....	Dates .....	Egypt.
Arika .....	Fermented Cow's milk, a variety of Koumis. ....	Tartary, Iceland.
Brandy <sup>1</sup> .....	Wine, figs, peaches, Persiman apple, mulberries, and sometimes other fruits....	Europe, Asia, N. & S. America : wherever wine is made.
Var. Lau .....	Rice....	Siam.
Rakia ... ..	Husks of grapes mixed with aromatics	Dalmatia.
Rossolio .....	A compound of brandy, Ros-solis, and other plants .....	Dantzic.
Troster .....	Husk of grapes, fermented with barley and rye .....	On the Rhine.
Sekis-kayavodka	Lees of wine and fruit .....	Scio.
Geneva <sup>2</sup> Hollands	Malted barley and rye, rectified on Juniper berries .....	Holland.
Var. Gin <sup>3</sup> .. ..	Malted barley, rye, potatoes ; rectified with turpentine .....	England.
Goldwasser.....	Wheat, barley, and rye, rectified with anniseeds, cinnamon, and other spices ..	Dantzic.
Kirchwasser .....	Machaleb cherry .....	Switzerland.
Maraschino .....	Macarska cherry .....	Zara, capital of Dalmatia.
Rum <sup>4</sup> .....	Cane Sugar and Molasses.....	West Indies and South America.
Var. Slatkaia trava	Maple sugar .....	North America.
Show-choo.....	A sweet grass.....	Kamschatka.
	The lees of Mandarin, a wine made from boiled rice.....	China.
Whiskey <sup>5</sup> .....	Malted and raw barley, rye, oats, and potatoes .....	Scotland & Ireland.
.....	Sloes .....	South of France.
Y-wer-a .....	The root of the Teeroot, baked, pounded and fermented .....	Sandwich Islands.

<sup>1</sup> The best Brandy is that of Cogniac, the next of Bourdeaux and Rochelle.  
<sup>2</sup> Named from *Genevre*, the French for Juniper.  
<sup>3</sup> The quantity made in England annually exceeds 3,000,000 gallons.  
<sup>4</sup> The appellation Rum is supposed to be derived from the terminal syllable of the word *Saccharum* ; but the native Americans called this liquor Rum.  
<sup>5</sup> 2,499,880 gallons were distilled in Scotland, in 1822, 1,341,978 gallons of which were sent to England, unrectified, for making Gin and compounds. In the same year, the quantity made in Ireland was 4,318,012 gallons. The name *whiskey* is supposed to be derived from *usque*, the two first syllables of *usquebagh*, the name it originally had in Ireland, whence the Scots appear to have derived their knowledge of it. In Ireland it was also called *buil-ceaun*, which literally sig-



The process of rectification is exceedingly simple. Any quantity of brandy, malt spirits, or rum, is diluted with an equal portion of water, and put into an alembic or still, to which a refrigeratory is united, and distilled with a very gentle heat. The first product is the strongest and purest, and when it has come over to the amount of one-fourth of the whole contents of the still, forms the rectified spirit. If the distillation be continued, the spirit continues to come over colourless, but weaker and weaker, till at length it is so watery as not to be inflammable. What remains in the alembic is water, the colouring ingredients, and any accidental impurities. When the ardent spirits which have been employed, contain much oil, the distillation requires to be repeated, and generally with the addition of alkali, lime, or other articles, before the empyreumatic flavour can be completely destroyed. When alkali is used, the spirit has an urinous taste; to free it from which it is again distilled with the addition of a little alum and charcoal, the acid of the former of which attracts the small portion of alkali which the spirit held in solution. Malt spirits, when properly rectified, yield as pure and as strong rectified spirit as brandy.

The strength of spirits is ascertained in common by several methods. The taste, and the degree of frothiness or size of the bubbles formed when spirit is shaken, is the least correct method; and the burning the spirit, and observing the quantity of water which remains after the combustion, although more accurate, is liable also to error, from the impossibility of performing the experiment always under the same circumstances. Pure Alcohol<sup>1</sup> leaves no water; rectified spirit of moderate strength, 25 per cent.; French brandy, 56; and common malt liquor, 65; and the like. Another test is the pouring a few drops of the spirit on gunpowder; but this is also very incorrect, and indicates two degrees of strength only; that which fires gunpowder, and that which cannot fire it. A more accurate test than any of these, and sufficient for common purposes, is to shake the spirit in a phial with very dry, pure sub-carbonate of potass, and observe the quantity of water attracted by the alkali, which indicates its strength. But the only certain mode of ascertaining the relative strength of spirits is by determining the specific gravity of the spirit, at a given temperature. Thus at 60° Fahrenheit the specific gravity of rectified spirit is .83599, at 65° it is .83362, and at 70° the gravity of the same spirit is .83124; while the gravity

---

nifies madness of the head. The best Scotch whiskey is *Glenlivet*; the best Irish, *Ennishowen*.

<sup>1</sup> By the term *pure alcohol* is meant alcohol of a specific gravity of 796 at 60° Fahrenheit, the strongest which can be procured.



of the proof spirit of the London College at the same degrees of temperature is  $\cdot 93002$ ,  $\cdot 92794$ , and  $\cdot 92580$ ; (see Table under the article *Alcohol* among the Preparations) the weakest spirit having the greatest specific gravity, and this diminishing as the temperature increases. The usual mode of ascertaining the relative gravity of different spirits is by the hydrometer<sup>1</sup>, of which there are different kinds in use. For ordinary purposes the relative strength of spirits may be known by weighing the sample to be tried in a phial capable of holding exactly 500 grains of distilled water. An equal bulk of rectified spirit weighs 418 grains, and of proof spirit 465: hence the number of grains above or below these sums will indicate the relative strength of the spirit.

*Qualities.* — Pure Rectified Spirit has a fragrant odour, and a hot, highly-pungent taste. It is colourless; always fluid; cannot be congealed at any known degree of cold; evaporates speedily at the ordinary temperature of the atmosphere; boils at  $163^{\circ}$  Fahrenheit; and is extremely inflammable, burning with a blue, lambent flame, without any sensible smoke. Like alcohol it combines with water in every proportion; and, on account of its affinity for water, precipitates many of the neutral salts from their aqueous solutions. It is capable of dissolving many saline bodies, and is the proper solvent of the greater number of the proximate principles of vegetables. Its constituents are 85 of pure alcohol and 15 of water, in 100 parts, when its specific gravity is 835 at a temperature of  $60^{\circ}$  of Fahrenheit; but 83 only of pure alcohol, and 17 of water, when it is 840 as designated by the Dublin College.

*Medical properties and uses.* — Rectified Spirit is a very powerful stimulant. In its undiluted state it is never exhibited as a remedy; and is employed only for forming the diluted spirit, and as a pharmaceutical agent.

Official preparations. *Alcohol*, L. D. *Spiritus Camphoræ*, L. E. *Spiritus Ammoniacæ*, L. D. *Spiritus Lavandulæ*, L. E. D. *Tinctura Aloes*, L. *Tinctura Assafoetidæ*, L. E. D. *Tinctura Benzoini composita*, L. E. D. *Tinctura Castorei*, L. E. D. *Tinctura Guaiaci*, L. E. D. *Tinctura Moschi*, D. *Tinctura Ferri Muriatis*, L. *Tinctura Muriatis Ferri cum Oxydo rubro*, D. *Liquor Hydrargyri Oxymuriatis*, L.

SPIRITUS TENUIOR, *Lond.* ALCOHOL DILUTUS, *Edin.*  
SPIRITUS VINOSUS TENUIOR, *Dub.* Weaker Spirit. Diluted Alcohol. Proof Spirit.

This is merely Rectified Spirit diluted with a certain proportion of water. According to the London and Dublin Colleges, its specific gravity should be to that of distilled water as 930 to 1000; while the Edinburgh College orders it of the gravity

<sup>1</sup> For a description of this instrument, see Part I.



of 935. The former may be formed by mixing four parts by measure of rectified spirit with three of water, and contains 44 parts of pure alcohol and 56 of water, in 100 parts; the latter is obtained from equal parts of rectified spirit and water, and contains 42 of pure alcohol and 58 of water in 100 parts. But very frequently, instead of being formed by mixing the pure rectified spirit with water, an impure spirit of a strength nearly similar is employed. The qualities of the diluted spirit are not different from those of the rectified spirit, except in degree, and in some instances it is better fitted for taking up the principles of vegetables submitted to its action. It is a curious fact, that diluted spirits, preserved in vessels covered with bladder only, become stronger: a circumstance that appears to depend on the solvent powers of the aqueous vapour acting on the bladder and relaxing its pores, so as to escape through them, whilst those of the alcohol, having the power of coagulating gelatin and animal albumen, tend to close their pores and are consequently detained.

*Medical properties and uses.*—Alcohol diluted to the degree of proof spirit is still a very powerful diffusible stimulant, and too strong for internal use. Externally applied, it is recommended in burns; to restrain bleeding in passive hæmorrhagies; and as a friction or fomentation to relieve muscular pains; and in a more diluted state it forms a good collyrium in the latter stage of ophthalmia. In cases of retention of the urine owing to paralysis of the bladder, bathing the hypogastric region with alcohol, has been found beneficial. Proof Spirit diluted with water is employed as a remedy in the form of tinctures and spirits; and the ardent spirits in common use may be regarded as nearly of the same nature. These taken in moderation increase the general excitement, communicate additional energy to the muscular fibres, strengthen the stomach, and exhilarate the mind. Hence they are often and advantageously used in cases of debility and the low stage of typhoid fevers, in which the use of wine is indicated; and, in habits disposed to create acidity, they are even preferable to wine; some of them, particularly brandy, proving gratefully stomachic, when wine is nauseated and rejected. As an article, however, of daily or dietetical use, particularly if taken in immoderate doses or long continued, ardent spirits, besides being the source of much moral evil, and debasing the human character nearly to a level with that of brutes, occasion disease, and are commonly the origin of dyspepsia, hypocondriasis, and hepatic and visceral obstructions.<sup>1</sup> The hurtful effects

---

<sup>1</sup> Besides those spirits which are distilled from wines, grain, and fruits, and the other articles mentioned in the Table, page 573., the Afghaniſtans make also



of ardent spirits, however, are obviated in a considerable degree, by diluting them with water, and adding lemon-juice and sugar to the mixture, so as to form what is generally known by the name of punch. Although all the varieties of ardent spirits may be regarded as diluted alcohol, yet each has a peculiarity of operation: thus, *Brandy* is simply cordial and stomachic; *Rum* heating and sudorific; *Gin* and *Whiskey* diuretic; and *Arrack* styptic, heating, and narcotic, and ill adapted to European constitutions.

Officinal preparations. *All the tinctures except those which are prepared with rectified spirit; and all the spirits.*

SPONGIA. *Syst. Nat. Gmelin. vi. 3817.*

Cl. 6. Ord. 4. Vermes Zoophyta.

G. 343. A flexile, fixed, torpid, polymorphous animal, composed either of reticulate fibres, or masses of small spires interwoven together, and clothed with a gelatinous flesh full of small mouths on its surface, by which it absorbs and ejects water.

Sp. 8. *S. officinalis*. Officinal Sponge. *Phil. Trans. lv. 283. t. 10.*

Officinal. SPONGIA, *Lond. Dub.* SPONGIA OFFICINALIS, *Edin.* Sponge.

*Syn.* Eponge (*F.*), Meerschwamm (*G.*), Spongie (*Dutch*), Svamp (*Dan.*, *Swed.*), Spugna (*I.*), Esponja (*S.*, *Port.*), Bodiaga (*Russian*), Isfunj (*Arab.*), Mo-ābādul (*H.*), Abeermoordele (*Pers.*), Uniwatta (*Jap.*).

This species of Sponge is found chiefly in the Mediterranean and Red Seas. In some of the islands of the Archipelago the inhabitants are trained from infancy to dive for sponges, which are generally found attached to the bottom of the rocks. Although the ancients had perceived something like sensation in sponges by their shrinking, they were long supposed to be of a vegetable nature, from their texture and the branched appearance which they assume, till Mr. Ellis's observations<sup>1</sup> first established the fact that they are animals sui generis, the mouths of which are the open ends of so many branched tubes opening on the surface, through which they receive their nourishment and discharge their fæces like polypes. These mouths are generally guarded with minute spines or points, and the tubes are filled with a gelatinous matter, and often with minute shells and sand.

*Qualities.* — Sponge cleaned and rendered fit for use is of a brownish yellow colour, soft, light, and very porous; absorbing rapidly by capillary attraction as much as it can contain

an intoxicating liquor from ewe's milk; and at Kamschatka a powerfully inebriating liquor is prepared from a species of mushroom, named Muchumer.

*Phil. Trans. lv. 284.*



of any fluid in which it is immersed, and again yielding it up on being compressed. When sponge is digested with boiling, distilled water it yields up to it a considerable proportion of gelatin, and the sponge loses much of its flexibility, and crumbles between the fingers when dry. Boiled with potass it forms a soap. Its principal constituents, according to the analysis of Mr. Hatchett, are animal gelatin, albumen, a small portion of common salt, and some carbonate of lime: and to these we may perhaps add iodine<sup>1</sup>; although some suppose that this constituent is present in burnt sponge only, and to be the result of a decomposition and recombination of principles produced by the combustion.

*Medical properties and uses.* — Sponge, in its usual form, is never employed as a remedy; but is an exceedingly useful instrument in the practice of surgery.

Officinal preparations. *Spongia usta*, L. D.

STALAGMITIS. *Spec. Plant. Willd.* iv. 980.

Cl. 23. Ord. 1. Polygamia Monœcia. *Nat. ord.* Tricoccæ.

G. 1888. *Hermaph.* Calyx four-leaved. Corolla four-petalled.

Stamens thirty, inserted into a fleshy four-angled receptacle.

Style thick. Stigma four-lobed. Berry one-celled, crowned by the style, three-seeded.

Male. Calyx, Corolla, and Stamens hermaphrodite.

Sp. 1. *S. Cambogioides*. The Gamboge-tree. *Murray App.*

*Med.* iv. 645. *Plencks Icones Plant. Med.* t. 421.<sup>2</sup>

Officinal. CAMBOGIA, *Lond.* GAMBOGIA, *Edin.* GAMBOGIA; GUMMI RESINA, *Dub.* Gamboge.

*Syn.* Gomme Gutte (*F.*), Gummigutt (*G.*), Gomma Gotta (*I.*), Guta Gamba (*S.*), Rom (*Port.*) Ossāra réwūd (*Arab.*), Gokoota (*Cingalese*), Mukki (*Tam.*).

The tree yielding this gum-resin is a native of the kingdom of Siam, and of Ceylon, where it is known by the names *Ghokata*, or *Gokkata*, or *Gohlatha*. It is of middling stature, and moderately branching. The leaves are on short petioles, ovate, opposite, entire, even, coriaceous, rigid, and of an obscure-green colour. The male flowers are either in distinct clusters, or mixed with the hermaphrodite, which are axillary. The calyx consists of four, ovate leaflets, the two exterior of which are smaller than the two interior: the petals are four, spreading, coriaceous, with ciliated margins, and of a yellow colour: the stamens are about thirty, affixed to a quadrangular, fleshy receptacle, and bearing subquadrangular, club-shaped anthers; sometimes there is the rudiment of a style, and an echinated, unequal, sterile stigma. The hermaphrodite flowers are in axillary whorls, or on the joints of the smaller branches, sometimes mixed with the male flowers, sometimes

<sup>1</sup> For an account of Iodine, see *Appendix* No. V.

<sup>2</sup> Decandolle says Gamboge is the produce of the *Garcinia Cambogia*.



in opposite gems. The calyx, corolla, and stamens are the same in both: the germen is globular, with a short solitary style, and a three or four-lobed stigma; with the lobes spreading, obcordate and persistent. The fruit is a smooth, globular, whitish or rosaceous berry, crowned by the lobes and style of the stigma, and containing several long, triangular seeds.<sup>1</sup>

In Siam the Gamboge is obtained in drops by breaking the leaves and young shoots, but in Ceylon from the bark of the tree wounded with a sharp stone, at the time the flowers should appear. It is collected first in cocoa-nut shells, and thence transferred into large earthen jars, where it remains until it is nearly dried to a cake, when it is formed into rolls and wrapped up in leaves. This cylindrical form, however, is said to be owing to its being run into the joints of the bamboo, when it is in a liquid state.<sup>2</sup> It is brought to Europe packed in cases and boxes. It was first brought to Europe by the Dutch, about the middle of the seventeenth century.

*Qualities.* — Gamboge is inodorous, and nearly insipid; opaque, brittle, breaks with a vitreous fracture, and is of an orange-yellow colour. Its gravity is 1.221.<sup>3</sup> When heated it melts, and by increasing the heat burns with a white flame, and leaves a very light, spongy charcoal. When wetted, it stains the fingers a brilliant-yellow; and when triturated with water about two-thirds are dissolved, and a turbid, yellow solution produced. Alcohol dissolves nine parts in ten, sulphuric ether six parts; and both form transparent deep golden-yellow tinctures. Gamboge is also soluble in a strong solution of pure ammonia, and potass, forming deep orange-red solutions.

The watery solution of Gamboge reddens tincture of litmus; is not precipitated by alcohol, but, on the contrary, is rendered transparent by it; oxysulphate of iron strikes with it a pale olive-brown hue, but is not precipitated, nor is it affected by solutions of any of the other metallic salts. The alcoholic tincture is rendered turbid and bright-yellow by the addition of water, but the precipitate is long of being deposited. The ethereal tincture, when evaporated in water, leaves a pellicle of beautiful orange-coloured brittle resin, which imparts no colour to water. When water is added to the alkaline solu-

---

<sup>1</sup> The *Cambogia gutta* of Linnæus, several species of *Hypericum*, *Chelidonium laciniatum*, and several other plants, yield a yellow juice; but that the tree we have described affords the true gamboge was clearly established by Koenig, who resided many years at Tranquebar.

*White's Voyage to the China Seas*, Boston, 1823, p. 250.

<sup>3</sup> Brisson.



tions they are not even rendered turbid; but, on the addition of acids, yellow precipitates are produced which are soluble in an excess of acid. These experiments confirm the analysis of Braconnot, from which he concluded that Gamboge contains one part of a gum resembling cherry-tree gum, and four of a brittle resin; but they do not throw any light on the nature of its cathartic property.

*Medical properties and uses.*—Gamboge is a powerful, drastic cathartic; and frequently excites vomiting, even in moderate doses. It is efficaciously used in obstinate costiveness, in dropsies and for the expulsion of *tæniæ*; but, from the violence of its operation, and the griping it occasions, it requires to be exhibited with caution. As a hydragogue it is usually combined with squill and supertartrate of potass; and for cathartic purposes, with calomel, soap, or rhubarb. The alkaline solution of it has also been administered in dropsy, in which form it is said to operate both by stool and urine.

The dose of the alkaline solution is from  $\mathfrak{m}$  xxx to  $\mathfrak{m}$  l, in a cupful of water twice a day. It is, however, more usually given in substance, in the form of pills, in doses of from grs. ij. to grs. vj, variously combined.

Officinal preparations. *Pilulæ Cambogiæ compositæ*, L. E.

STANNUM.<sup>1</sup> PLUMBUM ALBUM. Κασσιτερον. Tin.

*Syn.* Etain (*F.*), Zinn (*G.*), Tin (*Dutch, Dan.*), Tenn (*Swed.*), Cyna (*Polish*), Stagno (*I.*), Estanno (*S.*), Estanho (*Portug.*), Olowo (*Russ.*), Galai (*Turk.*), Resass (*Arab.*), Rangha (*H.*), Tágárum (*Tam.*), Urzeez (*Pers.*), Tima (*Malay.*), Trapso (*Sans.*).

This metal is not very diffusely spread over the surface of the globe; but is very abundant in the places where it occurs. It is plentifully procured in Cornwall<sup>2</sup>; and mines of it are also wrought in the Erzgebirje, on the borders of Bohemia; in the province of Gallicia, in Spain; and in the peninsula of Malacca, in Asia. It is said also to have been found in Chili.

It occurs

A. In its metallic state:

i. United with sulphur and copper.

SP. 1. *Tin pyrites.*

B. Oxidized:

ii. Combined with oxide of iron and silex.

1. *Tin stone.*

2. *Wood tin.*

The purest and best tin of Cornwall is procured from the

<sup>1</sup> Beckman is of opinion that the *stannum* of the ancients was neither our tin nor a peculiar metal; but the regulus of lead, or *werk* of the Germans, as procured by the first smelting of the ore, which always contains silver. *Hist. of Inventions*, vol. iv. p. 7.

<sup>2</sup> It is uncertain when the Cornwall mines were first wrought. The Phœnicians came to Britain for tin; and there is a tradition in Cornwall, that the very old forsaken works were those of the Jews, and are therefore called *Attal saraxim*. The Jews' cast off works. *Norden's Description of Cornwall*, 4to. 1728, 41.



tin-stone, which is found beneath the beds of streams, particularly at Carn near Perran. The ore is found under a stratum of clay about fifty feet thick, and a layer of roundish stones; occurring loose in lumps and grains of various sizes. It is first washed, then bruised and passed through wire sieves containing sixteen meshes in the square inch; and next smelted with charcoal, in a common blast furnace, through a hole in the bottom of which the reduced metallic-tin flows into a pit below. To purify it, the tin in the fused state is ladled from the pit into an iron boiler placed over a fire; and pieces of charcoal plunged, by means of an iron instrument, to its bottom, which occasions an ebullition, and causes any slag it contains to rise to the surface, whence it is skimmed off. The tin is lastly cast into bars or pigs, weighing from two to three cwt. each.<sup>1</sup>

*Qualities.* — Tin has a very slight, somewhat disagreeable, taste, and emits a peculiar odour when rubbed. It has the colour and the brilliancy of silver; is very malleable; but has little ductility and tenacity; is easily cut with a knife; and is flexible, producing a peculiar crackling noise when bent backwards and forwards. Its specific gravity, after fusion, is 7.291; and after being hammered, 7.299.<sup>2</sup> It melts at 442 Fahr., but is not volatilized, and on cooling crystallizes in rhombs. Tin exposed to the air soon loses its lustre, and when melted it is oxidized, a grey oxide being formed on its surface, which becomes yellow if the heat be continued; and if raised to a full-red heat takes fire, and acquires a pure-white colour. It is also oxidized by many acids; and combines readily with sulphur.

*Officinal.* STANNUM: STANNI LIMATURA, *Lond. Edin. Dub.* The filings and powder of tin.

Tin melted and agitated briskly in a heated mortar with a warm pestle till it cools, or shaken in a wooden box, is reduced to a kind of powder, consisting of small rounded particles, with very little lustre; but still in the metallic state. This is the powder of the pharmacopœias.

*Medical properties and uses.* — Tin is anthelmintic. It was formerly used in hysteria and hypochondriasis; but its efficacy in these complaints cannot extend beyond its power of dislodging worms. It is generally supposed to operate mechanically only; but it has been suggested, that “it is not improbable, it may act by generating hydrogen gas in the intestinal canal, which proves noxious to the animal; and its efficacy has been said to be increased by combination with sulphur, by which sulphuretted hydrogen gas will be evolved.”<sup>3</sup> For

<sup>1</sup> *Aikin's Dictionary of Chemistry.*

<sup>2</sup> *Brisson.*

<sup>3</sup> *Murray. System of Materia Medica*, i. 490.



the mode of exhibiting it, see *Pulvis Stanni* among the Preparations.

Official preparations. *Pulvis Stanni*, D.

STAPHISAGRIÆ SEMINA. Vide *Delphinium*.

STYRAX. *Spec. Plant. Willd.* ii. 623.

Cl. 10. Ord. 1. Decandria Monogynia. Nat. ord. Bicornes, Linn. Guaiacinae, Juss.

G. 874. *Calyx* inferior. *Corolla* funnel-shaped. *Drupe* two-seeded.

Sp. 1. *S. officinale*.<sup>1</sup> Official Storax. *Med. Bot.* 2d edit. 291. t. 101.

Sp. 3. *S. Benzoin*. Benjamin-tree. *Med. Bot.* 2d ed. 294. t. 102.

*Phil. Trans.* lxxvii. 307.

#### 1. STYRAX OFFICINALE.

Official. STYRACIS BALSAMUM, Lond. STYRACIS OFFICINALIS BALSAMUM, Edin. STYRAX CALAMITA; RESINA, Dub. Storax Balsam.

Syn. Storax (F.), Storax (G., Dan., Swed.), Stryaxboom, (Dutch), Storace (I.), Azumbar (S.), Estoraque (Port.), Usteruk (Arab.).

The officinal Storax-tree is a native of the south of Europe and the Levant; and flowers in July. It rises about fifteen feet in height, sends off many branches, and is covered with a rough, grey bark. The leaves, which are about two inches long, an inch and a half broad, and of a bright green on the upper surface, and hoary on the under, are petiolate, alternate, elliptical, pointed and entire. The flowers are in terminal clusters; the corolla is monopetalous, funnel-shaped, large, and of a white colour: the filaments are placed in a regular circle, and apparently adhere at the base, supporting erect oblong anthers; and the germ is oval with a slender style and simple stigma. The fruit is a juiceless drupe, of an ovate globular form, containing one or two compressed angular nuts.

From this tree, Storax is obtained in Asiatic Turkey. It issues from incisions made in the bark; and as it was formerly the custom to collect and export it in reeds, it was named *Styrax calamita*.<sup>2</sup> But only two kinds of Storax are now found in the shops, Storax in the tear, which is pure Storax; and Storax in the lump, or red Storax, which is mixed with sawdust and other impurities. Both kinds are brought from the Levant in chests and boxes.

*Qualities*.—Storax has a fragrant odour, and a pleasant, sub-acidulous, slightly pungent, and aromatic taste. It is of a reddish-brown colour, and brittle at the ordinary temperature of the air, breaking with a shining resinous fracture; but soon softens between the fingers, and melts at a low heat, exhaling a strong odour of benzoic acid. In a higher degree of heat it burns with a white flame, and leaves behind a light spongy

<sup>1</sup> *Στυραξ* Dioscoridis.

<sup>2</sup> The Dublin college retains this appellation, although no storax of this description is now to be met with in the shops.



charcoal. It is deprived of its red colour, and the little transparency it possesses by chewing; and becomes in a remarkable degree, more brittle. To water it imparts a yellow colour, and its odour and taste; but in distillation scarcely any oil is obtained. Alcohol and ether dissolve it completely, leaving only impurities, and the tincture is decomposed by the addition of water. Its constituents are resin, an empyreumatic oil, and benzoic acid.

*Medical properties and uses.* — Storax is stimulant, and in some degree expectorant. It was formerly much prescribed in asthma, catarrh, phthisis, and menstrual obstructions: but it is now scarcely ever employed, except as an adjunct on account of its fragrance; and, certainly, it cannot be recommended in the above complaints. The dose is from grs. x. to ʒss.

Officinal preparations. *Styrax purificata*, D. *Pilulæ e Styrace*, D.

## 2. STYRAX BENZOIN.<sup>1</sup>

*Officinal.* BENZOINUM, *Lond.* STYRACIS BENZOINI; BALSAMUM, *Edin.* BENZOE; RESINA, *Dub.* Benzoin; a balsam.

*Syn.* Benzoin (*F.*), Benzoë (*G.*), Benzoin (*Dutch, Dan., Swed.,*) Belzuino (*I.*), Benjui (*S.*), Bejoim (*Portug.*), Lubán (*H.*), Liban (*Arab.*), Malacca Sambranie (*Tam.*), Menian (*Jav.*).

The Benzoin or Benjamin tree, is a native of Sumatra. It is a tall tree sending off many round branches, which are covered with a whitish downy bark. The leaves are alternate, on round, striated, tomentose petioles, oblong, quite entire, pointed, veined, smooth above and tomentose beneath. The flowers are in compound, axillary clusters, nearly as long as the leaves; with the common peduncle tomentose; the partial alternate, spreading and tomentose; the pedicles short; and the flowers all hanging on the same side. The calyx is bell-shaped, short, and downy; the corolla consists of five, linear, obtuse petals, four times longer than the calyx, connected together at the base, and externally cineritious. The filaments are ten, a little shorter than the petals, connected at the base, and supporting linear, erect anthers: the germen is superior, ovate, and downy, with a slender style and double stigma.

Benzoin is obtained from this tree by wounding the bark near the origin of the lower branches. The tree is never wounded under six years of age; and cannot sustain these annual incisions above twelve years. A tree yields about three pounds of balsam annually. It is brought to this country in large masses packed in chests and casks. The

---

<sup>1</sup> Benzoin was long supposed to be the produce of a species of *Laurus*, a native of Virginia, which on this account was named *L. Benzoin*: this error was detected by Linnæus; but the real genus of the plant which yields it, was first assigned by Mr. Dryander. *Phil. Trans.* lxxvii. 307.



best is of a yellowish-brown colour, studded with white spots; the worst is blackish and full of impurities.

*Qualities.* — Benzoin has a very agreeable fragrant odour<sup>1</sup>; but scarcely any taste. Its specific gravity is 1·092. The mass is white and brownish, or yellowish, somewhat translucent, brittle, and breaks with a resinous fracture. When heated it exhales white fumes of a very fragrant, pungent odour, which are volatilized benzoic acid. Boiling it in water, lime and water, or solutions of the fixed alkalies, extracts the benzoic acid it contains, which can be afterwards separated by the addition of an acid. Sulphuric acid dissolves benzoin, and benzoic acid sublimes. Nitric acid assisted by heat dissolves it with violence, and the solution, on cooling, becomes turbid, and crystals of benzoic acid separate. In both these processes artificial tannin also is formed. Alcohol and ether readily dissolve this balsam, which is again precipitated from the tincture, in the form of a white powder, by the addition of water. According to Mr. Brande's analysis, 100 parts of benzoin distilled alone, yield of benzoic acid 9·0. Acidulous water 5·5. Butyraceous empyreumatic oil 60·0. Charcoal 22·0. Carburetted hydrogen and carbonic acid 3·5 parts.<sup>2</sup>

*Medical properties and uses.* — Benzoin is regarded as expectorant: and was formerly employed in asthmas and other pulmonary affections; and is still used for that purpose by the Tamool physicians; but it is scarcely ever ordered in modern European practice; and is principally used for preparing the acid.<sup>3</sup>

Officinal preparations. *Acidum benzoinum*, L. E. D. *Tinctura Benzoini composita*, L. E. D.

**SUCCINUM.**<sup>4</sup> *Lond. Edin. Dub.* Amber.

*Syn.* Succin (*F.*), Bernstein (*G.*), Barnsteen (*Dutch*), Bernsteen (*Dan.*), Bernsten (*Swed.*), Bursztyn (*Polish*), Ambra (*I.*), Sucino (*S.*), Alambar (*Port.*), Iantar (*Russ.*), Amber (*Tam.*), Karooba (*Pers.*), Kepoor (*H.*), Ambar (*Malay*), Kernulbehr (*Arab.*).

This substance is dug out of the earth in Ducal Prussia, near the sea coast; and is thrown in considerable quantity on the sea shore of Polish Prussia and Pomerania, particularly after tempestuous west or north-west winds.<sup>5</sup> It is evidently

<sup>1</sup> It is employed by the Malays to perfume their temples.

<sup>2</sup> *Nicholson's Journ.* x. 86. *Thomson's Chemistry*, 4th ed. v. 129.

<sup>3</sup> Dr. Paris (*Pharmacologia*) gives the following formula for preparing *Fumi, gating pastiles*, R *Benzoini* ʒj., *Cascarillæ* ʒss., *Myrrhæ* ʒj., *Olei nucis moschaet-olei Caryoph.* āā grs. x., *Potassæ Nitratiss* ʒss., *Carbonis ligni* ʒvj., *Muc. G. Tragacanthæ*, q. s.

<sup>4</sup> Ηλεκτρον Græcorum.

<sup>5</sup> It is found in small quantities on the east coast of Britain; and small pieces of it are occasionally found in the gravel-pits round London. The largest mass of amber ever found, was met with near the surface of the ground in Lithuania. It weighs 18lbs. and is preserved in the royal cabinet at Berlin.



of vegetable origin, and the lumps occasionally inclose small pieces of twigs and insects in their substance. The greater part of what is brought to this country comes from the Baltic; and a small quantity from Catania, in Sicily, packed in chests.

*Qualities.* — Amber is insipid, and also inodorous, except when heated, when it emits a fragrant odour. It is brittle, light, hard, and transparent, with a considerable degree of lustre; is commonly of a golden-yellow or brown colour, but occasionally colourless, and is electric. Its specific gravity is 1.065. It softens when heated, and in a strong heat burns, leaving few ashes. It is nearly insoluble in water, to which, however, it gives, when it is boiled with it, a sweetish odour, a yellow colour, and a bitter taste. Alcohol takes up about one-eighth part of amber, forming a tincture which is rendered milky by the addition of water, and precipitates a resin. The alcoholic solution contains a free acid. With boiling fixed-alkalies it forms a soap; and even a cold, weak solution of potass dissolves it, requiring, however, a considerable length of time. Sulphuric acid converts amber into a black resinous mass. Nitric acid, assisted by heat, acts violently upon it; nitrous gas is emitted, and the amber is ultimately entirely dissolved. Its constituents appear to be chiefly resin, an empyreumatic oil, and succinic acid. It is distinguished from resins by its higher refractive power, and the peculiar smell which it emits when rubbed or held near a hot iron.

*Use.* — Amber, although it was in high estimation among the ancients as a medicine, is now only used in pharmacy for the purpose of obtaining the oil and acid which it yields by distillation.

Officinal preparations. *Acidum Succinicum*, E. D. *Oleum Succini*, L. E. D.

SULPHUR. *Lond.* SULPHUR SUBLIMATUM. *Lond. Edin. Dub.* Sulphur. Sublimed Sulphur.

*Syn.* Soufre (*F.*), Schwefel (*G.*), Zwavel (*Dutch*), Zolfo; Solfo (*I.*), Azufre (*S.*), Euxofre (*Port.*), Svovel (*Dan.*), Svavel (*Swed.*), Siarka (*Polish*), Sera (*Russ.*), Kibreet (*Arab.*), Ghéndaguno (*Tam.*), Gandhaca (*San.*), Gowgird (*Pers.*), Blerong (*Malay*), Gundaka (*Cyng.*).

Sulphur is found native in the neighbourhood of volcanoes; and sometimes, although rarely, in veins traversing primitive rocks.<sup>1</sup> At the Solfatara near Naples, it is dug up in a state of comparative purity, being mixed with a white earth only, from which it is separated by sublimation, and the sulphur thus freed is melted, and cast into moulds forming the roll

---

<sup>1</sup> As for example in the sulphur mountain of Ticsan, in South America, where the sulphur is disseminated in small masses, having no contiguity with one another in the quartz which traverses the mica slate, parallel to its strata. The sulphur is dull, friable, and breaks with the slightest blow. *Ann. de Chim.* xxvii.



sulphur of commerce. It is imported into Britain chiefly from Sicily<sup>1</sup> and Naples; but a large proportion of what is used in this country is obtained from the roasting of pyrites. At the Pary's mines in Anglesea were works for this purpose on a large scale; where, in working the copper pyrites, the sulphur volatilized in the roasting, was collected in chambers, which were connected with the domes of the furnaces by means of horizontal flues. Each chamber had a door, by means of which it was cleared of the sulphur once in six weeks. This is the general mode of obtaining sulphur from pyrites, and thus procured, it is in rough, pulverulent, spongy crusts, of a dirty grayish-yellow colour. In order to purify it, the crusts are broken and thrown into a boiler, in which it melts; and after the impurities are separated by skimming and subsidence, it is cast into cylindrical moulds, forming roll sulphur; or into cones about two feet high, which form the loaf sulphur of commerce.<sup>2</sup>

The common English roll sulphur is said often to contain a full fifteenth part of orpiment, while the Sicilian sulphur contains seldom more than 3 per cent. of a simple earth; and therefore is justly preferred. Both of them are purified in the large way by conducting the vapour of melted sulphur into close chambers, where it concretes in the form of a fine powder: but for medicinal use, that which is sublimed by heating in a sand-bath, an earthen cucurbit, charged with roll sulphur, and conveying the vapours to be concentered into a set of alludels placed round the cucurbit, is to be preferred. Prepared in either mode, it is the *Sulphur sublimatum*<sup>3</sup> of the pharmacopœias. Sulphur, especially in the roll, contains many impurities, the amount of which may be detected by boiling 100 grains in four ounces of oil of turpentine, pouring off the solution while hot, and repeating the boilings, until no more is dissolved. The residue indicates the amount of the impurities.<sup>4</sup>

*Qualities.* — *Roll sulphur* is a crystallized, brittle, solid body of a yellow colour, has a peculiar well-known odour when rubbed or heated, and is insipid. It breaks from the heat of the hand, when held in it for a short time; and being a non-conductor of electricity, becomes electrical when rubbed. Its specific gravity is 1.99. *Sublimed sulphur* is in the form of a very bright-yellow powder, and contains a minute portion of sulphuric acid, from which it can be separated by washing it with water. Sulphur volatilizes under 220° Fahr., at which

<sup>1</sup> In Sicily it is procured from *Samattino, Gallati, Trabria, Pentellaria, Licati, Salato, Palmo, Tavera, Girgenti, and Falconara*.

<sup>2</sup> The *sulphur vivum* of the shops is the impure dregs of this process.

<sup>3</sup> Θειον πεπυρρουμενον Dioscoridis.

<sup>4</sup> *Brande's Manual*, p. 178.



it fuses, and, what is singular, by increasing the heat to  $320^{\circ}$ , it becomes thick and viscid, and if then poured into water, it assumes a red colour and ductility like wax<sup>1</sup>; while its specific gravity is increased to 2.325. At  $560^{\circ}$  it becomes an elastic fluid. When heated in the air it inflames at  $300^{\circ}$ , and burns with a pale blue flame, emits pungent suffocating vapours, and becomes acidified. It is insoluble in water; but soluble in a small degree in alcohol, ether, and oils; and combines with the alkalies, and many of the earths and metallic substances. The experiments of Davy led to the supposition that sulphur is a triple compound of oxygen, hydrogen and a peculiar base<sup>2</sup>, but it is still an undecomposed substance.

*Medical properties and uses.* — Sulphur is laxative, and a stimulating diaphoretic. From the gentleness of its operation on the bowels, it is one of the best means of keeping them lax in hæmorrhoidal affections; and the diaphoresis which it at the same time excites has rendered it serviceable in chronic rheumatism and catarrh, and in atonic gout, rickets, asthma, and other pulmonary affections not attended with acute inflammation. It is supposed that it combines with hydrogen in the stomach. It manifestly transpires through the skin, perhaps, however, in the state of sulphuretted hydrogen, which may be the cause that silver is blackened when kept in the pockets of those who take sulphur. It is specific in scabies and some other cutaneous affections, in which it is applied externally, and taken internally at the same time.

The dose may be from ʒj to ʒiij, mixed into an electuary with syrup or treacle, or in milk. To promote its purgative power it may be combined with super-tartrate of potass; and in hæmorrhoidal cases with magnesia.

*Officinal preparations.* *Sulphur lotum*, L. E. D. *Sulphur præcipitatum*, L. *Unguentum Sulphuris*, L. E. D. *Unguentum Sulphuris compositum*, L.

**SUPER-TARTRAS POTASSÆ IMPURUS**, *Edin.*  
**TARTARUM**, *Lond.* Impure Super-tartrate of potass.

*Syn.* Tartre (*F.*), Roher Weinstein (*G.*), Wynsteen (*Dutch*), Winsten (*Swed.*), Tartaro bianco (*I.*), Tartaro (*S.*).

This is a saline matter, which exists in the juice of the grape; and is deposited on the sides of casks of wine, in the form of a crust, during the continuance of the slow fermentation which goes on in wine, till it attains the greatest perfection age can give it. It is well known by the name of *tartar*, and is named red or white tartar, as it is more or less coloured, owing to the nature of the wine from which it is deposited. Besides colour-

<sup>1</sup> In this state it is kneaded under the water, and used for receiving the impressions of seals and medals.

<sup>2</sup> *Phil. Trans.* 1809.



ing matter, it contains extractive, potass combined with tartaric acid in excess, and tartrate of lime, It is only used for preparing the next article.<sup>1</sup>

SUPER-'TAR'TRAS POTAS'SÆ, *Edin.* POTASSÆ SUPERTARTRAS, *Lond.* TARTARUM CHRYSTALLI, *Dub.* Super-tartrate of potass, Crystals of tartar.

*Syn.* Tartrate acidule de Potasse (*F.*), Weinsteinrahm (*G.*), Wynsteenroom (*Dutch*), Cremore di tartaro; ossitartrato ossidulo di Potassa (*I.*).

This is the above-mentioned saline crust purified. It is first reduced to powder, then dissolved in boiling water in tubs, and the clear fluid poured off from the sediment. The clear solution is then allowed to remain at rest, when it deposits brown crystals of tartrate of potass, which are boiled in copper vessels with the mother liquor; and clarified by throwing in whites of eggs, and some finely sifted wood ashes. An effervescence immediately takes place, and a red scum is thrown up, which is carefully skimmed off with a perforated skimmer; and the throwing in of the wood ashes, with the subsequent skimming, are repeated for fourteen or fifteen times; after which the liquor is taken from the fire, and allowed to remain at rest for three days. On the fourth day, a dirty-white, saline crust is removed from the surface, and two thirds of the liquor ladled out. The crystals which now form are white and clean, and require no further preparation than drying on a wicker frame. In some places, instead of wood ashes, a portion of pure clay is diffused through the boiling solution. The exposure of the crystals on cloths to the air and light whitens them very considerably.<sup>2</sup>

*Qualities.* — Super-tartrate, or more properly bitartrate, of Potass is inodorous, and when allowed to dissolve in the mouth, which it does very slowly, and feeling gritty under the teeth, has a harsh, acid taste. Its crystals are small and irregular, generally run together into little masses, which are of a white colour, semi-transparent, brittle, and easily reduced to powder. Its specific gravity is 1.953. It requires for its solution 30 parts of boiling water, and 120 of cold water. The solution decomposes spontaneously by keeping; a mucous matter is deposited, and there remains a solution of carbonate of potass coloured with a little oil.<sup>3</sup> According to the analysis of Thenard, 100 parts of pure Supertartrate of Potass contain 57

<sup>1</sup> A large quantity of tartar comes from Sicily: the white is procured chiefly from *Palermo*, *Marsala*, and *Catania*; the red from *Mascali*, *Messina*, *Melazzo*, and *Vittoria*.

<sup>2</sup> Schaub says, it may be purified by simply boiling it with powdered recent charcoal, and very white crystals obtained. *Annales de Chimie*, xlix. 64.

<sup>3</sup> This decomposition was first described by Berthollet, in 1782. *Mem. Par. Thomson's Chemistry*, 4th edit. vol. iii. 95.



of tartaric acid, 33 potass, and 7 of water<sup>1</sup>; according to Burzelius, the proportions are, acid 70·45, potass 24·80, and water 4·75; while Mr. Brande states them to be acid 73·6, potass 26·4; and remarks that it is doubtful whether this salt contains any definite water of crystallization.<sup>2</sup>

*Medical properties and uses.* — This salt is purgative, diuretic, and refrigerant. As a purgative it is frequently employed, on account of its taste being less unpleasant than the generality of saline cathartics; but it is apt to excite too much absorption when long used, producing emaciation, and also disordering the digestive organs. This property, however, of exciting the action of the absorbents, is taken advantage of with great effect in the cure of dropsy, particularly ascites; in which the Supertartrate of Potass has been found extremely efficacious. It generally occasions a considerable discharge of serous fluid into the bowels, which is thrown off in the form of watery stools, at the same time that the discharge by urine is much augmented. The water in the cavity of the abdomen is thus rapidly carried off: and the chances of a return of the disease are supposed to be fewer than when other diuretics are employed. We are of opinion, however, that in cases complicated with hepatic obstructions the effects of this remedy are very uncertain. It may be advantageously united with squill; and, owing to the exhaustion it occasions, its use should be followed by preparations of iron, and other tonics. As a refrigerant, Super-tartrate of Potass dissolved in water, and the solution sweetened with sugar is a pleasant beverage in febrile diseases, when its purgative quality is not likely to prove injurious. As a purgative and hydragogue the dose is from ʒiv to ʒvj, in the form of electuary; and this dose for the latter purpose must be repeated until the kidneys are affected diluting freely during its use.

Officinal preparations. *Ferrum Tartarizatum*, L. D. *Potassæ Tartras*, L. E. D. *Antimonium Tartarizatum*, L. E. D. *Soda Tartarizata*, L. E. D.

SUS. *Syst. Nat. Gmelin*. i. 217.

Cl. 1. Ord. 6. Mammalia Belluæ.

G. 35. *Fore-teeth* four in the upper jaw converging, and six prominent in the lower jaw. *Tusks* two shorter in the upper jaw; and two in the under jaw displayed. *Snout* truncated, prominent, moveable. *Feet* cloven.

*Species* 1. *Sus Scrofa*.<sup>3</sup> The Hog. *Jonst. quadr.* 99. t. 47.

*Officinal.* A'DEPS, *Lond.* ADEPS SUILLUS, *Edin. Dub.* Fat Hog's lard.

*Syn.* Sain doux (*F.*), Schweineschmalz (*G.*), Swinister (*Swed.*), Szmalee (*Pol.*), Lardo (*I.*), Pingue (*S.*), Púnnie Colupoo (*Tam.*).

<sup>1</sup> *Annales de Chimie*, vol. xxxviii. p. 59.

<sup>2</sup> *Manuel*, p. 142.

<sup>3</sup> 'Tis Aristotle,



The hog is too well known to require a particular description. It is an inhabitant of the greater part of the temperate regions of the globe, the wild and the domestic being varieties of the same species; and of both there are several sub-varieties. The period of gestation of the sow is four months, and the offspring numerous, occasionally exceeding twenty at a litter, which the boar sometimes devours. The hog does not shed its teeth, and seldom lives beyond twenty-five or thirty years. It is much infested with vermin; and is subject to several diseases, particularly hydatid dropsy, scrophula, and scabies. Its food is of a vegetable nature; but it is asserted that pepper kills it. As an article of diet, the flesh of the hog, when the animal has been castrated and properly fed<sup>1</sup>, is very palatable, and not unwholesome; and when salted, keeps better than most other meats. But the frequent use of pork is said to favour obesity, and occasion disorders of the skin, particularly in the sedentary. The lard, which is the officinal part of the hog, is obtained chiefly from the flank of the animal. To free it from the membranes and vessels, it is cut in small pieces, then very well washed in water, until the water comes off colourless, and afterwards melted with a very gentle heat in a shallow vessel, continued on the fire till the whole of the water is evaporated. While in the melted state, it is run into bladders, in which it concretes; and is thus brought to market.

*Qualities.*—Lard is inodorous, tasteless, and white; soft, and nearly semifluid. Exposed to a heat of 97° it melts, and concretes again when cooled. It is insoluble in water, alcohol, and ether: but is dissolved by the strong acids, being at the same time decomposed; and, like the fixed oils, it combines with the alkalies and forms soap. It is oxidized, if, when melted, a little nitric acid be stirred into it; and assumes a greater degree of firmness, with a yellow colour. By destructive distillation it affords results very similar to those obtained from the analysis of fixed oil; and appears to be a compound of oxygen, hydrogen, and carbon in unknown proportions. From the experiments of Chevreul, it appears to consist of a mixture of two distinct oily substances; one of which is solid at the usual temperature, and has been named by him *stearin*; and the other liquid, which he has named *elain*. The proportion in 100 parts of lard, is, of stearin 38, elain 62 parts.<sup>2</sup> When lard is long exposed to a warm air, it becomes yellow, emits

---

<sup>1</sup> The qualities of the flesh depend much on the diet of the animal. Pork fed at a flour mill is always good: and Russel says, that which is fed near Aleppo on liquorice root, which grows in great abundance in the desert, is fat, delicious, and remarkably digestible.

<sup>2</sup> *Annales de Chimie*, t. 94. p. 129.



a foetid odour; and, owing to oxygen being attracted from the atmosphere, the sebatic acid is formed. This state of rancidity may, in some degree, be removed by washing it with very pure soft water; which during the operation becomes acid, and reddens litmus paper.

*Medical properties and uses.* — Lard is emollient; and owing to its softness and unctuousity is preferable to fat as a friction, but it is seldom used for this purpose; and is chiefly employed in the formation of ointments.

Official preparations. *Adeps preparata*, L. D.

SWIETENIA. *Spec. Plant. Willd.* ii. 557.

*Cl.* 10. *Ord.* 1. Decandria Monogynia. *Nat. ord.* Trihilatæ, Linn. Meliæ, Juss.

*G.* 843. *Cal.* five-cleft. *Pet.* five. *Nect.* cylindric, bearing the anthers at the mouth. *Capsules* five-celled, woody, opening at the base. *Seeds* imbricate winged.

*Sp.* 2. *S. Febrifuga*. Febrifuge Swietenia. *Roxburgh Coromandel Plants*, i. 18. t. 17.

*Official.* SWIETENIA FEBRIFUGA; CORTEX, *Dub.* The bark of Febrifuge Swietenia.

*Syn.* Rahuna (*H.*), Shemmairum (*Tam.*), Patranga (*San.*), Rohun (*Malay*).

This is a native of the East Indies, growing in various places, particularly in the vicinity of Omedurdah, not far from the cantonment of Hazareebagh; and at Rohun, from which it is known in India by the name of Rohun bark. It is also a native of Siam. It is a very lofty tree, with a large, shady head, and covered with a gray, scabrous bark. The leaves are alternate and abruptly pinnated; composed of three or four pairs of opposite, petiolated, oval, obtuse leaflets; each from three to five inches long, and from two to three broad, smooth, shining, and the lower side extending a little further down on the petiole than the upper side. The flowers are middle-sized, white, inodorous; and disposed in large, terminating, diffused panicles, furnished with minute bractes: the calyx is inferior; the nectary scarcely half the length of the petals and bellied; the filaments are very short, and inserted just within the mouth of the nectary; the germ is conical, bearing a thick tapering style, supporting a large, targeted stigma, which shuts up the mouth of the nectary; the capsule is large, ovate, and five-valved, with the valves gaping from the top. The bark is collected when the sap begins to ascend freely, at which period it readily separates. The bark of the small or middling sized branches are the best, and it is fit for use as soon as it is sufficiently dry for powdering.<sup>1</sup>

*Qualities.* — The bark of febrifuge Swietenia is very bitter

<sup>1</sup> Vide an Essay by Mr. Breton on this bark, in the *Medico-Chirurg. Trans.* vol. xi. p. 524.



and austere, yet it is not nauseous. It is brittle, compact, of a light-red colour internally, and externally covered with a rough, grey, inert epidermis. Water extracts its virtues both by infusion and coction; they are partially taken up by alcohol, and the aqueous and spirituous preparations do not suffer decomposition when mixed.<sup>1</sup>

*Medical properties and uses.* — The bark of this species of *Swietenia* is astringent and tonic. In India, it is used for the cure of intermittents with considerable advantage, and has also been found efficacious in most of the diseases in which the *Cinchona* bark proves serviceable. It is very little employed in Europe, and is not found generally in the shops. The dose, in substance, pulverized, is half a drachm.

**TAMARINDUS.** *Spec. Plant. Willd.* iii. 577.

*Cl.* 16. *Ord.* 1. Monadelphia Triandria. *Nat. ord.* Lomentaceæ, *Linn.* Leguminosæ, *Juss.*

*G.* 1250. *Calyx* four-parted. *Petals* three. *Nectary* of two short bristles under the filaments. *Legume* pulpy.

*Sp.* 1. *T. indica*.<sup>2</sup> The Tamarind tree. *Med. Bot.* 2d edit. 448.

*t.* 161. (Balam-pulli) *Rheede, Hort. Malab.* i. 39. *t.* 23.

*Officinal.* TAMARINDI PULPA, *Lond.* TAMARINDI INDICÆ FRUCTUS, *Edin.* TAMARINDUS; FRUCTUS, *Dub.* The pulp, or preserved fruit of the Tamarind.

*Syn.* Tamerins (*F.*), Tamarinden (*G.*, *Dutch*), Tamarindo (*I.*, *S.*), Tamarinho (*Port.*), Poollie (*Tam.*), Ambala (*Cyng.*), Timir hindie (*Pers.*), Assam Jaba (*Malay*), Umblie (*H. and Arab.*), Amlica (*San.*), Kamal Assam (*Jav.*).

This tree is a native of the East and West Indies, of Arabia, and Egypt. It is a large, beautiful, spreading tree.<sup>3</sup> The leaves are abruptly pinnate, composed of sixteen or eighteen pairs of sessile leaflets, half an inch only in length, and one-sixth of an inch broad, of a bright-green colour, downy, oblong, entire and obtuse; the flowers are in loose bunches of five or six, which come out from the sides of the branches; the calyx is of a straw-yellow colour and deciduous: the petals are also yellowish, and beautifully variegated with red veins; ovate, concave, acute, indented, and plaited at the edge; and the filaments purplish, bearing incumbent, brownish anthers: the pods are thick, compressed, and of a dull-brown colour when ripe; those from the West Indies are from two to five inches long, with two, three, or four seeds; those from the East Indies are twice as long, and contain five, six, or seven seeds; the seeds

<sup>1</sup> The wood of the tree, when treated in the same manner as the *Acacia catechu*, to obtain the Catechu, yields an extract which has the closest affinity to Kino, and may be substituted for it. *Medico-Chirurg. Trans.* vol. xi. p. 328.

<sup>2</sup> Ὁξύ φοίνικα Nicolai Myrepsici, the last of the Greek physicians.

The natives of India think that it is dangerous to sleep under a Tamarind tree, during the night.



in both are flat, angular, shining, and lodged in a dark, pulpy matter.

In the West Indies, the pods are gathered in June, July, and August, when fully ripe; and the fruit, being freed from the shelly fragments, is placed in layers in a cask, and boiling syrup poured over it, till the cask is filled; the syrup pervades every part quite down to the bottom, and when cool the cask is headed for sale.<sup>1</sup> The East India tamarinds are darker coloured and drier, and are said to be preserved without sugar. When tamarinds are good, they are free from any degree of mustiness; the seeds are hard, flat, and clean; the strings tough and entire; and a clean knife thrust into them does not receive any coating of copper. They should be preserved in closely-covered jars.

*Qualities.* — Tamarinds are inodorous, and have an agreeable, acid, sweetish taste. According to the analysis of Vauquelin, the pulp contains, independent of the sugar with which it is mixed, supertartrate of potass, gum, jelly, citric acid, tartaric acid, malic acid, and a feculent matter. The acid taste chiefly depends on the citric acid, the quantity being greater than that of the other; ℥xvj. of the prepared pulp containing ℥jss. of citric acid, but only ʒij. of tartaric acid, ℥ss. of supertartrate of potash, and ʒss. of malic acid.

*Medical properties and uses.* — Tamarind pulp is refrigerant, and gently laxative. The simple infusion of the pulp in warm water, or a whey made by boiling ʒij. of it in two pints of milk, and straining, form very grateful, refrigerant beverages, which are advantageously used in febrile diseases. The dose of the simple fruit required to act upon the bowels is so large, that it is seldom given alone as a purgative, but is generally combined with cassia or manna, the action of which it augments; or with such of the neutral purgative salts as are not decomposed by it; which is the case with those that have potass for their base, and are therefore incompatible in mixtures with this fruit. It forms an agreeable addition to infusion of senna; but the purgative power is weakened by it.

Officinal preparation. *Infusum Tamarindi cum Sennæ*, E. D.

TANACETUM. *Spec. Plant. Willd.* iii. 1809.

Cl. 19. Ord. 2. Syngenesia Superflua. Nat. ord. Compositæ Discoideæ, Linn. Corymbiferae, Juss.

\* *Discoid.*

G. 1472. *Receptacle* naked. *Pappus* sub-marginate. *Calyx* imbricate, hemispherical. *Calyx* rays obsolete<sup>2</sup>, trifid.

*Species* 18, *T. vulgare*.<sup>3</sup> Common Tansy. *Med. Bot.* 2d edit. 67. t. 27. *Smith Flora Brit.* 862. *Eng. Bot.* 1229.

<sup>1</sup> Long's *Jamaica*, iii. 729.

<sup>2</sup> *Radius calidiore æstate prodit.* Willdenow, l. c.

<sup>3</sup> Ἀρτεμισία λεπτοφύλλος Dioscoridis.



*Officinal.* TANACETI VULGARIS FLORES FOLIA, *Edin.* TANACE-TUM, FOLIA, *Dub.* The leaves of Common Tansy.

*Syn.* Tanassie (*F.*), Rheinfarn (*G.*), Wormkruid (*Dutch*), Rhreinfa (Dan.), Renfana (*Swed.*), Kiviat wrotyczowy (*Pol.*), Tanaceto (*I.*), Atanasia (*S.*), Tanasia (*Port.*) Dikaja riabina (*Russ.*).

This is an indigenous perennial plant, growing on hills, and by the sides of roads and fields; flowering in July and August; but it is generally cultivated for medicinal and culinary purposes. The root is creeping, sending up stiff, erect stems, about two feet in height, leafy, obscurely hexagonal and striated; with alternate leaves, doubly pinnatifid, acutely cleft, somewhat downy on the under side, eared at the base and embracing the stem; the flowers are in terminal, dense corymbs, of a bright-yellow colour, and flattish: the leaflets of the calyx are obtuse, with a dry, scaly margin: the florets are numerous; those of the disc hermaphrodite and five-cleft, those of the margin female and trifid; the seeds are small, uniform, inversely pyramidal, pentagonal, ribbed, of an ash-colour, and crowned with a narrow, marginate, membranous pappus.

*Qualities.* — Tansy has a strong, peculiar, fragrant odour, and an acrid, bitterish taste, somewhat resembling that of camphor. These qualities it yields both to water and alcohol; and in distillation with water affords a greenish-yellow essential oil, which has in perfection the odour of the plant, and probably contains camphor.

*Medical properties and uses.* — The leaves and flowers of Tansy are tonic and anthelmintic. It was formerly regarded as a powerful remedy in intermittents, dropsy, hysteria, and obstructed menstruation; but, experience, and the knowledge of better remedies, have set aside its use in these diseases. An infusion of the whole herb in boiling water has been highly extolled as a preventive of the return of gout<sup>1</sup>; but it is now scarcely ever used, except as an anthelmintic for expelling lumbrici, to which it has certainly some pretensions. The dose of the leaves in powder is from ʒj. to ʒj., twice a day.

TEUCRIUM. *Spec. Plant. Willd.* iii. 13.

Cl. 14. Ord. 1. Didynamia Gymnospermia. Nat. ord. Verticillatæ, *Linn.* Labiatæ, *Juss.*

G. 1093. Corolla no upper lip, but a fissure in place of it. Stamens protruded.

Species 12. *T. Marum*. Common Marum. *Med. Bot.* 2d edit. 324. t. 115.

Species 36. *T. Chamædrys*. Wall Germander. *Med. Bot.* 2d edit. 358. t. 130.

<sup>1</sup> Clarke, *Essays Physical and Literary*, iii. 438.



1. TEUCRIUM MARUM.<sup>1</sup>

*Officinal.* MARUM SYRIACUM; HERBA, *Dub.* The herbaceous part of Common Marum.

*Syn.* Herbe au Chat (*F.*), Arisberkraut (*G.*), Maro (*I., Port.*), Amaro (*S.*).

This plant is perennial, a native of Spain and Syria, but cultivated in our gardens.<sup>2</sup> It has a low, shrubby stalk, sending out many, woody, hoary branches; and, in its proper soil and climate, rises three or four feet in height. The leaves are small, and placed opposite at each joint, pointed, and sometimes nearly three-lobed; green on the upper surface, and hoary beneath: the flowers, which are in loose, terminal, whorled spikes, are very downy, and of a bright-red colour.

*Qualities.*—The leaves rubbed between the fingers have a volatile aromatic odour, which readily excites sneezing; their taste is bitterish, pungent and acrid, depending on a volatile oil, which can be obtained separate by distillation with water.

*Medical properties and uses.*—This plant is an useful er-rhine; and as it possesses no narcotic property, is in some cases preferable to tobacco. It is generally a component in sternutatory powders.

*Officinal preparation.* *Pulvis Asari compositus*, E. D.

2. TEUCRIUM CHAMÆDRYS.<sup>3</sup>

*Officinal.* CHAMÆDRYS HERBA, *Dub.* Wall Germander.

*Syn.* Petite Chêne (*F.*), Bathengel (*G.*), Gamander (*Dutch*), Ekeyräs (*Swed.*), Ozanka (*Pol.*), Camedrio (*I.*), Camedrey de agua (*S.*), Carvalhinha (*Port.*).

This is an indigenous perennial plant, growing on old<sup>4</sup> ruins and walls; flowering in June and July. It has a creeping root: the stems are nearly erect, branched, round, leafy and hairy: the leaves subovate, cut, crenate, hairy, veined and attenuated at the base: the flowers are axillary: the calyx is rough, with pointed segments: the corolla of a purple colour, bilabiate, with the upper lip short and cut in the middle, and the lower separated into spreading lobes, the central of which is large and roundish.

*Qualities.*—The recent leaves have a slight aromatic odour, which is dissipated by drying; their taste is moderately bitter. Water extracts its active matter completely; alcohol only partially.

*Medical properties and uses.*—Wall germander has been accounted tonic, stomachic, diuretic, and emmenagogue; and is said to prove efficacious in uterine obstructions, agues,

<sup>1</sup> Μάρων Dioscoridis.

<sup>2</sup> It appears to have been cultivated in Britain so early as 1640.

<sup>3</sup> Χαμαίδρυς Dioscoridis.

<sup>4</sup> Winchelsea Castle. Walls of Norwich, near Magdalen gate. *Smith.*



gout, and rheumatism<sup>1</sup>; but it is, perhaps, not improperly neglected, being scarcely ever ordered. The dose of the pulverized herb may be from grs. x. to ʒj, given three or four times a day.

**TOLUTANUM BALSAMUM**, *Lond. Dub.* **TOLUIFERÆ BALSAMI BALSAMUM**, *Edin.* Balsam of Tolu.

*Syn.* Beaume de Tolu (*F.*), Tolutanischer Balsam (*G.*), Balsama Tolutano (*I.*), Balsamo de Tolu (*S.*).

The tree which yields the Balsam of Tolu has been ascertained<sup>2</sup> to be the *Myroxylon Peruiferum*, the same from which the Balsam of Peru is procured; I have therefore rejected the *Toluifera balsamum*; which must henceforth be regarded merely as a synonyme of the *Myroxylon*. The Tolu Balsam is the white Balsam of Peru hardened, by exposure to the air. It is obtained from incisions of the bark, from which it flows freely in hot weather; and is afterwards put into mats and calabashes to condense and harden, in which state it is brought to this country.

*Qualities.* — Balsam of Tolu has an extremely fragrant, lemon odour, and a warm, somewhat sweetish taste. It is of a yellow, reddish-brown colour, and of a thick tenacious consistence, becoming solid and brittle by age. Exposed to heat it melts, easily inflames, and disperses, along with its peculiar odour, that of benzoic acid. In distillation with water, it yields a small portion of volatile oil, impregnates the water with its odour, and by continuing the process benzoic acid sublimes. It is soluble in alcohol, forming a tincture which is rendered milky by water, but no precipitate falls. Mr. Hatchett found, that when it is dissolved in the smallest quantity of solution of potass, its own odour is lost, and it acquires a permanent, fragrant smell, resembling that of the Clove-pink. When digested in sulphuric acid, a considerable quantity of pure benzoic acid sublimes; and the same occurs during its solution in nitric acid, which also evolves traces of prussic acid.

*Medical properties and uses.* — Tolu Balsam is a stimulating expectorant; and although less heating than the other balsams, is nevertheless improper in pulmonic affections attended with inflammation. It forms an elegant addition to more active medicines in cases of asthma and chronic catarrh: and on the whole is more employed on account of its agreeable flavour, than for any efficacy it possesses. The dose of the balsam may be from grs. v. to ʒss., suspended in water by means of mucilage or yolk of egg.

<sup>1</sup> Charles V. is said to have been cured of severe rheumatism by a vinous decoction of this plant, taken daily for sixty days.

<sup>2</sup> Vide *A Description of the tree named Quinquino in Peru, &c.* By Don Hipólito Ruiz; translated in *Lambert's Illustrations of the genus Cinchona*, 4to. Lond. 1821. p. 92.



Official preparations. *Tinctura Benzoini composita*, L. E. D.  
*Tinctura Toluiferæ Balsami*, E. D. *Syrupus Tolutani*, L.

**TORMENTILLA.** *Spec. Plant. Willd.* ii. 1112.

*Cl.* 12. *Ord.* 5. Icosandria Polygynia. *Nat. ord.* Senticosæ,  
*Linn.* Rosacæ, *Juss.*

*G.* 1001. *Calyx* three-cleft. *Petals* four. *Seeds* roundish,  
 naked, affixed to a small, juiceless receptacle.

*Species* 1. *T. erecta*.<sup>1</sup> Common Tormentil, or Septfoil (*officinalis*).  
*Smith, Flora Brit.* 552. *Eng. Bot. t.* 863. *Med. Bot.* 2d edit.  
 503. *t.* 181.

*Official.* TORMENTILLÆ RADIX, *Lond. Dub. Edin.* Tormentil  
 Root.

*Syn.* Tormentil (*F.*), Tormentilwurzel (*G.*), Meerwortel (*Dutch*), Blodrot  
 (*Swed.*), Kurzeziele (*Pol.*), Tormentilla (*I., S., Port.*), Sabiasnoi koren (*Russ.*).

This is a very common, indigenous, perennial plant, growing  
 in dry pastures and on heaths; flowering in June and July.  
 The root is woody: the stems are erect, branched, diffuse or  
 procumbent, round and leafy: the leaves are nearly sessile,  
 ternate, lanceolate, serrated, and hairy, accompanied by deeply-  
 incised stipules: the flowers are on long, capillary, opposite,  
 solitary, one-flowered peduncles: the calyx consists of ovate,  
 hairy, alternately larger and smaller segments, the latter of  
 which are exterior: the petals have short claws, are obcordate  
 and of a golden-yellow colour: the seeds are few, and wrinkled.

*Qualities.* — The root has a very slightly aromatic odour,  
 and an austere, styptic taste. It is knotty: externally blackish  
 and internally reddish. To boiling water it yields its active  
 matter, which appears to be chiefly tannin, as the infusion is  
 copiously precipitated by solution of isinglass, and strikes a  
 deep black with sulphate of iron. Except galls and catechu,  
 it contains more tannin than any other vegetable.

*Medical properties and uses.* — Tormentil root is a powerful  
 astringent. It has been employed with success in intermit-  
 tents, but more efficaciously in diarrhoeas; particularly those  
 attendant on phthisis, as it produces its astringent effects with-  
 out increasing the general excitement. As a local remedy it  
 may be advantageously used in the form of gargle and lotion  
 in ulcerations of the tongue and mouth, against spongy gums,  
 and as an application to foetid ill-conditioned sores; but it is  
 seldom used. It may be given in substance, or in the form of  
 decoction. The dose of the powdered root is from ʒss. to ʒj.

**TRITICUM.** *Spec. Plant. Willd.* i. 476.

*Cl.* 3. *Ord.* 2. Triandria Monogynia. *Nat. ord.* Gramina.

*G.* 152. *Calyx* two-valved, solitary, subtriflorous. *Flower* some-  
 what obtuse.

\* *Annual.*

<sup>1</sup> Πεντάφυλλον Dioscoridis.



*Species 2.* *Triticum hybernum*<sup>1</sup>, Winter Wheat. *Gartner de Fructibus.*

*Officinal.* FARINA. AMYL'UM, *Lond. Edin.* TRITICUM; SEMINUM FARINA; AMYLUM, *Dub.* Wheat Flour. Starch.

*Syn.* Farine du froment; Amidon (*F.*), Weizenmehl; Kraftmehl, Staerhe (*G.*), Abgoon (*Arab.*), Neshasté (*Pers.*), Geehoonkaheer (*H.*), Farina di Frumento l' Amido (*I.*), Acemite; almidon (*S.*), Godumbay mao (*Tam.*).

The country whence this valuable grain originally came is unknown; but it is certain that Sicily was the part of Europe where it was first cultivated. It will not vegetate beyond the 62° of northern latitude. It has two sets of roots; one set proceeding directly from the seed, and the other from what is denominated the *corona* of the plant, about two inches above the first: the *coronal* roots do not shoot till spring-time, and collect more nutriment than the *seminal* roots<sup>2</sup>; the ears or spikes are long, with the grain lodged in four rows, and imbricate: the chaff smooth, bellied, and terminated by very short awns, distinguishing it from Spring Wheat (*Triticum æstivum*), which has awns three inches long. Many varieties of wheat are cultivated in this country, of which the *white Dantzic* is considered the best. The grain is small and translucent, and yields flour which makes more bread in proportion to the quantity of flour than that of any other variety of wheat. After the operation of grinding, the farinaceous part of the seed is separated, by means of cloth sieves, into several, distinct portions of various degrees of fineness; but the whole may be resolved into two: 1. *flour*, which constitutes more than two-thirds of the whole; and, 2. *bran*, which consists chiefly of the husk of the seed.

*Starch* is manufactured by steeping either entire or coarsely bruised wheat in cold water, until it swells, and yields a milky juice when squeezed. It is then subjected to pressure in coarse bags placed in vats filled with water; and when all the milky juice is obtained, the bags are removed, and the fecula deposits itself. In a short time the supernatant liquor ferments, and alcohol and acetic acid are formed in it. The whole is now put into tubs called frames, in which the impure fecula is allowed to subside; and after the water is poured off, the upper part of the sediment which last subsided being dirty and discoloured, is scraped off from the starch below; this is then repeatedly well-washed, pressed in cloths, and dried by a gentle heat, during which it cracks into small, columnar masses, and is the finest white starch of the shops.<sup>3</sup>

<sup>1</sup> Πυρὸς χειμὸς πορόμμενος Dioscoridis.

<sup>2</sup> *Hunter's Georgical Essays*, Essay v.

<sup>3</sup> The ordinary blue starch, which is coloured with a solution of smalt and alum in water, is unfit for medicinal uses. The Chians first made starch.



*Qualities*.—*Flour* is inodorous and nearly insipid. Water with which it has been macerated acquires an opaline colour and a sweetish taste; affords precipitates with infusion of galls and the strong acids, and rapidly becomes sour. It appears to contain gluten, sugar, gum, albumen, and phosphate of lime; besides fecula or starch that remains insoluble. According to Vogel the constituents of flour are, in 100 parts, fecula 68, gluten 24, saccharine gum 5, albumen 1.50. The action of these principles on each other, when flour is kneaded with water, and yeast added to the mass, excites the panary fermentation, and produces bread, a little salt being added to give it sapidity. The large proportion of gluten in wheat flour renders it fitter for this purpose than any other kind of flour. During the process, a large quantity of carbonic acid gas is evolved, which swells up the mass, and gives it the sponginess and lightness that characterize well-baked bread.<sup>1</sup> For the purpose of baking bread a heat of 488° is required. When flour has been long kept, it becomes musty and undergoes the putrefactive fermentation, in which state the bread made with it is very unwholesome. Flour is fit for making bread only when all its constituents are entire, and as gluten is the most susceptible of decomposition among them, the ascertaining its presence is a proof of the goodness of the flour. M. Taddei has taught us that guaiac is a test of the presence of gluten, by striking with it a beautiful blue colour; flour, therefore, which exhibits this colour when rubbed with guaiac and a few drops of vinegar, may be pronounced good.

*Starch* is inodorous and insipid; in white columnar masses which are easily reduced to powder. It is insoluble in alcohol, ether, and cold water; but in the latter it falls into powder. Boiling water dissolves it, forming an insipid, inodorous, semi-transparent, opaline, gelatinous-like paste, which becomes brittle and opaque, when spread out in a dry air; but when exposed, without being spread out, it separates into a watery fluid, and an opaque paste; sours, and becomes mouldy. Alcohol precipitates starch white and tough from its solutions; acetate of lead and infusion of galls also throw it down, but the precipitate formed by the latter is re-dissolved by heating the liquid to 120°. Although potass dissolves starch, yet the solution of it is not altered by potass, carbonate of potass, nor ammonia: but a solution of potass in alcohol, and a solution of sulphuret of potass in alcohol, both produce precipitates.

---

<sup>1</sup> The method of making leavened bread was probably invented by the Egyptians; for it appears that the Israelites were acquainted with it after they sojourned in Egypt, but not before. It was known to the Greeks during the Trojan war: but the use of yeast or *barm* was discovered by the ancient Gauls.



From the products obtained from distilling starch *per se*, it appears to be a ternary compound of carbon, oxygen, and hydrogen.

*Medical properties and uses.*— The utility of *bread* as an article of diet requires no particular notice.<sup>1</sup> As a medicinal agent it is used for forming poultices, cataplasms, and for giving bulk and form to very active medicines which require to be given in minute doses, in the solid state, or as pills. When toasted and infused in water, it gives a pleasant flavour to the fluid, and renders it more acceptable as a diluent in febrile diseases, and as the ordinary beverage of the dyspeptic. *Starch* is less nutritive than bread, but is perhaps more digestible. It forms the greater part of the nutritive matter of the different farinaceous substances which are in general use as the diet of the sick, such as sago<sup>2</sup>, salep<sup>3</sup>, tapioca<sup>4</sup>, arrow root<sup>5</sup>, and gruel, which are only different modifications of

<sup>1</sup> Wheat flour is almost exclusively used for this purpose in England, part of Scotland, France, a part of Germany, Hungary, the Crimea and Caucasus, and some part of the middle of Asia. It is used also for the same purpose, but not so exclusively, in Spain, Portugal, Italy, Greece, Persia, Northern India, Arabia, Egypt, Nubia, Barbary, the Canary islands, North America, the Brazils, Buenos Ayres, Chili, the Cape of Good Hope, and the temperate zone of New Holland. Rye, barley, and oats, usurp its place in many parts of the North of Europe; rice in the East Indies, China and Japan, in Asia and Africa, in the torrid zone; and Maize in part of America and of Africa. Besides these grains, Yams, Casava, Batatas, the Banana, Doura (*Singhum*), Sago, the Bread fruit, some species of Arum, Chenopodium, Quinoa, *Acrastichum furtum*, and *Arachis hypogæa*, are used in different parts of the globe as substitutes for bread. *Schouw on the Geographic distribution of the Gramineæ.*

<sup>2</sup> Sago is the pith of various species of Palms. One of these, the Sagu-tree of Asia, *Metroxylon Sagu* (Roxburgh), when fifteen years old, will sometimes yield 600 weight of sago. It has been calculated, that one English acre of land will grow 435 Sagu-trees; which would yield 120,500lbs. avoirdupois of sago, or 8000 lbs. yearly; a produce triple that of wheat. *Hist. of the Ind. Archip.* i. p. 357.

<sup>3</sup> Salep is prepared from the bulbs of the *Orchis mascula*. The bulbs are first dipped in hot water and the skin rubbed off; after which, they are placed on a tin plate, and put into a heated oven for ten minutes; and lastly, dried in the sun. By this process, they acquire the appearance of horn, and, when pulverized, form the Salep of the shops.

<sup>4</sup> Tapioca is prepared from the roots of the *Iatropa manihot*. The roots are first freed from the rind; and then, are either held to a large wheel, which, on being turned round, soon reduces them to a fine pulp, or they are grated. The pulp is next put into bags, and pressed, to squeeze out the moisture, which contains a poisonous principle, and afterwards repeatedly washed. It is then pressed through plates full of round holes to granulate it; and, lastly, dried by means of heat in large flat pans.

<sup>5</sup> Arrow root is the pith of the *Maranta arundinacea*. The powder is prepared from roots of a year old, which, after being well worked, are beaten, and the fibrous part separated from the pulp. The farinaceous pulp is then thrown into a fresh quantity of water, and stirred until it becomes milky, when the fluid is passed through a sieve, and left at rest until the fecula is deposited. The supernatant fluid is now poured off, and the starch, after being well washed, is dried in the sun. In this state it is brought to Europe, and sold under the name of Indian arrow root.



starch. The solution of starch is employed medicinally as a demulcent; but as it is very readily acted on by the stomach, it cannot be of much service in involving acrid matters in the intestines when taken by the mouth. In the form of enema, however, it is often and advantageously used for allaying the effects of acrid bile on the coats of the rectum in bilious diarrhœa and dysentery; and for sheathing the rectum in cases of abrasion, and inflammation of the gut. It is the common vehicle for the exhibition of opium per anum.

Officinal preparations. *Mucilago Amyl.* L. E. D.

**TUSSILAGO.** *Spec. Plant. Willd.* iii. 1962.

*Cl.* 19. *Ord.* 2. Syngenesia Superflua. *Nat. ord.* Compositæ Discoideæ, *Linn.* Corymbiferæ, *Juss.*

*G.* 1483. *Receptacle* naked. *Pappus* simple. *Calyx* scales equal, as long as the disc, submembranaceous. *Corolla* female. *Florets* ligulate, toothless.

*Species* 12. *T. Farfara.*<sup>1</sup> Common Coltsfoot. *Med. Bot.* 2d edit. 45. *t.* 18. *Smith, Flora Brit.* 878. *Eng. Bot. t.* 429.

*Officinal.* **TUSSILAGO,** *Lond.* **TUSSILAGINIS FARFARÆ FOLIA ET FLORES,** *Edin.* **TUSSILAGO; FOLIA,** *Dub.* Coltsfoot leaves and flowers.

*Syn.* Tussilage; Pas d'Ane (*F.*), Huflattisch (*G.*), Hoefblad (*Dutch*), Tassilagem (*Port.*), Dwoje lisknik (*Russ.*), Farfara (*I.*), Unã de cabelo (*S.*).

Coltsfoot is an indigenous perennial plant, growing in moist, marley, and clayey soils. It flowers in March and April, and the leaves appear in May and June. The root is long and diffusely creeping, and sends up stems or scapes destitute of leaves, erect, five or six inches high, simple, unifloral, tomentose, with sparse, smooth, scale-like bractes of a brownish pink colour, lying close to the stem: the flower droops before it blows, but afterwards becomes erect, and is of a golden-yellow colour, the calyx is composed of linear, trinerved, plane, smooth, purplish scales, the length of the disc equal, uniform, and finely reflex: the florets of the ray are numerous, spreading, linear, twice the length of those of the disc, with a more slender stigma: the seeds are smooth, more frequently abortive, particularly in the disc: with the seed-down sessile, rough, white, and shining: the receptacle is pitted, flat at first, but finally convex: the leaves appear after the flower, are radical, petiolate, erect, cordate, angled and toothed; smooth, green above with reddish veins, but underneath white and woolly.

The leaves are more frequently employed than the flowers, and should be gathered and dried when they are fully expanded, before they have attained their greatest magnitude.

<sup>1</sup> Βηχιον Dioscoridis. The name is derived from βηξ, *tussis*, whence *tussilago*, showing the early opinion of the pectoral virtues of this plant.



*Qualities.* — The dried leaves are inodorous, and have a rough, mucilaginous taste. The mucus they contain is yielded to water by coction, and evolves by the boiling a peculiar odour.

*Medical properties and uses.* — Tussilago is demulcent, and has been regarded as expectorant from the earliest ages, having been smoked through a reed in the days of Dioscorides, with the view of relieving the chest from accumulated mucus in catarrh, asthma, and phthisis. It is still used as a demulcent in catarrhal and phthisical affections; but very little reliance is placed on its powers.<sup>1</sup> Cullen thought he perceived good effects result from the use of the expressed juice of the recent leaves in scrophula; but his observations have not been generally confirmed.<sup>2</sup>

The decoction of the leaves is the usual form of exhibiting tussilago. A handful of the leaves is boiled in Oij of water to Oj; and the decoction after being strained is sweetened with sugar-candy or syrup. The dose is a teacupful.

VALERIANA. *Spec. Plant. Willd.* i. 175.

Cl. 3. Ord. 1. Triandria Monogynia. Nat. ord. Aggregatæ, Linn. Dipsacæ, Juss.

G. 75. Corolla monopetalous, gibbous on one side of the base, superior. Seed one.

\* valerians, with a single downy seed.

Species 6. *V. officinalis*. Officinal, or great Wild Valerian.

Med. Bot. 2d edit. 77. t. 32. Smith, *Flora Brit.* 38.

Officinal. VALERIANÆ RADIX, Lond. Dub. VALERIANÆ OFFICINALIS RADIX, Edin. Wild Valerian root.

Syn. Valeriane (*F.*), Wilde Baldrianwurzel (*G.*), Wilde Valeriaan (*Dutch*), Wändelrot (*Swed.*), Kettull gunnung (*Jav.*), Valeriana Silvestre (*I.*), Valerian officinal (*S.*), Balder an (*Russ.*), Baldrian (*Danish*), Kozlki (*Pol.*).

This species of Valerian is an indigenous, perennial plant, flowering in June. There are two varieties of it; one growing in woods and marshy ground, the other on high pastures and heaths; and the sensible qualities of the second are considerably greater than those of the first. It has been often regarded as the φs of Dioscorides; but Sibthorp has proved this opinion to be erroneous, and has described the real Valerian of the ancients as a distinct species under the name of *Valeriana Dioscoridis*.<sup>3</sup> The roots of Valerian are long and slen-

<sup>1</sup> A vile, stimulant nostrum, consisting, according to Dr. Paris (*Pharmacologia*); of equal parts of *Balsam of Tolu*, and the compound tincture of *Benzoin*, with double the quantity of *rectified spirit of wine*, is sold under the name of *essence of Coltsfoot*, as a remedy for coughs

<sup>2</sup> *Mat. Med.* ii. 160.

<sup>3</sup> *Sibthorp, Flora Græca*, p. 24. t. 53. Dr Smith, the learned editor of *Sibthorp's Work*, says, "Hæc est vere φov Dioscoridis, a nemine botanicorum recentiorum ante Sibthorp detecta." Willdenow's 7th species, *V. phu*, which was supposed to be the plant of Dioscorides, does not accord with his description, whereas that of Sibthorp corresponds with it in every particular.



der fibres issuing from heads; the stems rise three or four feet in height; are round, grooving, hollow, and terminated with flowering branches, disposed crosswise; the leaves are larger at the base of the stem, decreasing in size towards the summit; opposite, connate, and bearded at the base below; pinnate, with a terminal leaflet a little larger than the rest; all the leaflets deeply veined and serrated, of a dark-green colour on the upper surface, and paler underneath. The flowers are small, in corymbs, odorous, and interspersed with lanceolate, connate, bearded, waved, pale bractes: the calyx is a slight margin at the top of the germen: the corolla tubular, white with a shade of pink, divided at the margin into five, reflected, obtuse segments: the filaments are spreading with the corolla, and support round, yellowish anthers: the style is shorter, with a trifid stigma; and the capsule is crowned with a feathery pappus, purplish at the base; and contains one oblong, ovate, compressed seed.

The roots should be dug up in autumn when the leaves decay, or in spring, before they expand; and be preserved in a dry place. Those which grow wild on a calcareous soil are preferable to those that are cultivated. They lose three-fourths of their weight by drying. Cats are allured and delighted with the odour.<sup>1</sup>

*Qualities.*—Valerian root has a strong, peculiar, unpleasant odour, and a warm, bitter, subacrid taste. Trommsdorff has chemically examined it. Its virtues appear to depend on a very liquid greenish-white-coloured volatile oil, which from its odour and taste seems to contain much camphor. Its specific gravity at 77° of Fah. is 0·9340; when exposed to light it becomes yellow; a small portion of nitric acid converts it into resin, and a larger dose into oxalic acid. The expressed juice of the root contains starch, extractive, and gum; while the roots deprived of this juice yield a portion of black-coloured resin, but consist chiefly of woody fibre.<sup>2</sup> The active matter of Valerian root is extracted by boiling water, alcohol, and solutions of the pure alkalies.

*Medical properties and uses.*—Valerian root is antispasmodic, tonic, and emmenagogue. It is advantageously employed in hysteria, symptomatic epilepsy, hemicrania, and other affections depending on a morbid susceptibility of the nervous system. We have also found it exceedingly serviceable in

<sup>1</sup> Mr. Lambert, has endeavoured to prove, that the *Valeriana iatamansi*, a Nepalese alpine plant, is identical with the spikenard of the ancients. This root is fusiform, about the thickness of the human finger, and bearing on the upper part, articulations covered with dense fibres, which give them somewhat of the appearance of the tails of animals. Vide *Illustrations of the genus Cinchona*, &c. 4to. Lond. 1821. 177.

<sup>2</sup> *Annales de Chim.* lxx. 95. Thomson's *Chemistry*, 5th ed. iv. 225.



hypochondriasis; and it is regarded as a useful adjunct to Cinchona in intermittents. It may be exhibited in substance combined with a small portion of *Mace* or *Cinnamon*; or in the forms of infusion or tincture. The extract is a bad form of preparation. The dose of the powdered root may be from ℥j. to ʒj., given three or four times a-day.

Officinal preparations. *Extractum Valerianæ*, D. *Infusum Valerianæ*, D. *Tinctura Valerianæ*, L. D. *Tinctura Valerianæ ammoniata*, L. E. D.

VERATRUM. *Spec. Plant. Willd.* iv. 895.

Cl. 23. Ord. 1. Polygamia Monœcia. Nat. ord. Coronariæ, Linn. Junci, Juss.

G. 1859. *Hermaphrodite*. Calyx none. Corolla six-petalled. Stamens six. Pistils three. Capsules three, many-sided.

————— Male the same. Rudiment of a pistil.

Species 1. *V. album*.<sup>1</sup> White Hellebore. *Med. Bot.* 2d edit. 753. t. 257.

Officinal. VERATRI RADIX, *Lond.* VERATRI ALBI RADIX, *Edin.*

HELLEBORUS ALBUS; RADIX, *Dub.* White Hellebore root.

*Syn.* Hellébore blanc (*F.*), Wiese Niesswurzel (*G.*), Zwartbloemige nieswortel (*Dutch*), Hevit Prustrot (*Swed.*), Elliboro blanco (*I.*), Veratro blanco (*S.*), Helliboro (*Port.*), Tschemeriza (*Russ.*).

*Veratrum* is a native of the mountainous parts of Greece, Italy, Switzerland, and Russia. Those specimens which are cultivated in our gardens flower in July. The root is perennial, fleshy and fusiform, beset with strong fibres gathered into a head: the stem is thick, round, hairy, erect, three or four feet in height, and branching; the leaves are oblong-ovate, about ten inches long, and five broad in the middle; plaited longitudinally, embracing the stem at the base, and of a yellowish green colour: the flowers are in long, terminal spikes, composed of small, alternate spikelets, each accompanied with a lanceolate bracte; the flower consists of six, persistent petals, of a pale-green colour; three of them oblong and lanceolate, with a membranous edge; and three calycinal, which enclose the other three in the bud, one half shorter and heart-shaped, with a small point at the top: the filaments closely surround the germen, diverge and bend down at the summit, and are terminated by yellow, quadrangular anthers: the germens are three in each hermaphrodite flower<sup>2</sup>, oblong, with erect, bifid, hairy styles, crowned with flat, spreading stigmas. The capsules contain many compressed, membranous seeds.

Although the root only is officinal, yet every part of the plant is extremely acrid and poisonous.

*Qualities.* — The recent root has a strong, disagreeable odour, and a bitterish, very acrid, permanent taste; but the

<sup>1</sup> Ἑλλέβορος λευκός Dioscoridis.

<sup>2</sup> The hermaphrodite flowers are generally on the upper, erect spike.



odour is lost by drying. The dry root, as found in the shops, is sliced, the thick part transversely, and the fibrous, longitudinally. The pieces have a dry, corrugated, yellowish-grey appearance, and break with a short, starchy fracture. They are inodorous, and have a slightly bitter taste. When very light and spongy, they must be rejected. The experiments of MM. Pelletier and Caventou have proved, that white Hellebore owes its medicinal properties to *Veratria*, the same alkaline principle which has already been described as the active ingredient in *Colchicum*. The following are the components of white Hellebore, according to their analysis; a fatty matter, composed of *elaine*, *stearine*, and *ammonia*, *acidulous gallate of veratria*, a yellow colouring matter, starch, gum and lignin.<sup>1</sup>

*Medical properties and uses.* — White Hellebore is a violent carthartic, emetic, and sternutatory. When taken internally, even in moderate doses, its operation is violent and dangerous; producing, besides hypercatharsis, with bloody stools and excessive vomiting, great anxiety, tremors, vertigo, syncope, sinking of the pulse, cold sweats, and convulsions, terminating, if the dose be large, in death. Its external application to an ulcerated surface also produces griping and purging. Notwithstanding these effects, *Veratrum* has been exhibited internally, and with advantage, in mania, epilepsy, scabies, lepra, and obstinate herpetic eruptions.<sup>2</sup> But the most ordinary use of white Hellebore is as a local stimulant; either as an adjunct to errhine powders in lethargic cases and gutta serena; or in the form of decoction as a wash, or mixed with lard as an ointment, in scabies and herpetic eruptions. In every form, however, it requires to be used with caution; and even as an errhine, its acrimony should always be obtunded by mixing it with some mild powder, as that of liquorice root or of starch. The dose of the powdered root should not exceed grs. ij; and for errhine purposes grs. ij or iij should be diluted with grs. xij of liquorice powder, and a pinch of it snuffed up the nose for several successive evenings. When taken internally as a poison, the best antidote is a strong infusion of nut-galls.

Officinal preparations. *Decoctum Veratri*, L. *Tinctura Veratri albi*, E. *Unguentum Veratri*, L. *Ung. Sulphur. comp.*, L.

VERONICA. *Spec. Plant. Willd.* i. 54.

Cl. 2. Ord. 1. Diandria Monogynia. Nat. ord. Personatæ, Linn. Pedicularis, Juss.

G. 44. Corolla, border four-cleft, with the lowest segment narrower. Capsule two-celled.

\*\* with corymbose racemes.

<sup>1</sup> Journ. de Pharm. Aout. 1820.

<sup>2</sup> Medical Communications, i. 297.



*Species* 30. *V. Beccabunga*.<sup>1</sup> Broad-leaved Brooklime. *Med. Bot.* 2d edit. 363. *t.* 132. *Eng. Bot.* x. 655. *Smith Flora Brit.* i. 20.

*Officinal.* BECCABUNGA; HERBA, *Dub.* The herbaceous part of brooklime.

*Syn.* Beccabunga; Veronique aquatique (*F.*), Bachbunge (*G.*), Beckeboom (*Dutch*), Bekkebunge (*Dan.*), Backabunga (*Swed.*), Potoeznik (*Polish*), Anagalide acquatica (*I.*), Beccabunga (*S., Port.*), Ibunka (*Russ.*).

Beccabunga is an indigenous, perennial plant, common in rivulets and clear ditches, flowering in July and August. The stem, which is procumbent or floating, and gives off from the joints long, simple, fibrous roots, is round, leafy, and, like every other part of the plant, smooth and shining. The leaves are opposite in pairs, on short petioles, oval, serrated, somewhat fleshy, punctured, and of a pale-green colour. The flowers are collected in opposite axillary clusters, and individually supported on delicate footstalks, accompanied by linear-lanceolate bractes. The calyx is divided into four, acute segments, shorter than the corolla, which is of a very beautiful sky-blue colour, with the tube white. The anthers are whitish, supported on filaments longer than the style; and the capsule cloven, almost twin. This plant is green throughout the year, but in greatest perfection in the spring.

*Qualities.* — It is inodorous and has, when much chewed, a bitterish, slightly astringent taste. The expressed juice reddens the more delicate vegetable blues in a small degree.

*Medical properties and uses.* — Although Brooklime was formerly considered as a good antiscorbutic, yet it is properly disregarded by modern practitioners; and, as Lewis observes, if it be expected to produce any good effect, it should be used as food.

VIOLA. *Spec. Plant. Willd.* i. 1159.

*Cl.* 5. *Ord.* 1. Pentandria Monogynia. *Nat. ord.* Campanaceæ, *Linn.* Cisti, *Juss.*

*G.* 446. *Calyx* five-leaved. *Corolla* five-petalled, irregular, horned at the back. *Anthers* cohering. *Capsule* superior, three-valved, one-celled.

\* *stemless.*

*Species* 12. *V. odorata*.<sup>2</sup> Sweet Violet. *Med. Bot.* 2d edit. 251. *t.* 89. *Smith, Flora Brit.* 245.

*Officinal.* VIOLÆ FLORES, *Dub.* VIOLÆ ODORATÆ FLORES, *Edin.* The recent flower of the violet.

*Syn.* Violette odorante (*F.*), Blaue veilchen (*G.*), Tamme, Viool (*Dutch*), Marts fioler (*Dan.*), Akta fioler (*Swed.*), Viola Mammola (*I.*), Violeta (*S.*), Violetta (*Port.*), Pachutschaja fialko (*Russ.*), Kiet tuong hoa (*Chinese*).

<sup>1</sup> The specific appellation is probably derived from the Flemish *beek-pungen*, mouth-smart. In Scotland, where it is often eaten like water cresses, it is called *Water-purpie*.

<sup>2</sup> Ἰσὺ πορφύρεον Dioscoridis.



This species of the violet is indigenous, growing in shady places; and flowering in April and May. It is a low creeping plant, giving out runners, which root at small intervals, and send up tufts of leaves and flowers. The roots are fibrous. The leaves heart-shaped, with crenated edges, on slender footstalks; the upper surface of a lively-green colour, the under paler and downy. The flowers are supported on delicate, quadrangular, channelled flower-stalks, about two inches long, furnished with two small bractes, and curved at the summit. The calyx consists of five green leaflets, the two posterior of which are separated by the spur of the corolla. The petals have a deep-violet colour, are white at the base, and irregular; the two lateral ones are bearded near the base; and the posterior, which is slightly keeled, has a large spur, enclosing glandular appendices of the corresponding anthers. The anthers are nearly sessile, whitish, flat, supporting orange-coloured, membranous expansions that cover the upper part of the germen; which is pyramidal, downy, and crowned with a falcated pistil.

For medicinal and chemical purposes, the sweet Violet is cultivated in great abundance at Stratford-on-Avon; but the London herb-shops are supplied chiefly from Kent. As the petals only, separated from the calyx, are brought to market, it is difficult to detect the admixture of the *Viola hirta*, an inodorous species, which is often practised. It is not, however, a matter of much importance.

*Qualities.* — Violets have an agreeable, sweet odour and a very slightly bitter taste. When chewed they tinge the saliva blue, and yield their colour and flavour to boiling water. The root, stem, leaves, flowers and seeds yield an alkaline principle not unlike *emetine*, which M. Boullay, its discoverer, has termed *violine*. It is, like *emetine*, a powerful poison.<sup>1</sup> It is united with malic acid in the Violet, as *emetine* is with Gallic in *Ipecacuanha*.

*Medical properties and uses.* — The petals of the Violet are gently laxative; and were formerly regarded as anodyne and pectoral: but they are now scarcely ever used, except for preparing the syrup, which is given occasionally as a purgative to infants. Their aqueous tincture, and the syrup, are useful and delicate tests of the presence of uncombined acids and alkalis: the former changing the blue colour to a red, the latter to a green. The infusion is not liable to change, if it be kept in a tin flask, well stopped. MM. Corte and Willemet, who employed the powdered roots to produce vomiting, found that that was fully effected by doses of two scruples.

---

<sup>1</sup> *Journ. de Pharmacie*, Jan. 1824.



Officinal preparation. *Syrupus Violæ*, E. D.

VITIS. *Spec. Plant. Willd.* i. 1180.

Cl. 5. Ord. 1. Pentandria Monogynia. Nat. ord. Hederaceæ, Linn. Vitis, Juss.

G. 453. *Petals* cohering at the apex, shrivelling. *Berry* five-seeded; superior.

*Species* 1. *V. vinifera*.<sup>1</sup> Common Vine. *Med. Bot.* 2d edit. 144. t. 57. *Duhamel Arb.* ii. t. 1—6.

Officinal. UVÆ PASSÆ, Lond. VITIS VINIFERA FRUCTUS, Edin.

UVÆ PASSÆ SOLE SICCATÆ, Dub. Raisins, Sun Raisins.

*Syn.* Raisins secs (*F.*), Rosinen (*G.*), Groote razynen (*Dutch*), Russin (*Swed.*), Uva passa (*I.*), Passas (*S.*), Uvas Passadas (*Port.*), Zabib (*Arab.*), Kishmish (*H.*), Dividatsipalavuttil (*Tam.*), Velit chamoodika gheddie (*Cyng.*), Mewuz (*Pers.*), Zebub (*Malay*).

The Vine is a native of Armenia, Georgia, and the Levant; but it is now found in most of the temperate regions of the earth, and is cultivated with care wherever its fruit can be brought to perfection. In France, the northern limit of the Vine is stated to be  $50^{\circ} 20'$ ; in Thuringia, Saxony, and Siberia, it is  $51^{\circ}$ ; but towards the east it is lower, for although Hungary has much wine, yet Galicia has none; and in the southern parts of the Russian Empire it ascends no higher than  $48^{\circ}$ . In America the vine is cultivated in the southern states only, extending no farther north than  $38^{\circ}$ . The limits southward in the northern hemisphere is properly  $15^{\circ}$ ; but in the high mountainous island of St. Thomas on the coast of Guinea, in Abyssinia, and in the Deccan, it is found almost under the equator. In the southern hemisphere, its southern limits are  $37^{\circ}$ . The greatest altitude is, in  $45^{\circ}$  latitude, 2460 feet; in the north of Switzerland, 1700 feet; on the Alpine range, 2000 feet; in Teneriffe, 2500 feet; and on the Apennines and in Sicily, 3000 feet.<sup>2</sup> Its culture is supposed to have been introduced from the East, where it was cultivated, and wine made from the fruit, in the earliest ages<sup>3</sup>; and afterwards to have extended to Italy, about 600 years after the foundation of Rome, and thence to Burgundy in the time of the Antonines. It was introduced into Madeira, from the island of Cyprus, in the fifteenth century. In Great Britain the Vine was cultivated before the year 731, when Bede finished his history; but although it was at one period brought to considerable perfection<sup>4</sup>, yet, from the greater value of the

<sup>1</sup> Ἀμπέλως Græcorum.

<sup>2</sup> See extract from Prof. Schoure's work on the Geography of Plants, *Edin. Phil. Journ.*

<sup>3</sup> We are told, that Noah, after coming out of the ark, planted a vineyard, and "drank of the wine, and was drunken." *Genesis*, chap. ix. ver. 20, 21.

<sup>4</sup> There were many vineyards in different parts of this country from which wine was made; and we are informed, that in the cellar at Arundel castle, in 1763, there were sixty pipes of excellent Burgundy, the produce of a vineyard attached to the castle. *Museum Rusticum*, i. 85.



ground for the cultivation of corn, and the wines produced in this country having never equalled those of the continent, vineyards are now scarcely known in Britain. The Vine, therefore, is cultivated here for the dessert only, no raisins are made, and scarcely any wine.

The Vine has a slender, twisted, climbing stem, covered with a rough, peeling, fibrous bark. The leaves are lobed, and sinuated, serrated, and placed alternately on long footstalks. The flowers, which appear in June and July, are small, and produced in clusters, attended by tendrils. The calyx is very minute; the petals are of a greenish-white colour, adherent at their apices, [and soon fall off, like a little cap, from the anthers, which then spread and shed their pollen. The fruit is a succulent, globular berry, one-celled when ripe; naturally containing five seeds; but in general only two, which are hard and of an irregular form. There are many varieties of the Vine; that which is called the Alexandrian Frontiniac yields the most delicious grapes for eating, and the Syrian the largest bunches.<sup>1</sup>

Raisins are made from the varieties named the *Black Raisin Grape*, and the *White-Raisin Grape*. They are cured in two methods; either by cutting the stalk of the bunches half through, when the grapes are nearly ripe, and leaving them suspended on the Vine till their watery part be evaporated, and the sun dries and candies them; or by gathering the grapes, when they are fully ripe, and dipping them in a ley made of the ashes of the burnt tendrils; after which they are exposed to the sun to dry. Those cured in the first method are most esteemed. They are brought to this country packed in boxes with sand.

*Qualities.* — *Grapes*, when recent and fully ripe, have an agreeable, cooling, sweet, subacid taste. They contain, besides water, sugar, mucilage, and jelly, albumen, gluten<sup>2</sup>, tannin, super-tartrate of potass, tartrate of lime, phosphate of mag-

<sup>1</sup> This is supposed to be the sort of grape which the spies, sent by Moses to examine Canaan, cut down at the brook Eshcol; "a branch with one cluster of grapes, and they bare it between two upon a staff." *Numbers*, chap. xiii. 23. Strabo relates, that in Margiana bunches of grapes were produced two cubits, or a yard long; and in some of the Archipelago islands, they weigh from thirty to forty pounds. The Syrian grape, in this country, has produced bunches weighing nineteen pounds and a half. *Martyn's edition of Miller's Dictionary*. There is a grape, cultivated in Madeira, as a dessert fruit, the clusters of which sometimes weigh twenty pounds.

<sup>2</sup> The gluten excites the vinous fermentation in the juice of the grape when expressed. Fabroni has shown, that it is lodged on the membranes that separate the cells of the grape; and becomes mixed with the saccharine part when the juice is expressed.



nesia, muriate of soda, sulphate of potass, and tartaric, citric, and malic acids; and a mucoso-saccharine principle, which Chaptal and Proust regard as the constituent on which the fermentative process in bruised grapes depends. *Raisins* differ from grapes chiefly in the quantity of saccharine matter being more abundant.

*Medical properties and uses.* — The ripe fruit of the Vine is cooling and antiseptic; and, when eaten in large quantities, diuretic and laxative. Grapes are very useful in febrile diseases, particularly in bilious and putrid fevers, dysentery and all inflammatory affections. In Syria the juice of ripe grapes inspissated is used in great quantity in these diseases.<sup>1</sup> Grapes have been strongly recommended as an article of common diet in phthisis<sup>2</sup>; and they certainly contain much bland nutritious matter, well fitted for phthisical habits. *Raisins* are more laxative than the fresh fruit, and are apt to prove flatulent when eaten in any considerable quantity. They are used as an adjunct to some officinal preparations; but add nothing to their efficacy.

#### VINUM. Wine.

*Officinal.* VINUM. *Vinum album Hispanicum*, Lond. VINUM ALBUM HISPANUM, *Edin.* Sherry Wine.

*Syn.* Vin d'Espagne (*F.*), Wein (*G.*), Wyn (*Dutch*), Win (*Swed.*), Vino (*I.*), Vino de Xerez (*S.*), Khumr (*Arab.*), Bāde (*Pers.*), Dakh ramudh (*H.*).

Although the London and Edinburgh Colleges have designated *Sherry* only, yet all the generous wines are occasionally used as medicinal agents, and, therefore, we shall take a general view of the manufacture, characters, and properties of wine.

In the wine countries, when grapes are fully ripe, they are gathered, and immediately subjected to the press, by which the juice is separated from the skins and seeds. In some places the grapes are previously picked from the stalks, the sound being separated from all the unsound with great care<sup>3</sup>; in some they are pressed just as they are gathered from the vines; and in other places they are almost converted into raisins before they are pressed.<sup>4</sup> The expressed juice is called *must*, and contains all the principles which we enumerated above as being present in the grape; these, when the vats holding the *must* are placed in a temperature of 70°, begin to act upon

<sup>1</sup> *Russel's Natural History of Aleppo*, vol. i. p. 85.

<sup>2</sup> *Moore's View of Society, &c. in Italy*, ii. letter 62.

<sup>3</sup> This is the case at Madeira; and at Epernay, where the best Champagne is made. In Madeira, every kind of grape which the island produces, except the Malmsey and the Sercial, are pressed together for making the wine which bears the name of the island.

<sup>4</sup> The wine of Chio, which was esteemed by the ancients for its strength, sweetness, and exquisite aromatic flavour, is made from nearly dried grapes.



one another, the liquor becomes turbid, an intestine motion is evident in it, its temperature increases, a scum collects on its surface, and carbonic acid gas is disengaged. This is the process of vinous fermentation. In a few days its activity gradually decreases, the scum and impurities subside to the bottom; and the liquor clears, having lost its saccharine taste, and become *wine*. It is then put into barrels, and in due time into bottles, in both of which kind of vessels the fermentation is continued, although in an imperceptible degree; nor is it altogether completed till the wine attains the utmost limits of its age, and passes into the acetous fermentation. All the principles of the *must* are perhaps required for the production of wine; but the saccharine matter, the gluten, and the vegetable acid, are essential; and on the proper quantity of the first in particular, and the manner in which the fermentation is conducted, depend the strength and goodness of the wine. When the sugar is in too great quantity, and not completely decomposed, or the fermentation is checked, the wine retains a sweet taste; a more proper proportion and perfect decomposition, with a brisker fermentation, render it strong and spirituous; but if the quantity of sugar be small, and at the same time there is a deficiency of tartar in the *must*, a thin and weak wine is produced. When it is bottled early it becomes brisk and sparkling; and it is rough and astringent when the fermentation has been conducted on the skins, particularly on those of the coloured grapes; which also gives colour to the wine; for when the juice only is fermented, white wines are produced from coloured grapes.

Wine that has been too long fermented before being put into the casks, is very apt to become sour; and frequently oxides of lead, as litharge, and white lead, are employed to correct the acidity. According to Fourcroy, these form a soluble, triple salt, an aceto-tartrate of lead, by uniting with the acetic and tartaric acids in the wine<sup>r</sup>; which, daily experience shows, produces violent cholic, and other deleterious effects on those who drink it. The fraud may be detected by means of a solution of sulphuretted hydrogen gas, as has been already explained (see *Plumbum*), or by the prussiate of potass, a few drops of which will produce a whitish precipitate, if lead be present. Port wine is often adulterated, and coloured with dye-stuffs. Liquid potash detects the artificial admixtures of wines. It precipitates the natural principles of wine, green; Berries of Yebb, precipitate them violet; Indian wood, violet; Red Mulberries, violet: Brazil wood,



red; Beet, red; Litmus, clear violet; Myrtle-berries, wine lees colour; Elder-berries, blueish. But besides those articles already mentioned, both arsenic and corrosive sublimate have been used for fining wines. Nitrous ether is sometimes employed to perfume wines.

*Qualities.* — Various circumstances, such as climate, soil, and the mode of conducting the fermentation, modify the flavour and taste of wine. The odour and flavour, in the more fully fermented wines, seem to depend on the vinous process, as it bears little resemblance to the natural flavour of the fruit; from which, however, in sweet and half fermented wines it is immediately derived; but flavouring ingredients, as Bitter Almonds and Orris root, are also used in the manufacturing of wines. Malaga, Frontignac, Tokay, Vino Tinto, Montifiuscone, Schiras, and the Malmsey wines of the Greek islands, are sweet to the taste, and consequently the result of imperfect fermentation; Champagne, Gooseberry, and all sparkling wines owe their briskness to carbonic acid gas; Hock, Rhenish, Mayne, Barsac, Burgundy, Claret<sup>1</sup>, and Hermitage contain a certain quantity of uncombined acid, and are termed light and dry; while Marsala, Madeira, Sherry<sup>2</sup> and Port, are dry and strong. The odour of *Sherry* is pleasant and aromatic; the taste warm, with some degree of the agreeable bitterness of the Peach kernel; the taste of *Port* is austere, and bitterish; *Claret* is less rough, thinner, slightly acidulous and higher flavoured; and *Hock* acidulous. Of the common white wines, *Marsala* is undoubtedly the strongest. But notwithstanding these and other differences, the essential components of all wines are the following: One or more *acids*; generally the malic, but in some the carbonic predominates, and all of them contain some tartaric; *extractive matter*, which in old wines is deposited with the tartar; a *volatile oil*, on which the flavour depends; *colouring matter*; and *alcohol*, the most important of the ingredients, and that one on which their dietetic and medical properties depend. Gay Lussac has proved that this principle is ready formed in wine, and not, as Fabroni supposed, the result of its distillation. The following table is intended to show the average quantity, by measure, of alcohol,

---

<sup>1</sup> The best claret is made from grapes grown at Chateau Margaux. *Vinum, tam album quam rubrum, claritum, medinum &c.* were claimed by the Monks of Winchester, on Festivals in 1285. *Regist. Priunt. S. Smith, Winton MS. quoted in Warton's Hist. of Poetry*, 4to. vol. i. p. 425.

<sup>2</sup> The sherry, or sherris-sack of Shakspeare's time, may mean, as Dr. Henderson supposes, the *dry* or *sec* wine of Xeres: or, if it was the first wine exported from Spain, the term sack, in my opinion, may have been given to it from that circumstance, *Vino Sacco*, in Spanish signifying export wine.



oily, unctuous, resinous and gummy matter contained in several wines, which were examined by Neumann.

<i>A Quart of</i>	<i>contains of</i>											
	<i>highly rectified Spirit.</i>		<i>thick, oily, unctuous, resinous Matter.</i>			<i>gummy &amp; tartareous Matter.</i>			<i>Water.</i>			
	℥.	℥.	℥.	℥.	grs.	℥.	℥.	grs.	lb.	℥.	℥.	grs.
Aland - -	1	6	3	2	0	1	5	0	2	5	3	0
Alicant - -	3	6	6	0	20	0	1	40	2	2	6	0
Burgundy - -	2	2	0	4	0	0	1	40	2	9	0	20
Carcassone - -	2	6	0	4	10	0	1	20	2	8	4	30
Champagne - -	2	5	0	6	40	0	1	0	2	8	3	0
French - -	3	0	0	6	40	0	1	0	2	8	0	20
Frontignac - -	3	0	3	4	0	0	5	20	2	4	6	30
Vin de Grave	2	0	0	6	0	0	2	0	2	9	0	0
Hermitage - -	2	7	1	2	0	0	1	40	2	7	5	20
Madeira - -	2	3	3	2	0	2	0	0	2	4	3	0
Malmsey - -	4	0	4	3	0	2	3	0	2	1	2	0
Vino de Monte Pulciano } - -	2	6	0	3	0	0	2	40	2	8	0	20
Moselle - -	2	2	0	4	20	0	1	30	2	9	0	10
Muscadine - -	3	0	2	4	0	1	0	0	2	5	4	0
Neufchatel - -	3	2	4	0	0	1	7	0	2	2	7	0
Palmsec - -	2	3	2	4	0	4	4	0	2	2	5	0
Pontac - -	2	0	0	5	20	0	2	0	2	9	0	40
Old Rhenish	2	0	1	0	0	0	2	20	2	8	5	40
Rhenish - -	2	2	0	3	20	0	1	34	2	9	1	6
Salamanca - -	3	0	3	4	0	2	0	0	2	3	4	0
Sherry - -	3	0	6	0	0	2	2	0	2	0	6	0
Spanish - -	1	2	2	4	0	9	4	0	1	10	6	0
Vino Tinto - -	3	0	6	4	0	1	6	0	2	0	6	0
Tokay - -	2	2	4	3	0	5	0	0	2	0	3	0
Tyrol, red - -	1	4	1	2	0	0	4	0	2	8	6	0
Red wine - -	1	6	0	4	40	0	2	20	2	9	3	20
White - -	2	0	0	7	0	0	3	0	2	7	0	0

Mr. Brande, Dr. Prout and M. Zez have prosecuted this enquiry with more accuracy. In the following table the quantity of alcohol of specific gravity .850, in 100 parts of Wine, is given on the authority of the experiments of these distinguished chemists.



TABLE of the PRINCIPAL KNOWN WINES, and of the QUANTITY of ALCOHOL in WINES.

Where produced.	Generic Names.	Varieties.	Quantity of Alcohol of spec. grav. .825 in 100 parts.	Qualities.
Portugal ..	<i>Red. Port</i> .....	..... (average)	22.96 B. <sup>1</sup>	Deep purple; rough; bitter sweet; spirituous.
		Vinho de Ramo.....	15.62 P.	
		Collares .....	19.75 P.	
	<i>White. Bucellas</i> .....	.....	18.49 B.	Pale straw; flavour delicate.
	Setuval ...	.....	—	
	Carcavellos .....	.....	18.65 B.	Amber-colour; sweet.
Spain .....	<i>White. Sherry</i> ...	.....	19.17 B.	Deep amber colour; nutty and aromatic.
		Amontillado .....		
		Paxarete.....	—	Amber colour; sweet and aromatic.
	Malaga... ..	..... (A. D. 1666)	18.94 B.	
		Pedro-ximenes .....	—	Amber colour; flavour delicate, rich, sweet.
		Lagrima de Malaga	—	
	Malmsey of Sitges Priory	.....	—	Resembles Malaga.
	<i>Red. Tent, Tintilla</i> .....	.....	13.30 B.	Purple; sweet; flavour strong, spicy.
	La Torre ...	.....		
	Peralez .....	.....	—	Sweet.
	Segorve ...	.....		
	Vinaroz ...	.....	—	Resembles Claret.
	Benicarlo...	.....		
	Carinena ...	.....	—	Resembles Sauterne.
	Val de Penas	.....		
Majorca ..	Manzanares	.....	—	Still, of an amber colour. Brisk or sparkling; delicate flavour and aroma; slightly acidulous; but some are still, or, at most, simply creaming; generally paler than Sillery.
	Ciudad Real	.....		
France ...	<i>White. Alba flor</i> .....	.....	17.26 B.	Good colour and body, and a high agreeable flavour.
	<i>White. Champagne</i> .....	Sillery .....	13.30 B.	
		Ay, Hautvilliers, Epernay, Dizy, Avenay, Avise, Oger, Pierry, Closet. Lemesnil, Cramont, Menil.	—	Colour and aroma of Burgundy, with lightness of Champagne.
	<i>Red. Champagne</i>	Verzy .....	11.93 B.	
		Verzynay, Mailly, Bouzy, St. Basle, Chamery, Ecueil, Villedemange .....	—	Inferior to Champagne, but resembling it in some of their qualities.
		Clos St. Thierry ....	—	
	<i>White. Arbois</i> ...	.....	—	
	Papillon .	.....		
	Chablis...	.....	—	

<sup>1</sup> B means on the authority of Mr. Brande; P of Dr. Prout; and Z of M. Zetz.



Where produced.	Generic Names.	Varieties.	Quantity of Alcohol of spec. grav. .825 in 100 parts.	Qualities.
France .....	<i>Red. Burgundy</i>	.....	14.57 B.	Beautiful, rich, purple colour; exquisite flavour, with a full body, yet delicate and light.
		Romanée Conti, Clos-Vougeot, Chamber-tin, Richebourg, Romanée de Saint Vivant, Tache, St. George .....	—	
		Volnay, Pomard, Corton, Vosne, Nuits, Beaune Chamboll, Morey, Meurseault, Savigny-sous, Beaune .....	—	
		Romaneche, Torins, Chenas, Tonnere, Auxerre .....	—	Excellent wines, but inferior to the former.
		.....	—	Strong, generous wines.
	<i>White. Burgundy</i>	Mont Rachet .....	—	High perfume and nutty flavour.
		La Perriere, la Combotte, la Goutte d'or, la Genevriere, les Charmes, Vaumorrillon, les Grisées, Valmur, Grenouilles, Vaudesir, Bougnereau, Mont de Milieu, Fuissey, Pouilly .....	—	Rich, high-flavoured wines.
		.....	—	Dark purple colour; flavour exquisite, and perfume resembling that of the raspberry.
	<i>Red. Hermitage</i>	Meal, Greffieux, Besas, Baume, Raucoule ...	2.32 B.	
		Crozes, Gervant, Mercureol .....	—	Less delicate in flavour.
	<i>White. Hermitage</i>	Vin de paille .....	17.43 B.	Amber colour; sweet, luscious.
		Côte Rotie .....	12.32 B.	Resemble Hermitage in flavour, but are weaker.
		Seyssuel ...	—	Violet perfume.
		Clarette of Die.....	—	Light, sparkling, delicate.
	<i>Red. Tavel.....</i>	.....	—	Bright rose colour; flavour and aroma delicate.
		Chuzlan ...	—	
		Beaueaire ..	—	Inferior.
		St. Geniez..	—	
		Lirac .....	—	Full rich colour; flavour of Ratafia.
		St. Laurence	—	
		St. Joseph .	—	Sprightly; colour of the violet.
		St. Georges	—	
		Cornas .....	—	Full rich colour; flavour of Ratafia.
		.....	—	
	<i>White. Vin de Cotillon.....</i>	.....	—	Sprightly; colour of the violet.
		St. Peray, St. Jean	—	



Where produced.	Generic Names.	Varieties.	Quantity of Alcohol of spec. grav. .825 in 100 parts.	Qualities.
France ....	White, Frontignan	.....	12.79 B. {	Luscious, flavour of the grape.
	Lunel ....	Clos-Mazet .....	15.52 B. {	Bright yellow colour; less luscious than Frontignan.
	Beziers ...	Cazoul, Bassan.....	—	Resembles Sherry.
	Red, Roussillon	..... (average) Bagnols sur Mer, Cosperon, Collioure, Torremila, Grenache, Terrato .....	18.13 B. {	Great body and colour, become tawny when old.
	White, Rousillon	Rivessaltes .....	—	Bright golden colour; fragrant aroma; flavour of the quince.
		Salces ( <i>Maccabac</i> ).....	—	Similar, inferior to Rivessaltes.
		.....	21.24 P. {	Red; somewhat rough; sweet.
	Red, Claret.....	..... (average) Lafitte, Latour, Leoville, Chateau Margaux, Rauzan .....	15.10 B. {	Deep purple; delicate flavour; violet perfume.
		( <i>Graves</i> ) Haut Brion, Haut Talanu, Merignac, (average) <i>Artimino</i> , <sup>1</sup> <i>Kissanos</i> <sup>2</sup>	13.37 B. {	Resemble the better sorts of Burgundy, but are rougher.
		Gorce, Larose, Brantmouton, Pichow, Longueville .....	—	Light wines; of good flavour.
		St. Emilion, Canon...	—	Harsh; odour of burning sealing-wax.
	White, Claret ...	Preignac, Beaumes, Langon, Cerons, Buzet .....	—	Secondary quality.
		St. Nessans, Sancé, Mont Basillac.....	—	Sweet.
	Barsac .....	.....	13.86 B. {	Amber colour; full; aroma somewhat like cloves.
	Sauterne ...	.....	14.22 B.	Amber colour; sweetish.
Germany .	White, Rhenish	Johannisberger (1788)	8.71 P. {	High flavour and perfume.
		Steinberg .....	—	Strongest of the Rhine wines; sweetish.
		Rüdesheimer (1811) Grafenberg .....	10.72	Like the former.
		Markebrune, Rothenberg .....	—	Soft and delicate flavour.
	Red, Rhenish...	(Hock) Hocheimer, (average) .....	13.68 B.	Light; acidulous.
		Amanshausen, Leibfrauenmilch, Scharlachberger .....	—	Considerable body.
		Laubenheim, Nierstein .....	—	Light; delicate perfume and taste.

<sup>1</sup> A Tuscan wine.

<sup>2</sup> A Candian wine.



Where produced.	Generic Names.	Varieties.	Quantity of Alcohol of spec. grav. .825 in 100 parts.	Qualities.
Germany	Red. Rhenish	Bodenheimer (1802)	13.96 Z.	Delicate perfume and taste.
	Moselle...	Braunenberg, Pisport, Zeltingen, Wehlen, Graach.....	} — }	Light pleasant flavour, high aroma.
Hungary	Tokay.....	.....	9.88 B. }	Brownish-yellow when new, greenish when old.
		Tokay Essence .....	—	Syrupy, thick, muddy.
		Ausbruch.....	— }	Thinner and more vinous.
		Maslas .....	— }	Inferior to the two former.
	Méneser .....	{ Ædinburg, Rusth, Ofen..... }	—	Sweet, resembles Tokay.
Italy .....	Montepulciano .	.....	—	Sweet, with high flavour.
		Aleatico ... ..	16.20 P. }	Brilliant purple; luscious aromatic flavour.
	Verdea .....	.....	— }	Greenish colour and high flavour.
	Trebbiano .....	.....	—	Golden colour; sweet.
	Albano .....	Montefiascone .....	—	Pale straw-colour; light.
	Orvieto .....	.....	—	Both red and white; light.
	Lacrima Christi	.....	19.70 B.	Red, luscious, sweet.
		Monte Somma, Gallite .....	} — }	The best Lacrima.
		Ischia, Nola, Ottajano, Novella, Torre de Greco, Pozzula .....	} — }	Second-rate wines.
	Vino Greco ...	.....	—	Sweet.
Sicily .....	Marzala .....	..... (average)	... 25.9 B.	
		Twenty-one years old, submitted to Soemmering's process five years .....	} 18.40 P. }	Resembles Madeira.
	Syracuse.....	.....	... 15.28 B.	Both red and white
	Etna .....	.....	... 30.00 P. }	Resembles Madeira, with the harsh flavour of Sicilian brandy.
	Lissa .....	.....	.. 15.90 P.	Resembles Claret.
Ithaca ...	Red wine of Ithaca.....	} .....	—	Hermitage flavour.
Cephalonia	Cephalonia .....	.....	—	A dry red wine.
Candia ...	Rithymo .....	.....	— }	A fine flavoured white wine.
Cyprus ...	Vino Santo.....	.....	—	Pale straw colour; sweet.
Tenos.....	Tenos .....	.....	—	Luscious, sweet.
Tenados...	Red Muscadine .....	.....	—	Resembles Tokay.
Smyrna ..	White Muscadine .....	.....	—	Luscious, sweet.
Madeira...	Madeira .....	..... (average)	... 22.27 B. }	Full; pungent, nutty, or bitter-sweet, rich, aromatic flavour.
		..... (West Indies)	... 21.20 P. }	
		Sercial..... (average)	... 20.32 B. }	
		Malmsey <sup>1</sup> .....	... 16.40 B.	Luscious, sweet.
Teneriffe .	Teneriffe.....	.....	... 19.79 B.	Resembles Madeira.

<sup>1</sup> This name is derived from *Malvasia*, a town in the bay of Epidaurus Limera, in the Morea, whence the grape was originally derived, but now producing no good wine.



Where produced.	Generic Names.	Varieties.	Quantity of Alcohol of spec. grav. 825 in 100 parts.	Qualities.
Cape of Good Hope	Constantia .....	.....	14.50 P.	Sweet, luscious, pungent. Resembles Rhenish. Sweet. Harsh, earthy taste. Yellow or Topaz colour; sweetish; resembles Madeira.
		Red Constantia .....	18.92 B.	
		White Constantia ....	19.75 B.	
	Steen wine .....	.....	10.60 P.	
	Cape Muschat	.....	18.25 B.	
Persia .....	— Madeira .	... (average)	20.51 B.	Resembles Tintilla, with a pitchy taste. Resembles Rhenish.
	Shiraz .....	White ..	19.80 P.	
		Red .....	15.52 B.	
England ..	Grape wine .....	.....	18.11 B.	Various. Brisk like Champagne. Thick, narcotic. Sweet, luscious, flavour of the fruit. Sweet.
	Raisin wine ....	..... (average)	25.12 B.	
	Currant wine ...	.....	20.55 B.	
	Gooseberry wine	.....	11.84 B.	
	Elder wine .....	.....	9.87 B.	
	Orange wine ...	.....	11.26 B.	
	Cyder .....	.....	9.87 B.	
	Perry .....	.....	7.26 B.	
	Mead .....	.....	17.32 B.	
	Sycamore wine {	<i>juice fermented with sugar.</i>		
Barbary ..	Usuph..... {	<i>water in which raisins are steeped.</i>		Strong and harsh.
Nepaul ...	Sihee .....	<i>a grape wine.....</i>	—	
Hindostan	Tari .....	<i>fermented juice of the Palmira tree. Borassus flabelliformis, Callu, Teildy, Saura.</i>		
	Sinday..... {	<i>fermented juice of Elate sylvestris, the wild date.</i>		
China .....	Cha.....	<i>nearly the same as Tari.</i>		
	Mandurin .....	<i>boiled rice fermented.</i>		
Tartary ...	Koumis .....	<i>fermented mare's milk.</i>		
	Airen .....	<i>fermented cow's milk.</i>		
Africa .....	Kanyangtsyen {	<i>the flesh of the lamb fermented with rice and other vegetables</i>		
	Millaffu .....	<i>fermented juice of the palm-tree, Congo.</i>		
	Pombie .....	<i>fermented millet, Caffres.</i>		
Brazil.....	Kooi .....	<i>fermented juice of Apples.</i>		
Mexico ...	Palque .....	<i>fermented juice of the Agave.</i>		
Norway ...	Birch wine... {	<i>juice of Betula alba fermented with sugar.</i>		



*Medical properties and uses.* — Wine when good, and of a proper age, is cordial and tonic; but when new it is flatulent, debilitating and purgative, and intoxicates sooner than old wine. In a dietetical point of view, the temperate use of wine promotes digestion, and gives additional energy to the action of the heart and arteries, strengthens the animal functions, exhilarates the spirits, sharpens the wit, and calls into action all the intellectual powers: but when taken in excess, it intoxicates, producing sickness, head-ache, vertigo, and diarrhoea, with nervous tremors, which continue for two or three days; and, like ardent spirit, its habitual, excessive use, extinguishes the faculties of both body and mind, producing dyspepsia, emaciation, and debility, hepatic and pulmonary inflammation, palsy, gout, dropsy, and a long train of diseases and wretchedness. We nevertheless hear of very extraordinary quantities of wine being drank with impunity by some individuals. I knew a man who had not retired sober to bed for twenty years; and yet lived to upwards of eighty years of age.<sup>1</sup> Drunkenness, however, is the vice of barbarians; and, as nations merge from that state, it evidently becomes less prevalent. In Britain it is now happily confined to the dregs of the people.

As a remedy, wine is stimulant, antiseptic, tonic, and antispasmodic. Its stimulating properties are less diffusible, but more permanent than alcohol; and thence its dose is more easily regulated, and its effects are more certain. In all diseases accompanied with much debility, as typhoid fevers, and in cases of extensive ulceration or gangrene, wine is not only the best addition to cinchona bark and opium, but is a remedy on which alone there is much reliance; in some convulsive affections, as symptomatic tetanus, and chorea, much benefit has been derived from its use; and in the convalescencies from all severe diseases, it is the most efficacious and the quickest mean we can employ for restoring the exhausted strength and vigour. Wine operates less powerfully on the system in a state of disease than in health; the quantity, however, to be given, and the proper period of exhibiting it, require to be regulated with much judgment. The skin being open, and not dry or hot, the strength sinking, and the ulcerations, if any exist, assuming a gangrenous appearance, indicate the use of wine; and when, in the event of the pulse being low and fluttering, wine restores its firmness without

---

<sup>1</sup> The Emperor Maximin could quaff six gallons of wine at a sitting; and a Mr. Vanhorn, of modern notoriety, drank in the course of twenty years, 35,688 bottles, or 50 pipes of red port. In the reign of Henry the Eighth, wine and beer were used at breakfast, and the quantity served to one person was a pint of each.



increasing delirium, and induces sleep, it may be given with a confidence of the greatest benefit. But if, on the contrary, it renders the pulse quicker, increases heat, thirst, delirium, or watchfulness, its exhibition ought immediately to be discontinued. The quantity to be given depends entirely on the nature of the disease, and the intentions for which it is administered. In typhus, the proper rule is to give it till the pulse fills, the delirium abates, and the extremities warm; and it should be repeated on the smallest appearance of stupor, quick and sinking pulse, or tremor.<sup>1</sup> A few glasses, and these even diluted with water, given in the space of twenty-four hours, will often produce all that is required from wine; but sometimes very large quantities are necessary. In a case of symptomatic tetanus, mentioned by Currie<sup>2</sup>, five bottles of Madeira wine were taken every day for some time, without producing the least symptoms of inebriety, or morbidly exciting the pulse; but on the contrary, with the utmost advantage in allaying irritation, and relieving the patient. In ordinary cases of fevers however, wine is, perhaps, in general too freely given, so as to occasion exhaustion instead of supporting strength. In a dietetical point of view, wine is useful or prejudicial, in proportion as the fermentation is more or less perfect. In wines containing the malic acid, when the fermentation has been imperfect, the fermentative state of the liquor is recommenced in the stomach and much carbonic acid and other gases are evolved, which, distending that organ, oppresses the individual; and, if he be dyspeptic, produces depression of spirits and all the horrors of hypochondriasm. When wine, however, is good, it stimulates gratefully the nerves of the stomach, and consequently, promoting a more healthy secretion of the gastric fluid, greatly assists the chymification of the food. *Sweet wines* are more apt to become acescent than the dry wines; but it is erroneously conjectured that the same objection applies to the use of the *Rhenish wines*; for, as these wines contain more free tartaric acid, and scarcely any malic or gallic acids, they are less liable to ferment than many of the stronger wines. Sparkling, brisk wines, such as Champagne, intoxicate more speedily than the stiller wines, but the morbid excitement is of short duration, and the subsequent exhaustion is comparatively trifling. They are said to induce gout in those strongly predisposed to that disease; but, probably, more is to be attributed to the luxurious dishes which generally accompany the use of Champagne, than to that exhilarating beverage. In febrile habits, Burgundy, Port

---

<sup>1</sup> *Moore's Medical Sketches*.

<sup>2</sup> *Reports on Water*, i. 174.



and the stronger white wines are to be avoided; but in diseases of debility, particularly where the stomach requires the aid of a tonic, these wines will prove beneficial. In convalescence, however, from acute diseases, if wine be admissible, Claret or some of the better kinds of Rhine wines, for example, Moselle or Hock, are preferable; and this is particularly the case, if with a low pulse and much exhaustion, the nerves are so excitable as to produce a febrile action in the arterial system, on the application of stimulants, either corporeal or mental. In those who have a disposition to obesity, the Rhine wines, on account of their diuretic properties, are preferable to every other kind for daily use. Where health abounds, wine is altogether unnecessary; but, as habit has rendered the use of it general, it is to be lamented that, in this country, the high prices of the more wholesome kinds force the great majority of the middling ranks to indulge in the use of those which contain too large a quantity of alcohol, and the lower classes to the abuse of spirituous liquors.

When wine is prescribed as a cordial in a state of convalescence from acute diseases, or in a weakened state of the habit, it should not be taken with dinner or any other meal, but at noon, upon an empty stomach.

ULMUS. *Spec. Plant. Willd.* i. 1324.

*Cl.* 5. *Ord.* 2. Pentandria Digynia. *Nat. ord.* Scabridæ, *Linn.* Amentaceæ, *Juss.*

*G.* 505. *Calyx* five-cleft. *Corolla* none. *Capsule* (samara) compressed, membranaceous.

*Species* 1. *U. campestris*.<sup>1</sup> Common Elm. *Med. Bot.* 2d edit. 710 t. 242. *Smith, Flor. Brit.* i. 281.

*Officinal.* ULMI CORTEX, *Lond.* ULMI CAMPESTRIS CORTEX, *Edin.* ULMUS; CORTEX INFERIOR, *Dub.* Elm Bark.

*Syn.* Orme (*F.*), Ulmrinde (*G.*), Olm (*Dutch*), Alm (*Dan.*), Kara Wiazowa (*Pol.*), Olmo (*I., S., Port.*), Ilim (*Russ.*).

The Elm-tree is indigenous, and very abundantly cultivated<sup>2</sup>, flowering in March or early in April, before the leaves are unfolded. It grows to a considerable height, sending off strong, spreading, lateral branches; with the bark of the trunk very rough and cracked, but that of the younger branches smooth and tough. The leaves are rough on both sides, villose beneath along the veins, doubly serrate, longer on one side of the midrib than on the other, about three inches long, two broad, and of a dark-green colour. The flowers, which appear before the leaves, are in distinct gems, clustered, scarcely peduncled, numerous, small, of a brownish-flesh

<sup>1</sup> Πτελεα Græcorum.

<sup>2</sup> The age of the Elm is, yet, unknown. Several stocks of it which were planted in France, by the orders of Sully in 1580, still survive. (1824.) They are from 80 to 90 feet high, and from 25 to 30 in circumference.



colour, and have a violet odour. The capsules are oval, oblong, bordered, and contain a single roundish seed.

The inner part of the bark of the younger branches, which is of a yellowish colour, is the part officinally used, and is sold freed from the epidermis.

*Qualities.* — Elm-bark is inodorous, and has a slightly bitter, slimy taste. When boiled, in a small quantity of water, it forms a thick dark-brown coloured decoction, which gelatinizes as it cools; and when evaporated leaves a brittle, semi-transparent substance, soluble in water, but insoluble in alcohol and ether, to which, however, it imparts a brownish colour. The brittle residue, when treated in the same manner as Klaproth treated the gum-like exudation from the *Ulmus nigra*, afforded nearly the same results<sup>1</sup>; and consequently it must be regarded as *Ulmin*: but from the effects of some re-agents (see *Decoction*), I am inclined to consider it a peculiar modification of mucus, combined with extractive, gallic acid<sup>2</sup>, and super-tartrate of potass, which Schæele detected in elm bark.

*Medical properties and uses.* — This bark operates as a diuretic. It has been given with seeming benefit in herpetic eruptions; and Dr. Lettsom<sup>3</sup> attributes the cure of a severe case of “lepra ichthyosis,” in which other remedies had failed, to the use of this bark. Other practitioners have also related cases of its efficacy; but Dr. Willan<sup>4</sup> thinks it is of little use. It is generally given in the form of decoction.

Officinal preparation. *Decoctum Ulmi*, L. D.

UVÆ PASSÆ. Vide *Vitis*.

UVÆ URSI FOLIA. Vide *Arbutus*.

WINTERA. *Spec. Plant. Willd.* ii. 1239.

Cl. 13. Ord. 4. Polyandria Tetragynia. Nat. ord. Oleraceæ, Linn. Magnoliaceæ, Juss.

G. 1063. Calyx three-lobed. Petals six or twelve. Germens club-shaped. Style none. Berries four or eight, obovate.

Species 1. *W. aromatica*. Winter's Bark-tree. *Med. Bot.* 2d edit. 647. t. 226. *Phil. Trans.* xvii. 923. t. 1. f. 1, 2.

Officinal. WINTERÆ AROMATICÆ CORTEX, *Edin.* Winter's Bark.

*Syn.* Cannelles de Winter (*F.*), Winterana (*L.*), Corteza Winterana (*S.*).

The tree is a native of the Straits of Magellan, growing in valleys which are exposed to the sun. It is a large ever-green tree; covered on the trunk with a grey, wrinkled bark, which on the branches is green and smooth. The leaves are petio- late, elliptical, obtuse, smooth, an inch and a half in length, an inch broad in the middle, and of a light-green colour.

<sup>1</sup> Thomson's *Chemistry*, 4th ed. iv. 695.

<sup>2</sup> Very little tannin is present, as it scarcely effects a solution of gelatine.

<sup>3</sup> *Medical Memoirs*, 152. <sup>4</sup> *Description, &c. of Cutaneous Diseases*, i. 139.



The flowers are axillary, two, three, or more together, on short peduncles of a milk white colour, with the odour of jasmine; the petals are unequal, oval, obtuse, concave and erect; the filaments shorter than the petals, supporting large, oval anthers, and the germens turbinated, with sessile, divided, flat stigmas. The berries are of a light-green colour, spotted with black, and contain several black, aromatic seeds.

This tree was discovered in 1577, by Captain Winter, the crew of whose ship used the bark as spice. It is not often found in the shops; and is frequently confounded with the *Canella alba*, from which it may be distinguished by being in larger pieces, and having more of a cinnamon hue.

*Qualities.* — Winter's bark has an aromatic odour: and a pungent, hot, spicy taste, slowly imparted, but very permanent. These qualities depend on a volatile oil, which can be obtained separate, in distillation with water. M. Henry obtained also resin, tannin,<sup>1</sup> a colouring matter, acetate of potass, muriate of potass, sulphate of potass, oxalate of lime and oxide of iron, in this bark.

*Medical properties and uses.* — This bark is stomachic and carminative. It has been found efficacious in scurvy, and may be used as an adjunct to simple bitters in dyspepsia; but it does not appear to be superior to *Canella alba*, and is very little used. On the continent it is given in the forms of powder, tincture, and infused in wine.

## ZINCUM. Zinc.

*Syn.* Zinc (*F.*), Zink (*G.*), Zinco (*I.*), Spiagter (*Russian*), Tootanāgum (*Tam.*), Sungbusrie (*Duk.*).

Zinc is a semi-ductile metal procured in great abundance in Britain, particularly in Derbyshire; and in most of the mining countries of Europe. It occurs in

### A. The metallic state.

i. combined with sulphur and iron. SP. 1. *Blende*.

Var. *a.* Yellow blende.

*b.* Brown blende.

*c.* Black blende.

### B. Oxidized:

ii. combined with silica.

1. *Red Zinc ore*.

iii. acidified by carbonic acid.

2. *Electric calamine*.

3. *Common calamine*.

Var. *a.* crystallized.

*b.* compact.

*c.* earthy.

As the fourth species of these ores is an article of the *Materia Medica*, we shall describe its characters and properties before we notice those of metallic zinc.

<sup>1</sup> *Canella alba* does not contain tannin.



## 1. COMMON CALAMINE.

*Officinal.* CALAMINA. *Carbonas Zinci impura*, Lond. CARBONAS ZINCI IMPURUS, *Edin.* CALAMINARIS, *Dub.* Calamine. Impure Carbonate of Zinc.

*Syn.* Pierre Calaminaire (*F.*), Galmey (*G.*), Kalmei (*Belg.*), Gallmeja (*Swed.*), Pietra Calaminare (*I.*), Calamina (*S.*), Piedra Calaminar (*Port.*), Madal tootum (*Tam.*).

This ore of zinc is found abundantly in Derbyshire, Somersetshire, Cumberland, and Flintshire, occurring in veins in secondary limestone, generally accompanied by galena, calcareous spar, quartz, and other ores of zinc. The three varieties are indiscriminately used; and consist, according to an analysis by Mr. Smithson<sup>1</sup>, of the following components: *var. a.* 65·2 oxide of zinc, 34·8 carbonic acid; *var. b.* 64·8 oxide of zinc, 35·2 carbonic acid; *var. c.* 71·4 oxide of zinc, 13·5 carbonic acid, 15·1 water,—in 100 parts of each variety. They are, however, generally calcined in a moderate heat, by which part of their carbonic acid is dissipated, before they are brought to the shops.

*Qualities.* — Calamine is usually in the form of grayish-yellow or reddish-yellow, friable lumps, without lustre, opaque, and breaking with an irregular, earthy fracture. The specific gravity of the two first varieties is 4·334: that of the last 3·584. Before the blowpipe, Calamine becomes yellow; and when exposed to its utmost heat, is sublimed. It dissolves in sulphuric acid with effervescence, but does not gelatinize. It is not used as a remedy till after it is prepared: and then only as an external application.

*Officinal preparation.* *Calamina preparata*, L. E. D.

## 2. METALLIC ZINC.

*Officinal.* ZINCUM, *Lond. Edin. Dub.* Zinc.

Although the method of extracting Zinc from its ores had been long known and practised in India and China, yet it was not known in Europe till about 1721, when Henke pointed out a method of extracting it from its ores. Von Swab first obtained it by distillation in 1742. The Greeks certainly were not acquainted with Zinc, although they employed Cadmia, which contained Zinc, in the manufacture of brass. At present it is well understood, and conducted in the following manner:—The sulphuret or blende, which is the ore usually employed, is first broken to pieces, and the galena and pyrites separated by hand; and is then roasted in a reverberatory furnace, by which the carbonic acid and part of the sulphur are driven off. The roasted ore being washed, to separate the metallic particles from the lighter parts, is now ground

<sup>1</sup> *Phil. Trans.* 1803. 17.



in a mill with one-eighth of its weight of charcoal; and put into large, earthen jars placed in a circular furnace, and through the bottom of each of which passes an iron tube which goes through the floor of the furnace into a vessel of water placed beneath. The cover of each jar is firmly and accurately luted on, so that the reduced zinc, as it is elevated by the strong heat of the furnace, not finding a vent to escape by the top, descends through the iron tube into the water, and is there condensed in small metallic drops, which are afterwards melted and cast into ingots, in which state it is brought to market<sup>1</sup>, under the name of *Speltre*.

*Qualities.* — Zinc when rubbed between the fingers, emits a very perceptible odour and has a peculiar taste. Its colour is brilliant, white with a shade of blue, and its fracture shining and lamellated; hard, yet staining the fingers black when rubbed upon them. Its specific gravity varies from 6·861 to 7·1. In any temperature between 212° to 400°, it is very malleable; but at a higher temperature, it can be pulverized in a mortar. It may be drawn into wire, but its ductility is not great. Zinc melts at 680° of Fahrenheit; if in contact with air, it is rapidly oxidized: and, at the temperature of ignition, burns with a white dazzling flame, and is volatilized in the state of a flocculent, white oxide. It is oxidized and soluble in all the acids; and decomposes water when aided by a small portion of sulphuric or of muriatic acid. It is used only for pharmaceutical purposes.

Officinal preparations. *Zinci Oxydum*, L. E. D. *Zinci Sulphas*, L. E. D.

### 3. IMPURE OXIDE OF ZINC.

*Officinal.* OXIDUM ZINCI IMPURUM, *Edin.* TUTIA, *Dub.* Impure OXIDE of Zinc. Tutty.

*Syn.* Tutie (*F.*), Tutia (*G.*), Tuzia (*I.*), Atutia (*S.*).

This substance is supposed to be an artificial compound of the sublimed oxide of zinc that collects in the chimneys of the furnaces in which the ores of this metal are roasted, mixed with clay and water and baked.

*Qualities.* — Tutty is inodorous and insipid, of a brownish colour on the outside, moderately hard and ponderous, and breaks with a smooth fracture. The internal colour is yellowish; and the pieces sometimes contain small globules of zinc. The oxide it contains consists of 85 zinc, 15 oxygen,—in 100 parts. It is not employed as a remedy until it is levigated and prepared.

Officinal preparations. *Oxidum Zinci impurum preparatum*, E. *Unguentum Oxidi Zinci impuri*, E. D.

<sup>1</sup> The principal works are near Bristol, and at Swansea.



**ZINGIBER.** *Trans. Linn. Society*, viii. 347.

*Cl.* 1. *Ord.* 1. Monandria Monogynia. *Nat. ord.* Scitamineæ, *Linn.* Cannæ, *Juss.*

*G. novum.* *Anther* double. *Filament* lengthened beyond the anther with a furrowed awl-shaped apex. *Style* received in the furrow of the anther.

*Sp.* 1. *Z. officinale*.<sup>1</sup> *Officinal Ginger.* *Jacquin Hortus Vindobonensis*, i. 31. t. 75. (*Amomum Zingiber*) *Willd. Spec. Plant.* i. 6. *Med. Bot. 2d edit.* 731. t. 250. *Rumph. Amb.* ii. t. 12.

*Officinal.* ZINGIBERIS RADIX, *Lond.* AMOMI ZINGIBERIS RADIX, *Edin.* ZINGIBER; RADIX CONDITA, *Dub.* Ginger root dried and preserved.

*Syn.* Gingembre (*F.*), Ingwer; Imber (*G.*), Zenzero (*I.*), Gengibre (*S.*), Gengivre (*Port.*), Gember (*Dutch*), Ingefer (*Dan.*), Ingefära (*Swed.*), Jembier (*Pol.*), Sont'h (*H.*), Sunt'hi (*Sans.*), Dshey (*Javanese*), Sookkoo (*Tam.*), Alia (*Malay*), Zungebell (*Pers.*), Siwe (*Amb.*).

The Ginger plant is a native of the East Indies<sup>2</sup>, and is particularly abundant in the mountainous district of *Gingi*, to the east of Pondicherry, whence it derived its name. It is now naturalized to the West Indies, where it flowers in September. The root is perennial, creeping, of a compressed roundish form, or tuberosc, fleshy, and sending off many long fibres and offsets; the stem is annual, rises about three feet in height, is an annual culm, solid, upright, round, and enclosed in an imbricate, membranous sheathing. The leaves are alternate, linear-lanceolate, six inches long, smooth, and on short embracing petioles. The flowers are in a dense spike close to the stem, composed of large, upright, ovate, subcuneate, coloured scales, half closing the flowers: the calyx is a small double spathe of a dingy-yellow colour, tubular, with the segments of the border conical, and nearly equal: the anther-bearing filament is extended beyond the anther, has an awl-shaped appendage with a groove to receive the style after it has passed between the lobes of the anthers, and is terminated with the stigma a little beyond the extremity of the filament: the capsule is smooth, containing many oblong seeds.

The herbaceous part of the plant withers in December, and the roots are dug up in January; but when the root is intended to be preserved in syrup, it is dug up when the shoots do not exceed five or six inches in height. For preparing the dried Ginger, after the roots are dug, the best pieces are selected, scraped, then washed, and dried in the sun with great care. This is called *White Ginger*; in contradistinction to which, the roots that are scalded in boiling

<sup>1</sup> *Ζιγγίβρις* Dioscoridis. The similarity between the Greek name of the plant, and the Sanscrit *Sringavera* (horn-shaped) is remarkable. *Asiatic Researches*, vol. xi. p. 346.

<sup>2</sup> It is named *alè* by the Brahmins.



water before being dried, are denominated *Black Ginger*. The confected or preserved Ginger is prepared by scalding the green roots till they are tender; then peeling them in cold water, and putting them into a thin syrup, from which in a few days they are shifted into the jars in which they come home, and a very rich syrup poured over them.

Dried Ginger is imported in bags, each containing about one hundred weight. The white kind brings the highest price, being more pungent and better flavoured. The external characters of goodness in either are soundness, or the being free from worm holes; heaviness and firmness: the pieces that are light and soft, or very friable and fibrous, should be rejected. The confected Ginger is nearly translucent when good.

*Qualities.* — Dried Ginger has a pungent, aromatic odour, and a hot, biting taste. Its odour appears to depend on a volatile oil, which can be obtained separate in distillation with water, and has all the flavour, but none of the pungency of the root. Water, alcohol, and ether extract its virtues. The greater part of Ginger root, however, is starch. To separate it, triturate the root with water, and strain through cloths; then after the fecula, which is suspended in the water, has subsided, separate it by decanting off the water, and macerate in alcohol; what remains undissolved is a tolerably pure, insipid starch. The pungency resides in a resino-extractive matter, which is combined with the fecula, but may be obtained separate by evaporating the ethereal tincture on the surface of water. M. Morin and Rouen have analysed Ginger, and give the following as its components:—Resin soluble in ether, resinous matter insoluble in ether, a blueish-green volatile oil, vegeto-animal matter, a substance analogous to osmazome, acetic acid, acetate of potass, starch, gum, sulphur, several oxides, and lignin.<sup>1</sup>

*Medical properties and uses.* — Ginger is stimulant, carminative, and sialogogue. It has been found useful in flatulent cholera, dyspepsia, and tympanitis; and in gout when it attacks the stomach. It is less frequently used alone than as an adjunct to other remedies to promote their efficacy, and give them warmth. The local stimulus of Ginger when chewed excites the salivary glands, and provokes a considerable flow of saliva: hence it has been found useful as a sialogogue in relaxations of the uvula and tonsils, and in paralysis of the muscles of the tongue and fauces.

The dose of powdered ginger may be from grs. x. to ℥j.

---

<sup>1</sup> *Journ. de Pharm.* June, 1823.



Officinal preparations. *Syrupus Zingiberis*, L. E. D. *Tinctura Zingiberis*, L. D. *Syrup. Rhamni*, L. *Tinct. Cinnamoni comp.*, L. *Acid. Sulphur aromat.*, E. *Confectio opii*, L. *Confect. Scammonii*, L. D. *Infusum Sennæ*, L. *Pulvis Cinnam. comp.*, L. E. D. *Pulv. Scammonii comp.*, L. D. *Pulv. Sennæ comp.*, L. *Pilula Scillæ comp.*, L. *Pilula Aloes*, D. *Vinum Aloes*, L. E. D.



## PART III.

### PREPARATIONS AND COMPOUNDS.

---

#### ACIDS.

THESE are substances which have a sour taste: and are capable of combining with alkalis, earths, and metallic oxides, while at the same time they lose their acidity, and form compounds named neutral salts, in which the properties of the acid, the alkali, the earth, or the oxide employed, are lost. They change to red, the blue, purple, and green colours of vegetables; and unite with water in almost every proportion. These circumstances, therefore, may be regarded as characteristic of this class of substances.

Acids were supposed by Lavoisier to be combustible bodies combined with oxygen, which he regarded as the principle of acidity; and this opinion was almost generally adopted until lately, when the discovery of *chlorine*, and the promulgation of Sir H. Davy's theory of the nature of muriatic acid, led to the revival of Berthollet's doctrine, "that no substance as an acidifying principle exists; but that the nature of a compound depends upon the way in which the constituents unite together." It is now, therefore, no longer questionable, whether there is any acid that does not contain oxygen as an essential component; and it is by no means true that the combination of every combustible with oxygen will constitute an acid.<sup>1</sup> Thus the combination of hydrogen with oxygen forms water which is not acid, and that of sulphur with hydrogen forms a body, which really possesses acid properties, but nevertheless contains no oxygen; and it is now known that the alkalis owe their alkaline properties to the presence of oxygen, as much as the acids owe to it their acid properties.<sup>2</sup> For the formation of an acid, however, although the union of the body with oxygen be not essential, yet it is not improbable, that some principle must be present in the body, common, in a greater or less degree, to all bodies capable of being formed into acids, which may be regarded as the principle of acidity, although it be yet unknown.

On the supposition that all acids are compounds of oxygen.

---

<sup>1</sup> Acids containing no oxygen—Muriatic, Prussic.

<sup>2</sup> See Davy's Discoveries, *Phil. Trans.* 1808.



with certain bases, the name of each has been derived from the base of which it is formed; from *sulphur*, for instance, comes *sulphuric acid*: but the same base being supposed capable of uniting with different proportions of oxygen, the terminations *ous* and *ic* were added to indicate the degree of acidification: thus, when sulphur is united with the smaller proportion of oxygen, the acid produced is named *sulphurous acid*; when with the full proportion *sulphuric acid*. One or two acids were moreover supposed to combine with a still larger proportion of oxygen, to denote which the syllable *oxy* (for oxygenized) was prefixed: thus muriatic (*hydrochloric*) acid, which was supposed to be combined with an excess of oxygen, became oxymuriatic acid.

It is unnecessary to enter into the consideration of the impropriety of these terms in a chemical point of view; they are those employed in the British Pharmacopœias, and are applied to well known substances, in daily use in the practice of medicine.

The stronger acids require to be kept in glass bottles, furnished with well ground glass stoppers, and having the name of the acid which each contains engraved on the glass. They should be dispensed also in glass stopped phials. The acids known to chemists are very numerous: but of these a small proportion only is employed for medical and pharmaceutical purposes. In the London Pharmacopœia, the arrangement of which we have adopted, they are placed in alphabetical order; but in this place it may be proper first to exhibit them according to the nature of their radicals or bases.

#### I. ACIDS COMPOSED OF A SIMPLE RADICAL AND OXYGEN.

SULPHUR

1. SULPHURIC ACID.

AZOTE

{ 2. NITROUS ACID.  
3. NITRIC ACID.

#### II. ACIDS COMPOSED OF A COMPOUND RADICAL AND OXYGEN.

CARBON AND HYDROGEN

{ 1. ACETIC ACID.  
2. CITRIC ACID.  
3. BENZOIC ACID.  
4. SUCCINIC ACID.

#### III. ACIDS COMPOSED OF A SIMPLE RADICAL AND CHLORINE.

HYDROGEN

1. MURIATIC ACID.

ACIDUM ACETICUM DILUTUM, Lond. *Diluted Acetic Acid.*

“ Distil the dilute Acetic acid from a glass retort, placed in a sand bath, into a glass receiver kept cool: throw away



the first pint, and preserve the six succeeding pints which are distilled.”

ACIDUM ACETICUM TENUE, Edin. *Weak Acetic Acid.*

“Distil eight pounds of Vinegar in glass vessels, with a gentle heat. The pound which first comes over, being too watery, is to be rejected; the five pounds that follow, will be the weak acetic acid. The distillation may be continued as long as a colourless acid comes over; but this, being too much burnt, and unfit for internal use, may be mixed with the pound which first came over, and preserved for various chemical purposes.”

ACETUM DISTILLATUM, Dub. *Distilled Vinegar.*

“Take of Wine Vinegar, *ten pints*; distil with a gentle heat *six pints*. The distillation must be performed in a glass vessel, and the first pound which comes over rejected.”

The specific gravity of this acid is to that of water as 1006 to 1000.

*Syn.* Vinaigre distillé (*F.*), Distillirter Essig (*G.*), Azynagtige zuur (*Dutch*), Aceto distillato (*I.*), Vinaigre distilado (*S.*).

Of the three appellations given to this preparation in the British Pharmacopœias, that of the Edinburgh College only is unobjectionable. It is the acetic acid in a more diluted state than that in which it exists in vinegar, but purer; being freed in a great degree from the mucilage, extractive, super-tartrate of potass, and other extraneous matters which vinegar contains.<sup>1</sup> The distillation on a large scale is often conducted in the common copper still, or in a tinned still with a pewter worm; but both these practices, as they are apt to give dangerous metallic impregnations to the product, are highly reprehensible, when the distilled vinegar is to be employed for medicinal purposes.

In performing the process, it is more important to avoid carrying the distillation too far than to reject the first eighth part which comes over; for although this is undoubtedly weaker, and contains a small portion of alcohol, yet it retains about one-twelfth<sup>2</sup> of the whole quantity of the real acid obtained: but by continuing the process a little too long, the whole product acquires an unpleasant empyreumatic flavour. This is avoided by changing the receiver rather before the quantity ordered has been obtained; and, if to the residue be added an equal quantity of hot water and half an ounce of recently burnt animal charcoal<sup>3</sup>, for every pint of fluid in the

<sup>1</sup> See *Acetum*, Part II.

<sup>2</sup> *London Medical Review*, No. x. 125.

<sup>3</sup> Animal charcoal is made by calcining bones in a crucible, to which is luted a perforated cover. When no more flame issues from the aperture, it must be closed, and the heat raised for half an hour.



retort, the distillation may be recommenced, and an additional portion of the diluted acid obtained, equally pure and strong as the former. At the end of the operation, when dilution has not been employed, the residue is a deep-red coloured liquor, strongly acid, very empyreumatic, and which deposits supertartrate of potass. Mr. Brande justly regards as an improvement in the distillation of vinegar, the employment of the heat of high pressure steam, instead of that of an open fire; as both the risk of empyreuma is prevented, and “a larger portion may usually be distilled off, before any foreign flavour is perceptible.”<sup>1</sup>

*Qualities.* — Distilled vinegar has a fainter and less agreeable odour than common vinegar; a grateful, not strong, acid taste: is limpid and colourless; and of a specific gravity varying from 1·007 to 1·0095. One fluid ounce of spec. grav. 1·007 should dissolve 13·8 grains of marble; or 1000 grains should dissolve 145 grains of crystallized subcarbonate of soda, for their saturation. It dissolves the gum resins and the active principles of plants, such as those of the squill and the meadow-saffron; and forms acetates with the alkalies and several of the metallic oxides. Darracq<sup>2</sup> ascertained that it differs from acetic acid in containing some uncombined mucilage and extractive matter, but that the acids are otherwise the same. According to Mr. Phillips, the acid of sp. gr. 1·007 contains of acetic acid 3·42, water 96·58, in 100 parts; and that of sp. gr. 1·009, acetic acid 4·73, water 95·27. It is owing to the extractive that, when distilled vinegar is boiled with potass, the solution has a deep reddish-brown colour, and, during evaporation, carbonaceous matter is deposited. It is sometimes adulterated. Sulphuric acid is detected by a precipitate being produced on the addition of a solution of acetate of barytes; lead and tin, by a solution of sulphuretted hydrogen, forming a dark-coloured precipitate; and copper, by the acid assuming a blue colour, when supersaturated with ammonia.

*Medical properties and uses.* — The same as those of common vinegar; but, as it is purer, and not liable to spontaneous decomposition, it is fitter for pharmaceutical purposes.

*Officinal preparations.* *Liquor Ammoniacæ Acetatis*, L. E. D. *Acetas Ferri*, D. *Liquor Plumbi Subacetatis*, L. D. *Acetum Colchici*, L. *Acetum Scillæ*, L. E. D. *Oxymel*, L. D. *Emplastrum Ammoniaci*, L. *Oxymel Colchici*, D.

ACIDUM ACETICUM, Dub. *Acetic Acid*.

“Take of Acetate of Kali (Potash), *six ounces*; Sulphuric acid, *three ounces by weight*. Pour the acid into a tubulated

<sup>1</sup> *Manual of Pharmacy*, p. 192.

<sup>2</sup> *Annales de Chimie*, xli. 264.



retort: then add to it in small portions and at different times, the Acetate of Potass, allowing the mixture to cool after each addition; finally, with a moderate heat, distil the acid until the residue be dry.

“The specific gravity of this acid is to that of distilled water, as 1070 to 1000.”

ACIDUM ACETICUM FORTE, Edin. *Strong Acetic Acid*.

“Take of dried Sulphate of Iron, *one pound*; Acetate (*superacetate*) of lead, *ten ounces*. Rub them together; then put them into a retort, and distil in a sand bath with a moderate heat, as long as any acid comes over.”

*Syn.* Acide Acetique (*F.*), Essigsäure (*G.*), Azynzuur (*Dutch*), Acido acetico (*I.*).

These processes furnish the same acid as that which is contained in distilled vinegar, but completely free from extractive and mucilage, and much stronger. In the process of the Dublin College, the sulphuric acid, by reason of its superior affinity for potass, decomposes the acetate of potass, and sets free the acetic acid; which must necessarily come over in a concentrated form, as the acetate does not contain much water, and the greater part of the water of the sulphuric acid is retained by the sulphate of potass which is formed in the retort. The Edinburgh process is that of Badollier, an apothecary of Chartres<sup>1</sup>, with the substitution of sulphate of iron for sulphate of copper. The affinity of the sulphuric acid of the sulphate of iron to the oxide of the acetate of lead, assisted by the heat, decomposes the acetate, and the sulphuric acid, uniting with the lead, the acetic acid is disengaged, and distils over. The dried state of the salts enables the acid to be obtained in a concentrated form.<sup>2</sup> The process of the Dublin College is the best; the acid obtained being stronger, not contaminated with any metallic impregnation, and free from sulphurous acid, a small portion of which the other generally contains.<sup>3</sup>

*Qualities.* — Acetic acid has a grateful, fragrant, pungent odour, a very sour, and acrid taste; and when applied to the skin, inflames it and raises a blister. It is limpid, colourless, highly volatile, and if much concentrated, takes fire when

<sup>1</sup> *Annales de Chimie*, xxxvii. 3.

<sup>2</sup> The process of the edition of the London Pharmacopœia of 1787, for obtaining this acid by the simple distillation of dried, subacetate of copper, afforded a much stronger acid than either of these processes. An excellent acetic acid is obtained by decomposing the acetate of lead at once, by means of sulphuric acid; putting into the retort, before distilling, a small portion of oxide of manganese, as directed by M. Baup. Vide *Journ. de Pharm.* Dec. 1816.

<sup>3</sup> It may be completely freed from this acid by re-distilling it from black oxide of manganese mixed with a small portion of carbonate of potass. *Nicholson's Journal*, xiii. 42.



heated in the open air. The sp. grav. of the strongest that can be procured is 1.063; in which state it contains 100 parts by weight of real acid and 14.78 of water.<sup>1</sup> It becomes solid and crystallizes at 28° of Fahrenheit, liquefying again at 40°; is capable of oxidizing iron, zinc, copper, nickel, and tin; combines with alkalies, earths, and metallic oxides, forming acetates; dissolves resins, gum-resins, camphor<sup>2</sup>, and volatile oils; and combines with alcohol, which, when aided by heat, it converts into a species of ether. The dry acid saturates its own weight of carbonate of lime, so that the strength of any sample of this acid may be readily determined.<sup>3</sup> With water it unites in any proportion, and, during the mixture, heat is evolved. From the experiments of Gay Lussac 100 parts of pure Acetic acid appear to consist of 50.224 of carbon, 5.629 of hydrogen, and 44.147 of oxygen.

*Medical properties and uses.* — Acetic acid is stimulant and rubefacient. It is principally employed as a refreshing scent, in syncope, asphyxia, and nervous head-achs; and for obviating the unpleasant smell of the confined air of crowded assemblies and of the sick-room. It may be used as a counter-irritant, when applied to the skin, in cynanche tonsillaris, croup, and other internal inflammations. It is, also, an excellent application to warts and corns, which it seldom fails to remove; but, in applying it, care must be taken to avoid eroding the surrounding skin. It is a useful and powerful substitute for a blister, when the effect of such an application is required to be rapidly obtained.

Officinal preparation. *Acidum aceticum camphoratum*, E. D.

ACIDUM BENZOICUM, Lond. *Benzoic Acid*.

“Take of Benzoin, *one pound*. Put the Benzoin into a glass vessel placed in a sand bath, and exposed to a heat of 300° gradually increased, and sublime until nothing more ascends; press the sublimed matter between bibulous paper, in order to separate it from the oil: then sublime it again with a heat not exceeding 400.°

*Edinburgh.*

“Take of Benzoin, *twenty-four ounces*; Subcarbonate of Soda, *eight ounces*; Water, *sixteen pounds*. Triturate the balsam with the subcarbonate; then boil them in the water for half an hour, stirring them constantly, and strain. Boil the residue of the balsam in other six pounds of water, and strain.

<sup>1</sup> *Ann. de Chim.* lxxiii. 88.

<sup>2</sup> *Henry's Aromatic Vinegar* is a solution of camphor, and some essential oil, in acetic acid.

<sup>3</sup> *Brande's Manual of Pharm.* p. 7.



Mix the strained liquors, and evaporate to two pounds: filter again, and drop in diluted sulphuric acid as long as any precipitation is produced.

Dissolve the precipitated Benzoic acid in boiling water: strain the liquor whilst it is hot through linen, and set it aside to crystallize. Wash the collected crystals with cold water; then dry, and preserve them for use."

*Dublin.*

"Take of Benzoin, *any quantity*. Liquefy it in a wide-necked retort, to which a receiver is adapted but not luted, and sublime. The sublimed matter must be now and then removed from the tube of the retort, lest it accumulate in too great quantity. If it be soiled with oil, press it between folds of blotting paper to separate the oil, and repeat the sublimation."

*Syn.* Acide Benzoïque (*F.*), Benzoessäure (*G.*), Benzoin zuur (*Dutch*), Acido Benzoico (*I.*).

The processes of the London and Dublin Colleges are nearly the same as the method recommended by Chaptal of separating the acid which the Benzoin contains: but although the quantity contained be greater, yet if the fire be not very nicely regulated, so large a portion of empyreumatic oil is also volatilized that it gives the acid a brown tinge, which cannot easily be entirely separated from it.<sup>1</sup> The Edinburgh process is that of Gren, which is a modification of that of Scheele published in 1775.<sup>2</sup> Subcarbonate of Soda separates the Benzoic acid from the resin with which it is united in the Balsam, combines with it, and forms a Benzoate which is dissolved in the water; whilst at the same time a small portion of resin is also dissolved, and gives the solution a yellow colour. This Benzoate is decomposed in its turn by the sulphuric acid, which combines with the alkali, and forms a sulphate; whilst the Benzoic acid that is set free, being insoluble in cold water, precipitates in the form of a brownish powder. The subsequent sublimation frees the acid of this colour, and gives it the crystallized form and brilliant appearance of the acid originally obtained by sublimation, without any adhering oil. The following are the quantities of acid obtained from one pound of Benzoin by these processes, according to Mr. Brande's experiments.<sup>3</sup>

---

<sup>1</sup> This acid was originally obtained by sublimation. It was described under the name of *flowers of benzoin* by Blaise de Vigenève, in 1608. *Thomson's Chemistry*, 4th edit. 289.

<sup>2</sup> *Scheele*, i. 124.

<sup>3</sup> *Nicholson's Journal*, x. 88.



	℥.	ʒ.	ʒ.	grs.
By Chaptal's ( <i>that of the London College</i> )	2	0	0	0
Gren's ( <i>the Edinburgh</i> ) - - -	1	6	2	19 <sup>1</sup>

Benzoic acid may also be extracted from the other balsams.

*Qualities.* — Benzoic acid is, when perfectly pure, inodorous; but as it is usually found in the shops, it has a slight aromatic odour<sup>2</sup>; its taste is pungent, sweetish, acrid, and acidulous. It is in feathery or flocculent crystals, soft to the touch and not pulverulent, of a beautiful whiteness, and a silky lustre. The crystals contain no water. Its specific gravity is 0·657. When heated, it melts, emits a suffocating, acrid vapour, and in a strong heat burns with a white flame. Benzoic acid is soluble in twenty-four times its weight of boiling water; but the water lets fall nineteen twentieths in cooling. Cold alcohol takes about one half its weight; boiling alcohol its own weight of this acid. With the alkalies, earths, and metallic oxides, it forms benzoates, which are not used in medicine. According to Berzelius it is a triple compound of 74·41 of carbon, 5·16 of hydrogen, and 20·43 of oxygen.

*Medical properties and uses.* — This acid is stimulant; but, although it is retained by all the Pharmacopœias, it is of little value as a remedy.

Officinal preparation. *Tinctura Camphoræ composita*, L. D. *Tinctura Opii ammoniata*, E.

#### ACIDUM CITRICUM, Lond. *Citric Acid*.

“Take of Lemon Juice, *a pint*; prepared Chalk, *an ounce*; or a quantity sufficient to saturate the juice; Diluted Sulphuric Acid, *nine fluid ounces*. Add the Chalk by degrees to the Lemon Juice heated, and mix them; then pour off the liquor. Wash the Citrate of Lime which remains, in repeated portions of warm water, and then dry it. On the dried powder pour the Diluted Sulphuric Acid, and boil for ten minutes; express the liquor strongly through a linen cloth, and filter it through paper. Evaporate the filtered liquor with a gentle heat, so that crystals may form as it cools. To obtain the crystals pure, dissolve them in water a second and a third time; filter each solution, boil it down, and put it apart to crystallize.”

*Syn.* Acide Citrique (*F.*), Acido Citrico (*I.*).

<sup>1</sup> M. Stoltze has obtained 180 parts of pure acid from 1000 of Benzoin by the following process. He dissolves the Benzoin in 5 parts of alcohol at 76°, filters the solution, introduces the liquor into a retort, and saturates it with neutral carbonate of soda, dissolved in 8 parts of water and 3 of alcohol. He then distils to obtain the alcohol; and to the decanted, remaining liquor, adds sulphuric acid to separate the Benzoic acid; which is afterwards purified by dissolving it in boiling water and crystallizing. *Berl. Jahrbuch*, 1823. Part II. p. 75.

<sup>2</sup> To free it from the oil on which its odour depends, dissolve it in alcohol, and precipitate by water. *Phil. Mag.* xiv. 331.



This process, which was contrived by Scheele, will seldom require to be performed by the apothecary, as the crystallized acid is now manufactured very pure, and sufficiently reasonable on the great scale.<sup>1</sup> The theory of the process is very simple. The lime of the chalk unites with the citric acid which exists ready formed in the lemon juice, and produces an insoluble citrate of lime, which precipitates united with some of the mucilaginous and extractive matter of the juice. These are separated by repeated washings; and the sulphuric acid, which is added to the dried citrate, decomposing it, owing to the superior affinity of the sulphuric acid for lime, an insoluble sulphate of lime forms, and is precipitated, while the citric acid is disengaged, remains in solution, and is crystallized by evaporation. The crystals of the first crystallization are dark coloured; which is partly owing to a portion of mucilage that still adheres to the citric acid, and partly to the excess of sulphuric acid acting on the citric acid and decomposing a portion of it. The repeated crystallizations free the crystals from this dark colour; but as it is of some importance to avoid any hurtful excess of sulphuric acid, and as the strength of lemon juice is variable and uncertain, it is better to determine the quantity of acid required by the quantity of chalk employed. For this purpose a portion of the sulphuric acid intended to be used must be previously saturated with the chalk, and the weight of the portion employed accurately ascertained; by the knowledge of which the exact quantity of sulphuric acid required to decompose the citrate may be found. According to the experiments of Proust<sup>2</sup>, 94 ounces of lemon juice saturate 4 ounces of chalk with citric acid, and produce  $7\frac{1}{2}$  ounces of dry citrate, which require for their decomposition, and the complete saturation of the lime they contain, 20 ounces of diluted sulphuric acid, composed of one part of the common acid, and three parts of water, or of a specific gravity of 1.15. To ascertain, however, the exact point of saturation of the lime with the sulphuric acid, take a little of the clear supernatant fluid, filter it, and add to it a few drops of acetate of lead; if no sulphuric acid be present, citrate of lead only will be formed, soluble in nitric acid, which does not dissolve sulphate of lead.<sup>3</sup>

*Qualities.* — Pure citric acid is in white, transparent, persistent, right rhomboic prisms; in general, however, variously

---

<sup>1</sup> The principal manufacturer in London, is Mr. Coxwell, of Fleet Street.

<sup>2</sup> *Journal de Physique*, lii. 366.

<sup>3</sup> Citric acid may be also obtained from the juice of the Craberry, *Vaccinium oxycoccus*; Bird cherry, *Prunus padus*; Bitter-sweet, *Solanum dulcamara*; and Dog-rose, *Rosa canina*.



modified. It is inodorous; has an extremely acid, almost caustic taste; and reddens strongly the vegetable blues. In a damp air, the crystals absorb moisture. One ounce of water at 60° Fahrenheit dissolves one ounce and a quarter of this acid; and at 212° twice its weight. The solution, when long kept, is liable to become mouldy and undergo spontaneous decomposition. Citric acid combines with the alkalies, earths, and metallic oxides, and forms citrates. The sulphuric and nitric acids decompose it. Its components, according to Berzelius, are 41·369 carbon, 3·800 hydrogen, and 54·831 of oxygen: and in its crystallized state contains Citric Acid 76·32, water 23·68 in 100 parts.<sup>1</sup> Citric acid may be adulterated with tartaric acid; or with citrate of lime. The first is discovered by adding to the solution of the acid to be tested a solution of nitrate, sulphate or muriate of potass, or saturating it with carbonate of potass, when an insoluble supertartrate, in small brilliant crystals, will be formed if the tartaric acid be present; the second is detected by dissolving the crystals in water, saturating the solution with ammonia, and adding to it some oxalate of ammonia, which will instantly precipitate the lime, if present. If it be mixed with sulphate of potass, the solution will yield a precipitate with muriate of barytes, which is insoluble in muriatic acid.

*Medical properties and uses.* — The solution of this acid in water, in the proportion of nine drachms and a half of the crystals to Oj. of water, answers nearly all the purposes of recent lemon juice; and is even preferable for forming the common effervescing draught with Carbonate of Potass. The following table shows the quantity of Citric Acid required to decompose one scruple of the alkaline salts mentioned in it.

<i>Alkaline salts.</i>	<i>Citric Acid.</i>
Carbonate of Soda.	gr. x.
Subcarbonate of Soda.	gr. xij.
Carbonate of Potass.	gr. xv.
Subcarbonate of Potass.	gr. xvij.
Subcarbonate of Ammonia	gr. xxvj.

A solution of ℥j. in Oj. of water, sweetened with sugar that has been rubbed on fresh lemon peel, forms a grateful refrigerant beverage, resembling lemonade; and which is equally useful in febrile and inflammatory complaints. It is said that

<sup>1</sup> *Phillips' Translation of Pharm.* p. 23.



the crystallized acid is not so useful in scurvy as the juice of the fresh fruit. Citric acid is incompatible in formulæ with the alkaline and earthy carbonates and acetates; the earthy and alkaline sulphurets; and tartrate of potass.

**ACIDUM MURIATICUM, Lond.<sup>1</sup>** *Muriatic Acid.*

“Take of dried Muriate of Soda, *two pounds*; Sulphuric Acid, (by weight,) *twenty ounces*; Distilled Water, *a pint and a half*. First mix the Acid with half a pint of the water, in a glass retort; and when the mixture is cold, add to it the Muriate of Soda. Pour the remainder of the water into the receiver; and having fitted it to the retort placed in a sand bath, distil over the Muriatic Acid into this water, with a heat gradually raised until the retort become red hot.

“The specific gravity of Muriatic Acid is, to that of Distilled Water, as 1·160 to 1·000.

“One hundred grains of this acid are saturated by one hundred and twenty-four grains of the crystals of Subcarbonate of Soda.”

*Edinburgh.*

“Take of Muriate of Soda, which has been previously exposed to a red heat, Sulphuric Acid, Water, of each *two pounds*. Pour the Acid, mixed with eight ounces of the Water and cooled, upon the Muriate of Soda, in a glass retort; to which adapt a receiver, containing the remainder of the Water, and distil from a sand bath with a moderate fire. In a short time the vessels may be luted together, and the distillation continued to dryness.

“The specific gravity of this acid is, to that of distilled water, as 1·170 to 1·000.”

*Dublin.*

“Take of Muriate of Soda dried, Sulphuric Acid, Water, of each *six pounds*. Dilute the Acid with the Water, and after it is cold add it gradually to the Muriate put into a glass retort: then distil the liquor until the residuum become dry.

“The specific gravity of this acid is, to that of distilled water, as 1·170 to 1·000.”

*Syn.* Acide Muriatique (*F.*), Kochsalzäure (*G.*), Zoutzuur (*Dutch*), Acido Muriatico (*I.*), Ooppoo trāvagam (*Tam.*).

The principal difference in these formulæ is in the quantity of Sulphuric Acid ordered for decomposing the Muriate.<sup>2</sup> The Sulphuric Acid is properly ordered to be diluted, to moderate the strong effervescence, and to prevent the too rapid disengagement of the muriatic acid gas, which would both endanger the bursting of the apparatus, and render the

<sup>1</sup> Spiritus Salis, P. L. 1720. Spiritus Salis Marini Glauberi, P. L. 1745.

<sup>2</sup> We use the word *muriate*, because it is the term employed by the Colleges, although *Chloride of Sodium* be that now generally and properly adopted.



process otherwise very unmanageable. The directions of the London and Edinburgh Colleges, to put part of the water into the receiver, is preferable to mixing the whole with the acid, and pouring it on the Muriate, as it facilitates very much the condensation: but Mr. Phillips justly remarks, that it would be better to put less water into the receiver and more into the retort. In the manufacturing laboratories, although the process is in principle the same as the above, yet the retort is generally of earthenware or of iron, which communicates the yellow colour that characterizes the common Muriatic acid, and which depends on a small portion of iron being raised, and brought over with the acid. Even when iron vessels are not employed the acid often assumes a yellowish colour, which depends either on a small portion of iron in the salt, or from the presence of some chlorine.

These processes were formerly explained, by saying that the decomposition of the muriate of soda is effected by the superior affinity of sulphuric acid for soda; aided by the affinity of the muriatic acid for the soda being weakened by the heat, which favours its tendency to assume the elastic form, in which state it passes over into the receiver, and is there condensed by the water. But as the doctrine of chlorine is now admitted, we must adopt the following explanation of Sir H. Davy, who regards dry common salt as a compound of 36 parts of chlorine and 24 of sodium, and consequently containing neither muriatic acid nor soda. In the processes of the Pharmacopœias, therefore, for obtaining muriatic acid, the water of the sulphuric acid is decomposed, its oxygen unites to the sodium, and forms soda, which combining with the sulphuric acid produces a sulphate of soda; while the hydrogen of the decomposed water combines with the chlorine and forms hydrochloric acid, or muriatic acid gas, a gaseous fluid consisting of equal volumes of hydrogen and chlorine, or by weight of hydrogen 2·7, chlorine 97·3, in 100 parts, which dissolving in the water contained in the receiver constitutes the liquid acid. The residue of the process is sulphate of soda with an excess of acid; to separate which, without breaking the retort, boiling water may be poured into the retort, after its contents have cooled down to 212°.

*Qualities.* — Liquid Muriatic Acid, thus obtained, is a colourless or very pale straw-coloured fluid: it has a strong pungent odour, an intensely sour, caustic taste; it reddens strongly the vegetable blues, emits white, suffocating fumes when exposed to the air, and erodes animal and vegetable substances. It unites with the alkalies, earths, and metallic oxides, forming muriates. According to Phillips, the statement of the London College as to the quantity of subcarbonate



of soda required to saturate 100 grains of the acid, is not quite correct.<sup>1</sup> The acid obtained by the process of the London College of the sp. gr. 1·160, consists of nearly 32·4 of muriatic acid, and 67·6 of water. The following part of a table, constructed by Mr. E. Davy, shows the quantity of real muriatic acid gas contained in 100 parts of fluid acid, of different densities, at the temperature of 60° :

Spec. Grav.	Real Acid Gas.	Spec. Grav.	Real Acid Gas.	Spec. Grav.	Real Acid Gas.
1·21	42·43	1·14	28·28	1·07	14·14
1·20	40·80	1·13	26·26	1·06	12·12
1·19	38·38	1·12	24·24	1·05	10·10
1·18	36·36	1·11	22·30	1·04	8·08
1·17	34·34	1·10	20·20	1·03	6·06
1·16	32·32	1·09	18·18	1·02	4·04
1·15	30·30	1·08	16·16	1·01	2·02

According to the new nomenclature, the muriatic acid of the shops is *Hydro-chloric acid*, or, to retain the common name, *Hydro-muriatic acid*. The real acid contained in the liquid acid, is a compound of equal volumes of chlorine and of hydrogen. The fluid muriatic acid found in the shops often contains sulphuric acid with small portions of iron, and sometimes copper; the first is detected by diluting the acid with 5 or 6 parts of distilled water, and adding a few drops of muriate of barytes<sup>2</sup>, which is precipitated white if sulphuric acid be present; iron is discovered by saturating the diluted acid with carbonate of soda, and adding prussiate of potass; if a blue precipitate be formed, it may be concluded that iron is present. A blue colour being produced by supersaturating the acid with ammonia detects copper.

*Medical properties and uses.* — This acid is tonic, and anti-septic. It has been efficaciously used in typhus fevers, and in some cutaneous eruptions. It is a common and useful adjunct to gargles, in the proportion of from fʒss. to fʒij. in fʒvj. of any fluid, in ulcerated sore throats, and scarlatina maligna; and, in a very highly diluted state, viz. mviij. in fʒiv. of water, it has been recommended as an injection in gonorrhœa.

This acid has even been regarded as an antidote in general syphilitic affections; but the observations of Mr. Pearson have showed this opinion to be erroneous; yet, by its salutary

<sup>1</sup> Philip's *Trans. Pharm.* p. 27.

<sup>2</sup> Mr. Hume discovered that muriatic acid precipitates muriate of barytes when no sulphuric acid is present; but this does not happen when the acid is much diluted.



effects on the stomach and general health, “it is a medicine capable of ameliorating the appearance of venereal ulcers, and of restraining for a time the progress of the disease,” where it is desirable “to gain a little time, previously to the entering on a mercurial course.”<sup>1</sup> The dose is from  $\text{m x.}$  to  $\text{m xx.}$  in a sufficient quantity of water, or in any bland fluid. It is incompatible in prescriptions with alkalies, most of the earths, oxides and their carbonates; sulphuret and tartrate of potass, tartarized antimony and iron, nitrate of silver and acetate of lead. In typhus, and fevers of a typhoid type, I have generally given it in the infusion of cinchona or cusparia bark. Dr. Paris states that he has found it a preventive of the generation of worms, when given after copious evacuations of the bowels.<sup>2</sup> Largely diluted in any mucilaginous fluid, and sweetened, it is a useful remedy in calculous cases depending on an excess of the phosphates.

When Muriatic Acid is taken as a poison, it may be detected by its sensible qualities; but if mixed with wine or other fluids, let a portion of it be distilled from a small retort over a candle, into a phial containing a solution of nitrate of silver. The precipitation of muriate of silver, which is soluble in ammonia, but not in nitric acid, will take place if the poison contain muriatic acid. The best antidotes, if exhibited in time, are soap and calcined magnesia suspended in water.

A very important property of muriatic acid, in the state of gas, is the power it possesses of neutralizing putrid miasmata, discovered by Morveau in 1773. It is therefore used as an agent for destroying infection in sick-rooms and hospitals, disengaged by pouring sulphuric acid on common salt.

Officinal preparations. *Murias Barytæ*, E. *Solutio Muriatis Calcis*, E. D. *Tinctura Ferri Muriatis*, L. E. D.

**ACIDUM MURIATICUM DILUTUM**, Dub. *Diluted Muriatic Acid*.

“Take of Muriatic Acid, and of Distilled Water, each one pound by weight. Mix them. The specific gravity of this acid is, to that of water, as 1·080 to 1·000.”

This formula is intended to render the dose of Muriatic Acid more easily apportioned: 100 parts contain about 14 of real acid.

**AQUA OXYMURIATICA**, Dub. *Oxymuriatic Water*.

“This is prepared by transmitting the superfluous gas of the process for making the solution of Oxymuriate of Soda, *Aqua Alcalina Oxymuriatica*, by means of a proper apparatus, through a pint of distilled water.


<sup>1</sup> *Pearson on Remedies for Lues Venerea*, 194.

<sup>2</sup> *Pharmacologia*.



“The specific gravity of this solution is, to that of water, as 1·003 to 1·000.”

*Syn.* Acide muriatique oxygéné (*F.*), Vollkomme Salzsäure (*G.*), Oerzuures Zoutzuur (*Dutch*), Acido muriatico ossigenato (*I.*).<sup>1</sup>



In the process by which this solution is prepared, Chlorine comes over in the gaseous form, and is condensed in the distilled water, which is placed in a Woolfe's bottle, or a globular vessel *b*, furnished with a safety tube *e*, and connected by a tube with a receiver *a*, that contains a solution of subcarbonate of potass. The gas first passes through the alkaline solution, part of it condenses and combines with the potass, forming a chlorate of potass, while the superfluous uncondensed portion passes on to the other vessels *c*, *d*, and there combines with the water, forming this solution, which is erroneously termed oxymuriatic acid.

The substance contained in this solution was discovered by Scheele in 1774, while making his experiments on manganese; and termed by him *Dephlogisticated Muriatic Acid*: but its nature was not perfectly understood until it was investigated by Sir H. Davy, who determined it to be an elementary substance, and named it *Chlorine*, from its colour, which is yellowish.

*Qualities.* — The saturated solution of Chlorine, or Oxymuriatic acid as it is still termed by the Dublin College, has a very offensive suffocating odour; and a harsh, styptic, but not acid taste. Its colour is a very pale-yellowish green; it destroys all the vegetable colours, rendering them white. It must be kept in opaque bottles or in a dark place; for by the action of the solar rays, part of the water is decomposed; the hydrogen of which, uniting with the Chlorine, forms muriatic acid, which remains in solution in the water, while the oxygen is set free. At a temperature of 50° this solution contains about twice its volume of Chlorine. The aqueous solution of *Chlorine* acts on almost all the metals, forming Murates or Chlorides.

<sup>1</sup> All these terms are now generally admitted to be erroneous, there being really no such substance as they imply: the composition alluded to is the solution of a simple elementary body, *Chlorine*, in water.



*Medical properties and uses.* — Fluid Oxymuriatic-acid (*Aqueous solution of Chlorine*) is stimulant and antiseptic. It has been strongly recommended in scarlatina and malignant sore throat; and as an antisyphilitic remedy. In the latter disease the same opinion may be given of it as of the simple muriatic acid; but in scarlatina and cynanche maligna more benefit has resulted from its use. From fʒss to fʒij, mixed in fʒviij of water, and sweetened with a little syrup, may be taken in the course of the day, in divided doses.

But the most important use of Chlorine is in its gaseous form, as a fumigation for neutralizing putrid miasmata, and correcting the infectious atmosphere of hospital wards and rooms in which have been cases of contagious fevers. For these purposes it is better adapted than the common muriatic acid gas; but as both of them are highly deleterious to animal life, they should be employed in such apartments only as the sick can be removed from while the gas is extricated. The Chlorine is easily procured by pouring fʒvj of strong sulphuric acid on a mixture of ʒiv of pulverized manganese, and ʒviij of dried common salt, in a china cup, which should be placed in a pipkin of hot sand. The doors of the room which is to be fumigated, must be kept shut for two hours after the cup or cups with this charge are placed in it; then be thrown open, and a free current of air permitted to pass through the apartment. By this process the offensive odour of the sick room is destroyed, the chemical constitution of the deleterious atmosphere altered, and its freshness completely restored.

For the more convenient application of this powerful agent, Morveau has invented what he terms disinfecting or preservative bottles. The apparatus consists of a strong glass bottle or phial, covered with a plate of glass, which is fitted by grinding so as to shut, accurately, the orifice of the vessel. The bottle is fixed in a wooden frame; and the plate of glass kept in its place, and closely applied by means of a screw. If the bottle be of 25 cubic inches of capacity, the charge to be put into it may consist of 372 grs. of black oxide of manganese in coarse powder, 3·5 cubic inches of nitric acid of 1·4 specific gravity, and an equal bulk of muriatic acid of 1·134 specific gravity. As soon as the charge is introduced, the glass plate must be firmly screwed down in its place. When the apparatus is to be used, the screw is to be turned so as to allow the gas which is extricated to escape from under the plate of glass; and this must be again screwed down, as soon as the smell of the Chlorine is perceptible in the distant corners of the apartment. Bottles of any dimensions may be used, but the charge must in no case occupy more than one-third part of the capacity of the vessel.



ACIDUM NITRICUM, Lond.<sup>1</sup> *Nitric Acid.*

“Take of Nitrate of Potass dried, and Sulphuric Acid, each by weight *two pounds*; mix them in a glass retort: and distil the Nitric acid from a sand-bath, until red vapours are produced. Then, having added an ounce of dry Nitrate of Potass, re-distil the acid in a similar manner.

“The specific gravity of this acid is, to that of distilled water, as 1·500 to 1·000. One hundred grains of this acid are saturated by two hundred and twelve grains of crystals of Subcarbonate of Soda.”

*Edinburgh.*

“Take of Nitrous Acid, *any quantity*. Put it into a retort, and having fitted a receiver, which must be kept cold, apply a very gentle heat until the reddest part shall have passed over, and the acid which remains in the retort, already almost free from colour, have become Nitric acid.”

ACIDUM NITROSUM, Edin. *Nitrous Acid.*

“Take of Nitrate of Potass bruised, *two pounds*; Sulphuric Acid *sixteen ounces*. Pour the acid upon the Nitrate of Potass in a glass retort, and distil from a sand-bath with a gradually augmented heat, until the iron pot becomes obscurely red hot.

“The specific gravity of this acid is, to that of distilled water, as 1·520 to 1·000.”

*Dublin.*

“Take of Nitrate of Kali, *six pounds*; Sulphuric Acid, *four pounds* by weight. Mix and distil until the residue becomes dry.

“The specific gravity of this acid is, to that of distilled water, as 1·500 to 1·000.”<sup>2</sup>

*Syn.* Acide nitrique (*F.*), Salpeter säure (*G.*), Zalpeterzuur Skerkwater (*Dutch*), Skedwatter (*Swed.*), Acido Nitrico (*I.*), Pottle Ooppoo trāvāgum (*Tam.*), Areki Shora (*Pers.*), Maulabker (*Arab.*).

In performing these processes it is advisable to use a Woolfe's apparatus, or a range of two or three globular receivers, the last of which should contain a small portion of water. The Nitric Acid is separated from its combination with potass in the nitrate by the superior affinity of the sulphuric acid for the potass, which, however, requires to be aided by quantity, a larger portion of sulphuric acid than is

<sup>1</sup> Acidum Nitrosum, P. L. 1787. Spiritus Nitri Glauberi. Aquafortis, P. L. 1745. Aquafortis simplex et duplex, P. L. 1720.

<sup>2</sup> For the preparation of this acid on a large scale in this country, rough nitre, with half the weight of sulphuric acid, is employed. These are put into a large glass or earthenware *body*, to which a glass pipe is luted communicating with an empty receiver, which is connected by means of pipes also, with several other receivers half filled with water.



necessary for saturating the potass of the nitrate being used : and also by heat, which volatilizes the Nitric Acid as it is disengaged. As soon as the materials are heated, orange-yellow vapours are disengaged, which in a short time, as the heat increases, become paler, and continue so until the ingredients in the retort are nearly dry, and the heat is augmented to  $500^{\circ}$ ; when, owing to a partial decomposition of the acid next disengaged, nitrous gas comes over in deep red fumes, with a quantity of permanently elastic pure oxygen, which may be collected in an inverted receiver filled with water, placed in a pneumatic trough, and connected with the last of the receivers by means of a bent tube. The nitrous gas combines with the condensed acid in the receiver, deepens its colour, and gives it that form which constitutes nitrous acid. It is with the view of preventing this, that the London College has ordered so large a portion of sulphuric acid to be employed, the principal use of which appears to be to contribute a sufficient portion of water to preserve the constitution of the Nitric Acid. The Edinburgh College orders the acid to be kept in its coloured state; and as a medical agent it answers the same purposes as the colourless acid; for, when both are diluted with water, they have the same appearance, and are brought to the same state, the addition of the water expelling completely the nitrous gas, which is only loosely united with the Nitric Acid to form the nitrous. The quantity of acid obtained by the Edinburgh process is about half the weight of the nitrate employed; and the residue is a white spongy saline cake of sulphate of potass with an excess of sulphuric acid, or a bisulphate of potass, which may be dissolved out of the retort by hot water.

By the London process the Nitric Acid is at first obtained tolerably free from nitrous gas; but in general the re-distillation, as directed, will be found necessary. In the expulsion of the nitrous gas, to change the nitrous acid into Nitric Acid, according to the directions of the Edinburgh College, a portion of the acid is carried over with the gas, as nitrous acid vapour; which should not be wasted, but be condensed by a small portion of water being put into the receiver, and thus form a diluted acid. Mr. Murray<sup>1</sup> justly observes, that the heat of a water-bath is best adapted for this operation, being sufficient for the purpose, and not too great to produce the decomposition of the acid. A completely colourless acid, however, is not obtained, unless the acid be re-distilled from

---

<sup>1</sup> *System of Materia Medica*, ii. 184.



a small portion of black oxide of manganese; but this is not at all necessary for medical purposes.<sup>1</sup>

As nitre sometimes contains a small portion of muriate of soda, Nitric Acid, in whatever method it has been procured, may be contaminated with a minute portion of muriatic acid, or of sulphuric acid, if a large proportion of this have been used for decomposing the nitre: the presence of the first is detected by dropping in nitrate of silver, which forms an insoluble muriate of silver; while the formation of a precipitate on the addition of nitrate of barytes discovers the second; but both the acid to be tested and the nitrate of barytes should be previously diluted with distilled water: if any undecomposed nitre be present, it will be detected by evaporating a portion of the acid, when, if it contain no nitre, it will have no residue. These contaminations do not affect the medicinal virtues of the acid.

*Qualities.* — *Nitrous Acid*, as the term is understood in the Edinburgh Pharmacopœia, is a yellow or orange-coloured fluid, emitting, when exposed to the air, deep orange-coloured, extremely suffocating fumes. In its chemical affinities and other qualities it agrees in many respects with Nitric Acid. It consists of nitrous gas loosely combined with Nitric Acid and water; and the colour varies according to the proportion of nitrous gas which is present. From experiments made by Sir H. Davy<sup>2</sup> on this subject, the following appear to be the proportions in the three states in which nitrous acid is usually procured for pharmaceutical purposes.

100 Parts of Acid.	Spec. Gravity.	Real Nitric Acid.	Water.	Nitrous Gas.
Pale yellow	1·502	90·5	8·3	2·00
Bright yellow	1·500	88·94	8·10	2·96
Dark orange	1·480	86·84	7·6	5·56

The Edinburgh College states the specific gravity too high, for it seldom exceeds 1·502, and scarcely ever 1·52.<sup>3</sup> When one part by weight of water is added to four parts of yellow nitrous acid, the colour is altered to a fine green; when equal parts of both are mixed, it becomes blue; and by another addition of water, or by allowing it to stand exposed to the air, it changes to a very pale straw-colour, or becomes nearly colourless.

<sup>1</sup> Nitric acid was first obtained by Raymond Lully, in the 13th century, by distilling a mixture of nitre and clay: a process still employed on the continent. The name *Nitric acid* was imposed in 1787, by the French chemists.

<sup>2</sup> *Researches*, 37.

<sup>3</sup> Rouelle states the specific gravity of the strongest nitric acid that can be procured to be 1·583; Kirwan makes it 1·5543 only, at 60° Fahrenheit.



*Liquid Nitric Acid* is a colourless, or very pale-yellow, limpid fluid, emitting, when exposed to the air, white suffocating vapours, and possessing strong acid properties. It is highly corrosive, and tinges the skin yellow, the tint remaining till the epidermis peel off. When exposed to the air, it attracts moisture and becomes weaker. It unites with water in every proportion, and while mixing, heat is evolved. One fluid ounce of the acid, prepared according to the formula of the London College, should weigh 11 drachms 1 scruple. The following table, constructed by Sir H. Davy<sup>1</sup>, shows the quantity of real acid and water contained in 100 parts of fluid acid of different densities.

100 Parts Nitric Acid of specific gravity.	Contain of		100 Parts Nitric Acid of specific gravity.	Contain of	
	True Acid.	Water.		True Acid.	Water.
1.5040	91.55	8.45	1.3186	52.03	47.97
1.4475	80.39	19.61	1.3042	49.04	50.96
1.4285	71.95	28.35	1.2831	46.03	53.97
1.3906	62.96	37.04	1.2090	45.27	54.73
1.3551	56.80	43.12			

Nitric Acid is volatilized by heat, and decomposed by light. It is also decomposed by all the simple combustibles, with great violence of action, which is accompanied with the evolution of copious red fumes. When poured on oils, particularly if these contain any essential oil, it sets them on fire; is capable of oxidizing all the metals; and combines with the earths, alkalies, and metallic oxides<sup>2</sup>, forming nitrates; one fluid ounce of specific gravity 1.500 should dissolve 476 grains of white marble. The constituents of Nitric Acid, independent of the water, which gives it the fluid form, are 25.93 azote, and 74.07 oxygen in 100 parts: or 1 volume of azotic gas and 2½ volumes of oxygen gas. The strongest fluid acid contains 25 per cent. of water, and 75 of dry acid.

*Use.* — Strong fluid Nitric Acid is seldom employed except for pharmaceutical purposes; but it has been lately recommended as an escharotic in sloughing phagedœnic ulcers by Mr. Welbank. To employ it properly, the surface of the ulcer must be well cleaned and dried, and, after applying a thick coating of lard to protect the surrounding sound skin,

<sup>1</sup> *Researches*, p. 41.

<sup>2</sup> In its concentrated state, it does not act upon the metals at ordinary temperatures; but when slightly diluted it quickly oxidizes them and dissolves the oxides. This is owing to the decomposition of the water, which yields up its oxygen to the metal, which in a state of oxide is readily soluble in the acid. It does not act on platina or gold.



a pledgit, moistened with the undiluted acid, must be pressed steadily on every point of the diseased surface, which, by this treatment, soon throws off a slough: and assumes a healthy aspect.<sup>1</sup> When extricated in the form of vapour, it is employed for destroying contagion. It is less powerful than the oxymuriatic acid, but is more generally useful, as it can be extricated in the chambers of the sick without proving deleterious to animal life.<sup>2</sup> For this purpose fʒij. of sulphuric acid may be poured over ʒiv. of coarsely powdered nitre in a china cup, and placed in a pipkin of hot sand. This quantity is sufficient for fumigating a room of ten feet square; and, where a larger portion is required, it is more advisable to multiply the number of pipkins, than to put a larger quantity of the materials into one vessel.

Officinal preparations. *Acidum Nitricum Dilutum*, L. E. D. *Argenti Nitras*, L. E. D. *Ung. Hydrargyri Nitratis*, L. E. D. *Hydrargyri Nitrico-oxydum*, L. *Spiritus Ætheris nitrici*, L. E. D. *Unguentum Acidi nitrosi*, E. D.

ACIDUM NITRICUM DILUTUM, Lond. *Diluted Nitric Acid.*

“Take of Nitric Acid, a fluid ounce; Distilled Water, nine fluid ounces. Mix.”

ACIDUM NITROSUM DILUTUM, Edin. *Diluted Nitrous Acid.*

“Take of Nitrous Acid, and of Water, equal weights. Mix, avoiding the noxious vapours.”

*Dublin.*

“Take of Nitrous Acid, and of Distilled Water, each one pound. The specific gravity of this mixture is, to that of distilled water, as 1280 to 1000.”

These processes are intended for the more convenient apportionment of the dose in the exhibition of this acid. In the former edition of the London Pharmacopœia the proportions of acid and water were equal by weight; but the alteration in the present edition makes a very important difference of strength, in a given measure of the diluted acid, prepared after the first, and the last of the above formulæ.

When prepared according to the directions of the London College, fʒi. contains about grs. 68 of Nitric Acid, of 1.500 specific gravity, and will saturate 144 grains of crystallized Subcarbonate of Soda; while the same measure of the same acid, prepared after the Edinburgh and Dublin, and the former London formulæ, contains grs. 390.5 of the same acid; a difference which may lead to errors in practice; and is

<sup>1</sup> *Medico Chirurg. Trans.* vol. xi. p. 69.

<sup>2</sup> *The effects of Nitrous Vapour, &c.* by J. C. Smyth, M. D.



therefore to be regretted, particularly as no reason is assigned for the change.

*Medical properties and uses.* — Nitric acid is tonic and antiseptic. When very largely diluted with water, it forms an agreeable and very useful beverage in fevers, particularly of the typhoid type. In larger doses, less diluted, it has been efficaciously administered in chronic hepatitis, even when dropsy has supervened; and has also been found serviceable in restraining violent sickness, in dyspepsia, asthma, and the majority of the cachexiæ. From some observations of Dr. Scott, published at Bombay, in 1796, this acid excited considerable attention as a remedy for syphilis; but after the most ample trials, by almost every practitioner of any eminence in the country, its antisiphilitic powers have not been found by any means to answer the accounts of them transmitted from India. The subsequent publications of Dr. Scott, however, have shown, that he did not employ Nitric Acid, but a mixture of three parts of muriatic acid, and two of nitric. It checks for a time the progress of the disease, but does not permanently remove the symptoms; and, as Mr. Pearson justly observes, “it would by no means be warrantable to substitute the Nitrous (or Nitric) Acid in the place of mercury, for the cure of venereal<sup>1</sup> complaints.” It is, however, in many cases of much benefit during a mercurial course, or prior to its commencement, when the constitution is impaired, and inadequate to support the effects of mercury; as by its tonic powers it promotes the general health, and lessens the action of the mercurial remedy on the mouth and fauces; yet, when it is pushed far, it affects the mouth, and produces ptyalism.<sup>2</sup> When dropsy supervenes on reiterated courses of mercury, which is not unfrequent in broken-down constitutions, this acid, Mr. Carmichael observes, given in as large doses as the stomach will permit, conjoined with digitalis, is productive of the utmost benefit. We have found it of considerable service, given at the same time with mercury, in old obstinate ulcerations of the legs, although no venereal taint could be suspected: and it is employed with benefit as a local stimulant, in the form of lotion, in the proportion of fʒij. of the diluted acid, to Oj of water, to fœtid ulcers, attended with a thin ichorous discharge, and in caries of the bones. In India, and in this country for some years past, Nitric Acid has been used combined with muriatic acid, in the form of a bath<sup>3</sup>, and

---

<sup>1</sup> *Pearson on Remedies for Lues Venerea*, 188.

<sup>2</sup> Blood drawn from the arm, after the daily use of this acid for some weeks, exhibits the buffy coat.

<sup>3</sup> The proportions are two parts of muriatic acid and one of nitric, so diluted with water as to taste as sour as vinegar. Experience has not justified the high encomiums bestowed upon them.



in this state produces a slight excitement of the skin, a peculiar taste in the mouth, and, in other respects, nearly the same effects as when it is taken internally; but the chief perceptible effect of the mixed acid is on the bowels, which it keeps moderately open. Diluted Nitric Acid has often been employed as a poison. It is detected by the orange-coloured spots, which are observed on the lips, chin, and hands of the patients; and, if death be the result, by the same colour being found in a large portion of the alimentary canal, the mucous membrane of which is converted into a fatty substance, and the stomach often perforated. If any of the fluid can be obtained, the extrication of orange-coloured fumes on boiling it over copper-filings, is a certain test of aquafortis. Soap, and calcined magnesia suspended in water, are the best antidotes.

The dose of the diluted acid is from  $\mathfrak{m}\text{x.}$  to  $\mathfrak{m}\text{xxx.}$  in  $\mathfrak{f}\mathfrak{z}\mathfrak{iij.}$  of water, given three or four times a day. When used as a bath, the mixed acid should be added to the water, until it is about as sour as weak vinegar. Nitric acid, even when diluted, is incompatible in prescriptions with oxides, earths, alkalies, the sulphurets, and the acetates of potass and of lead. It decomposes the two last named salts, and forms nitrates of lead and of potass.

**ACIDUM SUCCINICUM, Edin.** *Succinic Acid.*

“Take of Amber in powder, and pure Sand, *equal parts*; mix and put them into a glass retort, of which they may fill one half. Having adapted a large receiver, distil from a sand-bath, with a gradually raised fire. A watery liquor with a little yellow oil will first distil over: then a yellow oil with an acid salt: and lastly, a reddish and black oil. Pour the liquor out of the receiver, and let the oil be separated from the water. Press the acid salt, collected in the neck of the retort and on the sides of the receiver, between folds of bibulous paper, that it may be freed from the adhering oil; then purify it by solution, in hot water and crystallization.”

**ACIDUM SUCCINICUM, Dub.** *Succinic Acid.*

“Take of Amber, and pure Sand, each *a pound*. Distil, with a gradually increased heat, an acid liquor, an oil, and a salt discoloured with oil. Wrap up this salt in bibulous paper, and subject it to the press to separate the oil; then let it be again sublimed.”

*Syn.* Acide Succinique (*F.*), Bernsteinsäure (*G.*), Acido Succinico (*I.*).

The use of the Sand in these processes is to prevent the Amber, which swells very much, from passing over into the receiver. The heat which is necessary for the complete decomposition of the amber is very considerable; and therefore by following exactly the formulæ of the Colleges, this is



scarcely ever accomplished. The Succinic Acid is partly dissolved in the water which condenses in the receiver, but the greater part is sublimed in the neck of the retort, and is so much contaminated with the oil, that after repeated solution and crystallization, and even resublimation, it still retains a portion of it. According to Guyton Morveau<sup>1</sup>, it may be obtained perfectly pure by distilling from it a small portion of Nitric Acid, with a heat not strong enough to sublime the Succinic Acid.

*Qualities.* — The crystals of Succinic acid are minute triangular prisms. When pure, they are white, translucent, and shining; have a slight penetrating sour taste; redden infusion of litmus; and are volatile and inflammable, burning away without leaving any odour. They are soluble in twenty-four parts of water at 60°, and two parts at 212°; the greater part, however, crystallizing as the water cools. They are also soluble in alcohol, and in sulphuric and nitric acid, without suffering decomposition. With the alkalies, earth, and metallic oxides, succinic acid combines and forms succinates. It is a triple compound of 47·600 parts of carbon, 4·512 of hydrogen, and 47·888 of oxygen.<sup>2</sup>

This acid is often adulterated with tartaric acid, muriate of ammonia, and sulphate of potass. The first is detected by carbonate of potass; the second, by nitrate of silver; and the sulphate by barytic water. It is altogether discarded from practice.

ACIDUM SULPHURICUM DILUTUM, Lond. *Diluted Sulphuric Acid.*

“Take of Sulphuric acid, *a fluid ounce and a half*; Distilled Water, *fourteen fluid ounces and a half*. Add the acid gradually to the water, and mix.”

*Edinburgh.*

“Take of Sulphuric acid, *one part*; Water, *seven parts*; mix them.”

*Dublin.*

“Take of Sulphuric acid, *two ounces by weight*; Distilled Water, *fourteen ounces by weight*. Mix them gradually, and set the mixture aside to cool; then pour off the clear liquor. The specific gravity of this acid is, to that of water, as 1090 to 1000.”

*Syn.* Acidum Vitriolicum dilutum, P. L. 1787. Acide sulphurique étendu d'eau (*F.*), Verdüunte Schwefelsäure (*G.*), Verduintes Zwavelzuur (*Dutch*), Acido solforico diluito (*I.*), Zakab (*Arab.*).

It is very much to be regretted, that the London College, when it altered the proportions of acid and water in this mix-

<sup>1</sup> *Annales de Chimie*, xxix. 165.

<sup>2</sup> *Annals of Philosophy*, vol. v. p. 99.



ture, from those in the last edition of its Pharmacopœia, did not adopt the proportions ordered by the two other Colleges, so that, in this preparation at least, a standard strength might have been fixed for the whole kingdom. The reasons which induced it to adopt the present proportions are not easy to be conceived; for the puerile reason stated by Dr. Powell that “this mixture will be more conveniently made, and its dose more easily apportioned than the former one,” cannot surely have operated in causing the alteration. One fluid ounce of this diluted acid of the London Pharmacopœia, contains 80 grains of the strong acid; in the Edinburgh and Dublin pharmacopœias it constitutes an eighth part.

Owing to the great affinity of Sulphuric Acid for water, and the density of the mixture being much greater than the mean of the separated acid and water<sup>1</sup>, a very considerable increase of temperature is produced during their combination, sufficient to crack the glass vessels in which it is made, if the two ingredients be at once mixed together.<sup>2</sup> To prevent such an accident, the acid must be gradually added in small portions to the whole of the water, and the mixture agitated after every addition. It is of importance always to ascertain the specific gravity of the acid before the mixture be made. The mixture, when it has cooled down to the temperature of the atmosphere, lets fall a white precipitate, consisting of a small portion of sulphate of potass, and of sulphate of lead, which the strong acid always contains, but which the diluted acid is incapable of holding in solution. The diluted acid is thus purer than the strong acid, which suffers no other alteration except in point of strength: and hence the Dublin College properly directs the clear liquor to be poured off when the mixture has cooled.

*Medical properties and uses.* — Diluted Sulphuric Acid is tonic, antiseptic, and refrigerant. Its tonic and antiseptic powers render it extremely serviceable in low typhoid fevers, dyspeptic affections, diabetes, convalescencies, and in cutaneous eruptions. It restrains the colliquative sweats which attend hectic: locally applied, it is a common and useful adjunct to gargles in cynanche, and to check salivation; and as a refrigerant, it is given with certain benefit in passive hæmorrhages, from whatever part they may arise. Combined with mucilages, it has been beneficially given in passive diarrhœas,

---

<sup>1</sup> It is a curious fact that, after the mixture has cooled down to the temperature of the atmosphere, a considerable time elapses before it acquires its real density.

<sup>2</sup> If one part by weight of sulphuric acid of 1·845 specific gravity be mixed with one-fourth its weight of water, both being at the temperature of 50° Fah., the caloric instantly evolved is sufficient to raise the thermometer from 50° to 300°, and a still greater heat is produced by mixing 73 parts of the acid with 27 of water.



operating on the relaxed mucous coat of the intestine as an astringent. In the first-mentioned cases, the diluted acid may be combined with infusions of cinchona or other vegetable bitters, and aromatics; and in the latter with infusion of roses, mucilage, or simple water sweetened with syrup. It is certainly injurious to the teeth; and, therefore, should be sucked through a quill, when taken as an internal remedy. The usual dose is from  $\mathfrak{m}\text{x.}$  to  $\mathfrak{m}\text{xxx.}$ , but in malignant erysipelas, with a tendency to hæmorrhage, it has been given to the amount of  $\mathfrak{f}\mathfrak{z}\mathfrak{j.}$  in twenty-four hours; and we have given it, with evident advantage, to the same amount, in violent uterine hæmorrhages.

Officinal preparations. *Acidum sulphuricum aromaticum*, E. *Infusum Rosæ compositum*, L.

**ACIDUM SULPHURICUM AROMATICUM**, Edin.  
*Aromatic Sulphuric Acid.*

“Take of Alcohol, *two pounds*; Sulphuric Acid, *six ounces*. Drop the acid gradually into the alcohol. Digest the mixture in a covered vessel with a very gentle heat for three days; then add of Cinnamon-bark, bruised, *one ounce and a half*; Ginger-root, bruised, *one ounce*. Digest again in a closed vessel for six days; then filter through paper placed in a glass funnel.”

This preparation is generally regarded as an imperfect ether; but we are of opinion that the reciprocal action of the acid and alcohol during the digestion is scarcely sufficient to produce such a result; and the acid, undoubtedly, very much predominates. It is, therefore, a simple alcoholic solution of sulphuric acid, holding the essential oils of cinnamon and of ginger in solution.

*Qualities.*—The odour is peculiar and aromatic; the taste gratefully acid. It is limpid, and of a brownish colour.

*Medical properties and uses.*—This is an agreeable mode of exhibiting Sulphuric Acid in dyspepsia, chronic asthma, and most of the complaints for which the diluted acid has been found serviceable. The dose is from  $\mathfrak{m}\text{x.}$  to  $\mathfrak{m}\text{xxx.}$  in any convenient fluid vehicle; and may be given three or four times a day.

**ACIDUM TARTARICUM**, Lond. *Tartaric Acid.*

“Take of Supertartrate of Potass, *two pounds and a half*; boiling Distilled Water, *three gallons*; prepared Chalk, *one pound*; Sulphuric Acid *one pound*. Boil the Supertartrate of Potass with two gallons of the Distilled Water, and gradually add the prepared Chalk, until no more bubbles are produced: set the mixture apart that the Tartrate of Lime may subside; pour off the fluid, and wash repeatedly the Tartrate of Lime with distilled water, until it come off tasteless. Then pour upon the Tartrate the Sulphuric Acid diluted with a gallon



of the boiling distilled water, and set the whole apart for twenty-four hours, occasionally stirring it. Filter the liquor, and evaporate it in a water-bath to obtain the crystals.”

In this process, which is nearly the same as that of Scheele, the lime of the chalk separates the tartaric acid from the potass with which it was previously combined<sup>1</sup>; and again yields it up, in order to combine with the sulphuric acid, which is used in the second part of the process. It is quite unnecessary to boil the distilled water for diluting the sulphuric acid, as the degree of heat produced by the mixing the acid and water, even were both at the freezing point, is sufficient for every purpose of the process, and is not increased by using the distilled water in a boiling state. Before crystallizing, it is proper to test the liquor by dropping a little acetate of lead into a small portion of it, which throws down a precipitate insoluble in acetic acid, if any sulphuric acid be present. In which case a little more tartrate of lime should be added. The crystals are obtained in groups, and cannot be always procured of the same form under the most careful management: but their most frequent form is an oblique rhombic prism.

*Qualities.*—Tartaric acid in its crystallized state is white, imperfectly transparent, persistent in the air, inodorous, and very acid to the taste. The primary form of its crystal is an oblique rhombic prism.<sup>2</sup> It melts when heated a little above 212°, and boils at 250°, without losing its whiteness, and, unless the boiling be long continued, it loses little more than 4 *per cent.* of its weight; but the nature of the acid is changed; for on cooling, the semi-transparent mass into which it concretes is deliquescent. Tartaric acid is readily soluble in water, which dissolves one fifth of its weight, and the saturated solution is not liable to spontaneous decomposition. It is decomposed by a high temperature, and also by sulphuric and nitric acids, the latter of which converts it into oxalic acid. It combines with alkalies, earths, and metallic oxides, forming *tartrates*; and in its power of saturating alkalies closely resembles citric acid. According to the analysis of Berzelius, 100 parts of this acid, as it exists in tartrate of potass, are composed of 3.951 of hydrogen, 36.167 of carbon, and 59.882 of oxygen: but, in its ordinary crystallized state, or as a hydrate, it is composed of 88.16 parts of real acid, and 11.84 of water, in 100 parts. When carelessly made, it may contain sul-

---

<sup>1</sup> The components of supertartrate of potass are, tartaric acid 70.15, potass 25.13, and water 4.72, in 100 parts.

<sup>2</sup> *Phillips's Trans. of Pharm.* 1824.



phuric acid, which, however, can be detected by adding muriate of barytes to the solution, when a precipitate insoluble in an excess of muriatic acid will be thrown down, if sulphuric acid be present. One hundred grains of the crystallized acid saturate 200 grains of crystallized subcarbonate of soda.

*Medical properties and uses.* — Tartaric acid, largely diluted and sweetened, forms a cooling agreeable beverage in fevers and diseases connected with an increased secretion of bile. It is incompatible in prescriptions with solutions of salts containing potass, which it converts into a bitartrate, which being of difficult solution is thrown down in minute crystals. It also precipitates salts of lime and of lead. Dissolved in water, and added to an aqueous solution of carbonate of Soda, it forms a good substitute for soda-water.

## APPENDIX TO THE ACIDS.

IN the following Appendix I propose to give a short account of two acids, which, although they have not a place in any of the British Pharmacopœias, yet, are of great importance; the one being a remedy of considerable efficacy in a certain class of diseases, and the other demanding the attention of the medical practitioner, from the frequent employment of it as a poison.

### I. — ACIDUM HYDROCYANICUM.

#### *Hydrocyanic (Prussic) Acid.*

This acid is found in many vegetable productions; such as the bark of the *Prunus Padus*, or Bird-cherry; the leaves of the Peach and Nectarine trees, Bitter-almonds, and the kernels of many fruits; but, for the purposes of medicine it is artificially prepared.

*Preparation.* The best method of preparing Hydrocyanic acid, for medicinal use, is the following, which was first employed by Scheele.

Mix two ounces of Prussian-blue with six ounces of red precipitate of mercury, and add six ounces of water. Boil the mixture for some minutes, constantly agitating it, when the blue colour will disappear, and the mass assume a yellowish-grey hue. Pour the whole on a filter, and wash the residuum with a little hot water, which is to be added to the filtered liquor. Pour this upon an ounce and a half of clean iron filings, and add three drachms of strong sulphuric acid.



Shake this mixture well, and, after the powder subsides, pour the fluid into a retort, and distil one-fourth part of it over into a well luted receiver. This is the Hydro-cyanic acid, containing an admixture of a little Sulphuric acid, which is readily separated by means of Barytic water. La Planche recommends  $\frac{1}{6}$  only to be distilled over, and this to be rectified, by means of a gentle fire, over  $\frac{1}{200}$  of carbonate of lime; drawing off, afterwards,  $\frac{3}{4}$  only of the  $\frac{1}{6}$  of the whole, thus treated, by a second distillation. The acid is obtained of a uniform strength by this method.

In the above processes, the Iron filings and the Sulphuric acid added to the solution obtained from boiling the mixture of Prussian-blue and Red precipitate of Mercury in water, decompose the water; and the reduced Mercury combines with the Cyanogen, the base of the acid of the Prussian-blue, and forms a Cyanuret of Mercury. This new combination is again destroyed by the heat, and the Cyanogen acting upon the nascent Hydrogen of the decomposing water, forms Hydro-cyanic vapours, which are absorbed by the water in the receiver, and constitute the Hydro-cyanic Acid.

*Physical and chemical properties.* — Hydro-cyanic Acid, prepared in the above described manner, is a colourless, transparent liquid, with a peculiar odour, not unlike that of bitter almonds. It is at first bland and sweet to the taste, but ultimately impresses a pungent acrimony on the palate. It is very volatile; and, owing to this property, crystallizes if a drop of it fall upon paper. Its specific gravity is 0.996.

It is decomposed by a high temperature and by light; being resolved into Carbonic acid, Ammonia and Carburetted Hydrogen gas, which are dissipated, and leave behind a carbonaceous deposit. It should therefore be kept in an opaque stoppered bottle. It is very inflammable, burning with a blue flame: and is soluble both in water and in alcohol. According to Gay Lussac, it consists of a peculiar base, which he has named Cyanogen, acidified by Hydrogen. Cyanogen is a compound, consisting of two proportionals of Carbon and one of Nitrogen. But the acid employed in medicine contains one part only of the acid referred to by Gay Lussac, and eight parts and a half of water.

*Medical properties and uses.* — Hydro-cyanic Acid, when taken into the stomach, in a large dose, acts as an instantaneous and most powerful sedative, destroying completely the nervous energy and the irritability of the body, and consequently extinguishing life: but, in an animal thus killed, the action of the heart continues for some time after the animal has apparently ceased to live. The observation of this curious



fact led Professor Brera, in 1809<sup>1</sup>, to administer Prussic acid as a remedy in pulmonary inflammation; and he found that it quickly subdued the violence of the disease, “without having any recourse to more than preliminary bleeding.” British practitioners, however, were altogether unacquainted with this remedy, until after Dr. Majendie published his first essay on this subject, in 1815; when Dr. Granville, through the medium of the London Medical Repository, directed their attention to its powers: and I refer those who are desirous of tracing the introduction of Prussic acid into use, as a medicinal agent, to his work.<sup>2</sup>

Prussic Acid, *internally* exhibited, is a remedy of great efficacy in spasmodic coughs of every description, particularly asthma, chronic catarrh, and hooping-cough. It has also, been employed with success in palpitations of the heart.<sup>3</sup> In my own practice, I have witnessed its powers in that affection of the *trachea*, which is often mistaken for *phthisis pulmonalis*, and is not less fatal. In true tubercular phthisis, my own experience does not enable me to say much in favour of Prussic Acid; but the mass of evidence brought forward, in testimony of its beneficial influence in this disease, by Dr. Granville, should not be overlooked; and, as I have stated in another place<sup>4</sup>, the judicious exhibition of Prussic Acid in the early stage of pulmonary consumption may do much to bring that disease under the controul of art. Prussic Acid has been found extremely useful in the treatment of those epidemic catarrhs, with which this country is occasionally visited: and no remedy is so well adapted, as an adjunct to tonics, for removing those dyspeptic affections which are attended with acidity of the stomach, and accompanied with heat and soreness of the tongue. In these cases it reduces the morbid irritability of the stomach, and thereby enables the juices of that organ to be more slowly secreted and of a more healthy character.<sup>5</sup> Its beneficial effects in asthma and in hooping-

<sup>1</sup> Brera's work is entitled, “*Prospetti dei risultamenti ottenuti nella Clinica Medica dell' Imperiale R. Università di Padova, ne' sei anni scolastici, 1809 — 1815.*”

<sup>2</sup> Treatise on the internal use of Hydro-cyanic acid, &c. 2d edition, London, 1820.

<sup>3</sup> *Lond. Med. and Phys. Journ.* Nov. 1825.

<sup>4</sup> Vide Dr. Granville's Treatise, 2d. edit. p. 376.

<sup>5</sup> Dr. Elliotson has published a small volume containing the result of his practice with Prussic acid in dyspepsia; and has stated that accident led him to try the powers of the medicine in this class of diseases. Respect for my own character obliges me to say, that nothing could surprise me more than this statement of Dr. Elliotson; as he acknowledges having read the first edition of Dr. Granville's Treatise, which contains a letter from me, dated 20th February, 1819, stating my sentiments of the utility of Prussic acid, in dyspepsia, and the *modus operandi* of the remedy, previously to his having employed it!



cough are also well established. M. Heller recommends it in aneurism of the heart and aorta.<sup>1</sup> Cases are also on record in which this acid has proved serviceable in the treatment of painful and difficult menstruation, floodings, hemoptysis, and nervous diseases. It certainly is a very powerful sedative; and may be employed in all cases, in which sedatives and narcotics are indicated, with decided advantage.

As a *local remedy*, Prussic Acid is the only application which can be depended on for allaying the itching and tingling which are so distressing in impetiginous affections. I have lately employed it with unvarying success in these complaints, and having published my observations<sup>2</sup>, I am in hopes of seeing its value determined in the hands of others. I have found it useful, also, in combination with small doses of oxy-muriate of mercury in *acne rosacea*, and several other cutaneous diseases.

The dose of Prussic Acid is from *mij.* to *mvij.* It may be administered in distilled water, or in almond emulsion, or in infusion of cinchona bark, as circumstances may require. When an over dose has been taken, its deleterious effects are best counteracted by hot brandy and water, and the ammoniated tincture of iron. M. Heller recommends bleeding, ammoniacal frictions, and acidulated drink; but, more reliance is to be placed on stimulants. As a local application, it may be used in the form of lotion, in the proportion of a fluid drachm to a fluid ounce and a half of distilled or of rose water; or as a cataplasm composed of crumb of bread, soaked in a solution of *fʒjss.* of the acid in *fʒj.* of distilled water. It is incompatible in prescriptions with nitrate of silver, the salts of iron, and the mineral acids.

Although the instantaneous power of Prussic Acid, in destroying animal life, when it is taken in doses sufficiently large to operate as a poison, may, perhaps, always prevent medical art from proving beneficial in such cases; yet it is of importance to be able to ascertain in judicial enquiries, relative to suicide or to murder, that Prussic Acid has been administered as a poison. This may be done if the animal be opened from eighteen to forty-eight hours after death. The following means pointed out by Dr. Granville, for detecting its presence in the animal system after death, should be known. Collect the blood contained in the ventricles of the heart, a portion of the contents of the stomach, and of

---

<sup>1</sup> Vide *Traité de la nécessité de ne point insister sur l'usage interieur des excitans dans l'empoisonnement par l'acide Hydrocyanique.* Par H. S. Heller. Paris, 1824.

<sup>2</sup> Vide *Medical and Physical Journal.* Feb. 1822.



any fluid that may be found in the head, the chest, or the abdomen, agitate the mixture for some time with distilled water, and filter the liquid, taking care to preserve the whole at a low temperature. To a small quantity of the filtered liquid, add a few drops of a solution of Pure Potass in Alcohol; then add a few drops of a solution of sulphate of iron; and if a reddish precipitate of the colour of burnt Terra Siena now fall down, which on the addition of a little Sulphuric acid, changes to a blueish-green, and gradually, on exposure to the atmosphere, becomes a beautiful blue, we may conclude that the death of the individual has been occasioned by Prussic Acid. But the best test is that which M. Lassaigne has proposed. Suppose a practitioner is called to examine the body of a person who has been poisoned by Prussic Acid. The stomach is first to be examined entire, to ascertain whether the odour of that Acid is perceptible in it; after which it is to be cut in pieces under *distilled* water, and a portion of it distilled with an equivalent proportion of the water until one-eighth of the liquid has passed into the receiver. That liquid is to be rendered slightly alkaline with potass, then a few drops of a solution of Sulphate of Copper is to be added to a small portion of it, and afterwards, a sufficient quantity of Muriatic Acid to re-dissolve the excess of the oxide of copper. The liquid will appear more or less milky according to the quantity of Hydro-cyanic Acid present. This test will detect  $\frac{1}{20000}$  of Hydro-cyanic Acid in solution in water.<sup>4</sup>

## II. — ACIDUM OXALICUM.

### *Oxalic Acid.*

Oxalic Acid exists, ready formed, in many vegetable and some animal bodies. Combined with Potass, it is found in the leaves of *Oxalis acetosella* and *corniculata*, *Rumex acetosa* and *Geranium acidum*; and with Lime, in the roots of Rhubarb, Valerian, and of many other plants. Berthollet procured it from honey, hair, tendons, albumen, and some other animal matters; but that which is found in the shops, manufactured for the purposes of art, is produced artificially by the action of Nitric Acid on Sugar. The following process, which was first described by Bergman, is still adopted for the production of Oxalic Acid.

Into a tubulated retort, put one ounce of White Sugar, and

<sup>1</sup> *Ann. de Chim.* xxvii. p. 200.



pour over it three ounces of strong Nitric Acid of sp. gr. 1.567. When the whole is dissolved, boil the liquor until it cease to afford nitrous fumes, and acquire a reddish-brown colour; then add three ounces more of Nitric Acid, and continue the boiling until the fumes cease, and the colour of the fluid disappear. Empty the contents of the retort into a wide vessel, and, upon cooling, a crystallization will take place, which is *Oxalic Acid*. On boiling the lixivium with two ounces of Nitric Acid in the retort, until the red fumes almost disappear, a second supply of crystals will be obtained. One hundred grains of sugar, when properly treated, yield fifty-eight grains of crystallized Oxalic Acid. The rationale of the process is very obvious: the Nitric Acid is partially decomposed, and yields up a portion of its oxygen, which is one of its components, to the sugar, which is a compound of oxygen, carbon and hydrogen; and, by this addition of oxygen, is converted into Oxalic Acid. The relative proportion of these components of sugar is oxygen 49.4, carbon 44.5, and hydrogen 6.1, in 100 parts; that of Oxalic Acid, oxygen 64, carbon 32, and hydrogen 4, in 100 parts.

*Qualities.*—The crystals of Oxalic Acid, when they are properly prepared, are four-sided prisms, with the sides alternately larger, terminated at their extremities by dihedral prisms. They are white, transparent, inodorous, have a very acrid, sour taste, and redden all the vegetable blues except indigo. One grain of Oxalic Acid communicates a sensible acidity to 2633 grains of water. The crystals dissolve in twice their weight of water at 65°, and in their own weight of boiling water. Alcohol at a mean temperature dissolves 40 parts of them, and boiling alcohol 56 parts. They are sparingly soluble in ether. The crystals are in the state of a hydrate, 100 grains containing 52 only of acid, and 48 of water.

Oxalic Acid combines with alkalies, earths, and metallic oxides, forming oxalates. Muriatic and Acetic Acids dissolve it without alteration; but it is decomposed both by Sulphuric and Nitric Acids aided by heat.

*Medical properties and uses.*—Oxalic Acid in *small* doses, when it is dissolved in a large quantity of water sweetened with sugar, forms an agreeable, cooling beverage, which may be used in febrile diseases, in the same manner and with the same intention as lemonade. It may also be employed to check an external hemorrhagy, which it appears to do by charring the blood as it issues from the wound, and thereby mechanically stopping its flow. It is a virulent poison when swallowed in large doses; and from the resemblance of its crystals to those of Sulphate of Magnesia, many fatal accidents



have occurred from mistaking Oxalic Acid for that purgative salt. The acid taste of the one salt and the bitter of the other would always prevent such accidents, were individuals to taste their medicines before swallowing them; but besides the occurrence of accidents, Oxalic Acid has, of late, been too frequently employed by the wretched suicide for the purposes of self-destruction. It is, therefore, important, that every medical practitioner should be acquainted with the qualities of Oxalic Acid, its effects on the animal economy, and the means of counteracting these, when it has been taken in a sufficient dose to operate as a poison.

The exact extent of the dose of Oxalic Acid, which may be taken with impunity has not been determined; but its poisonous properties are more or less virulent according to the degree of dilution of the dose. From some experiments on animals, which I made ten years since, and the published details of the appearances in dissection of several fatal cases of poisoning by Oxalic Acid, I was led to form an opinion that “the primary morbid action of this poison is on the stomach itself, on the coats of which its chemical action occasions the organized animal solid to enter into new combinations, and thence produces a decomposition both of the acid and the part to which it is applied:” that the acid, however, also enters into the circulation by absorption; but “that the proximate cause of death from Oxalic Acid, is the suspension of the functions of the heart and brain, which are sympathetically affected by the local injury done to the stomach.”<sup>1</sup> The subject has been, since then, investigated with much care and great ingenuity by Dr. Christison and Dr. Coindet, and their observations published in the *Edinburgh Medical Journal*.<sup>2</sup> From the labours of these gentlemen I am induced to change my opinion as to the extent of the injury done to the living stomach, and to believe that the pultaceous state of that organ, which was found by me in my experiments on dogs, is to be attributed to the action of the acid on its coats after death. I am, however, still of opinion, that death is less to be attributed to the sedative power of the poison acting on the brain and spine, to which it is carried by absorption, than to the sympathetic action on the nervous system from the local injury of the stomach, an opinion according with their first conclusions, that “Oxalic Acid, when introduced into the stomach in large doses and highly concentrated, irritates it, or corrodes it, by dissolving the gelatin of its coats; and death takes place by a sympathetic injury of the nervous system.”<sup>3</sup>

<sup>1</sup> *London Med. Repos.* vol. iii. p. 386.

<sup>2</sup> See *Edin. Med. and Surg. Journ.* vol. xix. p. 163.

<sup>3</sup> Toland, p. 185.



The general symptoms attending poisoning by Oxalic Acid, are burning pain in the stomach, violent and incessant vomiting, the matter ejected being commonly dark-coloured and sometimes bloody; in some cases there have been violent gripings and purging; the pulse soon sinks and becomes almost imperceptible, and this state is followed by deadly coldness of the limbs, attended with lividity of the fingers and nails, and clammy sweats: convulsions, but not in every instance, and insensibility, precede death. With regard to the appearances after death, no particular change in the external state of the body has been noted. On opening the body, the *stomach* is found, generally, to contain a quantity of dark-coloured fluid, which is probably extravasated blood charred by the poison; in some instances, the coats of the stomach have been found greatly injured, presenting appearances of great vascularity, thickening of the mucous coat, the rugæ pultaceous and easily wiped off, and in some cases, the other membranes have been found tender, and even perforated, so that the contents of the organ have escaped into the cavity of the abdomen. The lungs and heart have not been often examined; but, in the lower animal skilled by Oxalic Acid, both have presented indications of inflammation having existed in them, particularly the lungs. The vessels of the brain have been found turgid.

The fatal effects of poisoning by Oxalic Acid are so rapid, that little opportunity is afforded for counteracting them by medical art. The first object certainly, in every case, is to evacuate the poison from the stomach, and, when the stomach-pump is at hand, it should be instantly employed. The vomiting which usually supervenes, precludes the necessity of employing emetics and copious dilution, which, in other cases of corrosive poison is advisable, is more likely to promote the absorption of the acid and, consequently, increase its powers. Before, therefore, emetics are employed, if they should be deemed necessary, the activity of the poison should be weakened by altering its nature by some substance with which, in chemically combining, its solubility is diminished. That chalk produces this effect, I discovered in making the experiments already alluded to, and several instances have since occurred in which its administration has saved the lives of individuals poisoned by Oxalic Acid. The lime of the chalk unites with the Oxalic Acid, and forms an oxalate, which is perfectly inert. Magnesia may be employed instead of chalk, and it has the advantage of not inconveniencing the patient by the extrication of carbonic acid gas, which is copiously evolved when the chalk unites with the acid; but, as the Oxalate of Lime is more insoluble than the Oxalate of Mag-



nesia, and consequently more inert, it may be questioned whether the inconvenience of the gas be equivalent to the greater security from the employment of chalk. A mixture of chalk and water, or of magnesia and water should, therefore, be instantly given when Oxalic Acid has been taken in a large dose; and, after the local effects of the poison have been counteracted, the system should be supported by cordials combined with opium, and the oxalate afterwards carried out of the system by the aid of purgatives.

To obtain legal evidence in cases of poisoning by Oxalic Acid, when none of the poison is found, we may be guided to suspect the nature of the poison by the symptoms and the post mortem appearances: but a correct opinion can be formed only by an analysis of the vomited matter, the contents of the stomach and its coats. For this purpose the vomited matter and the contents of the stomach should be, separately, diluted with distilled water; and the coats of the organ itself boiled in distilled water. These solutions should, then, be separately filtered and decolorized with chlorine, and subjected to the following tests. If Oxalic Acid be present, hydrochlorate (muriate) of lime will occasion a precipitate which is soluble in a small quantity of nitric acid, but not in muriatic, unless a very large quantity of the acid be used. Sulphate of copper throws down a bluish-white precipitate in any fluid containing free Oxalic Acid, which is insoluble in hydrochloric acid: and Nitrate of Silver occasions a heavy white precipitate, which, when dried and heated over the flame of a candle on the point of a spatula, becomes brown at the edge, then suddenly fulminates and is all dissipated in white fumes. This is a very delicate test; for Dr. Christison informs us that, from a quarter of a grain of Oxalic Acid dissolved in 4000 parts of water, he and Dr. Coindet procured enough of the oxalate of silver to show its fulmination twice.<sup>1</sup>

---

*Edin. Med. Journ.* vol. xix. p. 198.



ALKALIES<sup>1</sup> AND SALTS.<sup>2</sup>

THE general term ALKALI comprehends under it substances possessed of very important chemical properties, and capable of producing very powerful effects on the animal œconomy. Alkalies have an acrid, urinous taste; are caustic, or dissolve animal matter; change the blue vegetable colours to green; serve as the means of combining oil and water; are capable of being fused and volatilized by a strong heat; have a great affinity for water; and combine with acids, forming neutral salts, in which the qualities of both the components are lost. The discoveries of Sir H. Davy have clearly established that the greater number of them are compound bodies, with metallic bases. They are affected by the air, and require to be preserved in well stopped, glass bottles.

The Alkalies, in reference to their chemical properties, may be arranged nearly in the same manner as the acids.

## I. ALKALIES COMPOSED OF A SIMPLE RADICAL WITH OXYGEN.

POTASSIUM

1. POTASSA.

SODIUM

2. SODA.

LITHIUM

3. LITHIA.

## II. ALKALIES COMPOSED OF A COMPOUND RADICAL WITH OXYGEN.

HYDROGEN and CARBON

- |   |                |
|---|----------------|
| { | 1. ACONITA.    |
|   | 2. ATROPIA.    |
|   | 3. CINCHONIA.  |
|   | 4. BRUCIA.     |
|   | 5. DATURA.     |
|   | 6. DELPHIA.    |
|   | 7. EMETA.      |
|   | 8. HYOSCIAMIA. |
|   | 9. MORPHIA.    |
|   | 10. PICROTOXA. |
|   | 11. QUINIA.    |
|   | 12. STRYCHNIA. |
|   | 13. VERATRA.   |

<sup>1</sup> The words *Kali* and *Alkali* are of Persian origin, and derived from the terms *كلبا* and *السي*, or *Kalla* and *Alkalia*, signifying the ashes of marine plants.

Vide *Good's Nosology, Prelim. Disc.* p. xcv. The Rev. W. Palmer, the professor of Arabic at Cambridge, gives the following as the origin of the term.

“ From the Arabic root *قلع* (*Kala*), to dress any thing by fire, is derived the substantive *قلي* (*Kilyon*), the ashes produced by burning the *Salicornia* or any other plant of the same nature. Hence *Kali*, and with the article *Al-kali*.”

<sup>2</sup> The title of this section in the London Pharmacopœia is *Alkalies and their Salts*; but as these salts cannot be termed Salts of Alkalies, in strict language,



### III. ALKALIES COMPOSED OF A COMPOUND RADICAL WITHOUT OXYGEN.

AZOTE, HYDROGEN

1. AMMONIA.

Of these Potass, Soda, and Ammonia only are employed in an uncombined state in Pharmacy

NEUTRAL SALTS, strictly speaking, have neither acid nor alkaline properties : but salts exist, formed by the combination of acids with alkalies, in which the properties of the one or the other predominate ; and consequently, although these are *secondary* salts, yet they cannot, in strict language, be denominated *neutral* salts. When the acid predominates, the salt is designated by the syllables *bi* or *super* being added to the appellation of the neutral salt, formed with the same acid and alkali ; but when the alkali is redundant, the syllable *sub* is added ; thus, if to Carbonate of Potass be added an addition of acid, equal to that which it already contained, it becomes a bicarbonate or a supercarbonate of potass ; but if there be less acid than in the neutral carbonate, the salt is a subcarbonate of potass. When the acid has the term *oxy* prefixed to its name, the same syllables are prefixed to that of the salt ; thus, *oxymuriate of potass*, denotes a salt composed of the oxymuriatic acid and potass ; terms, the impropriety of which I have already noticed. When an alkali is united with an acid, the salt formed is named from the acid, and the alkali is regarded as the base : <sup>1</sup> thus, the salt formed by *sulphuric acid* and *soda* is named *sulphate of soda*, the soda being regarded as the base of the salt. Even when the acid is united with two bases, or is a triple salt, the compound is still named from the acid ; as, for instance, *tartrate of potass and soda*, which is a compound of *tartaric acid*, *potass*, and *soda*.

The neutral and secondary salts have very different degrees of solubility ; but that of almost all of them is increased by an augmented temperature, while their solution is, for the most part, accompanied with a diminution of temperature. They may be obtained unaltered from solutions by evaporation : and, if the process be slowly conducted, they form in regular, crystallized masses, which have more or less transparency according to the quantity of water which they retain in their composition. Exposure to air, heat, and moisture, variously

we formerly translated the phrase *Neutral Salts* : but, as this term, in strict language, applies to salts in which there is no excess either of acid or of alkalies, we, therefore, have adopted the simple term *salts*.

<sup>1</sup> Before Morveau corrected the chemical nomenclature, in 1781, the genera of salts were usually referred to their bases, and the species distinguished by the acids ; but the reverse of this was proposed by him, and has been since almost generally adopted.



affect the appearance of crystallized salts. When they lose their transparency, and are covered with a white crust, or fall to powder, on simple exposure to the air, such salts are said to *effloresce*; if, on the contrary, they attract moisture from the atmosphere, and become fluid, they are said to *deliquesce*; and they are *permanent*, when the air has no effect on their crystals. The circumstance of a salt first melting in a moderate heat, then becoming covered with a white crust, and ultimately being converted into a dry, opaque mass, is termed *watery fusion*; but when, instead of melting, it splits, and the fragments fly off with a crackling noise, this effect is termed *decrepitation*.

The efflorescent and deliquescent salts should be preserved and dispensed in well-stopped bottles; while those that are permanent will not suffer from being put up in paper.

The alkalies have been employed as poisons; in which case the practitioner ought to be able to distinguish them from other acrid or caustic poisons. The volatile alkali is readily known by its odour; but if any of the fixed alkalies have been taken, besides the characters already enumerated, and which merely demonstrate the fact that the poison has been an alkali, the simplest method of ascertaining which of the alkalies is the poison in question, is to evaporate the solution, or some of the filtered fluid contents of the stomach to dryness in a silver spoon or vessel; after which, by exposing the mass to the air, if the alkali be potass it will rapidly deliquesce, but it will remain dry if it be soda. Vinegar is the best antidote of the alkaline poisons, when given early enough after the poison has been swallowed.

#### PREPARATIONS OF AMMONIA.

AMMONIÆ SUCARBO'NAS<sup>1</sup>, Lond. *Subcarbonate of Ammonia.*

“Take of Muriate of Ammonia, *a pound*; prepared Chalk, dried, *one pound and a half*. Pulverize them separately; then mix them, and sublime with a gradually increased heat, until the retort become red hot.”

SUB-CARBONAS AMMONIÆ, Edin. *Subcarbonate of Ammonia.*

“Take of Muriate of Ammonia, *one part*: softer Carbonate of Lime, dried, *two parts*: Separately pulverize and mix them, and sublime from a retort into a receiver kept cold.”

*Dublin.*

“Take of Muriate of Ammonia, reduced to powder, and well dried, Carbonate of Soda, dried, each *half a pound*. Mix: then put them into an earthen retort, and sublime with a heat gradually increased, into a receiver kept cold.”

---

<sup>1</sup> Ammonia præparata, Sal cornu cervi, P.L. 1787.



*Syn.* Carbonate d'ammoniaque (*F.*), Kohlensäures ammonium (*G.*), Drooges Kohlenzuures Ammonium (*Dutch*), Sottocarbonato d'ammoniaco (*I.*).

This salt is produced by a double decomposition of the substances employed. The lime of the chalk attracts the muriatic acid of the muriate of ammonia, while the carbonic acid is attracted by the ammonia. The muriate of lime which is formed remains in the retort, while the subcarbonate of ammonia sublimes and concretes into a cake on the sides of the receiver.<sup>1</sup> The theory of the Dublin process, in which the carbonate of soda is ordered instead of chalk, is precisely the same, only less heat is required; but it is too expensive to be generally employed. The chalk, or the carbonate of soda, should be extremely well dried, and the ingredients very intimately mixed, that the decomposition may be as complete as possible. The retort should have a wide cylindrical neck; and the receiver have a nearly cylindrical form, to permit the concreted salt to be taken out without breaking the glass.<sup>2</sup>

*Qualities.* — Subcarbonate of Ammonia has a penetrating, pungent odour, and an acrid, yet cooling taste. It is usually in a white, semitransparent, hard mass, which breaks with a striated fracture; has the specific gravity of 0.966<sup>3</sup>; and is totally volatilized, when pure, in a moderate heat. It is soluble in less than three parts of water at 60°, and in an equal weight of warm water; but in the latter it effervesces, and is partially decomposed. It is insoluble in alcohol, which coagulates to a spongy mass a strong solution of the salt. It changes the vegetable blues to green. Exposed to the air, it gradually effloresces and loses its pungent odour, and becomes a bicarbonate, owing either to the volatilization of the superabundant ammonia it contains, or to the absorption of carbonic acid from the air.<sup>4</sup>

Bergman makes its constituents to be 45 parts of carbonic

<sup>1</sup> According to the new theory, however, this salt is “the result of more complicated affinities; muriate of ammonia consists of muriatic acid and ammonia, and chalk of carbonic acid and lime; but muriatic acid is itself a compound of chlorine and hydrogen, and lime of the metallic body calcium, and oxygen. When these substances act upon each other, the muriate of ammonia and the carbonate of lime are not only decomposed, but the muriatic acid and the lime also; the hydrogen of the former, and the oxygen of the latter combine and form water, which rises in vapour, and is condensed with the carbonate of ammonia, while the chlorine of the muriatic acid uniting with the calcium of the lime forms chloride of calcium, which remains in the vessel unacted on by the heat: the compound, which was formerly called dry muriate of lime, being now, according to more modern and correct views, termed chloride of lime.” See *Phillips's Trans. of Pharm.* 1824.

<sup>2</sup> This salt is prepared, on a large scale, by sublimation from an iron pot, to which the heat is directly applied, and which is connected with a large earthen or leaden receiver. *Murray's System of Materia Medica*, ii. 228. Large quantities, but very impure, are also produced in the distillation for the preparation of gas.

<sup>3</sup> *Annales de Chimie*, xxviii. 12.

<sup>4</sup> The neutral carbonate is inodorous.



acid, 43 ammonia, and 12 water, — in 100 parts; but this statement has been shown to be erroneous; and I am inclined rather to adopt that of Mr. Phillips, who justly regards it as a sesquicarbonate, and makes the proportions to be 55·93 carbonic acid, 28·81 ammonia, and 15·26 of water.<sup>1</sup> Sir H. Davy, however, has found that the quantity of alkali varies according to the temperature that has been employed in the preparation: thus, when it is formed at a temperature of 300°, it contains rather more than 50 per cent. of ammonia; but at a temperature of 60°, it contains only 20 per cent.

Subcarbonate of Ammonia is decomposed by the acids, the fixed alkalies and their subcarbonates, supertartrate of potass, sulphate of magnesia, the metallic salts, barytes, and lime, and partially by magnesia.

*Medical properties and uses.* — This salt is stimulant, antispasmodic, antacid, diaphoretic, and in large doses emetic. It is beneficially given in gout, hysteria and dyspeptic affections, when much acid is present in the stomach; and in infantine convulsions connected with dentition, or with acidity of the primæ viæ. As a diaphoretic it is occasionally exhibited in chronic rheumatism, in combination with guaiacum; and sometimes, although rarely, it is employed to produce vomiting in gouty and paralytic cases. From the ammonia it contains in excess, the subcarbonate is applied as a local stimulant to the nostrils in syncope, hysteria, and languors; and with the addition of a little scent, forms the common smelling-salts of the shops. One part of pulverized Subcarbonate of Ammonia, and three parts of Extract of Belladonna, spread on leather in the form of a plaster, is an excellent application for allaying rheumatic pains. The ordinary dose is from grs. v. to grs. xx. formed into pills, or dissolved in any aqueous vehicle; but to excite vomiting ʒss. may be given for a dose, and repeated if necessary, assisting its operation by plentiful dilution.

Officinal preparations. *Liquor Ammoniae Subcarbonatis*, L. *Liquor Ammoniae Acetatis*, L. E. D. *Linimentum Ammoniae Subcarbonatis*, L. *Cuprum ammoniatum*, L. E. D.

LIQUOR AMMONIÆ<sup>2</sup>, Lond. *Solution of Ammonia.*

“Take of Muriate of Ammonia, *eight ounces*; Lime newly burnt, *six ounces*; Water, *four pints*. Pour one pint of the Water upon the lime, then cover the vessel, and set it aside for an hour. Dissolve the Muriate of Ammonia in the remainder of the water previously heated, add to it the former mix-

<sup>1</sup> The bicarbonate contains, Carbonic Acid 55·70, Ammonia 21·52 and Water 22·78 in 100 parts.

<sup>2</sup> Aqua Ammoniae puræ, P.L. 1787.



ture, and again cover the vessel; after the liquor has become cold, strain it, and distil twelve fluid ounces of Solution of Ammonia into a receiver, the temperature of which does not exceed  $50^{\circ}$ ."

The specific gravity of Solution of Ammonia is to that of Distilled Water as 0·960 to 1·000.<sup>1</sup>

AQUA AMMONIÆ, Edin. *Water of Ammonia.*

"Take of Muriate of Ammonia, *one pound*; Lime newly burnt, *one pound and a half*: Distilled Water, *one pound*; Water *nine ounces*. Upon the Lime broken to pieces pour the Water in an iron or earthen vessel, cover it up, until the Lime have fallen into powder and become cold, then rub the muriate to a fine powder, and triturate it with the lime in a mortar; after which put them directly into a bottle glass retort. Place the retort in a sand-bath, and adapt to it a receiver furnished with a tube passing into a phial containing the Distilled Water: the phial, however, being sufficiently large to hold double the quantity of Water. Then apply the fire, gradually raising it, until the bottom of the iron pot be red hot, and as long as gas and vapour are produced. The specific gravity of this Solution of Ammonia is to that of Distilled Water as 0·939 to 1000. It should be preserved in small phials well stopped."

AQUA AMMONIÆ CAUSTICÆ, Dub. *Water of Caustic Ammonia.*

"Take of Muriate of Ammonia, *sixteen ounces*; Lime newly burnt, *two pounds*; Water, *six pints*. Effuse one pint of Water upon the Lime placed in an earthen vessel, and cover it up. Twenty-four hours afterwards, when the Lime is crumbled to powder, mix with it the salt, avoiding the vapours; then put the mixture into a retort, and pour upon it the remainder of the Water. Agitate them; and having luted carefully the joinings of the vessels, distil with a moderate heat into a cooled receiver twenty-one measured ounces of the liquor. The specific gravity of this Solution should be to that of Distilled Water as 0·934 to 1000.

*Syn.* Dissolution d'ammoniaque (*F.*), Atzender Ammonium-liquor (*G.*), Vlugtiger etzender Salammoniak geest (*Dutch*), Liquore di Ammoniac (*I.*).

In these processes, the Lime having a superior affinity for Muriatic Acid, decomposes the Muriate, from which the Ammonia is disengaged, and passes over in combination with the watery vapour. If the temperature of the Water rises to  $130^{\circ}$ ,

---

<sup>1</sup> This process, which differs from that of the former edition of the Pharmacopœia, was suggested by Mr. Phillips; but more water is employed than he proposed, which he conceives to be objectionable; both because more fuel and larger vessels are required; and the total product is weaker in the proportion of about 9·5 to 13. Vide *Remarks in the Editio Altera of the Pharm.* Lond. p. 18.



the Ammonia is again separated in the form of gas; and hence the necessity of keeping the receivers cold: but water at 60°, takes up 780 times its bulk of gaseous Ammonia, which increases the bulk of the liquid two thirds. The product thus obtained is an aqueous solution of Ammonia; while Muriate of Lime remains in the retort, and may be dissolved out by twice its weight of water.

*Qualities.* — Liquid Ammonia is a limpid colourless fluid. It has a very strong, pungent odour, an extremely acrid taste, and corrodes the skin. Obtained according to the London College, its specific gravity is .9060, to the Edinburgh .9039, while that of the Dublin College is .9310. The following table shows the strength of liquid ammonia of different degrees of specific gravity within a certain range: (Temp. 50° Fah. Pressure 29.8 Barom.)

100 parts Sp. Grav.	Contain of		100 parts Sp. Grav.	Contain of	
	Ammonia.	Water.		Ammonia.	Water.
.9000	26.00	74.00	.9513	12.40	87.60
.9054	25.37	74.63	.9545	11.56	88.44
.9166	22.07	77.93	.9573	10.82	89.18
.9255	19.54	80.46	.9597	10.17	89.83
.9326	17.52	82.48	.9619	9.60	90.40
.9335	15.88	84.12	.9684	9.50	90.50
.9435	14.53	85.47	.9639	9.09	90.91
.9476	13.46	86.54	.9713	7.17	92.83 <sup>1</sup>

But Mr. Phillips says that the Liquid Ammonia of the London College of sp. gr. 0.960 is a compound of Ammoniacal gas 10, Water 90 parts, in 100.<sup>2</sup> For ordinary purposes it is useful to know, that a phial capable of containing 224 grains of distilled water, can hold no more than 216 grains of the strong solution. At 46° Fah. the Ammonia in this solution crystallizes; and at 68° the fluid assumes the appearance of a thick jelly, and becomes almost inodorous

Liquid Ammonia assists the oxidizement of copper and zinc; dissolves many of the metallic oxides: and unites with all the acids without effervescence, forming neutral salts. It dissolves oils, resins, and many other vegetable principles. Its affinity for carbonic acid is so powerful, that it rapidly attracts it from the atmosphere: and hence the necessity of preserving it in well-stopped small phials, as directed by the Edinburgh College. The acids, the metallic salts, and alum are incompatible in formulæ with it. The constituents of the Am-

<sup>1</sup> *Davy's Researches*, p. 68.

<sup>2</sup> *Trans. of Pharm.* p. 38.



monia it contains, according to the latest experiments of Mr. Davy, are 74 measures by bulk of Hydrogen gas, and 26 of Azotic gas: or, according to the analysis of Dr. Henry, 100 parts of Ammonia consist of 80·36 of azote, and 19·64 Hydrogen by weight.<sup>1</sup> Mr. Phillips says 17·64 of Hydrogen, and 82·36 of Azote.<sup>2</sup> The presence of carbonic acid in solution of Ammonia may be readily detected by its effervescing with acids, and by adding to it muriate of lime, which forms a precipitate if carbonic acid be present.

*Medical properties and uses.* — This solution of Ammonia is stimulant, antacid, and rubefacient. It is usefully employed, when largely diluted, in paralysis, hysteria, and syncope; and is perhaps superior to all the other antacids in relieving cardialgia and other symptoms of acidity of the stomach. As a local stimulant it is applied to the nostrils in faintings: and a rag moistened with it, and laid over the scrobiculus cordis, sometimes raises an instantaneous blister, and, by quickly inflaming the skin, always proves useful in spasms and gout in the stomach. Combined with a small portion of oil, it forms a saponaceous rubefacient, which is beneficially applied to the throat in inflammatory sore-throat, and as a friction in deep-seated inflammation and rheumatism. The dose of the solution is from  $\mathfrak{m}\text{x.}$  to  $\mathfrak{m}\text{xx.}$  in a large cupful of cold water or milk. When taken as a poison, if death be not the immediate result, the best antidote is vinegar.

Officinal preparation. *Hydro-sulphuretum Ammoniae*, E. D. *Oleum ammoniatum*, E. *Spiritus Ammoniae*, L. *Linimentum Ammoniae*, L. D. *Spiritus Ammoniae succinatus*, L. *Lin. Camphoræ comp.* L.

AQUA AMMONIÆ DILUTA, Edin. *Diluted Water of Ammonia.*

“Take of Water of Ammonia, *one part*; Distilled Water, *two parts*. Mix them together.”

This preparation is probably intended for facilitating extemporaneous prescription; but as the solution of Ammonia is never given except in a diluted form, it appears to be superfluous.

LIQUOR AMMO'NIÆ ACETA'TIS<sup>3</sup>, Lond. *Solution of Acetate of Ammonia.*

“Take of Subcarbonate of Ammonia, *two ounces*: Diluted Acetic Acid, *four pints, or a sufficient quantity*. Add the acid to the Subcarbonate of Ammonia, until the effervescence cease.”

AQUA ACETATIS AMMONIÆ, Edin. *Water of Acetate of Ammonia.*

<sup>1</sup> *Philosophical Transactions*, 1809.

<sup>2</sup> *Trans. of Pharm.* p. 38.

<sup>3</sup> *Aqua Ammoniae Acetate*, P. L. 1787.



“Take of Carbonate of Ammonia in powder, *any quantity*. Pour upon it as much weak Acetic Acid as will exactly saturate the Ammonia.”

*Dublin.*

“Take of Carbonate of Ammonia, *two ounces*. Add by small portions, with frequent agitation, *three pints and a half* of Distilled Vinegar, or as much as will saturate the Ammonia, which may be ascertained by means of Litmus.”

*Syn.* Acetate d’ammoniaque liquide (*F.*), Essigsaures Ammonium liquor (*G.*), Liquore di Minderero (*I.*).

The Subcarbonate of Ammonia employed for this preparation, is decomposed by the Acetic Acid of the Distilled Vinegar; which combining with the Ammonia, forms an Acetate that remains dissolved in the water, while the disengaged carbonic acid flies off in the form of gas, exciting effervescence. In our experiments, Distilled Vinegar of a specific gravity of 1.007 required 320 grains of the Subcarbonate to saturate a pint; thence the proportion ordered by the London and Dublin Colleges is just sufficient for saturating three pints.<sup>1</sup> Owing, however, to the variable proportion of acid in distilled vinegar this preparation cannot be obtained of an uniform strength; and provided it be accurately neutralised, which is easily known by using litmus and turmeric paper, it is of little importance. If it be not accurately saturated, some of the metallic salts particularly those of antimony, which are often ordered in conjunction with it, are decomposed.<sup>2</sup>

*Qualities.*—This solution is inodorous; has a slightly nauseous taste; and, when made with pure materials, is limpid and nearly colourless. It is decomposed by the fixed alkalies, the strong acids, alum, magnesia, lime-water, sulphate of magnesia, oxymuriate of mercury, nitrate of silver, the sulphates of zinc, copper, and iron, and acetate and subacetate of lead, which are consequently incompatible in formulæ with it.<sup>3</sup>

*Medical properties and uses.*—As a diaphoretic it is in common use in febrile diseases; and may be combined with opium, camphor, antimonials, or nitrate of potass. It is necessary to assist its determination to the skin with plentiful dilution, and a moderate degree of external heat: for by free exposure to cool air it excites the kidneys, instead of opening the skin.

<sup>1</sup> The mercury of a thermometer, the bulb of which was immersed in the solution while effervescing, sunk five degrees.

<sup>2</sup> *Remarks on the Editio Altera of the Pharm. Lond.* p. 16.

<sup>3</sup> M. de Lassone obtained the salt crystallised by sublimation, in long, slender, flattened crystals, terminating in sharp points, an inch in length, and of a pearl-white colour. They are very deliquescent; impress on the tongue a sense of coldness and sweetness; melt at 170°, and sublime at about 250°. According to Richter, they consist of 68.77 acid, and 31.23 base. *Thomson's Chemistry*, 4th ed. vii. 62.



Externally it is employed as a discutient; as a lotion to inflamed surfaces; and when diluted with rose-water, holding in solution a small portion of opium, it is an excellent collyrium in chronic ophthalmia; and still more largely diluted, is occasionally used as an injection in the commencement of gonorrhœa. I have lately used it with the best effect as a lotion in porrigo, affecting the scalp. The ordinary dose is from fʒiv. to fʒxij, given every three or four hours.

**LIQUOR AMMO'NIÆ SUBCARBO'NATIS, Lond.**

*Solution of Subcarbonate of Ammonia.*

“Take of Subcarbonate of Ammonia, *four ounces*; Distilled Water, *a pint*. Dissolve the Subcarbonate of Ammonia in the Water, and filter through paper.”

**SOLUTIO SUBCARBONATIS AMMONIÆ, Edin.** *Solution of Carbonate of Ammonia.*

“Take of Subcarbonate of Ammonia, *one part*; Distilled Water, *four parts*. Dissolve the Subcarbonate in the Water, and filter through paper.”

**AQUA CARBONATIS AMMONIÆ, Dub.** *Water of Carbonate of Ammonia.*

“Take of Muriate of Ammonia, *a pound*; Carbonate of Soda, *twenty-eight ounces*; Water, *three pints*. Distil off two pints, with a fire gradually raised. The specific gravity of this liquor is, to that of distilled water, as 1095 to 1000.”

*Syn.* Soucarbonate d'ammoniaque (*F.*), Kohlensaures Amonium liquor (*G.*), Liquore Sotto carbonato di ammoniaco (*I.*).

The formulæ of the London and the Edinburgh colleges for this preparation are to be preferred, inasmuch as they obtain by simple solution the same result as is produced from the more complicated process of the other college. The theory of the Dublin process is the same as that of the preceding preparation. The Soda of the alkali employed unites with the Muriatic Acid of the Muriate, while the Ammonia combines with the Carbonic Acid, and the Subcarbonate of Ammonia, thus formed, is volatilized and carried over with the watery vapour.

*Qualities.* — This solution has the odour and taste of the concrete Subcarbonate; is limpid and colourless; and when shaken with twice its bulk of alcohol, a nearly uniform coagulum is formed. Its specific gravity should be 1150.

*Medical properties and uses.* — The same as those of the concrete salt. It suffers by keeping and therefore should not be kept in large quantities. It is given in doses of from fʒss. to fʒj. in any bland fluid.

**LIQUOR VOLATILIS CORNU CERVINI, Dublin.**  
*Volatile Liquor of Hartshorn.*

“Take of Hartshorn, *any quantity*; put it into a retort, and



distil with a gradually raised heat, a volatile liquor, a salt, and an oil; then repeat the distillation of the volatile liquor until it become as limpid as water, separating, after each distillation, the oil and salt by filtration. The Liquor will be more easily purified, if, after each distillation, except the first, there be added to it one-sixth part of its weight of charcoal, previously made red hot, then extinguished by covering it with sand, and powdered while hot. If a sufficient quantity of Hartshorn cannot be procured, the bones of any land animals may supply its place."

*Syn.* Alkali volatil fluor (*F.*), Wässrigter Kohlensaures Ammonium liquor (*G.*), Liquore di corno di Cervo (*I.*).

In this process the gelatine of the horns, or the bones, is decomposed, and its ultimate principles, which are carbon, nitrogen (*azote*), hydrogen, and oxygen, entering into new combinations, form Subcarbonate of Ammonia, empyreumatic Oil, and Water, which are the products of the process. The subcarbonate is obtained partly in a solid form, and partly dissolved in the water, which distils over; but in both states, it is contaminated with the empyreumatic oil. The subsequent distillations are intended to free it from this oil; which, although at one time it was supposed to add to the efficacy of the remedy, yet, is now conceived to be useless, and a disadvantage: nevertheless, when it is completely removed by the charcoal, this preparation does not differ from a simple solution of the subcarbonate of ammonia in water.

The Volatile Liquor of Hartshorn found in the shops is part of the product of the distillation of bones on a great scale; and is never completely free from the empyreumatic oil, which is very perceptible in its odour and taste, and gives it a slight yellow tinge. It is often adulterated by the addition of a considerable portion of water; and this cannot be known by its pungency, which is kept up by adding to it a small quantity of Liquor Ammoniae. The fraud, however, may be detected by mixing a small portion of the suspected Liquor with twice its bulk of alcohol; when, if no considerable quantity of salt be precipitated, it is certainly adulterated.

#### PREPARATIONS OF POTASS.

LIQUOR POTASSÆ<sup>1</sup>, Lond. *Solution of Potass.*

"Take of Subcarbonate of Potass, *a pound*; Lime fresh burnt, *half a pound*; Boiling Distilled Water, *a gallon*. Dissolve the Subcarbonate of Potass in two pints of the Water. Add the remainder of the Water to the Lime, mix the hot liquors together, then set the mixture aside in a covered vessel, and when it is cold, let it be strained through a cotton bag.

<sup>1</sup> Aqua Kali Puri, P. L. 1787.



If, on the addition of any diluted acid, effervescence be excited, more Lime must be added, and the filtration repeated. A pint of this solution ought to weigh sixteen ounces."

AQUA POTASSÆ, Edin. *Water of Potass.*

"Take of Lime fresh burnt, *eight ounces*; Subcarbonate of Potass, *six ounces*; Boiling Water, *twenty-eight ounces*. Let the Lime be put into an iron or earthen vessel, with twenty ounces of the Water. When the ebullition ceases, immediately add the salt, dissolved in the remaining eight ounces of the Water; and having thoroughly mixed the whole, cover the vessel till the mixture cool. The mixture being cooled, agitate it well, and pour it into a glass funnel, the tube of which is obstructed with a piece of clean linen. Cover the upper orifice of the funnel while its tube is inserted into another glass vessel, that the solution of potass may gradually drop through the linen into the lower vessel. When it first ceases to drop, pour a few ounces of water into the funnel, but cautiously, so that the fluid may swim above the matter. The Water of Potass will again begin to drop. The affusion of Water, however, must be repeated until three pounds have filtered, which will be in the space of two or three days; then let the upper parts of the solution be mixed with the lower by agitation, and preserve it in a well-stopped vessel."

AQUA KALI CAUSTICA, Dub. *Water of Caustic Kali.*

"Take of Lime fresh burnt, *eight ounces*; Subcarbonate of Kali, *six ounces*. Pour upon the Lime, put into an earthen vessel, two pints of Boiling Water; and when it is slacked, mix the Salt with the Lime, and cover the vessel. Pour the materials, as soon as they are cold, into a glass funnel, the tube of which is obstructed with a linen rag. Cover the funnel, and allow the lixivium to drop into a vessel placed below it, pouring water into the funnel occasionally, until three pounds are filtered. Let the solution be shaken, and preserved in a well-stopped green-glass bottle.

"If the ley be rightly prepared it will be colourless, inodorous, and will scarcely effervesce when mixed with an acid. If it effervesce considerably, let a small portion of fresh burnt Lime, in fine powder, be added; digest for twenty-four hours in a covered vessel, frequently agitating; and finally, filter the ley in the manner already directed.

"The specific gravity of this solution is, to that of distilled water, as 1100 to 1000."

*Syn.* Dissolution de Potasse (*F.*), Flüssiges ätzendes Kali (*G.*), Potachlog (*Dutch*), Liquore di Potassa (*I.*).

The rationale of these processes is very obvious: the lime attracts the carbonic acid of the subcarbonate of potass and leaves its alkaline base in a state of purity. In considering



the proportions of the two latter of these processes, there appears *a priori*, a much larger proportion of lime ordered than is necessary for the decomposition of the subcarbonate of potass: but if the theory of Berthollet<sup>1</sup>, as to the effect of quantity in influencing chemical affinities, be just, the superabundance is necessary to insure the more perfect separation of the carbonic acid from the potass. If the lime, however, be well burnt, and recent, the solution is obtained almost perfectly free from carbonic acid, by the quantity ordered in the London formula; but unless much care be taken to exclude the air during the filtration, it will be rapidly attracted from the atmosphere. Calico is the best substance for stopping the mouth of the funnel, and it should be supported on a rough pebble or siliceous stone, previously dropped into the funnel, and allowed to settle itself. It should be kept in small glass-bottles, fitted with ground stoppers, to prevent the absorption of carbonic acid from the atmosphere.

*Qualities.*—Solution of Potass is inodorous, and so caustic as not to admit of being tasted. It is limpid, colourless, dense, and has an oily appearance when agitated; does not effervesce with acids; and ought not to afford a precipitate with lime-water; but it is seldom procured so pure. It feels soapy when rubbed between the fingers, owing to the solution of the cuticle. When perfectly pure, it remains transparent on the addition of barytic water. Prepared according to the formulæ of the Pharmacopœias, it is not a simple solution of potass, but contains small portions of muriate, and sulphate of potass, silica, and generally some lime. The presence of muriates may be discovered by saturating a portion of the solution with nitric acid, then adding nitrate of barytes to precipitate the sulphates, if any; and lastly, adding a solution of nitrate of silver, which is precipitated if any muriate be present. Sulphates are discovered by saturating with muriatic acid, and adding muriate of barytes; and if lime be present, blowing into the solution through a tube will render it turbid; but these contaminations do not alter its effects as a remedy, nor as a pharmaceutical agent. One pint of it should weigh sixteen ounces; or the specific gravity be 1.056. It dissolves gum, extractive and resin, and converts fat and oils into soaps.

*Medical properties and uses.*—This solution is diuretic, antacid; and lithontriptic. The two first properties it certainly possesses in a considerable degree: but its continued use, even when much diluted, is said to debilitate, and otherwise injure the stomach. As a solvent of calculus, both in the kidneys

---

<sup>1</sup> *Chemical Statics*, vol. i.



and bladder, this alkali has long been celebrated: it acts, however, on calculi composed of uric acid, or of urate of ammonia, only; the presence of which in the habit is known by a red deposit in the urine of the patient. But although the continued use of Solution of Pure Potass certainly renders the urine alkaline, yet there is reason to believe that its solvent effects on calculous matter in the kidney or the bladder are not equivalent to the irritation which its long use excites both in the stomach and the bladder; and as a prophylactic, its place can be much better supplied by magnesia and the alkaline carbonates. Dr. Willan says, he has seen the most beneficial effects experienced from the internal use of this solution in lepra<sup>1</sup>: and from my own experience I can assert, that it may be almost regarded as a specific in the various species of psoriasis, which depend altogether on acidity of the primæ viæ, and a hasty and consequently imperfect formation of the juices of the stomach. It is also used as a local stimulant, much diluted, in the form of lotion, to the joints, in rachitis and gouty swellings; and, in its concentrated state, as a caustic, to destroy the poison introduced by the bite of rabid or venomous animals.

The dose of this solution may be from  $\mathfrak{m}\mathfrak{x}$ . to  $\mathfrak{f}\mathfrak{z}\mathfrak{j}$ . 'taken in chicken-broth, milk, or almond mixture<sup>2</sup>; but in cases of psoriasis, it should be gradually increased to  $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$ . In cases of acidity of the stomach, it may be administered in some bitter infusion.

Officinal preparations. *Potassa fusa*, L. E. D. *Potassa cum Calce*, L. E. D. *Liquor Sulphureti Kali*, D. *Antimonii Sulphuretum præcipitatum*, L. E.

**LIQUOR POTASSÆ SUBCARBONATIS**, Lond.  
*Solution of Subcarbonate of Potass.*<sup>3</sup>

“Take of Subcarbonate of Potass, *a pound*; Distilled Water, *twelve fluid ounces*. Dissolve the Subcarbonate of Potass in the water, and filter the solution through paper.”

**AQUA SUBCARBONATIS KALI**, Dub. *Water of Subcarbonate of Kali*.

“Take of Subcarbonate of Kali, *any quantity*; let it be put into a wide-mouthed glass funnel, the neck of which is obstructed with a piece of linen; then set it in a cellar that the moist air may liquefy the salt. Let the ley or solution be received into a vessel placed under it.”

*Syn.* Dissolution de Soucarbonate de Potasse (*F.*), Flüssiges Kohlensaures Kali (*G.*), Liquore di Sotto-carbonato di Potassa (*I.*).

<sup>1</sup> *Willan on Cutaneous Diseases*, p. 141.

<sup>2</sup> Table Beer is sometimes recommended as the vehicle for administering it, but that liquor is seldom so free from acid, as not to destroy the alkaline properties of the remedy.

<sup>3</sup> *Aq. Kali præparatum*, P. L. 1787.



Of these two formulæ, that of the London College is to be preferred, as it affords the preparation with greater facility, and always of a definite strength. The bulk of the fluid is increased rather more than one-third part. In the Dublin preparation, from the length of time which is required, the deliquescent salt attracts carbonic acid from the air, and becomes nearly a neutral carbonate.

*Qualities.*—This solution should be perfectly clear, colourless, and inodorous: and possess the properties of the subcarbonate from which it is prepared. It cannot enter into extemporaneous formulæ with vegetable infusions containing much tannin, or with lime-water, magnesia, sulphate of magnesia, or the metallic salts; as these substances decompose it, or are decomposed by it.

*Medical properties and uses.*—These are the same as those of the concrete salt. The dose may be from  $\mathfrak{m}x.$  to  $\mathfrak{f}3ij.$ , in any convenient vehicle.

POTASSA CUM CALCE, Lond. *Potass with Lime.*<sup>1</sup>

“Take of Solution of Potass, *three pints*; Lime, fresh burnt, *a pound*. Boil the solution of Potass down to a pint, then add the Lime, previously slaked by the water, and intimately mix them.”

POTASSA CUM CALCE; olim, CAUSTICUM COMMUNE MITIUS, Edin. *Potass with Lime*; formerly, *Milder common Caustic*.

“Take of the Water of Potass, *any quantity*. Evaporate it to one-third part in a covered iron vessel: then mix with it as much newly-slaked lime as will bring it to the consistence of a solid paste, which is to be preserved in a well-stopped vessel.”

KALI CAUSTICUM CUM CALCE, Dub. *Caustic Kali with Lime*.

“Evaporate water of Caustic Kali to one-third part; then add as much fresh burnt Lime in powder as will form a mass of a proper thickness, which is to be preserved in a well-stopped bottle.”

The addition of the Lime in these preparations renders the Potass less deliquescent, and consequently more manageable as an escharotic.

POTASSA FUSA, Lond. *Fused Potass.*<sup>2</sup>

“Take of Solution of Potass, *a gallon*. Evaporate the Water in a clean iron vessel over the fire, until the ebullition having ceased, the Potass melts, and then pour it out upon a clean iron plate into pieces of proper form.”

POTASSA; olim, CAUSTICUM COMMUNE ACERRIMUM, Edin. *Potass*; formerly *Stronger common Caustic*.

<sup>1</sup> *Calx e Kali Puro*, P. L. 1778.

<sup>2</sup> *Kali Purum*, P. L. 1787.



“Take of Solution of Potass *any quantity*. Evaporate in a covered very clean iron vessel, until, the ebullition being over, the saline matter flow smoothly like oil, which happens before the vessel becomes red hot. Then pour it out upon a clean iron plate; cut it into small masses before it hardens, and let it be preserved in well-stopped phials.

KALI CAUSTICUM, Dub. *Caustic Kali*.

“Take of Solution of Caustic Kali, *any quantity*. Evaporate it over the fire in a clean iron vessel, until, the ebullition having ceased, the saline matter, on increasing the heat, remain almost quiescent in the vessel. Pour out the melted salt upon a clean iron plate; and while it is concreting, let it be cut into proper pieces, which must be immediately put into a phial closely stopped. During the evaporation the operator must avoid the drops which may be thrown out from the vessel.”

*Syn.* Potasse fondue (*F.*), Trocknes ätzendes Kali (*G.*), Pietra caustica *I.*, Reine Potasche (*Dutch*).

The concrete Potass procured by these processes is a hydrate, sufficiently pure for medical purposes, but it still contains the same foreign ingredients as the solution. To procure it as pure and free as possible from carbonic acid, the evaporation should be performed in a silver vessel very quickly, the vessel should be deep, so that the watery vapour which rises may exclude the atmospheric air. It is generally run into moulds, and formed into solid cylinders, which are covered with paper, and kept in well-stopped bottles. The method of Berthollet<sup>1</sup> for obtaining it in perfect purity, which is usually described in chemical and pharmaceutical works, is too troublesome and expensive to be generally adopted. The following method proposed by Lowitz is more economical.

A solution of Potass must be evaporated till a pellicle forms on its surface, then allowed to cool; and the saline deposit, which consists chiefly of the foreign salts, carefully separated. The evaporation is then to be renewed, skimming off the pellicles that form on the surface of the fluid, which, as soon as these cease to be produced, and the ebullition is ended, must be removed from the fire, and constantly stirred till it is cold. The mass is next to be dissolved in twice its weight of distilled cold water, the solution filtered, and evaporated in a clean iron or silver basin<sup>2</sup> until crystals are deposited. If the heated fluid consolidate into a mass, in any degree, a small portion of water must be added, and the mass again heated to fluidity. The supernatant liquor is left of a brown colour, which, after being kept for some time at rest in well-stopped phials, deposits the colouring matter, and may be again evaporated and

<sup>1</sup> *Journal de Physique*, xxviii. 402.

<sup>2</sup> Lowitz orders the evaporation to be performed in a glass retort; but Pure Potass, when hot, dissolves glass.



crystallised as before. The crystals obtained in the various evaporations are colourless, Pure Potass.<sup>1</sup>

*Qualities.* — Concrete Pure Potass should be a white, brittle substance, having the peculiar odour of slaking quicklime, and a degree of causticity which prevents it from being tasted. It is usually, however, of a blueish colour. It attracts water and carbonic acid rapidly from the atmosphere, and is completely soluble in less than its own weight of that fluid at 60°. It dissolves readily in alcohol. When heated to 360° it melts, and at red heat is volatilised. It unites with sulphur, the acids, many of the metallic oxides, and the fixed oils. Its constituents, according to the analysis of Sir H. Davy, who first ascertained its compound nature, are, in 100 parts, 86 of a metal which has been named potassium, and 14 of oxygen.<sup>2</sup> It is, however, in fact, a compound of potass and water, or a hydrate, consisting of potass 84·2, and water 15·8, in 100 parts<sup>3</sup>; or, according to Mr. Brande, potassium 70·2, oxygen 14·0, and water 15·8.<sup>4</sup>

*Medical properties and uses.* — Concrete Potass is used only as an escharotic, for forming issues in diseases of the hip-joint, the spine, and in deep-seated inflammations. It erodes the skin and soft parts beneath it to a certain extent, destroying the life of the part, which is subsequently thrown off as a slough, and an ulcer is left. To prevent inconvenience from its deliquescent nature, the skin should be covered with a piece of calico, spread with adhesive plaster, and having a hole in its centre sufficient to bare the part only where it is intended to apply the caustic. It is much and justly recommended for the removal of strictures of the urethra.

POTASSÆ ACETAS, Lond. *Acetate of Potass.*

“Take of Subcarbonate of Potass, *a pound and a half*, of the stronger Acetic Acid, *two pints*; boiling Distilled Water, *two pints*. Mix the Acid with the Water, and pour it upon the Subcarbonate of Potass till all ebullition cease, after which filter. First evaporate the solution in a water-bath until no more bubbles rise; then expose it to a gradually augmented heat, and continue the evaporation until a pellicle form; which should be removed, and dried on blotting paper. Repeat the evaporation again and again, removing the pellicles as they form, and drying them in the manner already described.”

ACETAS POTASSÆ, Edin. *Acetate of Potass.*

“Take of very pure Carbonate of Potass, *one pound*; Weak Acetic Acid, *a sufficient quantity*. Boil the Subcarbonate in five pounds of the Acid; and add more Acid at different times, until the watery part of the former portion being nearly dis-

<sup>1</sup> *Nicholson's Journal*, 4to. i. 164.

<sup>2</sup> *Phil. Trans.* 1808.

<sup>3</sup> *Phillips's Trans. of Pharm.* 1824.

<sup>4</sup> *Manual of Pharm.* p. 222.



sipated by evaporation, the Acid newly added occasion no effervescence, which will be the case when about twenty pounds of it have been consumed; then evaporate slowly to dryness. Liquefy this impure salt with a gentle heat for a short time; then let it be dissolved in water, and filtered through paper. If the liquefaction have been properly performed, the filtered fluid will be limpid; but, otherwise of a brown colour. Afterwards evaporate this fluid in a shallow glass vessel, so that when removed from the fire, it may pass into a crystalline mass. Finally, the Acetate of Potass ought to be preserved in closely shut vessels."

ACETAS KALI, Dub. *Acetate of Kali.*

"Take of Subcarbonate of Kali, *any quantity*. Add to it at different times about five times its weight of Distilled Vinegar moderately heated. When the effervescence shall have ceased, and the fluid is somewhat evaporated, add, at intervals, more Distilled Vinegar, until the mixture entirely cease to effervesce: then evaporate to dryness, and having raised the fire a little, cautiously liquefy the mass. Dissolve the salt in water, after it is cold: filter the solution and let it be boiled, until, on being removed from the fire, it concrete into a crystalline mass, which should be very white. Put this mass, at the moment, into closely-stopped bottles."

*Syn.* Kali acetatum, P. L. 1787. Acetate de Potasse (*F.*), Easigsaures Kali (*G.*), Azynzuure Potasch (*Dutch*), Acetato di Potassa (*I.*).

In these processes, the Acetic Acid of the Distilled Vinegar combines with the Potass of the Subcarbonate, and expels the Carbonate Acid in a gaseous form, exciting effervescence. Owing to the largely diluted state of the acid in Distilled Vinegar, a very considerable quantity is required to saturate the Potass. The London College, therefore, in order to obviate this advantage, now order the Pyroligneous Acid diluted with an equal quantity of water, to be used instead of Distilled Vinegar, 100 grains of which saturate 48 grains of the Subcarbonate of Potass. Towards the point of saturation, the solution acquires a reddish-brown colour, and during the evaporation a quantity of carbonaceous matter is deposited, arising from the Distilled Vinegar retaining some of the extractive of the common vinegar; or if the liquor be evaporated to dryness, a brownish-coloured salt is obtained. The filtering the evaporated fluid, or fusing the salt, or keeping it for a little time fluid, then dissolving it in water, and filtering it, frees it almost entirely from colour, and a light carbonaceous matter remains on the filter. The filtered solution is nearly limpid and colourless; and when again evaporated forms a nearly colourless salt.<sup>1</sup> It is rendered still more

<sup>1</sup> A very pure and beautiful salt, but too expensive for common use, may be prepared by adding to a solution of two parts of Superacetate of Lead, a solution



colourless, if a portion of animal charcoal be added to the solution.<sup>1</sup>

This salt may also be prepared with the residue after the distillation of vinegar; but the process is not more economical than when distilled vinegar is used.<sup>2</sup>

*Qualities.*—Acetate of Potass has a slight, peculiar odour, and a warm, sharp taste. It is usually in white masses, of a foliated, soft texture, shining, and becoming soon moist if exposed to the air. Exposure to a red heat converts it into subcarbonate of potass. One fluid ounce of distilled water at 60° dissolves 504 grains; or 100 parts of it are soluble in 102 parts of water, and in twice its weight of alcohol. It is sometimes adulterated with tartrate of potass, which may be detected by adding to a solution of the salt a solution of tartaric acid, which, if tartrate of potass be present, will occasion a copious precipitate. The same adulteration may be detected by the acetate of lead forming a precipitate, soluble in acetic acid. Sulphates are detected by adding a solution of nitrate of barytes; and muriates, by adding nitrate of silver. In the watery solution it is spontaneously decomposed; and is also decomposed by the strong acids; by a decoction of tamarinds; the sulphates of soda and of magnesia: the muriate of ammonia; the tartrate of potass and soda; and by solutions of oxymuriate of mercury, and of the nitrate of silver; which consequently cannot enter into formulæ with it.

Its constituents, according to the experiments of Dr. Higgins, are 38·5 of acid, and 61·5 of alkali—in 100 parts<sup>3</sup>: but they are more correctly stated to be 51 of acetic acid, and 49 of potass.<sup>4</sup>

*Medical properties and uses.*—Acetate of Potass is mildly cathartic and diuretic. It is found to be occasionally beneficial in febrile affections and jaundice; but its principal use is in dropsies, and other diseases in which a copious discharge of urine is required. The manner in which this is effected, is endeavoured to be explained by Dr. Paris, by assuming as a fact, that the stomach possesses “the power of readily decomposing all saline compounds, into which vegetable acids enter as ingredients, and of eliminating their alkaline base, which

---

of one part of Subcarbonate of Potass, and after filtering the liquor, adding to it a small portion of Sulphate of Potass, and filtering again before evaporation. Vide *Journ. de Pharm. Maii*, 1818. p. 203.

<sup>1</sup> *Ann. de Chim.* vol. lxxxvi. p. 44.

<sup>2</sup> This salt was first described by Raymond Lully, and has been known by a great variety of names; as, for instance, *Arcanum tartari*, *secret foliated earth of tartar*, *essential salt of wine*, *regenerated tartar*, *diuretic salt*, and *digestive salt of Sylvius*.

<sup>3</sup> *Higgins on Acetous Acid*, p. 8.

<sup>4</sup> *Phillips's Translation of the Pharm.* 1824.



being in the course of the circulation carried to the kidneys, excites them into action and promotes the excretion of urine."<sup>1</sup> To produce the latter effect, the dose may be from ℥j. to 3j., given every three or four hours, in any bland fluid, or united with infusions of any of the lighter vegetable bitters, such as quassia or gentian. Doses of ʒij. or ʒiij. open the bowels.

Officinal preparations. *Acetas Hydrargyri*, E. D. *Tinctura Acetatis Ferri*, D. *Acidum aceticum*, D. E.

POTASSÆ CARBONAS, Lond. *Carbonate of Potass.*

"Take of the Solution of Subcarbonate of Potass, *a gallon*. By means of a proper apparatus transmit Carbonic Acid through the solution of Subcarbonate of Potass, until it is saturated, and filter. Evaporate the filtered solution until crystals form, being careful not to raise the heat above 120°. Separate the crystals from the fluid, and dry them on blotting-paper.

"Carbonic Acid is easily obtained from White Marble and Diluted Sulphuric Acid."

CARBONAS POTASSÆ, Edin. *Carbonate of Potass.*

"Take of Pure Subcarbonate of Potass, *two parts*; Water, *three parts*. Dissolve the salt in the Water, and by means of a proper apparatus, throw into it a stream of Carbonic Acid Gas. Filter the solution, when it ceases to absorb the acid, and then evaporate it by a heat not exceeding 180°, that crystals may form. The Carbonic Acid is easily obtained by pouring diluted Sulphuric Acid on pulverized Carbonate of Lime."

*Syn.* Carbonate de Potasse (*F.*), Kohlensaures Kali (*G.*), Koolenzuure Potasch (*Dutch*), Carbonato di Potassa (*I.*).

By these processes a pure and completely neutralized Carbonate of Potass is obtained; and any silex the Subcarbonate may have contained is completely separated. The present formula is an improvement on the last, which directed this salt to be prepared by heating a mixed solution of Subcarbonate of Potass and Subcarbonate of Ammonia, so as to drive off the Ammonia, produced by the decomposition of the Subcarbonate, in a gaseous form. But it involves some objections which have been properly pointed out by Mr. Phillips. Thus, by using sulphuric acid to evolve the carbonic acid from the marble, the sulphate of lime which is formed, envelopes a portion of the carbonate and prevents its decomposition, unless the vessel containing it be provided with an agitator, which is not required if muriatic acid be employed. Another objection is that the solution of the subcarbonate of potass is so strong, that owing to the bicarbonate being less soluble in water than the subcarbonate of the Pharmacopœias, that salt is inconveniently precipitated during the process. The alkaline solution should therefore be diluted with four parts of water.

<sup>1</sup> *Pharmacologia.*



*Qualities.* — This salt, prepared by these formulæ, is, properly speaking, a Bicarbonate; the Subcarbonate of the Pharmacopœias being really a Carbonate. It is inodorous, has a slightly alkaline taste, without any acrimony, and scarcely acts upon turmeric paper. It is in small tetrahedral, rhomboidal prisms, with dihedral summits, of a beautiful white colour, which are not altered by exposure to the air, are soluble in four parts of water at 60°, and  $\frac{5}{8}$ ths of their weight of boiling water, in which they are partially decomposed, carbonic acid gas being emitted during the solution. It is insoluble in alcohol. Its spec. grav. is 2.012, and its constituents, according to Pelletier, are 43 Acid, 41 of Alkali, and 16 of Water—in 100 parts;<sup>1</sup> but according to Mr. Phillips, of Carbonic acid 43.56, Potass 47.53, and Water 8.91. It is incompatible in formulæ with the acidulous salts, borax, muriate of ammonia, alum, sulphate of magnesia, lime-water, and all the metallic salts. Mr. Phillips remarks, that calomel, unless heat be applied, is not decomposed by it.

*Medical properties and uses.* — On account of the increased quantity of carbonic acid which this salt contains, it is preferable to the common subcarbonate for effervescing draughts; but does not differ from it in its properties as a remedy. The dose is from grs. x. to 3j.

POTASSÆ SUBCARBONAS, Lond. *Subcarbonate of Potass.*<sup>2</sup>

“Take of Impure Potass (*pearl-ashes*), reduced to powder, *three pounds*; Boiling Water, *three pints and a half*. Dissolve the Potass in the Water, and filter; then pour the solution into a clean iron pot, and evaporate with a gentle heat until the liquor thicken; lastly, being taken from the fire, stir assiduously with an iron spatula, until the salt concrete into small grains.”

“A purer Subcarbonate of Potass may be prepared in a similar manner from Tartar, previously burnt until it be of an ash colour.”

SUBCARBONAS POTASSÆ, Edin. *Subcarbonate of Potass.*

“Let Impure Carbonate of Potass be put into a crucible and exposed to a red heat. Then triturate it with an equal weight of Water. Pour the solution, after the impurities have subsided, into a clean iron pot, and boil it to dryness; stirring the salt constantly towards the end of the boiling, to prevent it from adhering to the vessel.”

SUBCARBONAS KALI, Dub. *Subcarbonate of Kali.*

“Take of Pot-ashes coarsely powdered, Cold Water, each *six pounds*. Mix them by trituration, and macerate them in a

<sup>1</sup> *Ann. de Chim.* t. xv. p. 55.

<sup>2</sup> *Kali præparatum*, P. L. 1787.



wide vessel for the space of a week, with frequent agitation; then filter the solution, and evaporate it to dryness; and towards the end of the process assiduously stir the saline mass with an iron spatula. In this manner having reduced it to a coarse powder, preserve it in well-stopped vessels. Previous to dissolving the ashes in the water, if they be very impure, roast them in a crucible until they become white."

*Syn.* Soucarbonate de Potasse (*F.*), Kohlensaures Kali (*G.*), Unvollmaakle Kohlenstoffzuure potasch (*Dutch*), Sotto-carbonato di Potassa (*I.*).

The Potass of commerce is a heterogeneous mass, consisting chiefly of Subcarbonate of Potass, with small portions of sulphate of potass, muriate of potass, siliceous earth, oxide of iron, and oxide of manganese, in various proportions. (See *Part ii.*) The above processes are intended to separate the Subcarbonate of Potass in as pure a state as possible; and by following the directions of any of the Pharmacopœias, it is obtained sufficiently pure for medicinal purposes; while the insoluble metallic salts, and the greater part of the siliceous earth, are left on the filter when the solution is strained. It may be obtained in a still purer state by using cold water instead of warm; evaporating the solution till a pellicle form on the surface, and allowing it to stand for some hours, in order that the muriate of potass and the sulphate of potass may be crystallized and separated; after which the solution of the subcarbonate can be evaporated, and treated as above.

*Qualities.* — The salt obtained by the above processes is really a Carbonate, being composed of one atom of each of its components. In its dry state it consists of carbonic acid 31.43, potass 68.57 in 100 parts; but as a hydrate, the state in which it usually exists, it contains 16 per cent. of water. It is in coarse white grains, which are so deliquescent, that they soon attract from the air as much water as dissolves them, forming a fluid of the consistence of oil; and thence the salt must be kept in well-stopped bottles. Its taste is acrid and urinous; it changes to green the vegetable blue and red colours, combines with oils, and forms soaps, and is decomposed by acids with effervescence.<sup>1</sup> It does not decompose tartrate of iron, with which it may be, therefore, ordered in prescriptions.

**SUBCARBONAS POTASSÆ PURISSIMUS, Edin.**  
*Pure Subcarbonate of Potass.*

"Take of Impure Supertartrate of Potass, *any quantity.* Wrap it up in moist bibulous paper, or put it into a crucible; and having placed it among live coals, let it be burnt to a black mass; which, after having reduced it to powder, expose

<sup>1</sup> *Nicholson's Journal*, 4to iii. 215.



in an open crucible to a moderate fire, until it become white, or at least ash-coloured, taking care that it be not melted. Then dissolve it in warm water; strain the solution through a linen cloth, and evaporate it in a clean, iron vessel, stirring constantly towards the end of the process with an iron spoon, lest any of it should adhere to the bottom of the vessel. A very white salt will remain, which is to be left a little longer on the fire, till the bottom of the vessel become red hot. Finally, when it is cold, let it be preserved in well-stopped glass vessels.

KALI E TARTARO, Dub. *Kali from Tartar.*

“Take of Crystals of Tartar, *any quantity*: Heat it to redness in a silver crucible lightly covered, until fumes cease to be emitted. Let the residue be reduced to a coarse powder, and in the same crucible left uncovered, roast it for two hours, stirring it frequently. Then boil it in twice its weight of water, during a quarter of an hour; and after due subsidence of the impurities, pour off the pure solution. Let this part of the process be three times repeated. Filter the mixed leys, and evaporate them in a silver vessel; then, while the residuary salt is drying, granulate it by brisk agitation, and expose it to an obscure red heat. Take it out of the vessel before it be quite cold, and let it be preserved in well-stopped phials.”

*Syn.* Soucarbonate de Potasse (*F.*), Sotto carbonato di Potassa (*I.*).

The product of these processes is a Carbonate of Potass. The degree of heat to which the crude Supertartrate is exposed decomposes its tartaric acid; and by the re-union of two of its components, oxygen and carbon, carbonic acid is formed, which combines with the potass, while the remaining carbonaceous matter produced by the decomposition is burnt out by the subsequent roasting. The resulting saline mass, besides Carbonate of Potass, contains also a small portion of carbonate of lime and some argil, which however are separated by the solution and filtration.

*Qualities.*—These are in every respect the same as those of the salt obtained from the potashes of commerce; it, however, contains fewer impurities. Its constituents, according to Berard, are 29·79 acid, and 70·21 alkali and water, in 100 parts.<sup>1</sup>

Subcarbonate of Potass (to use the name of the Pharmacopœias) is often adulterated, or very carelessly prepared. If one part of it be dissolved in eight parts of distilled water and saturated with pure nitric acid, the presence of siliceous earth will be indicated by the solution becoming turbid, and by weighing the precipitate, its quantity may be ascertained. A

<sup>1</sup> *Annales de Chimie*, lxxi. 55.



precipitate being formed on the addition of muriate of barytes indicates the presence of sulphates; a white preeipitate turning blueish on exposure to the light, on adding nitrate of silver, proves the presence of muriatic salts; and calcareous earth is rendered evident by dropping into a solution of the subcarbonate a few drops of a solution of oxalic acid or oxalate of ammonia.

*Medical properties and uses.* — Subcarbonate of Potass is deobstruent, diuretic, and antacid. In small doses, it is sometimes given in cases of glandular obstructions of the abdominal viscera, particularly hepatic obstructions, with seeming advantage; but it is not certain that the benefit does not arise from the effects of the remedy in correcting acidity of the primæ viæ. Its effects on the kidneys are considerable, when aided by plentiful dilution. The principal use, however, of this salt in medicine, is for the formation of saline draughts, for which purpose it is given in combination with a solution of citric acid, or with recent lemon juice, in the proportion of ℥j. of the salt to fʒiv. of the lemon juice, or of the acid solution, in febrile affections. When given as an antacid, its taste and acrimony are most perfectly covered with milk.

*Officinal preparations.* *Aqua Supercarbonatis Potassæ*, E. *Potassæ Acetas*, L. E. D. *Potassæ Carbonas*, L. *Potassæ Tartras*, L. E. D. *Liquor Potassæ*, L. E. D. *Liquor Potassæ Subcarbonatis*, L. D. *Sulphas Potassæ*, E. *Magnesiæ Carbonas*, L. D. *Potassæ Sulphuretum*, L. E. D. *Liquor arsenicalis*, L. *Sulphur antimoniatum fuscum*, D. *Alcohol*, L.

POTASSÆ SULPHAS, Lond. *Sulphate of Potass.*<sup>1</sup>

“Take of the salt which remains after the distillation of the Nitric Acid, *two pounds*; Boiling Water, *two gallons*. Mix them so as to dissolve the salt; and then add as much Subcarbonate of Potass as may be sufficient to saturate the acid. Next boil till a pellicle form on the surface, and after filtering the liquor set it aside to crystallize. Pour off the water, and dry the crystals on bibulous paper.”

SULPHAS POTASSÆ, Edin. *Sulphate of Potass.*

“Dissolve the Acidulous Salt, which remains after the distillation of Nitrous Acid, in hot water, and add as much Carbonate of Lime in powder, as will saturate the superfluous acid, and leave the whole at rest, until the fæces subside. Having poured off the fluid, filter it through paper, and evaporate until crystals form.”

SULPHAS KALI, Dub. *Sulphate of Kali.*

“Dissolve the Salt which remains after the distillation of Ni-

---

<sup>1</sup> This name was imposed by the French chemists in 1787. The following are some of its old names: *Nitrum fixum*, *arcantum duplicatum*, *sal de duobus*, *sal polychrestus*, *tartarum vitriolatum*, *kali vitriolatum*.



tric Acid, reduced to a powder, in a sufficient quantity of warm water. Add as much Potass as will saturate the superfluous acid. Let the filtered solution be evaporated with a gentle heat, that crystals may be formed."

*Syn.* Sulfate de Potasse (*F.*), Schwefelsaures Kali Vitriolweinstein (*G.*), Zwavelzuures Kali (*Dutch*), Solfato di Potassa (*I.*).

In the Edinburgh process, which is the best of these three modes of preparing this salt, the carbonate of lime, which is added to the solution of the salt, combines with the superfluous sulphuric acid, while the carbonic acid is expelled; and the residue is thus converted into Sulphate of Potass. Both the London and the Dublin formulæ are objectionable on the score of expense; the value of the salt, as obtained by the London process, at the price of the pure salt made on the large scale, being to its cost very nearly as 5 to 10.<sup>1</sup> The greater part of the Sulphate of commerce is prepared from the residue of the distillation of nitrous acid from nitre and sulphate of iron. This is a mixture of Sulphate of Potass, and red oxide of iron, from which the Sulphate is easily separated by boiling water, while the oxide remains undissolved.<sup>2</sup>

*Qualities.* — Sulphate of Potass has a nauseous, bitterish taste. It is usually procured in small, grouped, transparent crystals, of which the form is a pyramidal dodecahedron, with isoceles triangular faces<sup>3</sup>; but this form is subject to various modifications, according to the mode of conducting the evaporation. Their specific gravity is 2.4073.<sup>4</sup> The crystals are not efflorescent; they decrepitate when heated; and are soluble in 16 parts of water at 60° and 5 parts of boiling water. This salt is partially decomposed by the nitric, muriatic and tartaric acids; and in solution is completely decomposed by muriate of barytes, muriate of lime, oxymuriate of mercury, nitrate of silver, and acetate and subacetate of lead, which therefore cannot enter into formulæ with it. Charcoal also decomposes it at a high temperature. Its constituents, according to the analysis of Mr. Phillips<sup>5</sup>, are 45.45 of acid, 54.55 of alkali; to that of Berzelius 47.1 of acid, and 52.9 of alkali; and to that of Berard 42.76 of acid, and 57.24 of alkali.<sup>6</sup>

*Medical properties and uses.* — This salt is deobstruent and cathartic. It is given with great advantage in the visceral obstructions to which children are liable; and in combination

<sup>1</sup> Vide *London Medical Review*, April 1810, p. 135.

<sup>2</sup> This oxide, when dried, is of a deep red colour, and is the *colcothar* of commerce.

<sup>3</sup> *London Medical Review*, April 1810.

<sup>4</sup> Hassenfratz. *Ann de Chimie*, xxviii. 12.

<sup>5</sup> *System of Chemistry*, 4th edit. ii. 660.

<sup>6</sup> *Annales de Chimie*, lxxi. 47.



with rhubarb or with aloes, I have found it more useful than any of the other saline purgatives in jaundice and dyspeptic affections. On account of its sparing solubility, it is generally given in the form of powder, in doses of from grains x. to ʒj., according as it is intended to act as a deobstruent or purgative.

Officinal preparation. *Pulvis Ipecacuanhæ compositus*, L. E. D.

POTASSÆ SUPERSULPHAS, Lond. *Supersulphate of Potass.*

“Take of the Salt which remains after the distillation of the Nitric Acid, *two pounds*; Boiling Water, *four pints*. Mix them, so that the salt may be dissolved, and filter. Then boil the solution till one half is dissipated, and set it aside to crystallize. Pour off the water and dry the crystals on bibulous paper.”

This Salt is the *Sal enixum* of commerce; a bisulphate of Potass. The solution should not be filtered until it be cold, as a copious deposition of uncrystallized salt takes place when it is filtered while hot.

The excess of Sulphuric Acid is so very loosely combined with Sulphate of Potass, that great part of it may be washed off by water; but nevertheless, the crystallized salt differs in several respects from the neutral sulphate. It may be prepared, by heating together three parts of Sulphate of Potass and one part of sulphuric acid.

*Qualities.* — Its crystals are long, slender, hexangular prisms, impressing a sour and slightly bitter taste; and containing from 12 to 13 per cent. of water. It reddens the vegetable blues; is soluble in two parts of water at 60°; in less than an equal weight of boiling water; and effervesces with the carbonates of alkalies. The proportions of its constituents are stated to be, of potass 32·87, acid 54·80, and water 12·33 in 100 parts. When exposed to a red heat, the water of crystallization and half the acid are expelled, and simple Sulphate of Potass remains.

*Medical use.* — As a remedy its efficacy is as yet unknown; but we are informed<sup>1</sup> that it has been introduced into the pharmacopœia from an idea that it will afford “a useful means of producing the effects of Sulphuric Acid combined with those of an opening salt; and that it may be exhibited at once in a solid form, an indication which is often desirable.” Dr. Paris<sup>2</sup> says, it forms a grateful adjunct to rhubarb. The dose is from gr. x. to ʒij. exhibited in combination with infusion of rue or of some bitter or rhubarb.

SULPHAS POTASSÆ CUM SULPHURE, Edin. *Sulphate of Potass with Sulphur.*

<sup>1</sup> *Powel's Translation of the London Pharmacopœia*, 2d ed. 73.

<sup>2</sup> *Pharmacologia*.



“Take of Nitrate of Potass in powder, and of Sublimed Sulphur, *equal weights*. Mix them well together, and throw the mixture in small quantities at a time into a red-hot crucible. The deflagration being finished, let the salt cool, and preserve it in a well-stopped glass vessel.”

In this process the Sulphur is oxidized, and converted partly into Sulphuric Acid, and partly into Sulphurous Acid, by uniting with the oxygen afforded by the decomposition of the Nitric Acid of the Nitrate, which is effected by the degree of heat employed. During the deflagration, however, a part of the acid is volatilized in the form of nitrous oxide, and consequently the oxygen evolved is not sufficient to acidify all the sulphur, and the unaltered portion remains united with a portion of potass. The sulphuric and sulphurous acids combine with the remainder of the potass; and thence the product is a mingled mass, consisting of Sulphate and Supersulphate of Potass and Sulphuret of Potass. It is the preparation which was originally known under the name of *salpolychrest*.

*Qualities.* — This salt has a sensibly acid taste, and reddens infusion of litmus. It is almost wholly dissolved in eight parts of water, at 60°; and by exposure to the air it is altogether converted into Sulphate of Potass. In general I have found that it emits no sulphureous odour on the addition of sulphuric acid, and is not precipitated by acids; but in other specimens prepared with equal care sulphur was thrown down by the muriatic acid.

*Medical properties and uses.* — The same as those of Sulphate of Potass; and consequently it is scarcely ever used.

AQUA ALCALINA OXYMURIATICA, Dub. *Oxy-muriatic Alkaline Water.*

“Take of Muriate of Soda dried, *two pounds*; Manganese in powder, *a pound*; Water, Sulphuric Acid, each *two pounds*. Mix together the Muriate of Soda and the Manganese, put them into a matrass, and add the water; then by means of a proper apparatus, add gradually, and at intervals, the sulphuric acid; and transmit the disengaged gas through a solution consisting of *four ounces* of Carbonate of Kali, and *twenty-nine ounces*, by measure, of Water. Towards the end of the operation apply a moderate heat to the matrass. The specific gravity of this solution is to that of distilled water as 1087 to 1000.”

For this process Woulfe's apparatus is necessary; two-thirds of the alkaline solution being put into the second bottle, and the remainder into the third. In this process, Chlorine is extricated from the decomposition of the dried Common Salt; which is a chloride of Sodium, and, combining with a



portion of the oxygen extricated from the Oxide of Manganese, is changed into Chloric Acid, which passes over in the gaseous form into the alkaline solution, where it unites with the Potass of the Carbonate, expelling the Carbonic Acid, and forming a Chlorate of Potass, the greater part of which remains dissolved in the water, while a small part is precipitated crystallized. But, along with the Chloric Acid, a portion of simple Chlorine also passes over; and, as this cannot unite with the alkali, it is absorbed by the solution, and remains loosely combined with it. Part of the oxygen also of the Oxide of Manganese unites with the Sodium, and forms Soda, which combines with a portion of the Sulphuric Acid; while the remainder of this acid attaches itself to the Manganese, forming a Sulphate of that metal. To prevent the inconvenience likely to result from the extrication of the Carbonic Acid Gas, a solution of lime should be put into the last bottle of the range of receivers. The product in the receivers is a *Solution* of hyperoxymuriate (*Chlorate*) of Potass, containing some uncombined Chlorine: what remains in the matrass is a mixed Sulphate of Soda and of Manganese. When the alkaline solution is stronger, in the proportion of  $\text{℥} \text{ xvj.}$  of the subcarbonate to  $\text{Oiv.}$  of water, a sparingly soluble, crystallized salt is procured, which is regarded as hyperoxymuriate (*Chlorate*) of Potass, and is a more certain preparation than the solution.

*Qualities.* — This solution has, in a slight degree, the odour of Oxymuriatic Acid (*Chlorine*), and a cooling taste. It destroys the vegetable colours, owing to the Chlorine it contains.

*Medical properties and uses.* — Hyperoxymuriate of Potass is stimulant and diuretic. The supposition which prevailed, that the effects of the nitric acid in syphilis arose from its imparting oxygen to the system, brought forward this preparation as a remedy in the same disease; and its use was soon extended to scurvy and other complaints which were supposed to depend on a deficiency of oxygen. But although it evidently has the power of checking for a time the action of the syphilitic virus, yet it has scarcely in any case succeeded in effecting a cure; and consequently has fallen into disrepute, and is now seldom employed in any case.

POTASSÆ TARTRAS, Lond. *Tartrate of Potass.*

“Take of Subcarbonate of Potass, *sixteen ounces*; Supertartrate of Potass, *three pounds*; Boiling water, *a gallon*. Dissolve the Subcarbonate of Potass in the water, and add the Supertartrate of Potass reduced to powder till the effervescence cease. Filter the solution through paper; then boil it until a pellicle appear on the surface, and set it aside to crystallize.



Having poured off the water from the crystals, dry them on bibulous paper."

TARTRAS POTASSÆ, Edin. *Tartrate of Potass.*

"Take of Subcarbonate of Potass, *one part*; Supertartrate of Potass, *three parts*, or a sufficient quantity; Boiling Water, *fifteen parts*. To the Subcarbonate dissolved in the water add in small portions the Supertartrate of Potass, reduced to a fine powder, as long as it excites effervescence, which gradually ceases before three times the weight of the Subcarbonate of Potass be added; filter the solution when it is cold, and after due evaporation, set it aside that crystals may form."

TARTRAS KALI, Dub. *Tartrate of Kali.*

"Take of Subcarbonate of Kali, *a pound*; Crystals of Tartar, reduced to a very fine powder, *two pounds and a half*, or a quantity sufficient to saturate the kali; Boiling Water, *a gallon*. To the Subcarbonate of Kali dissolved in the water gradually add the Tartar; filter the solution through paper, evaporate, and set it aside that crystals may form as it cools."

*Syn. Kali Tartarizatum*, P. L. 1787. *Tartrate de Potasse* (F.), *Weinstein-saures Kali* (G.), *Tartrato di Potassa* (I.).

In these processes the superabundant acid of the Bitartrate of Potass is saturated by the Potass of the Subcarbonate, the carbonic acid gas of which is expelled, and a neutral Tartrate obtained. The quantity of alkali required for this purpose must necessarily vary, owing to the degree of dryness of the subcarbonate employed. To obtain regular crystals, a very slow, nearly spontaneous, evaporation is necessary; and, therefore, this salt as found in the shops, and prepared on a large scale, is in the form of a white granular powder, which is produced by the evaporation being continued to dryness with frequent stirring.<sup>1</sup>

*Qualities.* — This salt has a bitterish, cool taste. Its crystals are right oblique angled prisms, and in this state it is soluble in its own weight of water at 60°; but in the granular form, four parts of cold water are required for its solution. It attracts water from the atmosphere. When long kept in solution, its acid is decomposed, and its alkali remains in the state of a subcarbonate<sup>2</sup>: and the same circumstances occur when it is exposed to a red heat. Alcohol also dissolves it readily. It is partially decomposed by the weaker acids and the acidulous salts, and also by tamarinds and other sub-acid fruits, which reduce it to the state of supertartrate; and it is completely decomposed by lime-water, muriate of barytes, magnesia,

<sup>1</sup> It was formerly named *Soluble Tartar*, *Sal vegetabile*, *Tartarus tarturizatus*.

<sup>2</sup> *Murray's Chemistry*, 2d ed. iv. 329.



nitrate of silver, and acetate and superacetate of lead. Its constituents, abstracting the water of crystallization, are, in 100 parts, 58.26 of acid, and 41.74 of alkali.<sup>1</sup>

*Medical properties and uses.*—Tartrate of Potass is a valuable purgative, operating easily and without griping; and even correcting the griping properties of some other substances, as of senna and the resinous purgatives, with which it is, therefore, usually combined. It enters into Klein's *Pulvis lenitivus hypochondriacus*, which consists of the following articles — R̄ Flavédinis Corticis Aurantiæ, Radicis Rhei, Potassæ Tartratis, ā ā ʒss, Olei Cajeputi gutt. iij M. ft. pulvis una pro dosi.<sup>2</sup> The dose is from ʒj. to ʒj. in solution.

**AQUA SUPERCARBONATIS POTASSÆ, Edin.**  
*Water of Supercarbonate of Potass.*

“Take of Water, *ten pounds*; Pure Subcarbonate of Potass, *one ounce*: dissolve, and expose the solution to a current of Carbonic Acid gas arising from Carbonate of Lime in powder, *three ounces*, Sulphuric Acid, *three ounces*, and Water, *three pounds*, gradually and cautiously mixed. The chemical apparatus of Dr. Nooth is well adapted for this preparation. But if a larger quantity of the solution be required, an apparatus which will admit of a sufficiently great pressure should be employed. The solution must be preserved in well-stopped vessels.”

This water is seldom sufficiently impregnated with the acid, when made on a small scale; but in the great way, and with an apparatus from which a much greater pressure is obtained, a solution is prepared for sale, which contains a very large quantity of uncombined carbonic acid.

*Qualities.*—When properly prepared, this Solution has a pungent, acidulous taste, and reddens tincture of litmus; is perfectly transparent, sparkles when poured into a glass, and effervesces violently with all the acids.

*Medical properties and uses.*—This solution is tonic, diuretic, and antacid. It has also been regarded as lithontriptic, and is much used in calculous cases: but its properties as a solvent of calculi, if it possess any, must depend upon the potass it contains, and, thence, the more completely it is impregnated with carbonic acid, the more its lithontriptic powers must be diminished. There is, however, reason for believing, that even pure potass when taken into the stomach exerts no influence on ready-formed calculi, and consequently its operation, even as a palliative or preventive, is confined to the

<sup>1</sup> Phillips' *Trans. of the Pharmacopæia*, p. 56.

<sup>2</sup> Quoted by Brande, *Manual of Pharm.* p. 252.



stomach, where it neutralizes the acid that always prevails there in calculous affections, and relieves many of the uneasy symptoms it occasions. In this view the solution of the Supercarbonate is a grateful mode of exhibiting Potass, as its acrimony is destroyed by its combination with the acid, which is nevertheless so weak as not to interfere with its operation as an alkali. On the same principles it proves beneficial in dyspepsia and gout, and forms with lemon juice an effervescing draught still preferable to that prepared with the Carbonate. The dose in calculous affections is  $\text{f}\overline{\text{3}}\text{viij.}$ , taken three or four times a day.

#### PREPARATIONS OF SODA.

**SODÆ CARBO'NAS**, Lond. *Carbonate of Soda.*

“Take of Subcarbonate of Soda, *a pound*; Distilled Water, *three pints*. Dissolve the Subcarbonate of Soda in the Distilled Water. Then transmit Carbonic Acid through the solution, by means of a proper apparatus, until it be saturated, and set it apart to crystallise. Dry the crystals involved and compressed in blotting-paper. Evaporate the remainder of the solution by a heat not exceeding  $120^{\circ}$ , that more crystals may be obtained. These are to be compressed and dried in the same manner as the former.”

**CARBONAS SODÆ**, Edin. *Carbonate of Soda.*

“Take of Subcarbonate of Soda, *two parts*; Water, *three parts*. Dissolve the salt in the water, and subject it to a stream of Carbonic Acid gas, until the acid be no longer absorbed. Then filter the fluid and evaporate it in a heat not exceeding  $180^{\circ}$ , that it may crystallise. The Carbonic Acid is easily obtained from equal weights of pulverised Carbonate of Lime and of Sulphuric Acid largely diluted with water.”

*Syn.* Carbonate de Soude (*F.*), Kohlensaures Natrum (*G.*), Koolenzuure Soda (*Dutch*), Carbonato di Soda (*I.*).

The constituents of this salt, which is a bicarbonate, are according to Rose, 49.05 of acid, 29.85 of alkali, and 20.20 of water, in 100 parts: but according to Mr. Phillips, 39.76 of Carbonic Acid, 38.55 of Soda, and 21.69 of Water. It is sometimes regarded as a sesquicarbonate, owing to its containing one atom and a half of Acid, and one atom of Soda.<sup>1</sup> This salt does not appear to possess any advantages over the subcarbonate as a remedy, but it is less nauseous. The dose is from gr. x. to 3ss.

**MURIAS SODÆ SICCATUM**, Dub. *Dried Muriate of Soda.*

“Take of Muriate of Soda, *any quantity*. Roast it over

<sup>1</sup> *Trans. of Pharm.* p. 63.



the fire in an iron vessel slightly covered, until it cease to decrepitate, occasionally agitating."

*Syn.* Muriate de Soude sec (*F.*), Getrocknetes salzaures Natrum (*G.*), Muriato di Soda secco (*I.*).

One hundred parts of crystallized Muriate of Soda contain, according to Kirwan, 8.12 of water, which is nearly dissipated by the heat; and the salt was thus supposed to be rendered of a more uniform strength: but in this state it contains no muriatic acid, being in fact a Chloride of Sodium. It is employed chiefly for the distillation of Muriatic Acid, which is obtained colourless from the dried salt.

PHOSPHAS SODÆ, Edin. *Phosphate of Soda.*

"Take of Bones, burnt to whiteness and reduced to powder, *ten pounds*; Sulphuric Acid, *six pounds*; Subcarbonate of Soda, a sufficient quantity. Mix the powdered Bones with the Sulphuric Acid in an earthen vessel; then add nine pounds of Water, and again mix: keep the vessel in a vapour bath for three days: after which, dilute the matter with nine pounds more of Boiling Water, and strain through a strong linen cloth, pouring Boiling Water gradually over it until the whole of the Phosphoric Acid be washed out. Set the strained liquor apart that the impurities may subside, from which pour it off, and evaporate it to nine pounds. To this liquor, separated from its impurities, and heated in an earthen vessel, add a warm solution of Subcarbonate of Soda, until the effervescence cease: then strain, and set the liquor aside, that crystals may form. These being removed, add to the liquor, if necessary, a little Subcarbonate of Soda, that the Phosphoric Acid may be accurately saturated; and dispose it by evaporation again, to yield crystals, as long as these shall be produced. Finally, let the crystals be preserved in a well-closed vessel."

*Dublin.*

"Take of Burnt Bones reduced to powder, *five pounds*; Sulphuric Acid, *three pounds and a half*. Mix the powder with the Sulphuric Acid in an earthen vessel; add, gradually, five pints of Water, and agitate the mixture. Digest for three days, adding from time to time more water, lest the materials should become dry, and continue the agitation; then pour over them five pints of boiling water, and strain through a linen rag, pouring on, at intervals, boiling water, until all the acid be washed out. Set the liquor apart that the impurities may subside, from which decant it, and evaporate it to one half; then add three pounds ten ounces of Carbonate of Soda (dissolved in a sufficient quantity of warm water): filter, and obtain crystals by repeated evaporation and cooling. The crystals are to be preserved in well-closed vessels.



“ If the salt be not sufficiently pure, repeat the solution and crystallization.”

*Syn.* Phosphate de Soude (*F.*), Phosphorsäures Natrum (*G.*), Fosfato di Soda (*I.*).

When Bones are burnt to whiteness, the residue is chiefly Phosphate of Lime, 100 parts of which consist of 41 parts of acid, and 59 of base <sup>1</sup>; and with a small portion of Carbonate of Lime. The addition of Sulphuric Acid, as directed in the above formulæ, abstracts 0·40 parts of the Lime, so as to form an insoluble Sulphate of Lime, and, involved in its mass, a soluble Superphosphate of Lime, for the separation of which, the digestion in vapour and the repeated effusions of Boiling Water are ordered. The Soda of the Subcarbonate of Soda, which is added to the defecated and filtered solution, now unites with the superabundant Phosphoric Acid, by which means the Lime is again left combined with as much of this acid only as renders it a neutral Phosphate, which from its insolubility precipitates and is easily separated from the Phosphate of Soda, which being soluble, remains dissolved in the Water, and crystallizes on the subsequent evaporation of the filtered liquor.

There are some niceties in the manipulation of this process that require to be noticed. In the first place, if too much Sulphuric Acid be employed, Sulphate of Soda will be also produced; and as four parts only of Sulphuric Acid are required to decompose ten parts of Phosphate of Lime, both the above formulæ err in this particular; secondly, as the Phosphate of Soda does not crystallize well without an excess of base, a little more Subcarbonate of Soda must be added than is required simply to neutralize the excess of acid of the Superphosphate: and lastly, the evaporation must not be carried quite to the formation of a pellicle, as in this case the crystallization is indeterminate, and the whole often concretes into an irregular mass. <sup>2</sup>

*Qualities.* — This Salt has a purely saline taste, resembling very much that of Common Salt. Its crystals are large, regular, transparent, rhomboidal prisms, terminated by three-sided pyramids, having a specific gravity of 1·333, and efflorescing on exposure to the air. It is soluble in three parts

---

<sup>1</sup> Vauquelin,

<sup>2</sup> A cheaper mode of preparing this salt has been given by M. Funcke, a German chemist. He adds to the matter of calcined bones diffused in water, just enough dilute sulphuric acid to saturate the small portion of carbonate of lime it always contains. When the effervescence ceases, the whole is dissolved in nitric acid, and as much sulphate of soda added to the solution as of bone ashes used. The whole is then distilled to recover the nitric acid; and the phosphate of soda is separated from the residue, which is a mixture of sulphate of lime and phosphate of soda, by solution and crystallization.



of water at 60°, and in two parts of boiling water; and undergoes the watery fusion when heated. Its constituents, according to Thenard, are, in 100 parts, 19 of soda, 15 of phosphoric acid, and 66 of water: and according to Berzelius 20·33 of acid; 17·67 of base, and 62·80 of water. Muriate of Barytes, Lime, and Magnesia, decompose this salt; and by the strong acids it is converted into Biphosphate of Soda.

*Medical properties and uses.* — It is a mild cathartic, excellently adapted for children, and others who have a fastidious taste. It may be given dissolved in gruel, or broth, made without salt, by which its taste is very effectually covered. The dose is from ʒvj. to ʒij. Although Phosphate of Soda was known before 1740, at which time it was described by Haupt under the name of *Sal Mirabile Perlatum*, yet, it was not introduced into medical use as a purgative, until about thirty years since, when it was recommended by Dr. George Pearson of London.

SO'DÆ SUBCARBO'NAS, Lond. *Subcarbonate of Soda.*<sup>1</sup>

“Take of Impure Soda (*barilla*) in powder, *a pound*; Boiling Distilled Water, *four pints*. Boil the Soda in the water for half an hour, and filter the solution. Evaporate it to two pints, and set it apart that crystals may form. Throw away the liquor that remains.”

SUBCARBONAS SODÆ, Edin. *Subcarbonate of Soda.*

“Take of Impure Carbonate of Soda, *any quantity*. Bruise it, and then boil it in water until all the saline matter be dissolved. Filter the solution through paper, and evaporate it in an iron vessel, so that after refrigeration crystals may form.”

*Dublin.*

“Take of Barilla in powder, *ten pounds*; Water *two gallons*. Boil the Barilla in the water, in a covered vessel, for two hours, occasionally stirring; filter the liquor; then bruise the Barilla that remains with an equal quantity of water, and again boil it: this may be repeated a third time. The leys being filtered and mixed, evaporate them to dryness in a wide iron vessel, taking care that the saline mass, which will remain, be not again liquefied by too great a heat: stir it with an iron spatula till it become white; finally dissolve it in boiling water, and, after due evaporation, set it apart, that, as it cools, crystals may form. These will be purer if the barilla before each boiling be exposed for some time to the

<sup>1</sup> Mr. Phillips properly remarks, “that this salt, although it is termed a subcarbonate in the Pharmacopœia, yet, is a carbonate,” as it consists of one atom of acid and one of base. Vide *Trans. of the Pharm.* 1824.



air. The crystallization should be effected when the air is at the freezing temperature, and in a liquor the specific gravity of which is, to that of water, as 1220 to 1000. If the salt be not very pure, repeat the solution and crystallization."

*Syn.* Natron Preparatum, P. L. 1787. Soucarbonate de Soude (*F.*), Sotto carbonato di Soda (*I.*).

Barilla, besides the Subcarbonate of Soda, contains Sulphate and Muriate of Soda, charcoal, lime, magnesia, argil, and silex, from which these processes are intended to separate it. The earths being insoluble, are separated by the solution and filtration; while the foreign salts remain dissolved in the residuary liquor after the subcarbonate of soda has crystallised. In the London formula, the quantity of water ordered to be employed is much greater than is requisite; and it has been justly observed<sup>1</sup>, that the evaporation is directed to be too soon stopped. One pound of barilla yields, when properly managed, from  $\bar{z}$  iij. to  $\bar{z}$  v. of the crystallized subcarbonate.

A pure Subcarbonate of Soda is now manufactured on a great scale, by the decomposition of Sulphate of Soda, and of Muriate of Soda, which will probably supersede altogether the processes ordered in the pharmacopœias.<sup>2</sup>

*Qualities.*—Subcarbonate of Soda has a mild alkalescent taste, and changes the vegetable blue and red colours to green. Its crystals are large, transparent octahedrons, truncated at the summits of the pyramids, which contain a large proportion of water, effloresce when exposed to a dry air, and crumble down into a white opaque powder. It undergoes the watery fusion at 150° Fah.; is soluble in two parts of water at 60°, and in considerably less than its weight of boiling water, its abundant water of crystallization assisting the solution of the salt at that temperature. Its constituents, according to the late analysis of D'Arcet<sup>3</sup>, are, in 100 parts, 16·04 of acid, 20·85 of alkali, and 63·61 of water: Mr. Phillips makes them 14·38 of carbonic acid, 20·92 of soda, and 64·70 of water. By treating this salt in the method described under Subcarbonate of Potass, any muriates or sulphates it may contain are detected, while the tartaric acid, added to its solution, discovers potass, by forming a precipitate of the supertartrate.

*Medical properties and uses.*—This salt is antacid and deobstruent. It is less acrid than the Subcarbonate of Potass; and thence is in more general use in dyspepsia and acidities of the stomach, and in scrophulous affections. Its use has been

---

<sup>1</sup> *London Medical Review*, April 1808. 139.

<sup>2</sup> For experimental purposes, it is obtained, still purer from the crystallized acetate by ignition.

<sup>3</sup> *Annales de Chimie*, lxxi. 208.



lately strenuously recommended in whooping-cough, the protraction of which it is said to prevent. It is given, after the stomach and bowels have been duly evacuated, at first in combination with ipecacuanha and opium, and afterwards, when the violence of the cough has abated, with myrrh or cinchona.<sup>1</sup> The dose of this salt is from grs. x. to ʒj., given twice or thrice a day. M. Peschier has, lately, successfully employed it for the cure of bronchocele, and considers it much more efficacious than Iodine in this disease.<sup>2</sup> In bronchocele Dr. Peschier gave it to the amount of 118 grains in a day, dissolved in water.

Officinal preparations. *Sodæ Subcarbonas exsiccata*, L. D. *Sodæ Carbonas*, L. *Aqua Supercarbonatis Sodæ*, E. *Phosphas Sodæ*, E. D. *Soda tartarizata*, L. E. D.

**SODÆ SUBCARBONAS EXSICCATA**, Lond. *Dried Subcarbonate of Soda.*

“Take of Subcarbonate of Soda, *a pound*. Expose the Subcarbonate of Soda to a boiling heat in a clean, iron vessel, until it become perfectly dry, and at the same time stir it diligently with an iron spatula. Finally, rub it into a powder.”

**CARBONAS SODÆ SICCATUM**, Dub. *Dried Carbonate of Soda.*

“Liquefy the crystals of Carbonate of Soda in a silver crucible over the fire; then, in an augmented heat, stir the dissolved salt until by the evaporation of the water it become dry. Reduce it to a fine powder, and preserve it in stopped phials.”

*Syn.* Soucarbonate de Soude sec (*F.*), Getrocknetes Kohlensaures Natrum (*G.*), Sotocarbonato di Soda secco (*I.*), Unvollmaakte Koolenstoffzuure Soda (*Dutch*).

Owing to the great proportion of water of crystallization which the Subcarbonate of Soda contains, it readily undergoes the watery fusion, and is completely dried by continuing the heat; but its properties are not otherwise altered. The constituents of 100 parts, in this state, according to the analysis of Kirwan, are 40.14 of acid, and 59.86 of soda<sup>3</sup>, which nearly accords with the analysis of Dulong and Dalton.

*Medical properties and uses.*—The chief advantage obtained from drying the Subcarbonate of Soda is the facility of exhibiting it in the form of pills; for when the crystallized salt is used for this purpose, the pill formed with it falls to pieces as soon as the salt effloresces. Dr. Beddoes<sup>4</sup> has extolled it,

<sup>1</sup> *Medico Chirurgical Transactions*, vol. i.

<sup>2</sup> *Bib. Univer.* xxvii. p. 146.

<sup>3</sup> *Nicholson's Journal*, 4to. iii. 215.

<sup>4</sup> *Beddoes on the Nature and Cure of Calculus.*



in this form, as a remedy in calculous affections: and it certainly affords relief from the painful symptoms attending calculus in the kidneys, and other urinary affections: but its effects are palliative only; and depend on its destroying the prevalent acid in the stomach. The dose is from grs. x. to grs. xv., given three times a day. Beddoes directed it to be combined with soap and aromatics.

SO'DÆ SUL'PHAS, Lond. *Sulphate of Soda.*

"Take of the Salt which remains after the distillation of Muriatic Acid, *two pounds*; Boiling Water, *two pints and a half*. Dissolve the Salt in the Water: then add gradually as much Subcarbonate of Soda as will saturate the Acid. Boil the solution until a pellicle appear, and after having filtered it set it apart to crystallize. Pour the water from off the crystals, and dry them on bibulous paper."

SULPHAS SODÆ, Edin. *Sulphate of Soda.*

"Dissolve in Water the Acidulous Salt which remains after the distillation of Muriatic Acid, and having mixed with it Carbonate of Lime (*chalk*) in powder to remove the superfluous acid, set it apart until the impurities subside; then, having poured off the liquor, filter it through paper, and reduce it by evaporation, that crystals may be formed."

*Dublin.*

"Dissolve the Salt which remains after the distillation of Muriatic Acid in a sufficient quantity of Boiling Water. Evaporate the filtered solution to a proper point, and set it apart, that, as it slowly cools, crystals may form."

*Syn.* Natron Vitriolatum, P. L. 1787. Sulphate de Soude (*F.*), Krystalisirtes Natrum (*G.*), Solfato di Soda (*I.*).

The theory of the London process for preparing this Salt is analogous to that of the process for preparing the sulphate of potass; but, from the low price of the salt, manufactured on a great scale (see *Part ii.*) it is preferable to saturate the superabundant acid with chalk, and reject the sulphate of lime. The Salt obtained by the Dublin process has a slight acidulous taste, and contains a quantity of sulphuric acid, but so loosely combined as scarcely to entitle it to be regarded as a supersalt; nor do its crystals differ in form from those produced by the other two formulæ.<sup>1</sup>

*Qualities.* — The taste of this salt is at first simply saline, but afterwards very disagreeably bitter. Its crystals are transparent, six-sided, irregular, channelled prisms, with di-

---

<sup>1</sup> The crystals of the supersulphate formed by dissolving the sulphate in sulphuric acid, and crystallizing, are rhomboidal. The artificial salts, sold under the name of *Cheltenham Salts*, owe their activity to this salt. For an account of them, see Dr. Paris's *Pharmacologia*; art. *Sulphas sodæ*.



hedral summits; efflorescent and rapidly falling to an opaque white powder, when exposed to the air. It is soluble in 2·86 parts of water at 60°, and 0·8 of boiling water; undergoes the watery fusion when heated<sup>1</sup>, and in a strong heat is partially decomposed. According to Berzelius, 100 parts contain 24·64 of acid, 19·36 of alkali, and 56 of water; and in the dried state, according to Bulcholz, 54 acid, and 46 of alkali<sup>2</sup>: but according to Mr. Phillips', 55·55 of acid, and 44·45 of Soda.<sup>3</sup>

*Medical properties and uses.*— Sulphate of soda is a very common and useful purgative; but from its nauseous taste it is not very generally prescribed by the physician, although this may be readily disguised by a small quantity of lemon juice, or of cream of tartar, added to the solution. It is incompatible in prescriptions with subcarbonate of potass, muriate of lime, nitrate of silver, and acetate of lead. The dose is from  $\bar{3}$  ss. to  $\bar{3}$  ij., but, in the effloresced state, half of these quantities is sufficient.

AQUA SUPER-CARBONATIS SODÆ, Edin. *Water of Supercarbonate of Soda.*

“ Take of Water *ten pounds*; Subcarbonate of Soda, *two ounces*; dissolve and subject the solution to a stream of Carbonic Acid gas procured from *three ounces* of Carbonate of Lime, and the same quantity of Sulphuric Acid, with three pounds of Water, gradually and cautiously mixed together. It may be conveniently prepared in Nooth's apparatus. But if a large quantity of it be required, an apparatus capable of affording a greater pressure will be requisite. The fluid must be preserved in well-corked bottles.”

This preparation is milder and pleasanter than the water prepared with Subcarbonate of Potass. It is manufactured in large quantities on a great scale, of a much superior quality to any which the apothecary can prepare; and is in very general use as a cooling beverage. Half a pint of it poured over two table-spoonsful of lemon-juice, sweetened with a little sugar, forms an excellent and very agreeable effervescing draught; and the same quantity poured upon two ounces of boiling milk forms an excellent substitute for asses' milk.<sup>4</sup>

<sup>1</sup> Mr. Faraday found that, when it is heated in a flask, one part dissolves in the water of crystallization, while the other is thrown down in an anhydrous state: and that at 180°  $\frac{5}{9}$  of the salt take all the water, whilst  $\frac{4}{9}$  separate in the dry state. *Journ. of Science*, vol. xix. p. 153.

<sup>2</sup> *Nicholson's Journal*, 4to. iii. 215.

<sup>3</sup> *Trans. of Pharm.* p. 65.

<sup>4</sup> What are termed *Sodaic Powders*, are attempted to be passed upon the public as capable of answering in every respect the purpose of *soda water*; but the salt formed by the solution of these powders is a tartrate of soda, not a carbonate.



SODA TARTARIZATA, Lond. *Tartarized Soda.*

“Take of Subcarbonate of Soda, *twenty ounces*; Supertartrate of Potass in powder, *two pounds*; Boiling Water, *ten pints*. Dissolve the Subcarbonate of Soda in the Water, and add gradually the Supertartrate of Potass. Filter the solution through paper; then boil till a pellicle forms on the surface, and set it aside to crystallize. Pour off the water from the crystals, and dry them on bibulous paper.”

TARTRAS SODÆ ET POTASSÆ, Edin. *Tartrate of Soda and Potass.*

“Take of Subcarbonate of Soda, *one part*; Supertartrate of Potass, *three parts*, or a *sufficient quantity*; Boiling Water, *fifteen parts*. To the Subcarbonate dissolved in the water, gradually add the Supertartrate rubbed to a fine powder, as long as effervescence is excited, which generally occurs before three times the weight of the Subcarbonate is added: when the fluid is cold, filter it through paper, and after a proper degree of evaporation, set it aside that crystals may form.”

TARTRAS SODÆ ET KALI, Dub. *Tartrate of Soda and Kali.*

“Take of Carbonate of Soda, *twenty ounces*; Crystals of Tartar, reduced to a very fine powder, *two pounds*; Boiling Water, *ten pints*. Dissolve the Carbonate of Soda in the Water, and gradually add the Tartar; filter the solution through paper; evaporate it, and set it aside, that, as it slowly cools, crystals may form.”

*Syn.* Tartrate de Soude et de Potasse (*F.*), Natrum-weinstein (*G.*), Tartrato di Potassa e di Soda (*I.*).

In these processes the superabundant acid of the Supertartrate is saturated by the Soda of the Subcarbonate, the Carbonic Acid of which is dissipated in the gaseous form; and a triple salt is obtained by the evaporation, instead of two distinct salts being formed from the different alkaline bases.

*Qualities.* — This salt has a bitter saline taste. Its crystals are large, regular, transparent, hard, rhomboidal, six-sided prisms; very slightly efflorescent, and soluble in five parts of water at 60°. It is decomposed by the strong acids, muriate of barytes, lime, and by a red heat. The constituents of 100 parts of this salt, according to Schulze, are 41.3 of tartaric acid, 14.3 of potass, 13.3 of soda, and 31.1 of water<sup>1</sup>: according to Vauquelin, they are, tartrate of potass, 54 parts; tartrate of soda, 46 parts; and according to Mr. Phillips, (*Translation of the Pharmacopœia*,) tartaric acid 62.60, potass

---

The powders are packed in two distinct papers, the one blue and the other white; the blue containing ʒss. of carbonate of soda, the white gr. xxv. of tartaric acid.

<sup>1</sup> *Gehlen Journ.* iv. 210. *Thomson's Chemistry*, 4th edit. iii. 96.



22.44, soda 14.96; or tartrate of potass 53.74, and tartrate of soda 46.26.

*Medical properties and uses.*—Tartrate of Potass and Soda is a cooling and not very unpalatable cathartic. It was introduced into practice by M. Seignette<sup>1</sup>, an apothecary of Rochelle, and the preparation kept a secret until it was discovered and published by Boulduc and Geoffrey, in 1731. It operates moderately, and without exciting much irritation; hence it is well suited to nephritic and puerperal cases. The dose is from ℥j. to ℥jss., dissolved in any convenient vehicle.

---

## EARTHS AND EARTHY SALTS.

EARTHS possess peculiar properties, which distinguish them from other bodies, and constitute them a distinct class of natural productions. They are opaque, solid, unflammable, of very difficult fusibility, very sparingly soluble in water, and of a specific gravity not exceeding 4.9. Some of them resemble the alkalies in several particulars; are caustic, change to green the vegetable blues and reds, and neutralize acids; and have, therefore, been denominated ALKALINE EARTHS; but, nevertheless, are in truth metallic oxides. Of this division, three are medicinally employed; namely, LIME, MAGNESIA, and BARYTES; but the two former only are used as remedies in their pure state. Those earths which do not possess alkaline properties, although they, also, are metallic oxides, are denominated PROPER EARTHS; of which one only, ALUMINA, is a medicinal agent; and it is not used in its uncombined state.

Although some of the old chemists conjectured that the pure earths were metallic oxides, yet no direct proofs in support of the supposition were obtained, and they were generally supposed to be simple bodies, until the discoveries of Sir H. Davy revived the idea of their metallic nature.

The action of the Pure Earths on the animal economy is very similar to that of the alkalies.

The EARTHY SALTS<sup>2</sup> are compounds of the Acids with the Pure Earths, resembling the salts formed by the combination of Acids with Alkalies. Some of them are crystallizable, and soluble in water; others are nearly insoluble: some of them exert scarcely any action on the animal economy; while

---

<sup>1</sup> Hence its appellations of *Sal de Seignette*, *Sal Ruppellensis*.

<sup>2</sup> We prefer the title *Earths and Earthy Salts* to that of *Earths and their Salts*, which is the title of this section in the London Pharmacopœia, for the same reason that we preferred the term *Neutral Salts* to that of *Alkaline Salts*.



others are possessed of great activity, and produce very striking effects.

In extemporaneous prescription, it is absolutely necessary to avoid combining the earths or earthy salts with substances with which they form insoluble compounds.

The following Table shows the solubility of the above earths, and of the compounds which they form with acids.

PURE EARTHS.	Solubility in one part of Water.	ACIDS which, in combination with these Earths, form		
		Soluble Compounds.	Compounds scarcely soluble.	Insoluble Compounds.
Lime ....	0·002	Nitric Muriatic Acetic Benzoic Cinchonic Malic Gallic	Sulphuric Boracic Succinic Tartaric	Oxalic Phosphoric Carbonic Arsenic
Magnesia	0·000	Sulphuric Phosphoric Nitric Muriatic Carbonic Acetic Benzoic Succinic Malic Gallic	Tartaric Arsenic?	Boracic Oxalic?
Barytes .	0·050	Nitric Muriatic Acetic Benzoic Tartaric	Succinic Carbonic Citric	Sulphuric Oxalic Phosphoric Boracic Arsenic
Alumina	0·000	Sulphuric Nitric Muriatic Carbonic Acetic Benzoic Oxalic Tartaric Gallic	Boracic	Phosphoric Arsenic



## PREPARATIONS OF ALUMINA.

ALUMEN EXSICCATUM, Lond. *Dried Alum.*

“ Melt Alum in an earthen vessel over the fire, and increase the heat until the ebullition cease.”

*Edinburgh.*

“ Melt the Alum in an earthen or iron vessel, and let it be kept over the fire until it cease to boil; then rub it into a powder.”

ALUMEN USTUM, Dub. *Burnt Alum.*

“ Take of Alum *any quantity*. Expose to the heat of a strong fire in an earthen vessel until it cease to boil.”

*Syn.* Sulphate d'Alumine Séc (*F.*), Gebrannter Alaun (*G.*), Alume calcinato (*I.*).

In these processes the Alum loses its water of crystallization; but if the heat be too great, its acid is partly expelled, and partially decomposed. According to Kirwan, alum desiccated at  $700^{\circ}$  loses more than half its acid. By our experiments, English alum lost 0.43 in a moderate heat, and 0.46 in a red: Levant alum, 0.41 in a moderate heat, and 0.44 in a red heat. Chaptal found that in a red heat alum of his own manufacture lost 0.67; Roman alum, 0.50; English, 0.47; and Levant alum, 0.40.

*Qualities.* — Dried alum has a more astringent taste than the crystallized salt. It is obtained in the form of a light, opaque, white, spongy, friable mass, 100 parts of which consist of 36.25 acid, and 63.75 bases; when it has been exposed to a heat of  $700^{\circ}$ : but in general, the desiccation being conducted at a lower temperature the proportion of acid is greater.

*Medical properties and uses.* — It is chiefly used as an escharotic to destroy fungus in ulcers; but has also been given internally to the extent of ℥j. for a dose in cases of cholic, the pain of which it is said to allay, while at the same time it gently opens the bowels.

LIQUOR ALUMINIS COMPOSITUS, Lond. *Compound Solution of Alum.*

“ Take of Alum and Sulphate of Zinc, each *half an ounce*; Boiling Water, *two pints*. Dissolve the Alum and the Sulphate of Zinc in the Water, and then filter through paper.”

This powerful astringent solution has been successfully used in glects and leucorrhœa; and in some cases of ophthalmia. It requires to be diluted in all these cases, and to be employed with caution. Half an ounce of the solution and six and a half ounces of rose water form an excellent collyrium in ophthalmia, after local bleeding.



## PREPARATIONS OF LIME.

CALX, Lond. *Lime.*

“Take of White Marble, *a pound*. Break it into small pieces, and expose these in a crucible to a very strong fire for an hour, or until the Carbonic Acid be so completely expelled, that on the addition of Acetic Acid no air-bubbles are extricated.”

CALX E TESTIS, Lond. *Lime from Shells.*

“In the same manner lime is also prepared from shells.”

*Syn.* Chaux vine (*F.*), Kalk (*G.*), Brand ochoslächt kalk (*Swed.*), Calce (*I.*), Calviva (*S., Port.*).

Lime prepared on the great scale for the ordinary purposes of art, is sufficiently pure for medicinal use; but for some pharmaceutical purposes it is required to be more completely burnt than is usually the case of that which is obtained from the kilns; and perhaps it is with this view that the above preparations have been ordered by the London College. It may, however, be observed, that neither of the substances ordered affords Lime in a state of absolute purity; limestone frequently containing silex, alumina, magnesia, and marine shells; and a portion of phosphate of lime which is not decomposed by the fire. To obtain perfectly Pure Lime, dissolve white marble or clean oyster-shells in diluted muriatic acid, and to the filtered solution add solution of ammonia as long as any precipitate falls; then filter again, and decompose the muriate by a solution of pure carbonate of potass; wash the precipitate, and expose it to violent heat in a platina crucible, till it cease to lose weight. The result is pure lime, fifty-six parts of which should be furnished for every 100 parts of pure white marble used.

*Qualities.* — Well-prepared lime is of a white colour, moderately hard, sonorous and brittle. Its specific gravity is 2.3. It is inodorous; has a hot, pungent, bitter taste; on animal matter it operates as a most powerful caustic; changes the vegetable blues to green; and is infusible. Water poured on it is absorbed with a hissing noise, much heat is evolved, and the lime swells, falls to pieces, and is then said to be slaked; in which state, having combined with a portion of the water, it is converted to a *hydrate*, which readily combines with sulphur, forming a sulphuret, and is to a certain degree soluble in water. By exposure to the air it attracts carbonic acid, and again returns to the state of a carbonate or lime-stone. It appears to be an oxide of a peculiar metal which has been named *calcium*<sup>1</sup>, in the proportions of 71.42 of calcium, and

<sup>1</sup> This metal has the colour and appearance of silver, is solid, four times heavier than water, absorbs oxygen, and burns brilliantly in the open air, and by being oxidized is converted into quicklime. *Phil. Trans.* 1808.



28.58 of oxygen, in 100 parts : but slacked lime, or the hydrate consists of 75.68 of lime, and 24.32 of water.<sup>1</sup>

*Use.* — Lime in this state is chiefly employed for pharmaceutical purposes and for forming the solution.

Officinal preparations. *Liquor Calcis*, L. E. D. *Potassa cum Calce*, L. *Calcis Murias*, L. *Liq. Muriatis Calcis*, L. E. D.

LIQUOR CALCIS, Lond. *Lime-water*.

“Take of Lime, *half a pound*; Boiling Distilled Water, *twelve pints*. Pour the Water upon the Lime, and agitate them together; cover the vessel directly, and set it apart for three hours; then preserve the solution upon the undissolved Lime, in well-stopped glass bottles, and pour off the clear fluid when it is wanted for use.”

SOLUTIO CALCIS, sive AQUA CALCIS, Edin. *Solution of Lime, or Lime Water*.

“Take of Lime, fresh burnt, *half a pound*. Put it into an earthen vessel, and sprinkle upon it four ounces of Water, keeping the vessel covered until the Lime become hot, and fall into powder; then pour on it twelve pounds of Water, and mix the Lime with Water by agitation. After the Lime shall have subsided, repeat the agitation; and let this be done about ten times, the vessel being kept shut that the free access of the air may be prevented. Finally, let the Water be filtered through paper, interposing glass rods between the paper and the funnel, that the Water may pass through as quickly as possible. It is to be preserved in very well-stopped bottles.”

*Dublin.*

“Take of fresh burnt Lime *a pound*; Boiling Water, *a pint*. Put the Lime into an earthen vessel, and sprinkle the Water upon it, keeping the vessel shut until it become hot and fall into powder; then pour upon it *three gallons of Water*. The vessel being again shut, let the mixture be frequently shaken for twenty-four hours; and then filter the solution through paper placed in a covered funnel, and preserve it in well-stopped bottles.”

*Syn.* Eau de Chaux (*F.*), Kalkwasser (*G.*), Kalkwater (*Dutch*), Aqua di Calce (*I.*).

Of these formulæ, that of the London College is to be preferred: as by keeping the solution upon the Lime it is always in a completely saturated state, and the supernatant fluid is generally sufficiently clear to allow it to be decanted off without filtration. It is however adviseable, in making the solution, first to slack the Lime with a small portion of Water, before the whole quantity be added; as by this it is prevented from running into a paste, which confines the action of the Water. The direction of the Dublin College to use the water for slacking the Lime in a boiling state, is superfluous. Cold

<sup>1</sup> *Phillips's Trans. of Pharm.* p. 71.



water acts more powerfully on Lime than hot water. From Mr. Phillips's experiments, it appears that 10,000 grains of water at  $212^{\circ}$  dissolve 5·6 grains of lime, the same quantity of water at  $60^{\circ}$  dissolve 9·7 grains, and at  $32^{\circ}$  the quantity is increased to 11 grains.<sup>1</sup> By heating cold, saturated lime-water, a crystalline deposition of hydrate of lime was thrown down.

*Qualities.* — Lime-water is inodorous; has a strong, styptic, acrid taste; is limpid and colourless; and changes to green, the vegetable blue and red colours. It unites with oil, forming an imperfect soap. When exposed to the air, it attracts carbonic acid, which, combining with part of the lime held in solution, forms on its surface a pellicle of carbonate of lime, which thickens, cracks, and sinks to the bottom of the vessel, leaving its place to be supplied by another pellicle; and thus, by successive formations, the whole of the lime is abstracted from the water. Hence the necessity of preserving the solution in well-closed bottles; and, when it is filtered, in small bottles, containing such a quantity only as can be used at once. It is decomposed by the acids and sulphur, the alkaline carbonates, phosphates, borates, tartrates, and citrates; the infusions of orange-peel, columba, cinchona, rhubarb, and senna, which are consequently incompatible in formulæ with it.

*Medical properties and uses.* — Lime-water is tonic, antacid, anthelmintic, and externally detergent. It proves very useful in dyspepsia attended with much acidity of the stomach, by neutralizing the acid, and dissolving the sordid mucus with which it is often loaded in this disease; and has also been found efficacious in diarrhœa, diabetes, and leucorrhœa. It destroys intestinal worms, and dissolves the mucus which forms their nidus; and, for the same reason, proves serviceable in slimy bowels. Its internal use, however, should be occasionally suspended for a few days, as its long continued action on the stomach is apt to prove hurtful. Externally it is applied as a lotion to foul and cancerous ulcers, tinea capitis, and scabies, but with little advantage.

The dose is from fʒij to ʒss, alone, or diluted with milk.

Officinal preparations. *Oleum Lini cum Calce*, E. D. *Aqua Calcis Composita*, D.

MURIAS CALCIS, Lond. *Muriate of Lime*.

“Take of the Salt which remains after the sublimation of the Subcarbonate of Ammonia, *two pounds*; Water, *a pint*. Mix and filter through paper, then evaporate the solution until the salt become dry. Preserve the salt in a well-stopped bottle.”

This preparation is a Chloride of Calcium, and consists of Chlorine 64·29, and Calcium 35·71, in 100 parts.<sup>2</sup> Muriate

<sup>1</sup> *Translation of the Pharmacopœia*, p. 72.

<sup>2</sup> *Ib.* p. 74.



of Lime can exist only in a state of solution in water; and in evaporating such to dryness, the Muriatic Acid is decomposed and resolved into chlorine and hydrogen, while at the same time the Oxide of Calcium, forming part of the muriate, parts with its oxygen, the Chlorine attaches itself to the Calcium, forming a Chloride of Calcium, which is obtained as a dry salt, and the oxygen to the hydrogen forming water, which is evaporated.

*Qualities.* — Chloride of Calcium, or Muriate of Lime, as it is termed, is an inodorous, white, fusible, deliquescent, bitter, pungent salt. It is very soluble in both hot and cold water, and is changed into a muriate during its solution. It is very soluble also in alcohol. During its solution much cold is produced; and, if snow be used, a freezing mixture, capable of lowering the thermometer from  $32^{\circ}$  to  $50^{\circ}$ , is obtained.

*Medical properties and uses.* — It is sometimes, but rarely, administered in scrophulous affections, in the form of pill combined with Conium; but is chiefly used for preparing the *Liquor Muriatis Calcis*.

The dose, in the solid form, is from gr. i. to grs. iv. combined with a bitter or sedative extract.

Officinal preparations. *Liquor Muriatis Calcis*, L. E. D.

**LIQUOR MURIATIS CALCIS**, Lond. *Solution of Muriate of Lime.*

“Take of Muriate of Lime, *two ounces*; Distilled Water, *three fluid ounces*. Dissolve the Muriate of Lime in the Water, and filter the solution through paper.”

**SOLUTIO MURIATIS CALCIS**, Edin. *Solution of Muriate of Lime.*

“Take of the harder variety of Carbonate of Lime (namely White Marble) broken into small pieces, *nine ounces*; Muriatic Acid, *sixteen ounces*; Water, *eight ounces*. Mix the Acid with the Water, and gradually add the pieces of Carbonate of Lime. The effervescence being finished, digest for an hour. Pour off the fluid, and reduce it by evaporation to dryness. Dissolve the residue in its weight and a half of water, and filter the solution.”

**AQUA MURIATIS CALCIS**, Dub. *Water of Muriate of Lime.*

“Take of Chalk reduced to a coarse powder, *one ounce*; diluted Muriatic Acid, *two ounces*. Add gradually the Acid to the Chalk, and when the effervescence is finished, filter the solution.”

*Syn.* Dissolution de Muriate de Chaux (*F.*), Liquore di Muriato di Calce (*I.*).

In the processes of the Edinburgh and the Dublin Colleges, the Muriatic Acid unites with the Lime of the Carbonate, and disengages the Carbonic Acid, which is dissipated in the gaseous form, while the Muriate of Lime remains dissolved in the



water. The evaporation is unnecessary, if an acid of a determinate specific gravity be employed, as ordered by the Dublin College. According to the London process, the Chloride of Calcium, when dissolved in the water, decomposes that fluid, and forms a Muriate of Lime, which is dissolved in the remainder of the water. This preparation is therefore properly named, although the Muriate of Lime of the London College be a Chloride of Calcium.

*Qualities.* — This solution is colourless, and has a disagreeable, bitter, acrid taste. It is decomposed by the sulphuric, nitric, phosphoric, fluoric, and boracic acids; the neutral salts into which these enter; and the alkalies and alkaline carbonates, which precipitate the lime.

*Medical properties and uses.* — Muriate of lime is deobstruent and tonic. It was introduced into practice by Fourcroy, and has been much recommended as a remedy in scrophulous and glandular diseases. I have given it with evident advantage in bronchocele; and have witnessed more benefit result from its continued use in the varied forms of scrophula, than from any other remedy. Its operation is similar to that of Muriate of Barytes; but the danger of an over-dose is less to be dreaded, and its good effects are more uniform and certain. The dose of the solution is from  $\text{mxx.}$  to  $\text{f3j.}$ , increased gradually to  $\text{f3iv.}$ , in a sufficient quantity of water or milk, repeated twice or thrice a day.

CRETA PRÆPARATA, Lond. *Prepared Chalk.*

“Take of Chalk, *a pound.* Add a little Water to the Chalk, and triturate it to a fine powder. Throw this into a large vessel of water, stir it, and after a short interval pour off the supernatant turbid water into another vessel, and set it apart that the powder may subside: lastly, let the Water be poured off, and dry the powder.”

CARBONAS CALCIS PRÆPARATUS, Edin. *Prepared Carbonate of Lime.*

“Let Carbonate of Lime, triturated to powder in an iron mortar, and levigated with a little water, on a porphyry stone, be put into a large vessel: then pour Water upon it, which, after frequently shaking the vessel, is to be poured off loaded with the fine powder. The subtile powder which subsides, when the Water remains at rest, is to be dried. Let the coarse powder which the Water could not suspend be again levigated, and treated in the same manner.

CRETA PRÆPARATA, Dub. *Prepared Chalk.*

“Let it be triturated to powder in an earthen mortar, with the addition of a little Water; then mix this with a sufficiently large quantity of Water by agitation, and after a short interval, when the coarser particles have subsided, pour off the fluid.



This may be frequently repeated, always previously tritulating; and finally, collect the very fine powder, which after some time will subside, and dry it upon an absorbent stone, or paper."

*Syn.* Craie préparé (*F.*), Rein Kreide (*G.*), Carbonato di Calce preparato (*I.*).

By the suspension of the finer particles of the levigated Chalk in water, they are reduced to a more impalpable form, and are more effectually separated from the coarser particles than could be accomplished by any other mechanical means; but the Chalk is not freed from the foreign earths it generally contains, (see *Calx*, Part ii.) although it be sufficiently pure for medicinal use.

*Qualities.* — Pure Carbonate of Lime, in the form of Chalk, is a white, opaque, soft, light substance. It is a compound of 44 parts of Carbonic Acid, and 56 of Lime, in 100 parts.

*Medical properties and uses.* — Chalk is antacid and absorbent. It is exhibited advantageously in acidities of the primæ viæ; and in diarrhœas, after all irritating matters have been removed from the bowels by previous evacuation, (see *Mistura Cretæ*.) As an external application, it is sprinkled over ulcers discharging a thin ichorous matter, which is thus absorbed by the Chalk, and prevented from excoriating the neighbouring sound skin. In cases of burns it is applied in a similar manner, and a poultice laid over it, by which the skinning of the sore is much hastened.<sup>1</sup>

The dose of the chalk is from grs. x. to ℥ij., or more.

Officinal preparations. *Mistura Cretæ*, L. E. *Hydrargyrus cum Creta*, L. *Pulvis Cretæ compositus*, L. E. *Pulvis opiatus*, E. *Trochisci Carbonatis Calcis*, E. *Confectio aromatica*, L. E.

CRETA PRECIPITATA, Dub. *Precipitated Chalk.*

"Take of solution of Muriate of Lime, *any quantity*. Add to it as much (*sub*) Carbonate of Soda, dissolved in four times its weight of hot Distilled Water, as may be sufficient to precipitate the Chalk. Wash the precipitate three times in a sufficient quantity of water; then collect it, and dry it on a chalk-stone or on bibulous paper."

A double exchange takes place in this process; the Muriatic Acid separates from the Lime and unites with the Soda, while the Carbonic Acid of the Subcarbonate combines with the Lime: the Muriate of Soda thus formed remains dissolved in the water, but the Carbonate of Lime is precipitated in the form of a white powder. It is an expensive preparation, and the benefit to be derived from a great degree of purity in this substance is not very obvious.

Officinal preparations. *Hydrargyrum cum Creta*, D. *Electuarium aromaticum*, D. *Mistura Cretæ*, D.

<sup>1</sup> *Kentish on Burns*, passim.



MURIAS BARYTÆ, Edin. *Muriate of Barytes.*

“ Take of Carbonate of Barytes, Muriatic Acid, each, *one part* ; Water, *three parts*. To the Water and the Acid, mixed together, add the Carbonate broken into small pieces. The effervescence being finished, digest for an hour ; then filter, and after due evaporation set the solution apart that crystals may form. Repeat the evaporation as long as any crystals are formed.

“ If the Carbonate of Barytes cannot be procured, the Muriate may be prepared from Sulphate of Barytes in the following manner :

“ Take of Sulphate of Barytes, *two pounds* ; Charcoal in powder, *four ounces* ; Muriatic Acid, *a sufficient quantity*. Roast the Sulphate, that it may be the more easily reduced to a very fine powder, and mix it with the powder of Charcoal. Put the mixture into a crucible, and having fitted to it a cover, let it be exposed to a strong fire for six hours ; then, having well triturated the matter, put it into six pounds of Boiling Water, in a glass or earthen vessel, and mix by agitation, preventing, as much as possible, the action of the air.

“ Let the vessel stand in a vapour-bath until the undissolved part shall have subsided, and then pour off the liquor. Pour upon the residue four pounds of Boiling Water, which, after agitation and subsidence, add to the former liquor : and then, while it is still hot, or, if it shall have cooled, after it is again heated, let Muriatic Acid be dropped into it as long as any effervescence is excited. Then let the solution be filtered and evaporated, that crystals may be formed.”

*Syn.* Muriate de Baryte (*F.*), Salzsäure Schwererde (*G.*), Muriato di Barita (*I.*), Zoutzuure Zuaaraarden (*Dutch*).

The simplicity of the first of these processes, in which the superior affinity of the Muriatic Acid for Barytes effects the decomposition of the Carbonate, recommends its general adoption ; and we believe this mineral can now be procured without difficulty, and in abundance. The second, however, may sometimes be required to be performed : it is somewhat complicated, but the theory is sufficiently obvious.

The Charcoal, by the assistance of heat, decomposes the sulphuric acid of the sulphate of barytes, attracting its oxygen, and forming with it carbonic acid, which is dissipated in a gaseous form, while the sulphur remains united with the barytes. The boiling water added to this sulphuret dissolves it ; but, during the solution, the water is partially decomposed, a portion of the sulphur attracts the oxygen of the decomposed water, and forming sulphuric acid, unites with a little of the barytes, so as reproduce some sulphate which precipitates ;



while its hydrogen unites with another portion of the sulphur, and forms sulphuretted hydrogen, the combination of which with the remaining sulphuret, converts it into a hydroguretted sulphuret, and prevents its further decomposition. Lastly, the muriatic acid added to the hot aqueous solution of these sulphurets decomposes them, disengages the sulphuretted hydrogen in the form of gas, and precipitates the sulphur; while at the same time it unites with the earth, and muriate of barytes remains in solution.

Several other methods have been proposed for the preparation of this salt; the following is that recommended by Bouillon La Grange<sup>1</sup>. Pulverise together equal parts of sulphate of barytes and muriate of lime; project the mixture into a red hot crucible, and let the fire be continued till the whole be melted, which is then to be poured out on a heated tile. After it is cold, reduce the mass to powder; boil it for some minutes in six times its weight of distilled water, and filter the solution: then evaporate the liquor to a pellicle, and set it aside to crystallize. The crystals require to be redissolved and again crystallized, to free them from any of the calcareous muriate they may retain on the first crystallization. The Edinburgh process, however, is still preferable to this of La Grange, as the previous calcination reduces any metallic salts that may be combined with the sulphate; and being thus rendered insoluble, they are separated during the first solution of the sulphuret.<sup>2</sup>

*Qualities.*—Muriate of Barytes, or in more correct language *Chloride of Barium*, has an acrid, very nauseous, bitter taste. It crystallizes in grouped quadrangular tables, bevelled on the edges; transparent, white, and very brilliant; of a specific gravity of 2.8257: and not alterable from exposure to the air. When heated, it decrepitates, becomes opaque, and ultimately melts, but is not decomposed. One part requires three of water at 60° for its solution, and 2.20 of hot water. According to Berzelius, the constituents of 100 parts of this crystallized salt are 61.85 of barytes, 23.35 of muriatic acid, and 14.80 of water<sup>3</sup>: but, as it is now properly regarded as a chloride of Barium, its composition, according to Sir H. Davy, is of Barium 66.04, chlorine 33.96 in 100 parts. It is used only for forming the following solution.

---

<sup>1</sup> *Annales de Chimie*, xlvii. 151.

<sup>2</sup> Goetling advises muriate of soda to be added to the charcoal, by which a smaller quantity of charcoal is capable of reducing a larger quantity of sulphate of barytes. A mixture of one part of muriate of soda and two parts of muriate of lime is sufficient to decompose six of the sulphate.

<sup>3</sup> *Nicholson's Journal*, 4to. iii. 25.



SOLUTIO MURIATIS BARYTÆ, *Edin.* *Solution of Muriate of Barytes.*

“Take of Muriate of Barytes, *one part* ; Distilled Water, *three parts.* Dissolve.”

*Syn.* Dissolution de Muriate de Baryte (*F.*), Soluzione di Muriato di Barite (*I.*).

*Qualities.* — This solution is a real Muriate, as it is termed, the chloride decomposing a portion of the water, the hydrogen being applied to the chlorine, and the oxygen to the Barium; Muriate of Barytes is formed, and remains dissolved in the water. It is limpid, transparent, and colourless; but is rapidly decomposed by the earthy, metallic, and alkaline sulphates and nitrates; the alkaline phosphates, borates, and carbonates, being precipitated in the form of a white powder. Its affinity for sulphuric acid is so great, that, as a re-agent, it is capable of detecting 0·00009 of that acid in any fluid, forming with it an insoluble compound.

*Medical properties and uses.* — This solution is stimulant and deobstruent, and in large doses poisonous. It was introduced into practice by the late Dr. Crawford as a remedy for cancer and scrophula; and is strongly recommended by Professor Hufeland in the latter affection, when it attacks organs endowed with exquisite irritability, as the eyes and lungs. Its use was afterwards extended to syphilis. When taken in moderate doses, it appears to increase the secretion by the skin, augments the flow of urine, and improves the tone of the system; but in large doses, it causes violent vomiting, purging, and vertigo; and the most dangerous symptoms are produced. When death is the consequence, it is owing, as Mr. Brodie has ascertained, to the poison acting on the brain and heart. It has undoubtedly been found beneficial in several instances of scrophula, in some cutaneous affections, and in ulcerations connected with elephantiasis; while in syphilis it has the power of suspending some of the symptoms for a short period. But although it be a medicine of some efficacy, yet, to use the words of Mr. Pearson, in whose opinion of its deficient powers as an antisymphilitic we place implicit faith, its “good qualities are uncertain in their operation, and narrowly circumscribed; nor is it a preparation on which great confidence can be placed for the cure of any disease.”<sup>1</sup> The dose requires to be carefully apportioned, and very gradually increased, from *℥v.*, which are sufficient at first, until *℥xx.* are taken twice a day; or more, if nausea be not excited.

It is sometimes used externally as an escharotic to fungous ulcers and specks on the cornea.

<sup>1</sup> *Observations on Remedies for Lues Venerea*, 92.



As antidotes of Muriate of Barytes, when it has been taken as a poison, M. Orfila has proposed the soluble sulphates, “if administered before a quantity of the salt sufficient to exert its fatal influence on the nervous system be absorbed!”<sup>1</sup>

MAGNESIA, Lond. *Magnesia*.

“Take of Subcarbonate of Magnesia, *four ounces*. Burn the Subcarbonate in a very strong fire for two hours, or until no effervescence is excited when acetic acid is dropped on it.”

*Edinburgh.*

“Let (*sub*) Carbonate of Magnesia be exposed in a crucible to a red heat for two hours; after which preserve it in close-stopped bottles.”

MAGNESIA USTA, Dub. *Calcined Magnesia*.

“Take of Magnesia *any quantity*. Let it be put into a crucible, and subjected to a strong heat for two hours; and when it has cooled preserve it in a well-closed glass vessel.”

*Syn.* Magnesie (F.), Gebrannte Magnesia (G.), Gebrennte Bidderzoutard (Dutch), Magnesia (I.).

The carbonic acid and water are expelled by the heat, and the pure earth remains in the proportion of five-twelfths of the weight of the subcarbonate employed: or ℥j. leaves 200 grs. of magnesia.<sup>2</sup>

*Qualities.* — It is inodorous and insipid; in the form of a white, very light, soft powder, having a specific gravity of 2.3. It turns to green the more delicate vegetable blues; does not effervesce with acids; is infusible; and requires for its solution 2000 parts of water at 60°. It does not become hot when mixed with water as lime does; and water filtered through it does not affect the vegetable blues. When exposed to the air it attracts slowly carbonic acid. Sir H. Davy has ascertained that, like the other alkaline earths, it is a compound of 60 parts of a peculiar metal, which he has named *magnesium*, and 40 of oxygen.

*Medical properties and uses.* — The same as those of the Subcarbonate. It sometimes contains lime, which is discovered by a precipitate falling when oxalate of ammonia is added to its solution in sulphuric acid. Its dose is from grs. x. to ʒss. taken in water or milk.

7406 MAGNE'SIÆ SUBCARBO'NAS, Lond. *Subcarbonate of Magnesia*.

“Take of Sulphate of Magnesia, *a pound*; Subcarbonate of Potass, *nine ounces*; Water, *three gallons*. Dissolve separately the Subcarbonate of Potass in three pints of water, and the Sulphate of Magnesia, in five pints, and filter: then add the rest of the water to the solution of Sulphate of Magnesia, and boil it, adding to it while it is boiling the solution of the Subcarbonate, with constant stirring; and strain through linen.

<sup>1</sup> *Traité des Poisons*, &c. vol. i. p. 182.

<sup>2</sup> *Black on Magnesia Alba*, 28.



Lastly, wash the powder repeatedly with boiling water, and dry it upon bibulous paper with a heat of  $200^{\circ}$ ."

CARBONAS MAGNESIÆ, Edin. *Carbonate of Magnesia.*

"Take of Sulphate of Magnesia, *four parts*; Subcarbonate of Potass, *three parts*; Boiling Water, *a sufficient quantity*. Dissolve the salts separately in twice their weight of water, and strain, or otherwise free from impurities; then mix them, and instantly add eight times their weight of boiling water. Boil the liquor for a short time, stirring it; then let it remain at rest until the heat be a little diminished, and strain it through linen, upon which the Carbonate of Magnesia will remain. The Carbonate after being well washed with pure water, is to be dried with a gentle heat."

MAGNESIA, Dub. *Magnesia.*

"Take of Sulphate of Magnesia, Subcarbonate of Potass, each *two pounds*; Boiling Water, *twenty pints*. Dissolve the Sulphate of Magnesia and the Alkali, each in ten pounds of water. Mix together the defecated liquors; then boil the mixture for a short time, and strain it while it is hot through linen stretched in a proper manner for collecting the Magnesia. Wash away the Sulphate of Kali by repeated affusions of boiling water; and finally dry the magnesia."

*Syn.* Carbonate de Magnésie (*F.*), Kohlensaure Magnesia (*G.*), Carbonato di Magnesia (*I.*).

The product of these processes is a compound of Magnesia; in the Carbonate, 11.48 of Magnesia; in the Hydrate, 38.42 of Carbonic Acid, and 15.76 of water.<sup>1</sup> Mr. Phillips states the components to be Carbonic Acid 52.4 and Magnesia 47.6 in 100 parts.<sup>2</sup> In its production the salts are decomposed, and a double exchange takes place; the Sulphuric Acid separates from the Magnesia, and unites with the Potass of the Subcarbonate, disengaging the Carbonic Acid, which in its turn combines with the Magnesia. The success of the operation depends very much on the degree of attention which is paid to the following circumstances. The water employed in every part of the process must be very soft, either rain water or pure distilled water; the Subcarbonate of Potass should be previously freed as completely as possible from any admixture of silica, by passing through the alkaline solution a current of carbonic acid, or exposing it to the air for some time before it be used, and the mixing the salts in small portions of water; and after boiling the mixture, throwing it into a large quantity of water. The large proportion of water ordered, and the boiling, are necessary for dissolving the sulphate of potass, and for expelling any redundant carbonic acid which

<sup>1</sup> *Henry's Elements*, 9th edit. 1. 594.

<sup>2</sup> *Trans. of the Pharm.* p. 78.



might occasion the magnesia to crystallize, and render it gritty. Mr. Henry recommends to pour off the water by inclination, and to put the precipitate upon chalk-stones for a little time; after which it is to be wrapped up in sheets of white paper, and dried before the fire.<sup>1</sup>

The greater part, however, of the Subcarbonate of Magnesia found in the shops is prepared, on a great scale, from bittern, the liquor remaining after the crystallization of common salt from sea water. The Bittern is heated to  $212^{\circ}$ , a solution of impure subcarbonate of potass instantly added to it, and the fire withdrawn. The other steps of the process resemble those above detailed. It is frequently adulterated with chalk, and sometimes gypsum: the former is detected by adding a little diluted sulphuric acid, which converts the magnesia into soluble sulphate, but produces an insoluble salt with the lime of the chalk. Gypsum is detected by boiling a portion of the magnesia in distilled water, and adding to the solution muriate of barytes, which will produce an insoluble precipitate, if gypsum be present.

*Qualities.* — Subcarbonate of Magnesia is inodorous and insipid; perfectly white, very light, smooth to the touch, nearly insoluble in water, and effervesces with acids. Its specific gravity is 0.294.<sup>2</sup> It is decomposed by all the acids, the alkalies, the neutral and metallic salts, lime, barytes, alumina, and by a strong heat.

*Medical properties and uses.* — Subcarbonate of Magnesia is antacid. It is a useful remedy in acidity of the primæ viæ, particularly of children, in aphthous fever, and that which attends dentition. The compound formed by its union with an acid in the stomach is purgative; but if no acid be present, magnesia does not appear to increase in any degree the peristaltic motion of the bowels. It is preferable to chalk and other absorbents in heartburn, when the bowels are costive; and has been given with advantage in dysentery, combined with ipecacuanha and opium, and the dose followed by a draught of lemonade. In calculus, when the concretions are formed in the kidney, no remedy is so efficacious. The extrication of the carbonic acid in the gaseous state, when the Subcarbonate is decomposed by acid in the stomach, sometimes proves inconvenient from the distention it occasions; but more generally it is beneficial. The usual dose is from ʒss. to ʒij., taken in water or milk.<sup>3</sup>

<sup>1</sup> Henry's *Experiments on the Preparation, &c. of Magnesia*, 8vo. Lond. 1773.

<sup>2</sup> Hoffmanni *Op.* iv. 473.

<sup>3</sup> The empirical nostrum, sold under the name of DALBY'S CARMINATIVE, consists of carbonate of magnesia ʒij., oil of peppermint, m j., oil of nutmeg m ij., oil of anniseed m iij., tincture of castor m xxx., tincture of assafœtida m xv., spirit of pennyroyal m xv., compound tincture of cardamoms m xxx., and peppermint water f ʒij.



Officinal preparations. *Magnesia*, L. E. D. *Hydrargyrum cum Magnesia*, D.

## APPENDIX TO THE ALKALINE AND EARTHY SALTS.

### I.—HYDRIODAS POTASSÆ.

#### *Hydriodate of Potass.*

In noticing this substance, which, although it have not been admitted into any of the British Pharmacopœias, yet has been extensively used, I will take the opportunity of describing Iodine, the combination of which with Hydrogen forms the Hydriodic acid, the acid base of the Hydriodate.

Iodine is procured by first lixiviating powdered kelp with cold water; then evaporating the ley till a pellicle form, and setting it aside to crystallize. On separating the crystals, the mother water is to be evaporated to dryness, and to the mass, put into the flask of an alembic, is to be added half its weight of sulphuric acid, and the same weight of black oxide of manganese; and, after adapting a capital and receiver to the flask, the mixture is to be distilled with a gentle heat, as long as violet vapours arise, which condense chiefly in the capital, in the form of opaque crystals, with a metallic lustre. These are IODINE. Various other methods have been employed for procuring Iodine, the best is the following, proposed by Dr. Ure. Take eight fluid ounces of the brown liquid, which drains from the salt which the soap-makers, who employ kelp, boil up and evaporate to dryness, heat it to 230° Fahrenheit, and add one fluid ounce of sulphuric acid diluted with its own bulk of water. When the mixture cools, separate the crystals of the salts<sup>1</sup>, which will form in it by filtration through a woollen cloth, and add to the fluid poured into a mattress, 830 grains of black oxide of manganese in powder. A glass globe is then to be inverted over the mouth of the mattress, and the heat of a charcoal chaffer being applied, Iodine will sublime in great abundance. It must be washed out of the globe with alcohol, then drained and dried on plates of glass; and purified by a second sublimation from dry quicklime.<sup>2</sup>

Iodine has been procured from sponge by M. Straub of Hofwyl<sup>3</sup>; and from various sea plants; for instance, *Fucus saccharinus*,—*digitatus*,—*serratus*,—*vesiculosus*,—*siliquosus*,—*filum*,—*rubens*,—*cartilagineus*,—*membranaceus*, and—*filamentosus*; *Ulva pavonia*, and *U. linza*. Its presence in plants is detected by mixing with their aqueous infusions, starch and sulphuric acid, and then adding a small quantity of aqueous solution of chlorine. At the place where the fluids touch will be perceived a blue zone, which becomes more visible if the vessel be slightly agitated; but disappears if the whole be mixed.<sup>4</sup>

The discovery of Iodine is due to M. Courtais, a French chemist, who first obtained it in 1811; but its nature was not known to the philosophical world till 1813, when it was announced to the French Institute by M. Clement. Its properties and chemical affinities were afterwards determined by the experiments of Gay Lussac<sup>5</sup>, Sir H. Davy<sup>6</sup>, Vauquelin<sup>7</sup>, Colin, Gauthier de Claubry<sup>8</sup>, and M. Pelletier.<sup>9</sup>

*Qualities.* — Iodine, when properly prepared, is a crystallized substance of a greyish black colour, having a specific gravity of 4.946, and a metallic lustre: its smell is disagreeable, not unlike that of chlorine, and its taste acrid and hot. It

<sup>1</sup> These are sulphate of soda, sulphate of potass, hydriodate of soda and sulphur.

<sup>2</sup> *Phil. Magazine*, l. p. 161.

<sup>3</sup> *Journ. of Science and the Arts*, vol. x. p. 456. <sup>4</sup> *Ann. de Chimie*, xxviii. p. 179.

<sup>5</sup> Gay Lussac's Experiments were published in November 1813. Vide *Ann. de Chim.* t. lxxxviii. p. 319.

<sup>6</sup> Sir H. Davy's Experiments were published in December 1813.

<sup>7</sup> *Ann. de Chimie.* t. xci.

<sup>8</sup> *Journ de Phys.* Aout. 1814.

<sup>9</sup> *Bulletin de Pharmacie*, t. vi.



is usually obtained in romboidal plates, which show a lamellated fracture; are scarcely soluble in water, but more so in alcohol, and still more in sulphuric ether. It melts at 224° Fahrenheit, and is volatilized at a temperature between 347° and 556°. Its vapour is of a beautiful violet colour, (whence its name from *ιωδής* violet): It changes vegetable colours to yellow: is sparingly soluble in water; is soluble in alcohol; and when combined with hydrogen, forms an acid, the Hydriodic. It is sometimes adulterated with coal, but this is easily detected by alcohol, in which pure Iodine is completely soluble.

*Medicinal Properties.* — From the fact that burnt sponge forms the basis of all the remedies that have been productive of any benefit in the treatment of bronchocele, Dr. Coindet of Geneva, supposing that Iodine was the active principle of the sponge, proposed to employ it in different combinations for the cure of that disease. He gave it in the form of tincture, made by dissolving forty-eight grains of Iodine in a fluid ounce of alcohol; and, also, in the form of hydriodate of potass.

The *Hydriodate of Potass* is made by dissolving Iodine in a solution of pure potass. Both an iodate and a hydriodate are formed; the first of which being much less soluble than the second, falls to the bottom of the solution in the form of small grains; the second (the hydriodate) remains dissolved in the liquid, which assumes a bright-yellow colour if the Iodine be not in excess, but a deep, brownish-yellow, if it be in excess. Numerous cases of the beneficial results of the exhibition of both these preparations of Iodine in bronchocele and in scrophulous swellings, have been published by Dr. Coindet and others: but, in this country, the remedy has been too little used to determine its real value.

The Hydriodate of Potass is the form of the medicine now generally preferred. Professor G. Taddei (*Gionale di Fisica Chim.* &c. Dec. 11. Tome iv. p. 81. Pavia, 1823.) gives the following formula for its preparation. Dissolve Iodine in Alcohol, and add to the tincture some Hydrosulphate of Potass; the fluid becomes first turbid, then assumes a deep red passing to flesh colour, and finally becomes white, like milk. If the addition of some drops of hydrosulphate of potass do not produce turbidness, the operation is complete: after a time the sulphur, which gives it the white colour, falls in flakes to the bottom, and may be separated by decantation or filtration. Distil the solution to regain the alcohol employed in the process, and the residue to dryness to obtain the *Hydriodate of Potass*.

Much caution is requisite in the administration of Iodine. In delicate, nervous habits, it is apt to bring on palpitations, dry cough, tremors, especially of the arms, and other febrile symptoms; and occasionally diarrhœa and priapism. M. Hufeland has recorded an instance of the use of the Tincture of Iodine for six months completely obliterating the mamma of a young healthy female of twenty years of age. What would be the result of its use in cancer of the mamma? <sup>1</sup> Professor Ullman, of Marburg, has published several cases of cancerous ulcer, which he treated successfully by the application of Hydriodate of Potass, used both as an ointment and a lotion.

The dose of the Tincture of Iodine is from ten to fifteen drops for an adult, given in a glassful of sugared water, or of syrup of capillaire and water, three times a day<sup>2</sup>, and of that of the Hydriodate, from six to ten drops, in the same vehicle.

Besides the use of Iodine as a curative agent, it has been employed as a test of the presence of *oxide of arsenic* and of *corrosive sublimate*. Brugnatelli, who first proposed its employment for this purpose, directs as much Iodine to be added to recently boiled starch as will give it a blue colour, and then as much distilled water to be mixed with this coloured starch as will bring it to the state of an aqueous solution. A few drops of an aqueous solution of oxide of arsenic added to this solution, changes its colour at first to a reddish hue, which gradually disappears, leaving the mixture colourless; but the blue colour is restored by the addition of a few drops of sulphuric acid. The same effects, with the exception of the restoration of the blue colour by the acid, are produced by Iodine on a solution of corrosive sublimate. <sup>3</sup>

<sup>1</sup> *Journ. de Hufeland*, June 1824.

<sup>2</sup> The Tincture has been successfully used in leucorrhœa by Dr. Sublairoles. See *Journ. Univers. des Sciences Med.* Oct. 1823. p. 124.

<sup>3</sup> *Jiorn. di Fisica*, ix. p. 465.



## No. 2.

TABLE presenting a synoptical View of the Neutral Salts with Alkaline and Earthy Bases, employed as remedies, or for pharmaceutical purposes<sup>1</sup>, retaining the Names in the British Pharmacopœias.

Salts.	Taste.	Figure of Crystals.	Action of Air.	Solubility in 100 parts of Water.		Action of Heat.
				at 60°	at 212	
Sulphate of Barytes	None	Rhomboidal prisms	None	0	0·002	Decrepitates.
— Potass	Bitter	Six-sided prisms	None	6·25	24	Decrepitates.
— Soda	Bitter	Six-sided prisms	Effloresces	35	125	Watery fusion.
— Magnesia	Bitter	Four-sided prisms	None	100	133	Watery fusion.
Alum	Astringent	Octahedrons	Little	20	33	Watery fusion.
Nitrate of Potass	Cooling	Six-sided prisms	None	14·3	100	Fuses.
Muriate of Barytes	Astringent	Four-sided prisms	None	20	30	Decrepitates.
— Soda	Salt	Cubes	None	35·46	36·16	Decrepitates.
— Lime	Bitter	Six-sided prisms	Deliquesces	400		Watery fusion.
— Ammonia	Acrid	Four-sided pyramids	Subdeliquesces	31	50	Sublimes.
— Magnesia	Bitter	Needles	Deliquesces	151		Watery fusion.
Hyperoxymuriate of Potass	Cooling	Rhomboidal plates	None	6	40	Gives out oxygen.
Phosphate of Lime	None	Six-sided prisms	None	0	0	Vitrifies.
— Soda	Salt	Rhomboidal prisms	Effloresces	25	50	Watery fusion.
Borax	Styptic	Six-sided prisms	Effloresces	0·025	16·8	Watery fusion.
Carbonate of Barytes	None	Rhomboidal prisms	None		0·043	Little.
— Lime	None	Various	None	0	0	Decrepitates.
— Potass	Alkaline	Four-sided prisms	None	25	83½	Watery fusion.
— Soda	Alkaline	Octahedral truncated	Effloresces	50	100 +	Watery fusion.
— Magnesia	None	Six-sided prisms	Effloresces	2		Decrepitates.
— Ammonia	Urinous	Irregular	None	50 +	100	Evaporates.
Acetate of Potass	Hot	Plates	Deliquesces	99		Melts.
— Ammonia	Cool	Slender prisms	Deliquesces	Very soluble		Melts and sublimes.
Tartar	Acid	Irregular prisms	None	1	3½	Melts.
Tartrate of Potass	Bitter	Four-sided prisms	None	25	50	Melts.
Tartrate of Potass and Soda	Bitter	Eight-sided prisms	Effloresces	20	30	Melts.

<sup>1</sup> We have formed this Table from the more general Table of Dr. Thomson, correcting some of the proportions by experiment. See *System of Chemistry*, 4th ed. iii. 368.



## METALLIC PREPARATIONS.

THE pure metals exert no action on the animal system ; for, although Iron be administered in its metallic state, yet it must be changed by acid in the stomach to an oxide before it can prove active as a remedy. Tin operates only by mechanical attrition ; and Mercury, which has also been given internally in the metallic form, on mistaken principles, cannot act otherwise than as a mechanical substance : but when metals suffer oxidizement, or are changed by acids to the state of salts, they constitute a class of remedies of great activity and importance. The following are employed as remedies :

a. in a metallic state,

TIN, MERCURY ;

b. variously combined with oxygen, acids, sulphur, &c.

SILVER,	IRON,	BISMUTH,
MERCURY,	LEAD,	ANTIMONY,
COPPER,	ZINC,	ARSENIC.

The union of oxygen with a metallic base is denominated oxidizement, and the resulting compound an *oxide*. This combination for medicinal purposes, is effected in four ways : 1. By the action of atmospheric air, aided by an increased temperature ; 2. By deflagration with Nitrate of Potass ; 3. By the action of water : and, 4. By solution in an acid, the acid being afterwards abstracted by an alkali, or some substance for which it has a greater affinity than it has for the oxide of the metal. In whatever manner the oxidizement is effected, metals in changing to oxides lose their lustre, tenacity, inflammability, and other metallic properties ; and are gradually converted into earthy-like substances, the weight of which is greater than that of the portion of metal employed. Different metals combine with different quantities of oxygen, which is even the case with the same metal ; and as a striking alteration of properties, particularly of that of colour, marks the maximum and minimum of oxidizement, colour is taken advantage of in naming the oxides : thus *Black Oxide of Iron* is Iron in its lowest degree of oxidizement ; *Red Oxide of Iron*, the metal in its highest degree of oxidizement. There are intermediate degrees, however, which cannot be correctly expressed in language from the colour alone ; and consequently the nomenclature of this division of preparations is defective.<sup>1</sup>

<sup>1</sup> Dr. Thomson has endeavoured to remedy this defect, by introducing the term *protoxide* to signify the lowest degree of oxidizement ; *peroxide* the highest ; and *deutoxide*, *tritoxide*, &c. the intermediate degrees.

→  
 protoxide  
 deutoxide  
 tritoxide  
 peroxide



Some metals are capable of so high a degree of oxidizement as to acquire acid properties, which is so particularly the case with the White Oxide of Arsenic, that it is regarded as an acid by several chemists. The activity of the oxides of metals on the animal system appears to be regulated, with a few exceptions, by the quantity of oxygen with which they are combined; and, therefore, as Dr. Murray has justly observed, “when a process for the preparation of any metallic oxide has once been established, and practitioners have become accustomed to its powers and strength, the process ought not to be varied nor changed, from the idea of some trivial improvement; as an alteration of circumstances, apparently of little importance, may give rise to a very important change in the result. And it is nearly demonstrable, that the oxides of a metal formed by different processes, as, for example, by a process conducted in the humid way, or by one with the application of heat, cannot be precisely the same.”<sup>1</sup>

Besides the above effects of oxidizement on metals, it renders them capable of uniting with acids, and forming soluble salts. The METALLIC SALTS, therefore, are oxides combined with acids: and this is the case, whether an oxide previously prepared be dissolved in an acid, or whether the salt be the product of the direct solution of a metal in an acid. In the latter case, the metal first gains oxygen either from a part of the acid itself, or from the water, or the air, which it decomposes; and the oxide thus formed is then dissolved by the remainder of the acid. The properties of the metallic salts are much varied by the previous degree of oxidizement of the metals; and this is a point, the fixing of which in pharmaceutical operations is of the first practical importance; for, if in all the indefinite degrees of oxidizement the metallic oxides combine with acids, the resulting salts must vary in as many shades as exist between the maximum and minimum of oxidizement. In the preparation of the metallic salts, therefore, the same strict attention is requisite in following one established and approved process.

No part of chemical and pharmaceutical language is so faulty as the nomenclature of the metallic salts. Thus, although there is no instance of a direct combination of a metal with an acid, yet we have *Sulphate of Iron*, *Nitrate of Silver*, *Muriate of Mercury*, &c.; and to express the combination of the metallic oxides containing a maximum of oxygen, with acids, the syllables *oxy* are prefixed, as *Oxysulphate of Iron*, *Oxynitrate of Silver*, *Oxymuriate of Mercury*, &c. generic

---

<sup>1</sup> *System of Mat. Med.* ii. 253



terms, which can be properly applied only to denote the compounds of Oxymuriatic Acid with salifiable bases; admitting the existence of such an acid.

The prefixing the terms *sub* and *super* to denote the quantity of acid below or above the point of perfect neutralization in any salt, is not objectionable in a chemical point of view: but for medicinal purposes this mode of distinguishing salts which have the most marked difference in their active properties, by the alteration or the addition of a syllable only, may be productive of the worst consequences; and, therefore, it is the more remarkable, that the latter terms are employed in all the British pharmacopœias to denote preparations betwixt which there is very little relationship, and which cannot be converted into each other by any subtraction or addition of acid. The illustration of these observations will be found under the individual preparations.

Many of the metallic salts are altered by exposure to the atmosphere; some effloresce and attract oxygen; some are altered in their properties by moisture; and others are reduced by the action of light: thence, all of them ought to be kept in well-stopped glass bottles; and perhaps these always should be either made of green glass, or otherwise rendered opaque. In compositions which require these salts to be dissolved in water, *distilled* or *filtered Rain-water* should always be employed, and much attention is requisite to avoid combining them with incompatible substances, which may either chemically decompose them, or alter their medicinal properties.

Sulphur also combines with the metals and their oxides; but its affinity for the former is greater, and thence there are more *Metallic Sulphurets* than *Sulphuretted Metallic Oxides*. Metallic sulphurets are also formed, when sulphuretted hydrogen gas is thrown into the acid solutions of those metals which have a weak affinity for oxygen: and, as the metallic solutions differ greatly in the degree of facility with which they are thus decomposed, sulphuretted hydrogen gas may be employed, as Proust has shown, for separating different metals held together in the same solutions. The metallic sulphurets are more used for pharmaceutical purposes than as remedies, their dose not being easily appreciated, and their effects uncertain.

### PREPARATIONS OF ANTIMONY.<sup>1</sup>

SULPHURETUM ANTIMONII PRÆPARATUM.  
Edin. *Prepared Sulphuret of Antimony.*

---

<sup>1</sup> The London College have wisely discarded the *oxide of Antimony* from the list of their preparations. As a remedy it was too violent and uncertain in its operation to be generally employed; and for the composition of any other antimonial preparation it was utterly useless.



“ Put Sulphuret of Antimony, rubbed to powder in an iron mortar and levigated with a little water upon a porphyry stone, into a large vessel; then pour water on it, and, after frequently agitating the vessel, pour it off loaded with the fine powder.

“ The coarse powder, which the water cannot suspend, is to be again levigated, and treated in the same manner.”

*Dublin.*

“ Let it be reduced to powder, and separate for use the very fine particles, in the manner directed for the preparation of chalk.”

*Syn.* Sulphure d'Antimoine (*F.*), Schwazer Schwefelspeiss-glanz (*G.*), Solfuro d'antimonio depurato (*I.*), Kohul (*Arab.*), Surmah (*Hind.*).

This mechanical preparation is intended to fit the Sulphuret for internal use.

*Qualities.* — Prepared Sulphuret of Antimony is an inodorous, insipid, blackish, or deep leaden-gray, dull powder, which stains the fingers, and is insoluble in water.

According to the analysis of Doctor Thomas Thomson, Sulphuret of Antimony is a compound of 100 parts of Antimony and 35.572 of sulphur, which is nearly a mean of the analysis by Proust, Vauquelin, Dr. John Davy, Bergman, and Berzelius. Mr. Phillips, however, makes the quantity of the sulphur to be greater; and gives the following as the proportions: Antimony  $73\frac{1}{3}$ , Sulphur  $26\frac{2}{3}$ , in 100 parts; or, if we adopt the atomic theory, sulphuret of antimony consists of one atom of antimony = 44, and one atom of sulphur = 16, the weight of the atom of the sulphuret being = 60.<sup>1</sup>

*Medical properties and uses.* — Sulphuret of Antimony is inert, unless it meet with acid in the stomach, in which case it usually operates either as a diaphoretic or a mild cathartic, but occasionally produces excessive vomiting and purging; and thence it is proper to evacuate the stomach and bowels previous to its use. It has been found efficacious in scrophula, gout, chronic rheumatism, and herpetic eruptions; but its beneficial effects are very slowly produced, and consequently the use of the remedy, in order that it may prove useful, must be continued for a considerable length of time. The dose is from grs. v. to ℥j., mixed with honey or any convenient vehicle.

Officinal preparations. *Antimonii Sulphuretum præcipitatum*, L. E. D. *Pulvis antimonialis*, L. E. D.

ANTIMONII<sup>2</sup> SULPHURETUM PRÆCIPITATUM, Lond. *Precipitated Sulphuret of Antimony.*

<sup>1</sup> Translation of the Pharmacopœia, 1824.

<sup>2</sup> Formerly, *Sulphur antimonii præcipitatum*, *Sulphur auratum Antimonii*. In strict compliance with the principles of the new nomenclature, the present name should be *Hydrosulphuretum Oxidi Antimonii* — Murray.



“ Take of Sulphuret of Antimony in powder, *two pounds* ; Solution of Potass, *four pints* ; Distilled Water, *three pints*. Mix them, and boil the mixture over a gentle fire for three hours, assiduously stirring it, and occasionally adding distilled water, so that the same measure may be kept up. Strain the solution directly through a doubled linen cloth ; and, while it is still hot, drop in gradually as much Sulphuric Acid as may be necessary for precipitating the powder ; then wash away the Sulphate of Potass with hot water, dry the precipitated Sulphuret of Antimony, and rub it to a fine powder.”

SULPHURETUM ANTIMONII PRÆCIPITATUM, Edin.

“ Take of Solution of Potass, *four parts* ; Water, *three parts* ; prepared Sulphuret of Antimony, *two parts* ; Diluted Sulphuric Acid, *a sufficient quantity*. Mix the Sulphuret with the Solution of Potass and the Water, then boil them in a covered iron pot over a gentle fire for three hours, frequently stirring with an iron spatula, and adding water as it may be required. Strain the hot liquor through a doubled linen cloth, and add to it when strained as much diluted Sulphuric Acid as may be necessary for precipitating the Sulphuret, which must be well washed with warm water.”

SULPHUR ANTIMONIATUM FUSCUM, Dub. *Brown Antimoniated Sulphur*.

“ Take of Subcarbonate of Kali, prepared Sulphuret of Antimony, each *one ounce*. Having mixed them, melt the mixture in a crucible and, when it is cold, reduce it to powder. Put it into a mattress with four pints of water, and boil for a quarter of an hour ; then remove the vessel from the fire, and cover it ; let it rest a little ; and when the liquor becomes limpid, cautiously decant it from the sediment. The Antimoniated Sulphur will partly separate as the liquor cools ; add as much diluted Sulphuric Acid as will precipitate the whole of it, which takes place with an excess of acid ; then agitate the mixture, in order that the latter precipitate (which is of an orange colour) may be mixed with the rest ; and after allowing it to subside, pour off the liquor from the sediment, which is to be washed with cold water as long as litmus indicate the presence of acid in the effused fluid. Finally, dry it upon bibulous paper.”

*Syn.* Soufre doré d'antimoine (*F.*), Gelber Spiessglanzschwefel (*G.*), Zolfo dorato di antimonio (*I.*).

Although the last of these formulæ differs from the two former, the products of all of them are the same, — a Sulphuretted Hydrosulphuret of Oxide of Antimony. The following is the theory of its formation. During the boiling, the Potass combines with the Sulphur of the Sulphuret of Antimony, and forms Sulphuret of Potass ; which decomposing part of the



water, and attracting its disengaged Hydrogen, is partly converted into a Sulphuretted Hydrosulphuret of Potass, while its Oxygen, aided by the Sulphuretted Hydrogen, oxidizes the Antimony, which is dissolved by the Sulphuretted Hydrosulphuret of Potass. The Sulphuric Acid, which is now added to the strained solution whilst it is hot, and which, in part, contains Potass, Oxide of Antimony, Sulphur and Hydrogen, combines with the Potass, disengaging Sulphuretted Hydrogen gas, and the Oxide of Antimony is precipitated, combined with the disengaged Sulphur and the remaining Sulphuretted Hydrogen. In the Dublin process, the precipitate thrown down whilst the decanted liquor cools is a powder of a brick-red colour, the well-known *Kermes Mineral*<sup>1</sup>, which is the Oxide of Antimony in union with such portions of Sulphur and Sulphuretted Hydrogen only as it can attract; while the precipitate afterwards thrown down by the acid, is the old *Sulphur auratum Antimonii*, or a Hydrosulphuret of Antimony with an excess of Sulphur; and thence by agitating the mixture, a compound, or intermediate product, is obtained, which is the Sulphuretted Hydrosulphuret of the Oxide, as in the former cases. According to Thenard, the Oxide in these two powders is in a different state of oxidizement; an opinion, however, which is at least very problematical. The following are the proportions of their constituents given by him: *Kermes Mineral* consists of 72·760 parts of Brown-Oxide of Antimony, 20·298 of Sulphuretted Hydrogen, 4·156 of Sulphur, and 2·786 of Water and loss: *Golden Sulphur of Antimony* contains 68·30 of Orange-Oxide of Antimony, 17·877 of Sulphuretted Hydrogen, 12·00 of Sulphur, and 1·823 of Water and loss — in 100 parts.<sup>2</sup> But the real difference appears to consist in the larger portion of Sulphur thrown down with the *Golden Sulphur*; the base being the same in both, as stated by Trommsdorff.<sup>3</sup>

*Qualities.* — The Precipitated Sulphuret of Antimony, as it is called, is an orange-coloured powder, slightly styptic to the taste, inodorous, and insoluble in water. It readily catches fire, and burns with a blue and greenish flame, exhaling the odour of sulphurous acid, and leaving the metal after the combustion, in the form of a greyish-white oxide. It is said to be

<sup>1</sup> This powder, although now discarded from the Pharmacopœias, was long a celebrated remedy. It was discovered by Glauber, and hence named *Panacea Glauberiana*; and the process kept secret until the French Government published it in 1720, having purchased it from one La Legerie, a surgeon, to whom it had been communicated by a pupil of Glauber.

<sup>2</sup> *Annales de Chimie*, xxxii. 268.

<sup>3</sup> *Annales de Chimie*, xxxiv. 132. The quantity of the sulphuretted hydrosulphuret is much increased, by adding to the sulphuret of antimony a small portion of sulphur.



frequently sophisticated. When pure it does not effervesce with acids, a fact which enables its adulteration with chalk to be readily discovered.

*Medical properties and uses.* — This preparation of antimony is diaphoretic, expectorant, and emetic according to the dose. It was formerly much employed in asthma, and in catarrhal affections; but it is uncertain in its operation, and is not much employed in modern practice; unless when combined with mercurials, when it forms a useful alterative in herpetic and other eruptions. It is also occasionally employed in conjunction with opium or other narcotics, and Sarsaparilla in cases of chronic rheumatism. During its use, the patient must avoid the use of acids and acidulous salts, as these increase very powerfully the emetic properties of the preparation. The dose is from gr. j. to grs. iv. in a pill, twice a day.

Officinal preparation. *Pilulæ Hydrargyri submuriatis comp.*, L.  
ANTIMONIUM TARTARIZATUM, Lond. *Tartarized Antimony.*

“Take of Glass of Antimony, finely pulverized<sup>1</sup>, Supertartrate of Potass, in powder, of each *one pound*; Boiling Distilled Water, *one gallon*. Mix the Glass of Antimony and the Supertartrate of Potass; and add them by degrees to the Boiling Distilled Water, constantly stirring the mixture with a spatula; then boil for a quarter of an hour and set it aside. When the solution is cold, filter it, and evaporate the filtered fluid so that crystals may form.

TARTRAS ANTIMONII; olim, TARTARUS EMETICUS, Edin.  
*Tartrate of Antimony, formerly Tartar Emetic.*

“Take of Sulphuret of Antimony, and Nitrate of Potass, of each an *equal weight*; Supertartrate of Potass, a *sufficient quantity*. Triturate separately the Sulphuret and the Nitre, and having mixed them well together, throw them into a red-hot crucible. When the deflagration is finished, separate the red matter from the white crust, and rub it into a very fine powder, which must be washed with several effusions of warm water; and afterwards dried.

“This powder is now to be rubbed together with an equal weight of Supertartrate of Potass, and the mixture boiled in a glass vessel, with four times its weight of Distilled Water, for an hour; then strained through paper, and the strained solution set aside to deposit crystals by evaporation.”

TARTARUM ANTIMONIATUM sive EMETICUM, Dub. *Antimoniated or Emetic Tartar.*

“Take of Nitro-muriatic Oxide of Antimony, *two ounces*;

---

<sup>1</sup> For the method of preparing Glass of Antimony, see Part I. p. 177.



Crystals of Tartar, rubbed to a very fine powder, *two ounces and a half*; Distilled Water, *eighteen fluid ounces*. Boil the Water in a glass vessel; then gradually throw into it the Oxide and the Tartar previously mixed together, and boil the mixture for half an hour; then filter the solution through paper, and allow it to crystallize by slow cooling."

*Syn.* Tartrate de potasse antimonié (*F.*), Spiessglanz-weinstein (*G.*), Tartaro Antimoniato (*I.*).

By all these methods, a little modified, crystallized Tartar Emetic may be prepared. With regard to the formula of the London College, the process has the advantage of being simple. When the Glass of Antimony, which is a protoxide combined with sulphur and a small portion of silica, is boiled with Supertartrate of Potass, the excess of acid dissolves the protoxide leaving the other components unacted upon. The Tartrate of Potass which remains, being however more soluble than the supertartrate, is held in solution with the Tartrate of Antimony; thence the product is a double salt, the Tartrate of Potass and of Antimony. Mr. Phillips recommends the employment of about one-tenth more of Glass of Antimony, and to boil the mixture for a longer time than is directed.

The theory of the other two processes is sufficiently obvious. The superabundant acid of the Supertartrate of Potass combines with the Oxides of Antimony, forming a double salt, or a Tartrate of Antimony, and of Potass; which, on the principles of the reformed nomenclature, should be the pharmaceutical name of this salt.

It must be regretted that all the colleges have not concurred in adopting the same preparation of antimony for the formation of this important salt.

*Qualities.* — Tartrate of Antimony and Potass is procured in crystals, the general character of which is an octahedron with a rhombic base. They are white, inodorous, very slightly styptic and metallic, and efflorescent. It should be bought in crystals, being, when in powder frequently adulterated with Supertartrate of Potass and Tartrate of Lime. If a few of the crystals be put into a dilute solution of *Ammoniæ Sulphuretum*, their goodness may be judged of by the quantity of orange-coloured precipitate that forms. When it is adulterated with Supertartrate of Potass, it is precipitated from its solution in water by spirit. If the crystals deliquesce, its purity may be suspected. It is soluble in about 15 parts of water at 60°, and in 2 parts at 212°, forming a perfectly clear, transparent solution. It is spontaneously decomposed when kept in aqueous solution; and is also decomposed by heat, the strong acids, the alkalies and alkaline carbonates, the earths, hydrosulphurets, some of the metals, and their oxides, lime-water, muriate of



lime, and acetate of lead, and by the decoctions or infusions of many bitter and astringent vegetables, as those of Cinchona bark, rhubarb, galls, and catechu; with which, therefore, it ought never to be conjoined in prescriptions. It is insoluble in alcohol. According to the analysis of Thenard<sup>1</sup>, its constituents are 35·4 of tartaric acid, 39·6 of oxide, 16·7 of potass, and 8·3 of water; and to a more recent analysis by Dr. Göbel, of Jena, protoxide of antimony 42·6, tartaric acid 45·0, potass 9·8, and water 5·75<sup>2</sup>: but these analyses are far from being satisfactory.

*Medical properties and uses.* — This triple salt is emetic, diaphoretic, expectorant, alterative, rubefacient, and sometimes cathartic. It is certainly the most important of the antimonial preparations, and in proper doses may supersede the use of all the others. It is given as an emetic in the commencement of fevers, in doses of from one to two grains dissolved in distilled water. To obtain its diaphoretic effect, the general dose is from one-sixteenth to one-fourth of a grain. But Laennec in France, and M. Rasori in Italy, prescribe it in very large doses, in acute rheumatism, inflammation of the chest, chorea, hydrocephalus, and apoplexy. The former orders from grs. iv. to grs. vj. in a glass of infusion of orange leaves, sweetened, and gradually increases the dose. Rasori begins with grs. xij. during the day, and the same quantity at night, dissolved in a pint of barley water, to be taken in divided doses; and carries it to doses of ii. grs. and a half.<sup>3</sup> I have given it in doses of grs. iij. in pleurisy: but with less evident benefit than is produced by smaller doses combined with submuriate of mercury and opium, after blood letting. In smaller doses combined with squill, ammoniacum, and camphor, repeated every three hours, it operates as an expectorant. In minute doses also combined with calomel, it is a powerful alterative in many cutaneous diseases; and when ʒij. of it are triturated with ʒj. of lard, and applied to the skin, it occasions a pustular eruption, which has proved very serviceable in mania, white swellings, and deep-seated inflammations as a counter-irritant. Dr. Joseph Tonelli supposes, that independent of the local irritation on the skin, which this ointment produces, it exerts a specific effect on the lungs; owing as he conjectures to its abstracting a portion of the oxygen which the lungs furnish to the blood.<sup>4</sup> This opinion is purely hypothetical: for we may inquire—Do the lungs furnish any

<sup>1</sup> *Annales de Chimie*, xxxviii. 39.

<sup>2</sup> *Schweigger's Journal*, b. vii. p. 71.

<sup>3</sup> *Arch. Gen. de Med.* May 1824.

<sup>4</sup> *Annali. Univ. di Medic.* July 1824.



oxygen to the blood? The effects of this ointment on the lungs, however, is very wonderful, as I have had many opportunities of proving. On some skins it does not always produce the required effect, and requires to be aided by the addition of white sugar, as in the following formula: R. Antimonii Tartarizati ʒij. Sacchari albi ʒj. Unguenti Cetacei ʒix. Tere ut fiat Unguentum.

When taken in very large doses, tartar-emetic acts as a corrosive poison, producing violent vomiting, hiccough, a sensation of burning in the stomach, colic, hypercatharsis, syncope, difficult respiration, convulsions, and death. The treatment consists in evacuating the poison by bland, oily liquids freely taken; after which decoction of yellow bark should be administered, with opium and local bleedings.<sup>1</sup>

Officinal preparations. *Vinum Antimonii tartarizati*, L. *Vinum Tartratis Antimonii*, E.

VINUM ANTIMONII TARTARIZATI, Lond. *Solution of Tartarized Antimony.*

“Take of Tartarized Antimony, *a scruple*; Boiling Distilled Water, *eight fluid ounces*; Rectified Spirit, *two fluid ounces*. Dissolve the Tartarized Antimony in the Boiling Distilled Water; then add the Spirit to the filtered solution.”

VINUM TARTRATIS ANTIMONII<sup>2</sup>, Edin. *Wine of Tartrate of Antimony.*

“Take of Tartrate of Antimony, *twenty-four grains*; Spanish White Wine, *one pound*. Mix, so that the Tartrate of Antimony may be dissolved.”

In the former edition of the London Pharmacopœia, this preparation was termed a solution, *Liquor*, and wine was employed; in the present edition, the old name *Vinum* is restored, although the wine has been rejected!

These solutions, when newly made, are equal in point of strength, fʒj. of each containing grs. ij. of Tartarized Antimony. When wine is used, a slow decomposition of the salt occurs; and, therefore, the intention of the process (the obtaining a solution of a determinate strength to administer Tartarized Antimony in very minutely divided doses) is thus, in some respects, frustrated. The precipitate appears to be an Oxide of Antimony, with a portion of Supertartrate of Potass; arising, perhaps, from the Potass attracting Tartaric acid from the wine, and thus breaking the affinity which retains it as a component of the antimonial salt. Dr. Paris remarks, that when good Sherry Wine is employed, no decomposition of the salt takes place; and if any precipitate occur, it is Tar-

<sup>1</sup> One fluid ounce of infusion of yellow bark, is sufficient to completely decompose ʒj. of emetic tartar.

<sup>2</sup> *Vinum Antimonii tartarizati*, P.L. 1787.



trate of Lime, arising from an accidental impurity in the Bitartrate of Potass of the preparation.<sup>1</sup>

*Medical properties and uses.* — These solutions are diaphoretic or emetic, according to the extent of the dose. In doses of m̄x. to fʒj. in any proper vehicle, repeated every three or four hours, they usually excite diaphoresis, but they are principally used as emetics for infants, a tea-spoonful being given every five minutes until vomiting be excited.

PULVIS ANTIMONIALIS, Lond. *Antimonial Powder.*

“Take of Sulphuret of Antimony in powder, *a pound*; Hartshorn Shavings, *two pounds*. Mix, and throw them into a broad, iron pot heated to whiteness, assiduously stirring until vapours cease to arise. Rub what remains to powder, and having put it into a proper crucible, expose it to a fire, which is to be gradually raised, so as to keep it at a white heat for two hours. Triturate the residue into very fine powder.”

OXIDUM ANTIMONII CUM PHOSPHATE CALCIS; olim, PULVIS ANTIMONIALIS, Edin. *Oxide of Antimony with Phosphate of Lime*; formerly *Antimonial Powder*.

“Take of Sulphuret of Antimony in coarse powder, Hartshorn Shavings, each *equal parts*. Mix and throw them into a wide, iron pot, heated to redness, and stir them assiduously until they are burnt into a matter of a grey colour, which remove from the fire, rub to powder, and put into a coated crucible, over which another crucible having a small hole in its bottom is to be inverted and luted: then apply the fire, which is to be gradually raised to a white heat, and kept at this increased heat for two hours. Finally, reduce the matter when it is cold to a very fine powder.”

PULVIS ANTIMONIALIS, Dub. *Antimonial Powder.*

“Take of Sulphuret of Antimony in coarse powder, Hartshorn Shavings, each *two pounds*. Boil the Hartshorn in a sufficient quantity of water to separate the gelatin; then dry it, and mix it with the Antimony. Throw the mixture into a wide iron pot heated to redness, assiduously stirring until the sulphurous vapours cease to be extricated, and the matter acquire a grey colour. Rub the mass to powder when it is cold, and put it into a coated crucible; over which invert another crucible having a small hole in its bottom, and lute the two firmly together. Roast the matter with a heat gradually raised to whiteness for the space of two hours; and lastly, when it is cold, grind it to a very fine powder.”

In these processes, by the first exposure of the materials to

<sup>1</sup> *Pharmacologia.*



the action of heat, the gelatin and the other principles of the hartshorn, except the phosphate of lime, are decomposed and dissipated; the sulphur of the sulphuret of antimony is at the same time expelled, and the metal is partially oxidized, the oxidizement being favoured by the shape of the vessel and frequent stirring. By the subsequent application of heat, the oxidizement of the metal is rendered more complete, and the oxide is partially vitrified; but whether the phosphate of lime is merely mechanically mixed with the oxide, or the lime yields up part of the phosphoric acid to it, and a ternary compound of phosphate of lime and of antimony be thus produced, is uncertain. From the experiments of Chenevix, however, the former supposition seems to be more probable. In the Dublin formula the boiling of the hartshorn shavings ordered is unnecessary, as the heat effectually decomposes the gelatin, which is the only part of them that can be extracted by the boiling. There is great uncertainty in this preparation. In two specimens analysed by Mr. Phillips, the quantity of Peroxide of Antimony in one was 35 parts, and in the other 38 parts, in 100.

From this uncertainty of uniformity in a preparation by the agency of fire, Mr. Chenevix proposed the substitution of a powder prepared according to the following formula:—Let equal parts of white oxide of antimony and of phosphate of lime be dissolved in the smallest possible quantity of muriatic acid, and pour the solution into a sufficient quantity of distilled water containing pure ammonia in solution. A powder precipitates, which is a mechanical mixture of submuriate of antimony and phosphate of lime.<sup>1</sup> The process by heat, however, is still continued in the pharmacopœias, from a desire of imitating, as closely as possible, the celebrated empirical preparation of Dr. James, “James’s Powder,” as a substitute for which this preparation was first introduced; and which, according to the analysis of Dr. Pearson, consists of 43 parts of phosphate of lime, and 57 of oxide of antimony, in 100 parts.<sup>2</sup>

*Qualities.*—The antimonial powder of the pharmacopœias is inodorous and insipid, of a dull white colour, insoluble in water, and only partially soluble in acids; in this particular differing from the powder of Chenevix, which is soluble in every acid that can dissolve either of its components.

<sup>1</sup> *Phil. Mag.* xi. 110.

<sup>2</sup> *Phil. Trans.* lxxxi. 317. Another analysis of this powder has been published lately by M. Pully, an Italian chemist, who gives the following as its constituents: seven parts of protoxide of antimony, four of phosphate of lime, four and a half of sulphate of potass, and three and a half of potass, holding in solution protoxide of antimony. *Annales de Chimie*, lv. 74. *Thomson’s Chemistry*, 4th ed. iii. 315.



*Medical properties and uses.* — Antimonial Powder is intended to operate as a diaphoretic, alterative, emetic, or purgative, according to the extent of the dose, and the state or habit of the patient to whom it is administered. It is the preparation of antimony which was until lately most commonly employed in the commencement of fevers, and in inflammatory affections; being generally given with a view to its diaphoretic effect: and it is true that when a copious perspiration is early induced, after having previously evacuated the stomach and bowels, fevers of the most threatening aspect are often cut short by it; but too often it fails in producing this effect, and according to Dr. Elliotson's experiments, 100 grains have been given without producing any effect. That the specimens which I have employed have sometimes produced the intention expected, I can bear testimony: but justice obliges me to acknowledge that I have been as often disappointed: and every object for which antimonial powder can be prescribed is more certainly obtained from the employment of small doses of Tartarized Antimony. Supposing that it is properly prepared and acts like true James' Powder, I should say that those labouring under inflammatory diseases, who can bear considerable discharges by stool, experience the most benefit from the use of the Antimonial Powder, particularly when venæsection has been previously employed. In acute rheumatism it is advantageously given, combined with camphor, calomel, and opium; and with calomel and guaiacum in several cutaneous affections. As it is insoluble in water, it is given either in the form of a powder, or made up in pills. The dose is from grs. iij. to grs. viij., repeated every fourth hour, diluting freely in the intervals, until its diaphoretic effects are obtained.

## PRÆPARATUM EX ARGENTO.

### PREPARATION OF SILVER.

ARGENTI NITRAS, Lond.<sup>1</sup> *Nitrate of Silver.*

“Take of Silver, *an ounce*, Nitric Acid, *one fluid ounce*; Distilled Water, *two fluid ounces*. Mix together the Nitric Acid and Water, and dissolve the Silver in the mixture on a sand-bath. Then gradually increase the heat, that the Nitrate of Silver may be dried. Melt this in a crucible on a gentle

---

<sup>1</sup> Argentum nitratum, P. L. 1787.



fire, until, the water being evaporated, the ebullition cease: then directly pour it into proper moulds.”

NITRAS ARGENTI, Edin. *Nitrate of Silver.*

“Take of Pure Silver flatted into plates and cut, *one part*; Nitric Acid diluted, *two parts*; Distilled Water, *one part*. Dissolve the Silver in the Acid and Water previously mixed together, in a phial with a gentle heat, and evaporate the solution to dryness. Then put the mass into a large crucible, and place it on the fire, which must be at first gentle, and gradually increased until the mass flow like oil; then pour it into iron pipes previously heated and rubbed with grease. Finally, let the preparation be preserved in a well-stopped glass vessel.”

*Dublin.*

“Take of Silver flatted into plates and cut; Nitrous Acid, each *one ounce*; Distilled Water, *two fluid ounces*. Put the Silver in a glass vessel placed in a sand-bath; and pour over it the acid previously diluted. Then dissolve the metal with a gradually raised heat, and evaporate the solution to dryness. Put the mass which remains into a crucible, and dissolve it over a slow fire; finally, let it be poured into proper moulds, and preserve it in a well-stopped, glass vessel.”

*Syn.* Nitrate d’Argent (*F.*), Salpetersaures Silber (*G.*), Nitrato di argento (*I.*).

In this process the Acid is partly decomposed by the Silver which is oxidized, and the oxide dissolved as it forms in the remaining acid. The effervescence is very violent, owing to the extrication of the Nitrous-gas of the decomposing acid, which flies off in orange-coloured fumes; part of them, however, is retained in the solution, and gives it a blue-greenish colour, which goes off as it cools. In this stage of the process, the Silver held in solution is in the state of an Oxynitrate, which, by due evaporation, may be obtained in brilliant, irregular, thin, six-sided plates, having an intensely bitter taste: and although by the subsequent melting a part of the acid is expelled, yet it is probable, that the product is not reduced to the state of a subnitrate.

The difference in the quantity of acid ordered in the different formulæ does not alter the nature of the product; but it is of some consequence, in an economical point of view, to know, that even in the Dublin formula, which orders equal parts of Silver and Acid, the quantity of acid is too great, ten fluid drachms being amply sufficient for the solution of two ounces of Silver. Several minute particulars are necessary to be attended to in conducting the process. The silver must be perfectly free from any alloy of copper, which renders the salt always more or less deliquescent. Its presence is indicated



when the solution remains of a permanently greenish-blue colour: in which case it may be purified by repeated solutions and crystallizations, as long as tabular crystals are produced, the Nitrate of Copper being left in the mother-water. The acid employed must also be pure; for, if muriatic or sulphuric acids be present, the solution is rendered turbid by the formation of a precipitate of sulphate and muriate of silver; which, however, when only in small quantity, does not impede the process, and is easily separated by simple subsidence, after the nitric acid is fully saturated. For the same reasons the water must be pure; and therefore distilled water, or filtered rain water, should be employed. The granular form of the silver is preferable to the laminated form ordered by the Colleges. For the subsequent evaporation and melting, a porcelain crucible should be used, as the fused silver is apt to sink into the substance of the common crucibles: and it should be of ample size to allow of the swelling and ebullition. The heat must not be continued after the fusion is complete; for by continuing the application of heat, the Nitric Acid is expelled, and the Silver partially reduced; but it should be directly run into the moulds, which may be made of iron; or, in a mass of well-tempered pipe-clay, holes of the size required may be perforated by means of a greased quill, and the fused Nitrate run into them. The moulds should be previously heated to prevent the sticks of Nitrate of Silver from becoming too brittle; and when cold, each piece must be cleaned from the grease, and separately rolled up in clean, white paper.

*Qualities.*—Fused Nitrate of Silver is in small, solid cylinders of a dark grey colour, and presenting, when broken across, a crystallized structure. It is inodorous, has an intensely bitter, metallic, caustic taste, and tinges the skin and hair black wherever it touches, owing to the reduction of the nitrate by the extension of it on the cuticle. It is not deliquescent; and the presence of copper in the preparation may always be suspected when this occurs. Copper may also be detected by putting a piece of the suspected Nitrate into Liquor Ammoniaë, which will become blue if Copper be present. It is soluble in an equal weight of water at  $60^{\circ}$ , and is also soluble in alcohol. It is blackened and reduced by exposure to light or a strong heat, by phosphorus, hydrogen gas, and the hydrosulphurets; is precipitated from its aqueous solution by mercury, copper, and some other metals; and is decomposed by the alkalies, with the exception of ammonia, by the alkaline earths, sulphuretted hydrogen, the hydrosulphurets, the sulphuric, muriatic, and arsenious acids, the majority of the neutral salts, and by astringent vegetable solutions and hard



water. The constituents of 100 parts are, 31·39 of oxide of silver, and 68·39 of nitric acid.<sup>1</sup>

*Medical properties and uses.*—Nitrate of silver is tonic, antispasmodic, and escharotic. It was introduced as an internal remedy by Angelus Sala, in the commencement of the 17th century. It is said to prove efficacious in epilepsy, in angina pectoris, and in chorea. In these cases it is given in doses of one eighth of a grain, gradually increased to grs. iv. or more, three times a day; but little advantage is gained, unless its use be preceded by a course of purgatives. The chief objection to the internal administration of Nitrate of Silver is the discoloration of the skin which it sometimes produces: but M. Sementini says this may be averted by the patient avoiding the light of sunshine.<sup>2</sup> How far is it useful also, in the same cases to avoid the use of muriate of soda? The best form of administering it is that of pill made with crumb of bread, or any vegetable extract. It may be combined with Musk, Camphor, and Opium. But the chief use of Nitrate of Silver is as an external application to destroy strictures of the urethra, warts, fungous excrescences, and incipient chancres. In solution, in the proportion of gr. ij. to fʒj. of distilled water, it forms a good injection in fistulous sores: and a lotion in an apthous state of the mouth; and in that disease of the gums generally denominated scurvy, in which the gum becomes spongy, and its edges hang loosely about the necks of the teeth. When this latter disease, however, rises to a great height, the sore edges of the gum should be touched with a hair pencil dipped in a much stronger solution, in the proportion of ʒj. of the nitrate of silver to fʒj. of distilled water.<sup>3</sup> A solution of one part of the nitrate in 1000 parts of water, is recommended by Hahnemann<sup>4</sup> as an application to old sores, and for healing the ulcers of the mouth produced by the use of mercurials. In the proportion of grs. x. to fʒj. of distilled water, it was introduced by Dr. Ridgeway as a remedy in the Egyptian Ophthalmia; and has been found very beneficial. One drop or two drops are inserted into the eye every second day.<sup>5</sup>

When given in too large doses, it acts as a poison on the system, producing symptoms resembling those induced by the other corrosive poisons. M. Orfila regards common salt as the antidote of this poison, when given sufficiently early to prevent the specific action of the Nitrate on the coats of the stomach<sup>6</sup>: and so completely is it decomposed by a solution of

<sup>1</sup> Phillips.

<sup>2</sup> *Giornale di Fisica*. xi. p. 355.

<sup>3</sup> Fox on the Natural History and Treatment of Diseases of the Teeth.

<sup>4</sup> *Annales de Chimie*, iii. 308.

<sup>5</sup> *Med. and Phys. Journ.* Feb. 1825. p. 122.

<sup>6</sup> *Traité des Poisons*, &c. tom. i. p. 49.



common salt out of the body, that when a saturated solution is mixed with a saturated solution of common salt and filtered, the fluid which passes is perfectly inert. When the antidote has not been administered very early, local and general bleeding, tepid baths, and emollient fomentations and glysters must be employed, if any symptom of abdominal inflammation be perceived.

---

## PRÆPARATA EX ARSENICO.

### PREPARATIONS OF ARSENIC.

ARSENICUM ALBUM SUBLIMATUM, Lond. *Sublimed White Arsenic.*

“Rub White Arsenic to powder; then put it into a crucible, and, applying heat, sublime it into another crucible inverted over the first.”

*Syn.* Oxide d'arsenique pure (F.), Weisses Arsenick (G.), Arsenico blanco (I.).

The greater part of the White Arsenic found in the shops, is in the form of semivitreous cakes, which are the product of a second sublimation of the Oxide, after it is obtained from roasting ores of Cobalt. Although prepared on a great scale, yet it is as pure as sublimation can make it, and, therefore, this process is superfluous; and as it is also not devoid of risk to the operator, it should be altogether rejected.

LIQUOR ARSENICALIS<sup>1</sup>, Lond. *Arsenical Solution.*

“Take of Sublimed White Arsenic rubbed to a very fine powder, Subcarbonate of Potass from Tartar, of each *sixty-four grains*; Compound Spirit of Lavender, *four fluid drachms*; Distilled Water, *a pint*. Boil them together in a glass vessel until the arsenic be entirely dissolved. Add to the solution, when it is cold, the Compound Spirit of Lavender, and, finally, as much Distilled Water as will make the whole up to a pint.”

SOLUTIO ARSENICALIS, Edin. *Solution of Arsenic.*

“Take of Oxide of Arsenic rubbed to very fine powder, very pure Subcarbonate of Potass, of each *sixty-four grains*; Distilled Water *fourteen ounces*. Boil them together in a glass vessel until all the oxide be dissolved. Add to the solution, when it is cold, *half an ounce* of Compound Spirit of

---

<sup>1</sup> This appellation is certainly very objectionable, as it conveys an erroneous idea of the preparation, even admitting that the term *arsenic* may be used to designate the *white oxide*: it should have been *Liquor Arsenialis Potassæ*, or perhaps, more properly, *Liquor alcalinus Oxidi Arsenici*.



Lavender, and as much Distilled Water as will make the whole up to sixteen ounces.

The White Arsenic possesses properties in some respect similar to those of an acid. It combines with alkalies, is soluble in water and the solution reddens tincture of litmus: but it also combines with and neutralizes the acids; so that, while some<sup>1</sup> regard it as an acid, others<sup>2</sup> consider it only as a highly oxidized oxide. In the above process, by combination with the potass, its solubility is much increased, and a solution obtained of an uniform strength, by which very minute doses can be correctly and easily apportioned. It was introduced by Dr. Fowler of Stafford, whose formula the London College has adopted, altering only the proportions of the water and the spirit of lavender, to make up the pint of the solution.

*Qualities.* — This Solution, one fluid drachm of which contains half a grain of White Arsenic, has the odour, taste, and colour of the compound spirit of lavender. It is decomposed by lime-water, hydrosulphuret of potass, nitrate of silver, the salts of copper, and instantly forms a copious precipitate when dropped into infusion or decoction of cinchona bark; with which, therefore, it ought not to be conjoined in extemporaneous prescriptions.

*Medical properties and uses.* — The Arsenical solution, as it is termed, is a powerful tonic, useful in all the cases in which the white oxide can be employed. (See *Arsenici Oxydum*, Part ii.) It was introduced by Dr. Fowler as a substitute for the celebrated empirical remedy known under the name of “The Ague Drop,” which owes its efficacy to the Oxide of Arsenic. In addition to the account we have already given of the medicinal use of the oxide, we have to add, that we have given this solution with decided advantage after cupping and purging, in threatened apoplexy when the strength was little and the complexion pale. The dose is  $\mathfrak{m}\text{iv}$ . gradually increased to  $\mathfrak{m}\text{xx}$ ., given twice a day.

ARSENIAS KALI, Dub. *Arseniate of Kali.*

“Take of White Oxide of Arsenic, Nitrate of Kali, each *an ounce*. Reduce them separately to powder; then having mixed them, put them into a glass retort, and place it in a sand-bath exposed to a gradually raised heat, until the bottom of the retort becomes obscurely red. The vapours arising from the retort should be transmitted through Distilled Water, by means of a proper apparatus, in order that the Nitrous Acid extricated by the heat may be disengaged. Dissolve the residue in four pounds of Boiling Distilled Water, and after due evaporation set it apart, that crystals may form.”

<sup>1</sup> Fourcroy.

<sup>2</sup> Berthollet.



In this process the nitrate of potass is decomposed by the heat; part of the oxygen of the nitric acid with the whole of its nitrogen escape in the form of nitrous gas, while the remainder of the oxygen is attracted by the oxide of arsenic, which is thus converted into arsenic acid, and combines with the disengaged potass of the nitrate, forming a superarsenate of potass: this remains in the retort in the form of a white saline mass, and is afterwards dissolved and crystallized. The nitrous acid is not worth condensing, as the process is not likely to be performed on a great scale.

*Qualities.* — Arseniate, or rather Superarsenate<sup>1</sup>, of Potass crystallizes in beautiful, transparent, tetraëdral prisms, having an excess of acid. They are soluble in water; and the solution reddens the vegetable blues.

*Medical properties and uses.* — This salt may be used exactly in the same manner, and in the same cases as the white oxide. It was discovered by Macquer, and long known under the name of “The Arsenical Neutral Salt of Macquer.” The dose is from one-sixteenth to one-eighth of a grain, formed into a pill with crumb of bread.<sup>2</sup>

## PREPARATUM E BISMUTHO.

### PREPARATION OF BISMUTH.

BISMUTHI SUBNITRAS, Lond. *Subnitrate of Bismuth.*

“Take of Bismuth, *one ounce*; Nitric Acid, *one fluid ounce and a half*; Distilled Water, *three pints*. Mix six fluid drachms of the Distilled Water with the Nitric Acid, and dissolve the Bismuth in the diluted acid: then filter the solution. Add the remainder of the water to the filtered solution, and set it apart that the powder may subside. Next, having poured off the supernatant fluid, wash the Subnitrate of Bismuth with Distilled Water; and, having wrapped it in bibulous paper, dry it with a gentle heat.”

<sup>1</sup> The arseniate is uncrystallizable and deliquescent, and altogether a different salt; the Dublin College, therefore, has improperly named it *Arsenias Kali*.

<sup>2</sup> Since Part II. of this edition was printed, a case of poison by arsenic has occurred, in which the life of the individual was apparently saved by the frequent exhibition of carbonate of magnesia and vinum opii; (see *Med. and Physical Journ.* vol. xvi. p. 466.,) and although this result is at variance with our experimental enquiries on this subject, yet the statement of the fact, as it occurred, is sufficient to arrest the attention of the profession to try the remedy.



*Syn. (Sub)* Nitrate de Bismuth, Blanc de Ford (*F.*), Salpetersaurer Wis-  
muth (*G.*), Termassido Bianco de Bismuto (*I.*).

In this process, the Nitric Acid is partially decomposed, and the Bismuth oxidized; and being thus rendered soluble, in the remainder of the acid a solution of the Nitrate of Bismuth is obtained. This solution, when filtered, is colourless and transparent; reddens the tincture of litmus, and has a harsh caustic taste; and where the process stopped here, crystals of the nitrate might be readily obtained by evaporation. The affinity of the Acid for the Oxide in the Nitrate is, nevertheless, so slight, that it can be separated from it by water; and this is effected in the second part of the process; for the water then added combining with the greater part of the acid, and depriving the fixed oxide of its solubility, this is precipitated, in combination with some water, and the remainder of the acid, forming what is termed Subnitrate of Bismuth.<sup>1</sup> It is, in fact, a Hydrated Oxide of Bismuth, combined with a small proportion of the Nitric Acid. The supernatant fluid contains in solution a Supernitrate of Bismuth, or a Nitrate with an excess of acid. The Oxide of Bismuth consists of 90 parts of bismuth and 10 of oxygen in 100 parts<sup>2</sup>, but the proportion of this in the Subnitrate has not been determined.

*Qualities.*—The Subnitrate of Bismuth is a pure white, inodorous, insipid power. It is soluble in the strong acids, from which it is readily precipitated by water, thence it is insoluble in that fluid, and, for the same reason, in very dilute acids. It is soluble in Ammonia, although this salt precipitates it from the Nitrate; but is very sparingly soluble in potass and soda. It is blackened by Sulphuretted Hydrogen gas, its solution in water, and all the Hydro-sulphurets. When mixed with charcoal and exposed to a strong heat, the Subnitrate is decomposed, and the Bismuth reduced to its metallic state.

*Medical properties and uses.*—Subnitrate of Bismuth is tonic and antispasmodic. It has been advantageously administered in spasmodic affections, palpitations of the heart, and epilepsy. We have found it extremely beneficial in pyrosis, gastrodynia, and some other varieties of dyspepsia; in which cases we have usually combined it with extract of hops; and when there have been merely atony of the digestive organs, without organic mischief, it has proved almost uniformly successful.<sup>3</sup>

<sup>1</sup> It was formerly known by the name of *Magistery of Bismuth*.

<sup>2</sup> *Phillips' Trans. of the Pharm.* 1824.

<sup>3</sup> It was first employed in these cases by professor Odier of Geneva. See *Manuel de Medecine pratique, &c. par Louis Odier*.



The dose of the Subnitrate is from one grain to twelve or fifteen grains. In every large doses, however, the Subnitrate of Bismuth acts with great violence, producing nausea, vomiting of a white ropy matter, pains in the stomach, acute colic, heat of the chest, and alarming anxiety: and when these symptoms are accompanied with rigors, sighing, vertigo, and convulsions, the poison always proves fatal. Post mortem examinations exhibit appearances of inflammation and ulceration of the stomach; inflammation of the duodenum and jejunum; the lungs gorged with blood of a deep-red colour, so as to resemble liver. It is easy to explain the inflammatory and corrosive effects of the Subnitrate on the coat of the stomach into which it is introduced; but the effect on the lungs is less obviously explained; unless we suppose the poison to act sympathetically through the nerves as well as locally on the animal fibre. In cases of poisoning by Subnitrate of Bismuth, both general and local blood-letting must be resorted to; whilst the patient should drink freely of milk and mucilaginous fluids. Fomentations and emollient glysters are also necessary; the degree of inflammation being such as to render the employment of ordinary purgatives hazardous.

When Subnitrate of Bismuth has been employed as a poison, and the whole has not been taken, it is detected by the chemical properties, and the effects of re-agents upon it, which we have already noticed; but, when the whole has been taken, the only certain method of ascertaining the nature of the poison is to dry a portion of the vomited matter, or of the contents of the stomach if it have proved fatal; and to calcine the mass, mixed with charcoal, in a covered crucible, so as to reduce the metal, in which state it is easily recognised.

---

## PREPARATA E CUPRO.

### PREPARATIONS OF COPPER.

ÆRUGO PRÆPARATA, Dub. *Prepared Verdegris.*

“Let the Verdegris be reduced to powder, and the more subtile parts be separated in the manner directed for the preparation of Chalk.”

*Syn.* Vert-de-gris (*F.*), Grünspan (*G.*), Acetato di Rame (*I.*), Cardenillo (*S.*).

By this process the Subacetate of Copper is obtained in a state of very minute mechanical division, better fitted for in-



ternal use, in the cases for which it is sometimes prescribed. (See *Ærugo*, Part ii.)

CUPRUM AMMONIATUM, Lond. *Ammoniated Copper.*

“Take of Sulphate of Copper, *half an ounce*; Subcarbonate of Ammonia, *six drachms*. Rub them together in a glass mortar until the effervescence cease; then wrap up the Ammoniated Copper in bibulous paper and dry it with a gentle heat.”

AMMONIARETUM CUPRI, Edin. *Ammoniuret of Copper.*

“Take of pure Sulphate of Copper, *two parts*; Subcarbonate of Ammonia, *three parts*. Rub them thoroughly together in a glass mortar, until all effervescence is finished and they unite in a violet-coloured mass, which wrap up in bibulous paper and dry, first on a chalk stone and afterwards with a gentle heat. Let it be preserved in a well-stopped glass phial.”

CUPRUM AMMONIATUM, Dub. *Ammoniated Copper.*

“Take of Sulphate of Copper, *an ounce*; Carbonate of Ammonia, *an ounce and a half*. Rub them in an earthenware mortar, until all effervescence having ceased, they unite into a mass, which is to be dried, wrapped up in bibulous paper and preserved in a phial closed with a glass stopper.”

*Syn.* Sulphate de Cuivre et d'Ammoniaque (*F.*), Schwefelsaures Kupfer mit Ammonium Kuppersalmiak (*G.*), Ammoniuro di Rame (*I.*).

The product of these processes is either a triple salt, a Subsulphate of Oxide of Copper and Ammonia, or a mixture only of subsulphate of copper, and sulphate of ammonia<sup>1</sup>: but the former is the more probable state of the compound, from the difference of capacity which it has for water being so great as to render the resulting mass extremely moist. During the trituration, the Sulphate of Copper is partially decomposed, and part of its acid yielded up to the Ammonia, which is consequently freed from the Carbonic Acid, the effervescence being the effect of the dissipation of this acid disengaged in the gaseous form. The action of the affinities, which produce these changes, is, perhaps, aided by the water of crystallization of the ingredients becoming fluid. In drying the product it must be very carefully excluded from the air.

*Qualities.*—This preparation has the odour of Ammonia, a hot, styptic, metalline taste, and a rich blue colour. By exposure to the air the blue colour is lost, and the salt acquires a greenish hue.

*Medical properties and uses.*—Ammoniated copper is tonic and antispasmodic. It has been principally employed in epi-

---

<sup>1</sup> It is certainly not an ammoniuret, although so designated in the Edinburgh Pharmacopœia.



epsy, as a remedy for which, it was first proposed by Dr. Cullen; and has since his time been frequently employed with evident advantage — although we must confess, that in our trials of it the event has not been such as to encourage us to place much dependence on its powers for relieving this severe disease. It has also been given in chorea after a course of purgatives. It is less apt to excite nausea than the other preparations of copper. Cullen, however, recommends its use not to be continued for more than a month at a time: and adds, that after the first interval, if the disease continue, the most benefit will be derived from giving the medicine “only for some days before an expected accession.”<sup>1</sup> It has been given with advantage in chorea, after a course of purgatives, combined with digitalis and myrrh.

The dose is one-fourth of a grain gradually increased to grs. v. given twice a day, either simply made into pills with crumb of bread, or combined with Valerian.

**LIQUOR CUPRI AMMONIATI, Lond.** *Solution of Ammoniated Copper.*

“Take of Ammoniated Copper, *a drachm*; Distilled Water, *a pint*. Dissolve the Ammoniated Copper in the Water, and filter the solution through paper.”

**AQUA CUPRI AMMONIATI, Dub.** *Water of ammoniated Copper.*

“Take of Lime-water, *eight fluid ounces*; Muriate of Ammonia, *two scruples*; Prepared Verdegris, *four grains*. Let them be mixed together, and digested for twenty-four hours; then pour off the clear liquor.”

As nearly the same result follows whichever of these processes is adopted: that of the London Pharmacopœia, from its simplicity, is undoubtedly to be preferred, although too much water is ordered, it being a curious fact that it is more soluble in a smaller quantity of water, owing to the larger quantity decomposing the Subsulphate of Copper, and leaving an insoluble oxide of copper which is precipitated. In the Dublin process, the lime decomposes the Muriate of Ammonia, and combines with its Muriatic Acid, forming a Muriate of Lime, while the disengaged ammonia unites with the oxide of copper of the verdegris, and forms a soluble compound. It differs from the simple solution of ammoniated copper, in holding also the muriate of lime in solution; and is a stronger preparation, for nearly one-half of the oxide of the Ammoniated Copper in the simple solution is precipitated by the excess of water.

*Medical properties and uses.*—This solution is detergent,

<sup>1</sup> *Mat. Med.* ii. 25.



and mildly escharotic. It forms an useful local stimulant for cleaning foul indolent ulcers, and disposing them to heal; and is also employed, still more largely diluted, for removing specks from the cornea.

**SOLUTIO SULPHATIS CUPRI COMPOSITA,** Edin. *Compound Solution of Sulphate of Copper.*

“Take of Sulphate of Copper, Sulphate of Alumina, each *three ounces*; Water, *two pounds*; Sulphuric Acid, *one ounce and a half*. Boil the Sulphates in the water, to dissolve them; and then to the liquor filtered through paper add the acid.”

This preparation is a simple solution of the sulphates. It is sometimes used as a styptic for stopping hæmorrhages; and largely diluted as a lotion in ophthalmia tarsi, and the purulent ophthalmia of infants.

As has been already noticed (Part ii.), sugar is the best antidote for the salts of copper, when these have been taken as poisons.

## PRÆPARATA E FERRO.

### PREPARATIONS OF IRON.

**LIMATURA FERRI PURIFICATA,** Edin. *Purified Filings of Iron.*

“Having placed a sieve over the filings, apply a magnet, so that it may draw the filings upwards through the sieve.”

*Syn.* Limaille de Mars (*F.*), Gepülvertes Eisen (*G.*), Limatura di Ferro (*I.*), Eerumboo Podia (*Tam.*).

The Iron filings obtained from the workshops are always mixed with many impurities, and often with filings of Copper and other metals. It requires some address to purify them by this process; at least the sieve must not be placed too close upon the filings, but as distant as the sphere of attraction of the magnet will admit of, so that the iron only may be raised.

**OXIDUM FERRI NIGRUM PURIFICATUM,** Edin. *Purified Black Oxide of Iron.*

“Let the Scales of the Black Oxide of Iron, found at the anvil of the blacksmith, be purified by the application of the magnet; for the magnet attracts the thinner and purer scales only, leaving the larger and less pure.”

**OXIDUM FERRI NIGRUM,** Dub. *Black Oxide of Iron.*

“Let the Scales of Iron, found at the blacksmith’s anvil, be purified by the application of the magnet. Then reduce them



to a powder, the finest parts of which are to be separated in the manner ordered for the preparation of chalk.”

*Syn.* L'oxide noir de Fer (*F.*), Schwarzes gesäuertes Eisin (*G.*), Ossido nero di Ferro (*I.*).

The Scales struck off from red-hot Iron by the hammer of the blacksmith are imperfectly oxidized, but still retain their magnetic quality in a sufficient degree to admit of being purified in the above manner.

*Medical properties and uses.*—This imperfect Oxide is tonic, deobstruent, and anthelmintic. It is efficaciously administered in general debility, dyspepsia, chlorosis, and worm cases. Its utility is determined by its meeting with acid in the stomach, which is known to be the case by the disagreeable eructations it produces and the black colour of the alvine evacuations. The dose is from grs. v. to ℥j., combined with any aromatic powder, or formed into an electuary with honey and taken twice a day.

FER'RUM AMMONIATUM, Lond. *Ammoniated Iron.*<sup>1</sup>

“Take of Subcarbonate of Iron, Muriatic Acid, Muriate of Ammonia, each *a pound*. Pour the Muriatic Acid upon the Subcarbonate of Iron, and set it aside until bubbles cease to arise. Filter the solution through paper, and boil it to dryness. Mix the residuum intimately with the Muriate of Ammonia; then instantly sublime, by the application of a strong heat; finally, reduce the sublimed matter to powder.”

MURIAS AMMONIÆ ET FERRI, Edin. *Muriate of Ammonia and of Iron.*

“Take of Red Oxide of Iron washed and again dried, Muriate of Ammonia, each *equal parts by weight*. Mix them well together, and sublime by a quick fire. Reduce the sublimation to powder and preserve it in a well-stopped phial.”

MURIAS AMMONIÆ ET FERRI, Dub. *Muriate of Ammonia and of Iron.*

“Take of Red Oxide of Iron, Muriate of Ammonia, each *equal parts by weight*. Having mixed them well together, sublime them with a sudden and sufficiently strong heat.”

*Syn.* Fleurs de Mars ammoniacales (*F.*), Eisenhaliges Salzaures Ammonium (*G.*).

In these processes of the Edinburgh and Dublin colleges, the theory of the operation is obvious: the sudden application of an intense heat enables the Oxide of Iron to decompose the Muriate of Ammonia, and to unite with part of its Muriatic Acid, and at the same time it probably enters into that degree of combination with the Ammonia, which exists in triple salts, the product being either a Muriate of Iron and

---

<sup>1</sup> Ferrum ammoniacale, P. L. 1787, Flores Martialis, Ens Martis, Flores auri, Calendulæ minerales.



Ammonia, or a mixed mass of Submuriate of Ammonia and Submuriate of Iron: some difference, however, takes place in that ordered by the London College. The subcarbonate of iron is intended to be converted into dry muriate, but the whole is not dissolved in the acid, thence the formation of *ferrum ammoniatum* is never the same. The preparation is a mixture of muriate of ammonia and permuriate of iron. Mr. Phillips justly remarks, that the evaporation of the solution of Muriate of Iron is an extremely unpleasant operation and should be conducted under a chimney.<sup>1</sup>

*Qualities.*—Muriate of ammonia and iron has an odour, resembling, in some degree, that of saffron, and a styptic taste. It is in crystalline grains of an orange-yellow colour: soluble in two parts of water, and also very soluble in alcohol, and deliquescent; on which account this salt requires to be preserved in very well-stopped phials. It is a mixture of Muriate Ammonia and Permuriate of Iron.

*Medical properties and uses.*—This preparation of iron is tonic, emmenagogue, and aperient. It was formerly much used in epilepsy, hysteria, chlorosis, scrophula, and rickets: but on account of the uncertainty of the preparation it is now seldom prescribed. The dose is from grs. iij. to grs. xv. given twice or thrice a day.

Officinal preparation. *Tinctura Ferri ammoniati*, L.

SUBCARBONAS FERRI PRÆPARATUS, Edin.  
*Prepared Subcarbonate of Iron.*

“Let purified filings of Iron be frequently moistened with Water, till they fall into rust, which is to be rubbed to powder.”

FERRI RUBIGO, Dub. *Rust of Iron.*

“Take of Iron-wire, *any quantity*. Cut it into small pieces, which are to be exposed to the air and frequently moistened with Water, until they be converted into rust; let this be rubbed in an iron mortar and, by pouring water on it, wash over the finest part of the powder, which is to be dried.”

*Syn.* Carbonate de Fer (*F.*), Kohlensaures eisen Rost (*G.*), Ossido Carbonato di Ferro (*I.*), Sudud ul hedeed (*Arab.*), Eerumboo Lohayka zung (*Duk.*), Mandura (*Sans.*), Kith (*Hind.*), Zafran ahun (*Pers.*), Tuppoo (*Tam.*).

In these processes the Iron is oxidized at the expence of the Water which is decomposed, while at the same time Carbonic Acid is attracted from the atmosphere, and combined with the oxide. The product is a Subcarbonate of Oxide of Iron; for the quantity of acid is not equivalent to the neutralization of the Oxide.

According to my experiments, it consists of eighty-five parts

---

<sup>1</sup> *Trans. of the Pharm.* p. 101.



of Oxide of Iron, and fifteen of Carbonic Acid; but these proportions must necessarily vary from variations in the mode of conducting the process.

*Qualities.*—It is inodorous, has a styptic taste, and a reddish-brown colour; dissolves in acids with effervescence; and is decomposed by heat.

*Medical properties and uses.*—The Rust of Iron is tonic and emmenagogue. Next to the Black Oxide it is the least active of the preparations of this metal. It has lately been recommended with much confidence, both as an internal remedy and an external application in cancer<sup>1</sup>; and I have sufficient proof of its power to at least suspend the symptoms of that inveterate disease. It is, perhaps, one of the best remedies that has yet been tried in tic douloureux. I have had an opportunity of observing its powers in several well-marked cases of this disease; in which it proved successful: but its use required to be long persisted in, as the disease returned in two of the instances, when its use was too soon discontinued. In large doses it often occasions uneasiness at the stomach; yet Cullen says, “We have always found the simple rust as effectual as any other preparation; and the stomach bears it better than any other.”<sup>2</sup> It is given in the form of pills, combined with aromatics and bitter extracts. The dose is from grs. iv. to ʒj. given twice a day.

Officinal preparation. *Tinctura Muriatis Ferri*, D.

FERRI SUBCARBONAS, Lond. *Subcarbonate of*<sup>3</sup>  
*Iron.*

“Take of Sulphate of Iron, *eight ounces*; Subcarbonate of Soda, *six ounces*; Boiling Water, *a gallon*. Dissolve separately the Sulphate of Iron and the Subcarbonate of Soda in four pints of Water, then mix together the solutions, and set the mixture aside, that the powder may subside; then pour off the supernatant fluid, wash the Subcarbonate of Iron in hot water, and dry it, wrapped up in bibulous paper, with a gentle heat.”

CARBONAS FERRI PRÆCIPITATUS, Edin. *Precipitated Carbonate of Iron.* CARBONAS FERRI, Dub. *Carbonate of Iron.*

“Take of Sulphate of Iron, *four ounces*; Subcarbonate of Soda, *five ounces*; Water, *ten pounds*. Dissolve the Sulphate in the Water: then add the Subcarbonate previously dissolved

<sup>1</sup> Carmichael on the Use of Carbonate of Iron in Cancer.

<sup>2</sup> It is the carbonate of iron which is contained in chalybeate waters, held in solution by the excess of carbonic acid. By exposing these waters to the air, the carbonic acid flies off, oxygen is attracted, and the carbonate falls down in the form of a yellowish sediment.

<sup>3</sup> Chalybs præparatus è aceto, et sine aceto, P. L. 1720, Chalybis rubigo præparata. P. L. 1745. Ferri rubigo, P. L. 1787.



in the Water, and mix them together. Let the Carbonate of Iron, which is precipitated, be washed with Tepid Water, and afterwards dried."

This preparation is also a Subcarbonate of Iron. By mixing the solutions together, a double decomposition is effected; the Sulphuric Acid of the Sulphate of Iron combines with the Soda, while the Iron attracts the disengaged Carbonic Acid of the Subcarbonate of Soda; and thence the products are an insoluble Subcarbonate of Iron, and a soluble Sulphate of Soda, which are easily separated by washing and filtration. When first precipitated, the Subcarbonate of Iron has a deep-green colour, and is at a minimum of oxidizement; but while drying, it attracts oxygen rapidly from the atmosphere, and is converted into the red oxide, or a peroxide, containing, according to Proust, 48 per cent. of oxygen: on which it should be very rapidly dried for the peroxide is incapable of remaining in combination with Carbonic Acid. I have found that the precipitate combines with the largest proportion of Carbonic Acid, when the solutions are mixed at a temperature of 150° of Fahrenheit; and filtration is necessary for separating it, the decantation of the clear fluid being very difficult, owing to the lightness of the precipitate. The great solubility of the Sulphate of Soda renders much subsequent washing unnecessary; and the precipitate, after being washed, should be dried in the paper on which it is filtered, by a heat not exceeding 200°.

*Qualities.* — Precipitated Subcarbonate of Iron is inodorous, has a slightly styptic taste; and when properly prepared is of a chocolate-brown colour. It is insoluble in water, but acids dissolve it with effervescence, disengaging the carbonic acid in the gaseous form. It is decomposed by heat, and converted into the black oxide of the metal. In my experiments, ten grains of the dried Subcarbonate, prepared with effloresced subcarbonate of soda, lost 2·3 grains, when dissolved in muriatic acid; and the same quantity, prepared with the crystallized alkali, and dried with great care, lost 1·4; so that prepared in the former method, it contained 23 per cent. of carbonic acid, and in the latter 14 per cent. Mr. Phillips says, he has found that it contains 15 per cent. when prepared with the greatest care, which indicates its composition to be Carbonate of Iron 40, Peroxide of Iron 60, in 100 parts.<sup>1</sup>

*Medical properties and uses.* — This preparation differs little from the former preparation in its effects, except that it sits easier on the stomach. The dose is from grs. iv. to grs. xxx.: given three times a day, combined with myrrh, or aromatics.

---

<sup>1</sup> *Trans. of the Pharm.* p. 98.



But in tic douloureux, it is given to the extent of a drachm for a dose.

Officinal preparations. *Ferrum Ammoniatum*, L. *Tartarum Ferri*, D. *Tinctura Ferri Muriatis*, L.

FERRI SULPHAS, Lond. *Sulphate of Iron*.<sup>1</sup>

“Take of Iron, Sulphuric Acid, each *eight ounces*; Water, *four pints*. Mix the Sulphuric Acid with the Water in a glass vessel, and to these add the Iron; then, when the effervescence is over, filter the solution through paper, and evaporate it over the fire, so that crystals may form as it cools. Pour off the water, and dry the crystals upon bibulous paper.”

SULPHAS FERRI, Edin. *Sulphate of Iron*.

“Take of Iron, Sulphuric Acid, of each, *by weight, eight ounces*; Water, *four pints*. Mix the Sulphuric Acid with the Water in a glass vessel and the Iron to them, when the effervescence is over, filter the solution through paper, and after due evaporation set it apart, that crystals may form. Having poured off the liquor, dry these upon bibulous paper.”

SULPHAS FERRI, Dub. *Sulphate of Iron*.

“Take of Iron-wire, *two ounces*; Sulphuric Acid, *three ounces and a half*; Water, *one pint*. Mix gradually the Acid with the Water; then add the wire cut into pieces, and digest the mixture that the metal be dissolved, after which filter the solution through paper; finally, after due evaporation, set it apart, that crystals may form by slow cooling.”

*Syn.* Sulfate de Fer (*F.*), Schwefelsäures Eisen (*G.*), Zwalvelzuures yzer (*Dutch*), Gron Vitriol (*Danish*), Solfato di Ferro (*I.*), Vitriola verde (*S.*), Caparosa verde (*Port.*), Unua Baydie (*Tam.*), Casis (*Hind.*), Heera Cashish (*Duk.*), Taroosee (*Malay*), Zungarmadenee (*Arab.*), Tootya Subz (*Pers.*).

In these processes part of the water is decomposed, the Iron is oxidized by combining with its oxygen, while its hydrogen is dissipated in the gaseous form; and the Oxide thus produced unites with the acid, and forms Sulphate of Iron, or rather Sulphate of Oxide of Iron; which is dissolved in the undecomposed portion of the water. Concentrated Sulphuric Acid, nevertheless, scarcely exerts any action on iron at a low temperature, and water alone is very slowly decomposed by it, so that the rapid decomposition of the diluted acid by the Iron must be ascribed to the sum of the affinities of the base of the Acid for oxygen, and of the Iron for oxygen being superior to that of the oxygen to the hydrogen of the water, which is therefore decomposed. The solution is of a pale-green colour and, when evaporated directly, yields crystals of Sulphate of Iron<sup>2</sup>; but, if it be exposed for some time

---

Old names of this salt: — *misy*, *sory*, *calchantum*, (Pliny) *sal martis*, *sal chalybis vitriolum ferri*, *vitriolum martis*.

<sup>2</sup> This salt, which is known in commerce by the name of *green vitriol*, is prepared on the great scale from native sulphurets of iron, by exposing them to the



to the atmosphere, it attracts oxygen, becomes turbid, a Sub-sulphate is precipitated and the salt obtained is an Oxysulphate.

*Qualities.*—Sulphate of Iron has a strong, styptic taste: it crystallizes in transparent, rhomboidal prisms, of a fine-green colour, which redden the vegetable blues; are soluble in two parts of water at 60° and three-fourths of their weight of boiling water, and are insoluble in alcohol. When exposed to the air, the crystals become opaque and are covered with a yellow powder, owing to the attraction of the oxygen of the atmosphere by the salt, during its efflorescence. Exposed to heat, sulphate of iron undergoes the watery fusion; and in an increased heat the acid, partly in the state of what is termed *Glacial oil of Vitriol*, is driven off, and the base remains in the state of a red oxide, the *colcothar of vitriol* of commerce. According to Dr. Thomson<sup>1</sup>, 100 parts of the Green Sulphate consist of 26·7 of sulphuric acid, 28·3 oxide of iron, in the state of protoxide, and 45·0 of water.<sup>2</sup> Mr. Phillips makes the proportions to be of Acid 28·8, Protoxide of Iron 25·9 and Water 45·3. The following substances decompose sulphate of iron: the earths, the alkalies and their carbonates; lime-water, borate of soda, phosphate of soda, muriate of barytes, nitrate of silver, acetate of lead, and every salt the base of which forms an insoluble compound with sulphuric acid and soaps: thence these are incompatible in formulæ with this salt. It is also decomposed by all infusions of vegetable astringents.

*Medical properties and uses.*—Sulphate of Iron is tonic, emmenagogue, and anthelmintic.<sup>3</sup> It is a useful remedy when exhibited with due caution, in all cases in which preparations of iron are indicated; but in improper doses it occasions pain of the bowels, nausea, and vomiting, and often proves hurtful by being too long taken. It has been given with advantage in diabetes, in the latter stage of phthisis, and in amenorrhœa depending on a weakened action of the blood-vessels. The dose is from gr. j. to v. combined with ammoniacum, rhubarb, myrrh, or bitter extracts. It has lately been used dissolved in water as a lotion to cancerous and phagedenic ulcers.<sup>4</sup>

air and moistening them, till a crust of sulphate of iron is formed on their surface, which is afterwards obtained in crystals by solution and evaporation.

<sup>1</sup> *System of Chemistry*, 4th edit. iii. 225.

Of this quantity of water, 8 parts are water of composition, the oxide being in the state of a *hydrate*.

<sup>3</sup> It was used as an anthelmintic in the time of Pliny, who says, “Sumitur ad depellenda ventris animalia drachmæ pondere cum melle.” *Nat. Hist.* lib. xxxiv. cap. 12.

<sup>4</sup> *Edinburgh Med. and Surg. Journal*, ii. 373.



Officinal preparation. *Tinctura Ferri Muriatis*, D. *Ferri Subcarbonas*, L. E. *Mistura Ferri Composita*, L. *Pilulæ Ferri Composita*, L.

SULPHAS FERRI EXSICCATUS, Edin. *Dried Sulphate of Iron.*

“Take of Sulphate of Iron, *any quantity*. Let it be heated in an unglazed earthen vessel, on a moderate fire, until it become white and perfectly dry.”

SULPHAS FERRI EXSICCATUM, Dub. *Dried Sulphate of Iron.*

“Take of Sulphate of Iron *any quantity*. Let it be whitened by exposing it in an unglazed earthen vessel, to a high temperature.”

In these processes the degree of heat should not exceed  $212^{\circ}$  of Fahrenheit. The salt is merely deprived of its water of crystallization, without undergoing any chemical change.

Officinal preparation. *Oxydum Ferri rubrum*, E. D.

OXIDUM FERRI RUBRUM, Edin. *Red Oxide of Iron.*

“Let Dried Sulphate of Iron be exposed to a violent heat, until it is converted into a red-coloured substance.”

OXYDUM FERRI RUBRUM, Dub. *Red Oxide of Iron.*

“Let Dried Sulphate of Iron be roasted in a strong fire until it is converted into a red substance; then let this be washed till the water poured from it does not indicate, by the test of litmus, the presence of any Acid; and lastly, dry it upon bibulous paper.”

*Syn.* Oxide de fer rouge (*F.*), Eissenoxyd (*G.*), Perossido rosso di Ferro (*I.*).

By the degree of heat employed, the Sulphuric Acid of the Sulphate is partly driven off in a highly concentrated state, and partly decomposed, Sulphurous Acid being disengaged, and the Oxide more highly oxidized. The residue is the Red Oxide of Iron, combined with a portion of the Red Sulphate, which renders it deliquescent; and which should therefore be separated by washing, as directed by the Dublin College. According to Proust, the Red Oxide at the highest degree of oxidizement consists of 48 parts of Oxygen, and 52 of Iron.

*Medical properties and uses.*—This preparation is possessed of the same medicinal properties as the other oxides of iron.

Officinal preparation. *Murias Ammoniacæ et Ferri*, E. D.

FERRUM TARTARIZATUM, Lond. *Tartarized Iron.*

“Take of Iron, *a pound*; Supertartrate of Potass, in powder, *two pounds*; Water, *five pints, or a sufficient quantity*. Rub the Iron and the Supertartrate of Potass together, and expose the mixture in an open glass vessel with a pint of water, to the action of the air for twenty days, stirring them daily, and keeping the mass always moist by additions of distilled water. Then boil it in four pints of Distilled Water, for fifteen minutes, and filter the solution. Evaporate in a



water-bath, until the Tartarized Iron be perfectly dried; reduce it to powder, and preserve it in a stopped bottle.

TARTRAS POTASSÆ ET FERRI, Edin. *Tartrate of Potass and of Iron.*

“Take of Purified Filings of Iron, *one part*; Supertartrate of Potass powdered, *two parts*; Water, *one part*. Rub them together and expose them to the air in a shallow earthen vessel for fifteen days, stirring the mass daily with a spatula, and keeping it moist by frequent additions of water. Then boil the whole for a short time in four times its weight of water, and pour off the solution from the other fæces. Evaporate the solution to dryness in a water-bath, and having rubbed the mass into powder, preserve it in a well-stopped bottle.”

TARTARUM FERRI<sup>1</sup>, Dub. *Tartar of Iron.*

“Take of Carbonate of Iron, *half an ounce*; Crystals of Tartar in very fine powder, *an ounce*; Distilled Water, *a pint*. Let them be put into a glass vessel, then boiled for an hour over a slow fire, and the liquor filtered through paper; when this is cold, filter it again and evaporate it until a pellicle appear on the surface: the liquor, as it cools, will concrete into a saline mass, which is to be reduced to powder and preserved in closely stopped phials.”

*Syn.* Tartrate de Fer et de Potasse (*F.*) Eissen Weinstein (*I.*), Tartrato di Potassa e di Ossido di Ferro (*G.*).

Of these three processes, that of the Dublin College is to be preferred, as it affords a perfect triple salt; whereas much of the iron employed in the London and the Edinburgh processes remains unaltered, or is at least only in the form of a simple oxide. In the London process, the Iron is first oxidized to a protoxide by the partial decomposition of the water, aided by the action of the air, and the oxide thus formed unites with the superabundant acid of the Supertartrate of Potass: hence the dried mass consists of tartrates of Potass and of Iron, mixed with oxide of iron, and, when it is carelessly prepared, some metallic iron. The subsequent addition of water and re-exposure to the air, are intended to render the oxidizement complete, and convert the whole to the state of the triple salt; but, as this is not effected, it is probable that the proportion of Supertartrate of Potass ordered, is insufficient for the large quantity of the metal directed to be used. In the Dublin process, the superabundant acid of the Supertartrate of Potass dissolves as much of the Oxide of the Carbonate of Iron as it can take up; and a clear solution is ob-

---

<sup>1</sup> It is remarkable, that both the London and Dublin Colleges should err in giving a name to this triple salt so dissonant to the principles of the reformed nomenclature of their pharmacopœias.



tained, which, by evaporation, yields a true Tartrate of Potass and of Iron. As it is almost impossible to procure this salt in crystals, the solution may be evaporated to dryness.

*Qualities.* — Tartrate of Potass and of Iron is inodorous, has a slightly styptic taste, and is of a brownish-green colour. It is very soluble in water; and deliquesces, in some degree, when exposed to the air, so as to require to be kept in closely stopped phials. The cold solution of the alkalies and their subcarbonates do not decompose this salt; but it is instantly decomposed when boiled with any one of them except ammonia and its subcarbonate, which in neither state affect it. The strong acids, lime-water, hydro-sulphuret of potass and infusions of astringent vegetables decompose it, and are, therefore, incompatible in formulæ with it.

*Medical properties and uses.* — This salt possesses the same medicinal powers as the other preparations of Iron; but from its mildness, slight taste and ready solubility, it is a more convenient form for the administration of Iron to children; and in many cases in which the other saline preparations of it prove nauseating, and sit uneasy on the stomach. It is advantageously given in all the cases in which chalybeates prove useful: and is also extolled as a remedy in dropsy, in which it is supposed to exert both a diuretic and a tonic power. The dose is from grs. x. to ʒss., given either in a state of solution, or in the form of powder or bolus, combined with an aromatic, or a bitter.

LIQUOR FERRI ALKALINI. Lond. *Solution of Alkaline Iron.*<sup>1</sup>

“Take of Iron, *two drachms and a half*; Nitric Acid, *two fluid ounces*; Distilled Water, *six fluid ounces*; Solution of Sub-Carbonate of Potass, *six fluid ounces*. Mix together the Acid and the Water, pour the mixture over the Iron, and when the effervescence has ceased, pour off the acid solution. Add this gradually, and at intervals, to the solution of Sub-carbonate of Potass, frequently agitating until it become of a brownish-red colour, and no more effervescence is excited. Finally, set it aside for six hours, and pour off the liquor.”

*Syn.* Teinture martiale alkaline de Stahl (*F.*) Tintura di marte alkalina de Stahl (*I.*).

Although this composition has been long known, yet it is not well understood. The Diluted Acid<sup>2</sup> acts violently upon the iron and oxidizes it, while heat is evolved, and red fumes are extricated, consisting of nitrous gas and nitrous oxide gas. If this action be moderated by the iron being in a lump,

<sup>1</sup> This name has been justly criticised, as implying an unknown substance, “alkaline iron;” it should have been, *Liquor alkalinus ferri oxidi*.

<sup>2</sup> The concentrated acid scarcely acts on iron.



and putting the vessel in which it is dissolving into cold water, the solution is of a pale-green colour, and the Iron is at the minimum of oxidizement; but if heat be applied, or the effervescence be allowed to proceed with violence, the solution is of a reddish-brown colour, and contains Oxynitrate of Iron; a great excess of acid being present in both cases. On each addition of the solution of the Subcarbonate of Potass to either of these solutions, effervescence is excited by the disengagement of Carbonic Acid, and a red precipitate is instantly produced, which is, however, kept suspended by agitation, and ultimately re-dissolved by the excess of alkali. By allowing the mixture to stand for six hours as directed, particularly if the weather be cold, the whole becomes involved in a spongy mass of acicular crystals of Nitrate of Potass, from which the alkaline, metallic solution is to be poured off: it is clear, and of a deep brownish-red colour, if the acid solution contain the metal at the minimum of oxidizement: but if at the maximum it is turbid and of a redder hue. The first of these is the preparation of the London College.

*Qualities.*— This preparation has an alkaline, slightly styp-tic taste, and excites the sensation of coldness in the mouth produced by Nitrate of Potass, but in an inferior degree. The metallic part is precipitated by water<sup>1</sup>; and the clear supernatant fluid, when evaporated, yields crystals of Nitrate of Potass, proving that the solution contains this salt mixed with the Alkaline Oxide of Iron, if not a Nitrate of Potass and of Iron. On the addition of alcohol to this solution, the whole of the solid ingredients are thrown down in a con-creted mass; it is also decomposed by the strong acids and the pure alkalies; and after being kept for some time, if not very well excluded from the air, it becomes gradually turbid, perhaps owing to the abstraction of oxygen from the atmosphere more completely oxidizing the metal, and much of the oxide is deposited.

*Medical properties and uses.*— This solution very probably agrees with the other preparations of Iron in its medicinal properties; but, setting aside the difficulty of procuring it always of an uniform strength, and the many circumstances in conducting the process, that may alter altogether the nature of the product, we do not know in what mode it can be given; as water, and consequently all infusions and decoctions, decompose it, and precipitate the metal. It is, therefore, difficult to conceive for what purpose it has been introduced into the London Pharmacopœia.

---

<sup>1</sup> This precipitate, when the iron has been rapidly dissolved in the acid and heat employed, is a combination of peroxide of iron and carbonic acid. It effervesces strongly with muriatic acid, and gives off carbonic acid during the solution.



**TINCTURA FERRI AMMONIATI**, Lond. *Tincture of Ammoniated Iron.*

“Take of Ammoniated Iron, *four ounces*; Proof Spirit, *a pint*. Digest and filter.” This being merely a spirituous solution of Ammoniated Iron, it seems to be unnecessarily introduced into the Pharmacopœia by the London College.

**TINCTURA FERRI MURIATIS**, Lond. *Tincture of Muriate of Iron.*

“Take of Carbonate of Iron, *half a pound*; Muriatic Acid, *a pint*; Rectified Spirit, *three pints*. Pour the Acid over the Carbonate of Iron, in a glass vessel, and shake them occasionally for three days. Set apart the liquor, that the fæces, if there be any, may subside; then pour off the solution, and add the spirit to it.”

*Edinburgh.*

“Take of Black Oxide of Iron, purified and reduced to powder, *three ounces*; Muriatic Acid, about *ten ounces*, or as much as may be sufficient to dissolve the powder. Digest with a gentle heat, and the powder being dissolved, add as much alcohol as will make the whole liquor amount to two pounds and a half.”

*Dublin.*

“Take of Rust of Iron, *half a pound*; Muriatic Acid, *three pounds*; Rectified Spirits of Wine, *three pints*. Put the Rust into a glass vessel, pour on the Acid, and agitate it occasionally; then set it aside that the fæces may subside, and pour off the liquor: evaporate this slowly to *one pint*, and when it is cold, add the spirit.”

**TINCTURA MURIATIS FERRI CUM OXIDO RUBRO**, Dub. *Tincture of Muriate of Iron, with the Red Oxide.*

“Take of Red Oxide of Iron, *an ounce*; Muriatic Acid, *four fluid ounces*; Rectified Spirit of Wine, *a sufficient quantity*. Digest the Oxide with the Acid for twenty-four hours; then boil the solution for half-an-hour: evaporate the filtered solution to the thickness of honey, and, when it is cold, add Rectified Spirits of Wine, with frequent agitation, until the specific gravity of the tincture be, to that of distilled water, as 1050 to 1000.”

*Syn:* Koch salzaure Eissentinktur (*G*), Tinctura di muriato di Ferro (*I*).

Of the formulæ given for the preparation of this tincture, those of the London and the Dublin Pharmacopœias are to be preferred. The metal, as ordered in them, is in a higher state of oxidizement, and forms at once, by its combination with the acid, an uniform compound soluble in alcohol; whereas, by following the Edinburgh process, the solution is a mixture of the above Muriate, and of the less soluble or green Mu-



riate<sup>1</sup>, the Black-oxide not being all completely oxidized; and it is not till after the exposure to air, and by attracting oxygen, that it is converted altogether into the more soluble Muriate. Hence the Edinburgh preparation cannot be always of an uniform and fixed strength, which, for an active medicine, is a matter of much importance. The introduction of the last preparation by the Dublin College is superfluous. When it is properly prepared according to the London formula, Mr. Phillips found that its spec. grav. is 0·994, and it contains 16·8 grains of Peroxide of Iron in half a fluid ounce.<sup>2</sup>

*Qualities.* — The alcoholic solution of Muriate of Iron is of a brownish-yellow colour, has a peculiar ethereal odour, and a very styptic taste. It contains the iron in the state of a chlorate; and when it is distilled, a black oxide of iron remains in the retort. With the alkalies and their carbonates it gives a red precipitate; strikes a black colour with infusions of astringent vegetables; and forms with mucilage of acacia gum an orange-coloured jelly. Hence these substances cannot enter into compositions with this tincture.

*Medical properties and uses.* — This is an active and elegant preparation of iron, well adapted for all the diseases in which chalybeates prove serviceable. I have found it more useful than any other tonic, in scrophula; when it is given in doses, gradually increased, until one hundred and twenty drops be taken for a dose, twice a day. It is also recommended in dysury depending on spasmodic stricture of the urethra, in which case it is given in doses of five or six drops, repeated every ten or fifteen minutes, until nausea be induced. It is used externally, as a styptic in cancerous and loose fungous sores. The usual dose is from ℥x. to ℥xxx. in a glass of water: but it may be gradually increased to ℥cxxx.

ACETAS FERRI, Dub. *Acetate of Iron.*

“Take of Carbonate of Iron, *half an ounce*; Acetic Acid, *three fluid ounces*. Digest for three days and filter.”

*Syn.* Acetate de Fer (*F.*); Acetato di Ferro (*I.*).

This preparation is a mild and efficacious chalybeate: but if the variety of forms in which iron is ordered to be prepared for medicinal purposes be considered, it will obviously appear to be superfluous.

TINCTURA ACETATIS FERRI, Dub. *Tincture of Acetate of Iron.*

“Take of Acetate of Kali, *two ounces*; Sulphate of Iron, *one ounce*; Rectified Spirit of Wine, *two pints*. Rub together the

<sup>1</sup> When iron filings are dissolved in muriatic acid, and completely excluded from the air, a muriate is formed insoluble in spirit of wine. *Davy's Researches*; p. 180.

<sup>2</sup> *Trans. of the Pharm.* p. 100.



Acetate of Kali, and the Sulphate of Iron, in a stone-ware mortar, until they unite into a soft mass; dry this with a moderate heat, and triturate it with the spirit; then put the mixture into a phial; cork it closely, and digest for *seven days*, frequently shaking it: finally, pour the clear tincture from off the *fæces*."

TINCTURA ACETATIS FERRI CUM ALCOHOLE, Dub. *Tincture of Acetate of Iron with Alcohol.*

"Take of Sulphate of Iron, Acetate of Kali, each *an ounce*; Alcohol, *two pints*. Rub together the Acetate of Kali and the Sulphate of Iron in a stone-ware mortar, until they unite into a soft mass; then dry this with a moderate heat, and when it is cold, triturate it with the alcohol. Put the mixture into a phial, cork it closely, and digest for *twenty-four hours*, frequently shaking it: finally pour the clear tincture from off the *fæces*."

*Syn.* Teinture de l'Acetate de Fer (*F.*), Tinturo di marte astringente (*I.*).

These two preparations differ in scarcely any thing except in strength; the theory of the formation of the Acetate of Iron being the same in both cases. During the process a double decomposition takes place; the Sulphuric Acid of the Sulphate of Iron leaves the Iron and unites with the Potass of the Acetate of Potass, while the disengaged Acetic Acid of the latter salt combines with the Iron, forming Acetate of Iron, which is soluble in the Alcohol. It is also probable that the oxide of Iron absorbs oxygen during the trituration, and being thus more completely oxidized, the mass must contain, instead of an Acetate, an Oxyacetate of Iron, which is more readily dissolved in the alcohol. The Sulphate of Potass remains undissolved in both processes, with a small portion of uncombined oxide of iron, in the form of a brownish precipitate.

*Qualities.* — These tinctures have a peculiar odour, a warm styptic taste, and a reddish-brown colour. They are decomposed by the alkalies and their carbonates, and the strong acids, and by infusions of astringent vegetables, which are, therefore, incompatible in formulæ with them.

*Medical properties and uses.* — These spirituous solutions of Acetate of Iron possess the same properties as the other preparations of this metal; but if the introduction of the simple Acetate be superfluous, the double form of its spirituous solution is still more objectionable. Indeed every advantage that can be expected from any of these forms of the acetate can be equally obtained from the Tartrate of Iron and of Potass (*ferrum tartarizatum*). And we cannot conceive that any particular benefit can result to practice from loading the list of remedies with all the multifarious states of combination of which the same substance is susceptible. The dose of either of these



tinctures may be from  $\mathfrak{m}\mathfrak{x}$ . to  $\mathfrak{f}\mathfrak{z}\mathfrak{j}$ . given in a sufficient quantity of water, or any other proper vehicle.

VINUM FER'RI, Lond. *Wine of Iron.*

“Take of Iron, *one drachm* ; Supertartrate of Potass powdered, *six drachms* ; Distilled Water, *two pints, or as much as will be required* ; Proof Spirit, *twenty fluid ounces*. Rub together the Iron and the Supertartrate of Potass, and expose them in an open glass vessel, with *one fluid ounce of water*, to the air for six weeks, stirring daily with a spatula; and adding frequently as much distilled water as may be necessary to keep the mass moist. Then dry it with a gentle heat; rub it to powder, and mix it with *thirty fluid ounces* of distilled water. Filter the solution and add the spirit.”

*Dublin.*

“Take of Iron-wire cut in pieces, *four ounces* ; White Rhenish Wine *four pints*. Sprinkle a little of the wine over the iron filings, and expose them to the air, until they be covered with rust, then add the remainder of the wine; digest for seven days, frequently agitating; and, lastly, filter.”

*Syn.* Vinum chalybeatum, P. L. 1745. Eissenwein (G.).

The Iron is oxidized and dissolved in the Acid of the Supertartrate of Potass. It is, therefore, a solution of Tartrate of Iron and of Potass, with an excess of Supertartrate of Potass: and when prepared in the mode ordered by the London College, each pint should contain about 22 grains of Peroxide of Iron; but owing to a portion of the tartarized iron being rendered insoluble by drying, and the precipitation by the spirit of a part of what was dissolved by the water, the usual quantity is grs. xvi. of peroxide of iron. The strength of the Dublin preparations must altogether depend upon the state of the wine: and it is to be regretted that the college did not order a given portion of *ferrum tartarizatum*, which readily dissolves in wine, and forms a permanent solution.

*Medical properties and uses.* — This is the least unpleasant of the preparations of iron. It is chiefly employed in chlorosis, and the relaxed habits of young females. The dose is from  $\mathfrak{f}\mathfrak{z}\mathfrak{j}$ . to  $\mathfrak{f}\mathfrak{z}\mathfrak{vj}$ . given twice or three times a day. As far as regards the formula of the London College, it is absurdly denominated a wine.

TABLE SHOWING HOW MUCH OF EACH PREPARATION OF IRON OF THE LONDON PHARMACOPŒIA CONTAINS ONE GRAIN OF OXIDE.

Ferri Sulphatis.....	gr. 5·8	Liquoris Ferri Alkalini .....	$\mathfrak{m}\mathfrak{x}\mathfrak{x}\mathfrak{x}$ .
Ferri Subcarbonatis.....	1·2	Misturæ Ferri compositæ.....	$\mathfrak{f}\mathfrak{z}\mathfrak{j}\mathfrak{s}\mathfrak{s}$ .
Ferri Ammoniaci .....	66·0	Tincturæ Ferri Ammoniaci.....	$\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$ .
Ferri Tartarizati.....	5·0	Tincturæ Ferri Muriatis.....	$\mathfrak{m}\mathfrak{x}\mathfrak{i}\mathfrak{v}$ .
Pilulæ Ferri compositæ.....	20·0	Vini Ferri.....	$\mathfrak{f}\mathfrak{z}\mathfrak{j}$ . <sup>1</sup>

<sup>1</sup> *Phillips's Trans. of the Pharm.* 1824.



## PRÆPARATA EX HYDRARGYRO.

## PREPARATIONS OF MERCURY.

HYDRAR'GYRUM CUM CRETA, Lond. *Mercury with Chalk.*<sup>1</sup>

“Take of Purified Mercury (by weight), *three ounces*; prepared Chalk, *five ounces*. Rub them together until globules are no longer visible.”

HYDRARGYRUM CUM CRETA. Dub. *Mercury with Chalk.*

“It is prepared in the same manner as the Mercury with Magnesia, only instead of Magnesia employing precipitated Chalk.”

In these processes the mercury is generally supposed to be oxidized during the trituration, and in the state of the protoxide, 100 parts of which, according to Fourcroy, contain, when well prepared, about 4 of oxygen: but Mr. Phillips justly says, that the mercury in this preparation is not a protoxide, being insoluble in acetic acid: but he adds, that when the chalk is taken up by acetic acid the mercury does not run together, but remains in minute globules, a circumstance which induces the belief that it is not merely mechanically divided. Eight grains contain three of mercury.

*Medical properties and uses.*—It is alterative, and is occasionally prescribed in mesenteric affections, in porrigo, and other cutaneous affections. It merits attention as a mild alterative for children. The dose may be from grs. v. to ʒss. given twice a day, mixed in any viscid substance.

HYDRARGYRUM CUM MAGNESIA, Dub. *Mercury with Magnesia.*

“Take of Mercury, Manna, of each *an ounce*; Magnesia, *half an ounce*. Triturate the Mercury with the Manna in an earthen mortar, adding as many drops of water as will give to the mixture the thickness of syrup, and continue the rubbing until the metallic globules completely disappear; then add, still triturating, *a drachm* of Magnesia; and after the whole is well mixed together, add *a pint* of hot Water, and agitate the mixture. Allow the mixture to remain for some time at rest, in order that the sediment may subside, from which the fluid is to be decanted. Repeat the washing a second and a third time, that the whole of the manna may be removed; and add the remainder of the Magnesia to the sediment while

<sup>1</sup> Mercurius alkalizatus, P. L. 1745.



it is still moist. Finally dry the powder upon bibulous paper."

The addition of the manna in this process, and in the former preparation with chalk of the Dublin College, is intended only to facilitate the oxidizement of the mercury; and, therefore, it is afterwards removed by the subsequent washings, so that the product is supposed to remain a grey or black oxide of mercury, mixed with magnesia. It is a preparation which might well be rejected.

HYDRAR'GYRI NITRICO-OX'YDUM, Lond.<sup>1</sup> *Nitric Oxide of Mercury.*

"Take of Purified Mercury (by weight), *three pounds*; Nitric Acid (by weight), *a pound and a half*; Distilled Water, *two pints*. Mix them in a glass vessel, and boil until the Mercury be dissolved and a white mass remain after the evaporation of the Water. Rub this into powder, and put it into another vessel very shallow; then expose it to a gentle heat and gradually raise the fire until it cease to emit red vapours.

OXYDUM HYDRARGYRI RUBRUM PER ACIDUM NITRICUM, Edin. *Red Oxide of Mercury by Nitric Acid.*

"Take of Purified Mercury, *three parts*; Diluted Nitrous Acid, *four parts*. Dissolve the Mercury, and evaporate the solution over a gentle fire to a white, dry mass, which, being rubbed to a powder, is to be put into a glass cucurbit, and covered with a thick plate of glass. Then adapt a capital to the vessel; and, having placed it in a sand-bath, let the contained matter be roasted with a fire gradually raised until it pass into small very red scales."

OXYDUM HYDRARGYRI NITRICUM, Dub. *Nitric Oxide of Mercury.*

"Take of Purified Mercury, *ten ounces*; Diluted Nitrous Acid, *ten fluid ounces*. Let them be mixed in a glass, and the Mercury dissolved with a gradually raised heat; then increase the fire until the residuary matter in the bottom of the vessel be converted into red scales."

*Syn.* Oxyde mercure rouge par l'acide nitrique (*F.*), Rother präcipitatur (*G.*) Mercurio Precipitato Rosso (*I.*).

In this process, the mercury is first oxidized at the expense of part of the acid employed; and the oxide, which is in a high state of oxidizement, combines with the undecomposed acid, so as to form a nitrate of mercury. By augmenting the heat, this nitrate is decomposed, the acid and water are nearly altogether expelled and the oxide is left of a bright-red colour, combined with a small portion of acid. However simple the

<sup>1</sup> Hydrargyrus nitratus ruber, P. L. 1787.



process may appear to be, yet it has been always found difficult to produce the bright-red scaly appearance, which the product should have when it is properly prepared. Much of the success of the process appears to depend on the purity of the acid; the proper regulation of the heat, which, at the utmost, should not be  $600^{\circ}$ <sup>1</sup>; and the scale on which it is formed, the heat being more steadily maintained, and acting with more uniformity, on a large than on a small quantity of materials. Hence the Red Precipitate prepared in Holland, where it is manufactured largely, has always been considered better than any prepared in this country. The proportions used by the Dutch chemists are fifty pounds of pure mercury, and seventy of pure nitrous acid of a specific gravity 1.3. The decomposition is conducted in very large flat vessels, the fire being raised when the gaseous nitrous acid ceases to be sensibly disengaged; and the test of its perfection is the inflammation of a match which has been just blown out, by introducing it into the vapour arising from the decomposing oxide.<sup>2</sup>

*Qualities.* — When properly prepared, this is a peroxide mixed with some nitrate of mercury. It is in small scales of a bright-red colour, very acrid and corrosive: insoluble in water, but totally soluble in nitric acid without effervescence. It is completely volatilized in a red heat, and at the same time decomposed. We have found the observation of Dr. Murray correct, that “if the preparation be boiled for a short time with five or six times its weight of water, the liquor, when filtered, has the styptic, metallic taste, and gives a white precipitate with water of ammonia or carbonate of potass; a plain proof that it holds dissolved nitrate of mercury.”<sup>3</sup> According to Payssè, 100 parts contain 82 of mercury, and 18 of oxygen: but according to Mr. Phillips<sup>4</sup>, the proportions are mercury 92.6, oxygen 7.4. It is sometimes adulterated with red oxide of lead, which may be detected by dissolving one part of the oxide in four of acetic acid: if lead be present, the solution has a sweetish taste; and when sulphuretted water is dropped into it, a dirty, dark precipitate is thrown down. When pure it is perfectly volatilized when thrown on a red hot iron.

*Medical properties and uses.* — Nitric oxide of mercury is stimulant and escharotic. It is an external application only,

<sup>1</sup> Higgins, *Essays*, i. 133.

<sup>2</sup> M. Payssè, *Annales de Chimie*, li. 202.

<sup>3</sup> *System of Materia Medica*, ii. 329. Dr. Murray suggests, that it should have been named *Subnitrus Hydrargyri ruber*, but improperly, as the undecomposed salt which it contains is a nitrate.

<sup>4</sup> *Trans. of Pharm.* p. 110.



being used, when rubbed into a fine powder, as a stimulant to old sores, and for destroying fungus. As a powder, in the proportion of gr. ss. to grs. iv. of sugar, it is blown into the eye to remove specks on the cornea; and formed into an ointment with lard, it is an useful application to ulcerations of the eyelids, and to chancres.

Officinal preparation. *Unguentum Hydrargyri Nitrico-oxidi*, L. E. D.

ACETAS HYDRARGYRI, Edin. *Acetate of Mercury.*

“Take of Purified Mercury, *three ounces*; Diluted Nitrous Acid, *four ounces and a half, or a little more than may be required for dissolving the mercury*; Acetate of Potass, *three ounces*; Boiling Water, *eight pounds*. Mix the Mercury with the Acid; and, towards the cessation of the effervescence, digest, if necessary, until the Mercury be completely dissolved. Then dissolve the Acetate of Potass in the Boiling Water; and immediately to this solution, still hot, add the former, and mix them together by agitation. Set the mixture aside to crystallize; then wash the crystals placed in a funnel with cold Distilled Water; and finally dry them with a very gentle heat.

“In preparing Acetate of Mercury, it is necessary that all the vessels which are used, and the funnel, be of glass.”

ACETAS HYDRARGYRI, Dub. *Acetate of Mercury.*

“Take of Purified Mercury, *three ounces*; Diluted Nitrous Acid, *three fluid ounces*; Acetate of Kali, *three ounces*; Boiling Distilled Water, *eight pints*. Add the Acid to the Mercury, and when the effervescence is over, digest upon hot sand, that the metal may be dissolved; mix this solution immediately with the Boiling Water in which the Acetate of Kali has been previously dissolved, and then let the mixture be passed as quickly as possible through a double linen cloth. Cool it, that crystals may form; wash these with cold Distilled Water, and dry them upon paper with a very gentle heat.

“In the whole of this process glass vessels must be used.”

*Syn.* Acetate de Mercure (F.), Essigsaures Quecksilber (G.), Acetato di Mercurio (I.).

Acetic acid scarcely acts on mercury, but by either of the above processes the Acetate may be formed. Nitrate of mercury is first obtained by the action of the nitric acid on the mercury; and this is decomposed by the acetate of potass, the alkali of which unites with the acid of the metallic salt, and forms nitrate of potass, which remains in solution; while the disengaged acetic acid combines with the oxide of mercury, and forms the Acetate of Mercury, which readily crystallizes and is thus easily separated. By preparing the solution of the nitrate of mercury with a gentle heat, when there is an



excess of acid, the portion of mild Acetate of Mercury produced is considerable; but if the quantity of acid be sufficient for the saturation only of the oxide, a sudden decomposition of the solution is effected by the hot water which contains the acetate of potass in solution, independent of the action of the acetate, and a subnitrate of mercury of a yellow colour is precipitated. Hence the propriety of the direction of the Edinburgh College to use more acid “than is required for dissolving the mercury.” It is of much importance also that the degree of heat be low; for if a high temperature be employed, the metal is oxidized to a maximum, and the product of the subsequent part of the process is an oxyacetate, which is very acrid and soluble, instead of the salt intended to be produced. For the success of the process, which often fails, the solution of the acetate of potass should not be used immediately after it is made, but should be scarcely more than tepid when it is mixed with the solution of nitrate of mercury; and to the water employed for washing the salt, should be added about fʒj. of distilled vinegar for every Oss. of water; which prevents the partial decomposition of the acetate, and the consequent yellow colour of the crystals, that sometimes occur in the washing.

*Qualities.* — This salt, when properly prepared, is in small flat crystals of a silvery whiteness, acrid to the taste, soluble in hot water, but scarcely soluble in cold water. It is insoluble in alcohol. The alkalies decompose it, and it is readily decomposed by heat. Light also has this effect, blackening the salt. The proportion of its constituents has not yet been ascertained.

*Medical properties and uses.* — Acetate of Mercury is anti-syphilitic and alterative; but it is scarcely ever used, unless as the active ingredient of Keyser’s pills. As an external application, a solution of it in the proportion of grs. ij. in fʒij. of rose-water, is used in some cutaneous affections. The internal dose is gr. j., night and morning.

HYDRARGYRI OXYDUM CINEREUM, Lond.  
*Grey Oxide of Mercury.*

“Take of Submuriate of Mercury, *an ounce*; Lime-Water, *a gallon*. Boil the Submuriate of Mercury in the Lime-Water, stirring it assiduously, until the grey oxide of mercury subsides. Wash this with distilled water, and then dry it.”

OXYDUM HYDRARGYRI CINEREUM, Edin. *Grey Oxide of Mercury.*

“Take of Submuriate of Mercury, *half an ounce*; Lime-Water, *five pounds*. Boil the Submuriate in the solution, for a quarter of an hour in a slightly covered vessel. Pour off



the supernatant fluid, then wash the oxide with distilled water; and dry it."

PULVIS HYDRARGYRI CINEREUS, Dub. *Grey Powder of Mercury.*

"Take of Mercury, *two ounces*; Diluted Nitrous Acid, *two fluid ounces*; dissolve the Mercury in a slow heat, and dilute the solution with *eight fluid ounces* of Cold Water; then drop into it *one ounce and a half* of the Water of Carbonate of Ammonia, or as much as may be sufficient for precipitating the whole of the metal, which is to be washed with boiling Distilled Water, until the poured-off fluid yield no sediment when Water of Sulphuret of Ammonia is dropped into it. Finally, let the precipitate be dried."

*Syn.* Oxide gris de Mercure (*F.*), Schwarzes gesauertes Quecksilber (*G.*), Protossido cinereo di Mercurio (*I.*).

The appellations given to these preparations would lead to the supposition that they were essentially the same, although the Dublin process differs from the other two; and scarcely any difference, indeed, exists between the products of the three processes, when they are properly conducted. In the London and Edinburgh processes, the lime-water decomposes the calomel; its lime unites with the acid of the mild muriate, and the insoluble protoxide remains of a greyish colour, while the muriate of lime which is formed, being soluble, is easily separated by washing: but if we admit the calomel to be a proto-chloride of mercury, we must admit that the water of the lime-water is decomposed, and its hydrogen uniting with the chlorine, forms muriatic acid, which converts the lime into a muriate, while its oxygen changes the mercury into the protoxide. In the Dublin process it is intended, first, to produce a nitrate of mercury with the metal, at a low state of oxidizement; so that by the addition of the carbonate of ammonia a decomposition may be effected, and grey oxide of mercury, and nitrate of ammonia formed, which are to be separated by the subsequent washings. But if the nitrate of mercury be formed, with the assistance of heat, as ordered by the Dublin College; or even if the solution be quickly made without heat, the metal becomes too highly oxidized, and the result is not the grey oxide, which the College intends should be produced, but is a mixture of the grey oxide, and a triple compound of oxide, mercury, ammonia, and nitric acid.<sup>1</sup> The directions of this formula are not sufficiently distinct to produce the effect intended; and, therefore, the following directions given by Hahneman for this preparation are absolutely necessary to be followed for obtaining the grey oxide in a

<sup>1</sup> *Gren's Chemistry*, (translation) ii. 230.



purser form. Dilute the acid with two parts of water, and add the mercury in small quantities at a time, placing the vessel in cold water to moderate the rise of temperature during the solution, which thus proceeds very slowly. When the acid has taken up as much of the metal as will saturate it, dilute the solution with twenty parts of distilled water, and drop in solution of ammonia as long as any precipitate is produced. Wash the precipitate immediately in water, and dry it on bibulous paper before the fire.<sup>1</sup> The same effect is produced if subcarbonates of ammonia be used, the carbonic acid being disengaged.

*Qualities.* — Grey Oxide of Mercury, properly prepared, is in the form of an impalpable, blackish-grey coloured powder, which becomes paler if exposed to air and light. It is inodorous, insipid and insoluble in water. In the state in which it is usually found in the shops, it is of a light grey colour, almost approaching to a white, in which state it contains the triple salt above mentioned. When prepared according to the London formula, it has been supposed<sup>2</sup>, that the lime not being able to abstract the whole of the acid, the product is strictly a submuriate of mercury mixed with the protoxide. We find from experiment, that this is actually the case, and it is not improbable that a minute portion of the lime may also be precipitated in the state of carbonate. The constituents of the protoxide are 96 parts of mercury, and 4 of oxygen, in 100 parts.<sup>3</sup> It ought to dissolve totally in acetic acid and be insoluble in muriatic; if it contain a peroxide of mercury, therefore, the muriatic acid will dissolve it: and this salt is also detected by Liquor Ammonia producing a white precipitate, and Liquor Potasse a yellow one.

*Medical properties and uses.* — The Grey Oxide of Mercury, when well prepared, may be used as a substitute for the oxide prepared by trituration; and as it is more likely to be always of an uniform strength, it may of course be more depended on than that preparation. It has been, however, objected to for forming ointment for the purposes of mercurial frictions; (see *Ung. Oxydi Hydrargyri cinerei*) — but perhaps the objections have originated from that form of the preparation having been used which contains the triple salt. We have seen it used with advantage for fumigation, both locally applied to assist the healing of venereal ulcers; and, generally, to bring the habit under the influence of mercury, when it could not be introduced by the ordinary mode. The dose of this Oxide is from gr. i. to grs. iij. given in the form of pill twice a day.

<sup>1</sup> Murray's *Chemistry*, 2d edit. iii. 178.

<sup>2</sup> Murray's *System of Materia Medica*, ii. 326.

<sup>3</sup> Fourcroy.



Official preparations. *Unguentum Oxidi Hydrargyri cinerei*, E.  
**HYDRARGYRI OXYDUM RUBRUM**, Lond.<sup>1</sup> *Red Oxide of Mercury.*

“Take of Purified Mercury (by weight), *a pound*. Put the Mercury into a tall glass vessel with a narrow mouth and broad at the bottom. Expose this vessel open to a heat of 600°, until the Mercury be converted into red scales; then rub these to a fine powder.”

OXYDUM HYDRARGYRI, Dub. *Oxide of Mercury.*

“Take of Purified Mercury, *any quantity*. Put it into an open, glass vessel with a narrow mouth and a broader bottom, and expose it to a heat of about 600°, until it be converted into red scales.”

*Syn.* Oxide de Mercure rouge par le feu (*F.*), Roth's Quecksilberoxyd (*G.*), Perossido rosso di Mercurio (*I.*).

In this process the mercury is brought nearly to the boiling point<sup>2</sup>, so as to be volatilized, in which state it decomposes atmospherical air attracting its oxygen, and is converted into a red oxide. A small quantity of mercury requires several weeks to be thus oxidized; and, therefore, as much only is introduced into the vessel as can cover its bottom; and both on this account, and in order to prevent the dissipation of the volatilized metal, the shape of the vessel is of some importance. It should have a wide bottom and a long neck, the extremity of which is extended almost to a point; and it should be heated in a sand-bath, the sand not rising higher round the vessel than the mercury stands within it. By maintaining a steady heat, a constant circulation of the mercurial vapour is kept up in the upper part of the matrass, and as it combines with oxygen, a dull film first forms on the surface of the mercury, which is next converted into a black powder, and then into red shining scales. A part of the mercury is always lost; and as the process requires much attention, and so long a time for its completion, the preparation is necessarily expensive.

*Qualities.*—*Red-oxide of Mercury* is obtained in the form of minute, crystalline, very brilliant, sparkling, deep-red scales, inodorous, but acrid and caustic, although less so than the former preparation. It is a peroxide, is soluble in several of the acids without decomposition: potass and soda decompose the solution, and precipitate the peroxide of an orange colour. It is slightly soluble in water also, and the solution changes to green the syrup of violets. When rubbed with running mercury, both are changed into black oxide; and when heated to ignition in a glass retort, it is decomposed; very pure

<sup>1</sup> Hydrargyrus calcinatus, P. L. 1787.

<sup>2</sup> Irvine makes the boiling point of mercury to be 672°; Crichton, 655°; and Dalton, 660°.



oxygen being obtained, and the metal again returns to the state of running mercury. According to Lavoisier, 100 parts of this oxide contain 7 of oxygen; Mr. Phillips makes the proportion 7.4; Fourcroy 8; and Chenevix, 15 parts. The light partially decomposes the red oxide, hence it should be kept in opaque bottles.

*Medical properties and uses.* — This is a very active preparation of mercury, and has been employed by some very celebrated practitioners<sup>1</sup> as an internal remedy in syphilis. It is, however, very apt to occasion vomiting, purging, and otherwise violently to affect the stomach and bowels; consequently it is now scarcely ever exhibited internally, or employed as an antisymphilitic. The dose may be one-eighth of a grain to gr. j. combined with gr. ss. of opium, in the form of pill, night and morning. It is chiefly used as an external stimulant and escharotic, in the same cases as the nitric oxide; being previously rubbed to a fine powder, and either sprinkled over the ulcers; or united with lard, and applied as an ointment.

HYDRARGYRI OXYMURIAS, Lond. *Oxymuriate of Mercury.*

“Take of Purified Mercury (by weight), *two pounds*; Sulphuric Acid, *thirty ounces* (by weight); Dried Muriate of Soda, *four pounds*. Boil the Mercury with the Sulphuric Acid in a glass vessel, until the Sulphate of Mercury become dry: rub this, when it is cold, with the Muriate of Soda in an earthenware mortar; then sublime it in a glass cucurbit with a gradually raised heat.”

MURIAS HYDRARGYRI CORROSIVUS, Edin. *Corrosive Muriate of Mercury.*

“Take of Purified Mercury, *two pounds*; Sulphuric Acid, *two pounds and a half*; Dried Muriate of Soda, *four pounds*. Boil the Mercury with the Sulphuric Acid in a glass vessel placed in a sand-bath, until the matter becomes dry. Mix this, when it is cold, in a glass vessel, with the Muriate of Soda; then sublime in a glass cucurbit with a gradually raised heat. Separate the sublimed matter from the scorixæ.”

MURIAS HYDRARGYRI CORROSIVUM, Dub. *Corrosive Muriate of Mercury.*

“Take of Purified Mercury, *two pounds*; Sulphuric Acid, *three pounds*; Dried Muriate of Soda, *two pounds and a half*. Dissolve the Mercury in the acid, and gradually increase the heat until the matter become almost dry; let this when it is cold, be rubbed with the muriate of soda in an earthenware mortar; and then sublime it in a proper vessel with a gradually raised heat.”

<sup>1</sup> John Hunter.



*Syn.* Muriate de Mercure corrosif (*F.*), Azzendes alssaures Quecksilber (*G.*), Mercurio sublimato corrosivo (*I.*).

Sulphuric acid does not act upon mercury at a low temperature; but when three parts of this acid are boiled upon two of mercury, the metal decomposes the acid, and is oxidized, sulphurous gas being emitted with effervescence; and there remains a dry mass of a fine white colour, which is a hypersulphate of mercury combined with an excess of acid. By triturating this salt with dried muriate of soda (chloride of sodium), and exposing the mixture to heat, a double decomposition is effected. According to the old doctrines, the muriatic acid leaves the soda, and combines with the oxide of mercury of the hypersulphate, while the sulphuric acid unites with the soda, thus forming muriate of mercury and sulphate of soda, the former of which, being easily volatilized is separated from the latter by sublimation. But, according to the latest doctrines, the chlorine of the common salt leaves the sodium, and uniting with the mercury of the hypersulphate, forms a *bichloride* of mercury, which sublimes; while the oxygen of the oxide of mercury combining with the sodium converts it into soda, which unites with the sulphuric acid and forms sulphate of soda which remains in the bottom of the cucurbit. This process was first proposed by Kunkel, but no salt has been prepared by a greater variety of methods; and as it is now generally manufactured on the large scale, the proportions of the ingredients ordered by the Colleges are perhaps but seldom adopted. Of the three formulæ of the British Colleges, however, that of the Dublin College is to be preferred, as by the larger proportion of sulphuric acid, and the smaller of muriate of soda, a more complete decomposition of the muriate of soda is effected; and consequently, a greater quantity of chlorine being evolved, a larger proportion of the mercury must necessarily be converted into chloride. Sixteen ounces of mercury should yield about  $\frac{3}{4}$ xx. of corrosive muriate. The most simple process, and perhaps the best, is the direct solution of the red oxide of mercury in muriatic acid, by which the salt is obtained by spontaneous crystallization<sup>1</sup>; but it is too expensive for general purposes.

*Qualities.* — Corrosive Muriate of Mercury<sup>2</sup> is obtained by the above processes in the form of a white, shining, semi-trans-

<sup>1</sup> *Annales de Chimie*, xxviii. 12.

<sup>2</sup> This appellation, which is that of the Edinburgh and Dublin Colleges, is certainly preferable to that of the London College. But as the name *oxymuriate* is improper in a strictly chemical sense, perhaps the name *deutochloride*, or *perchloride of mercury*, which most justly designates its chemical character, would be less exceptionable than any other. The old names were *Hydrargyrus muriatus*, *Mercurius sublimatus corrosivus*.



parent, easily pulverized mass, made up of small, acicular crystals. When the process is very carefully and slowly conducted, the crystals are separate, regular, tetrahedrons, compressed and pointed. On exposure to the air the mass effloresces on the surface. Its specific gravity is 5.1398. It is inodorous, and has a very acrid, disagreeable, metallic taste and changes to green several of the vegetable colours; Water at 60° dissolves one-twentieth of its weight, at 212° one-third, and alcohol at 60° four parts. It is soluble also in the sulphuric, nitric, and muriatic acids, and may be again obtained unaltered by evaporating the solutions. The fixed alkalies and alkaline earths decompose it, precipitating it from its solution of an orange-yellow colour, which becomes brick-red. It is also partially decomposed by exposure to light, and by some metals; and changed into calomel.<sup>1</sup> The carbonates of the fixed alkalies precipitate it of a fixed, yellow hue, and ammonia forms with it a white, triple, insoluble compound, containing muriate of ammonia and oxide of mercury. When triturated with olive oil, the oil is whitened; and when boiled with it, a small portion of calomel is thrown down; and the same is the case when it is boiled with sugar. The volatile oils reduce it. It is also decomposed by solutions of tartrate of potass and antimony; nitrate of silver and acetate of lead; and forms precipitates in infusions and decoctions of the following vegetable substances; chamomile flowers, horseradish root, columba root, catechu, cinchona bark, rhubarb root, senna leaves, simaruba bark, oak bark, tea, and in the almond mixture: consequently, it is incompatible in extemporaneous formulæ with these substances. The components of this salt are 25.46 parts of chlorine, and 74.54 of mercury in 100 parts: taking the analysis of Zaboada, and correcting it so as to make it correspond with the real composition of the per-chloride.<sup>2</sup> According to Mr. Phillips, the proportions are of chlorine 26.48, of mercury 73.52 parts in 100.

*Medical properties and uses.* — This salt has been long known to chemists.<sup>3</sup> It is a powerful stimulant and alterative; and in large doses is one of the most violent of the metallic poisons. As an antisyphilitic it was early much extolled, and is the active ingredient of many a celebrated empirical nostrum; but modern practice has fixed its real merits much lower than they were formerly placed. When taken in over-doses, either by

---

<sup>1</sup> Mr. Chenevix found, "that if a bit of copper be put into a solution of corrosive sublimate, a white powder shortly falls to the bottom, and that powder is calomel. When washed, it does not contain an atom of copper, nor of corrosive sublimate."

<sup>2</sup> *Thomson's Chemistry*, 5th ed. vol. i. p. 465.

<sup>3</sup> The preparation is said to have been long known to the Chinese, and it is mentioned by Rhazis and Avicenna. *Bergman*, iv. 281.



mistake, or designedly as a poison, the best antidote is white of egg *diluted* with water, and given in large, frequently repeated doses. The albumen decomposes the corrosive muriate, reducing it to the state of the mild muriate, while the compound which it forms with it, exerts no deleterious effect on the stomach.<sup>1</sup> M. Taddei has found that gluten and rye-meal are equally serviceable as antidotes to this poison. He ascertained that gr. j of sublimate is neutralized by 13 grains of dried gluten, and 500 to 600 grains of rye-meal.<sup>2</sup> The presence of corrosive sublimate in any solution suspected to contain it, may be detected by putting into the fluid a small piece of clean polished copper, which, if the poison be present, will be covered with a white coating, or white streaks that acquire a metallic lustre when rubbed. As it is of importance to be able to distinguish calomel from corrosive sublimate in the stomach or intestines of a person supposed to be poisoned; (for the latter salt may be reduced almost to the state of calomel by the contents of the stomach,) Orfila has laid down the following rules for our guidance in determining this point. If the salt be found in the form of a white powder, easily separable from the contents of the alimentary canal or its membranes, and acquire a black colour when put into lime-water, we pronounce that calomel has been taken; but if the salt be not found as a white powder, but intimately combined with the contents of the stomach, and these suffer no change of colour when mixed with lime water, while the presence of a mercurial salt is nevertheless ascertained, then we may conclude that the poison has been corrosive sublimate.<sup>3</sup> The hydriodate of zinc is a still better test. A very ingenious galvanic mode of detecting it has also been proposed by Mr. Sylvester. Bend a piece of iron wire three inches long into this shape  $\Pi$ ; and tie the two ends to a common, gold, wedding ring, on a plate of glass placed horizontal, drop some sulphuric acid, diluted with six parts by weight of water, till it spread the size of a halfpenny; and then at a little distance, some of the solution supposed to contain the sublimate, till the two edges of the liquids join, and place the wire and the ring in such a manner, that the wire may touch the acid and the ring the solution. If any sublimate be present, the ring will in a few minutes be coated with mercury where it touches the solution. Corrosive Muriate of Mercury sometimes succeeds in curing the primary symptoms of syphilis, but it as often fails; and although it check the progress of the secondary symptoms, relieving venereal pains, and

<sup>1</sup> Orfila. *Traité des Poisons*, t. i. part i. 101.

<sup>2</sup> *Girnale di fisica et Isis*. 4. l. 1823. p. 419.

<sup>3</sup> *Nouveau Journ. de Medecine*, tom. x. p. 147



healing ulcers of the throat, “yet, even in these cases,” says Mr. Pearson, “it never confers permanent benefit; for new symptoms will appear during the use of it; and on many occasions it will fail of affording the least advantage to the patient from first to last.”<sup>1</sup> It is given with more advantage in some other affections, as old ulcers, chronic rheumatism and some cutaneous diseases, particularly lepra, in which Willan says, it is the only useful preparation of mercury, “its operation being promoted by giving at the same time an antimonial<sup>2</sup>,” and the decoction of the woods. Its sensible operation is by urine; but sometimes it occasions the most violent nausea, griping, and purging, in which cases it should be combined with opium; and it is always necessary to take, during its use, some mucilaginous fluid, to moderate the irritation it is apt to induce. It is also used as an external application; in which case the best vehicle is the bitter almond emulsion; and it has the property of preventing this emulsion from fermenting. The dose is from gr. one-eighth to gr. one-fourth twice a day, made into a pill with crumb of bread or extract of poppies.

Officinal preparations. *Liquor Hydrargyri Oxymuriatis*, L. *Hydrargyri Submurias*, L. E. D. *Hydrargyri Precipitatus albus*, L.

**LIQUOR HYDRARGYRI OXYMURIATIS**, Lond.

*Solution of Oxymuriate of Mercury.*

“Take of Oxymuriate of Mercury, *eight grains*; Distilled Water, *fifteen fluid ounces*; Rectified Spirit, *a fluid ounce*. Dissolve the Oxymuriate of Mercury in the water, and add to it the spirit.”

This solution is intended to facilitate the administration of minute doses of oxymuriate of mercury, each fluid ounce of the solution containing half a grain of the salt. It ought not to be long kept or exposed to a clear light, as the oxymuriate is gradually decomposed, and calomel precipitated. It is, however, the most safe and convenient form of administering this active salt; and may be given as an antisymphilitic in doses of from fʒss. to fʒij. in fʒij. of linseed infusion, or of water and syrup, and in more minute doses, when its alterative effects only are required. As a local application, this solution, diluted with two parts of water, forms a useful gargle in venereal sore throat; and without dilution we have found it serviceable as a gargle for breaking the abscess in cynanche tonsillaris, when suppuration takes place. Diluted with an equal quantity of water, it is employed as a wash against tetter and scabies; and very largely diluted, it may be used as an injection in gonorrhœa: and given in the form of enema, when the stomach will not receive it.

<sup>1</sup> *Pearson on Remedies for Lues Venerca, &c.* 116.

<sup>2</sup> *Willan on Cutaneous Diseases*, 140.



HYDRAR'GYRUM PRÆCIPITATUM ALBUM<sup>1</sup>,  
Lond. *White Precipitated Mercury.*<sup>2</sup>

“Take of Oxymuriate of Mercury, *half a pound*; Muriate of Ammonia, *four ounces*; Solution of Subcarbonate of Potass, *half a pint*; Distilled Water, *four pints*. Dissolve first the Muriate of Ammonia, then the Oxymuriate of Mercury, in the distilled water, and add to the mixed solution, the solution of Subcarbonate of Potass. Wash the precipitated powder until it become tasteless, and then dry it.”

SUBMURIAS HYDRARGYRI AMMONIATUM, Dub. *Ammoniated Submuriate of Mercury.*

“Add to the fluid which has been poured off from the precipitated Submuriate of Mercury a quantity of Water of Caustic Ammonia sufficient to precipitate the whole of the metallic salt. Wash the precipitate with cold distilled water, and dry it upon bibulous paper.”

*Syn.* Muriate de Mercure précipité (*F.*), Salzsaures Quecksilber präzipitat (*G.*), Precipitato bianco di mercurio (*I.*).

As the products of these two processes are precisely the same, that of the Dublin College is to be preferred, both on account of its economy, and its greater simplicity. The fluid, which it orders to be used, is that which is decanted from the precipitated, mild muriate of mercury, prepared by heat; and which, as we have already observed, holds the corrosive muriate in solution: so that the oxide of this salt is precipitated by the ammonia, combined with a portion of acid and also of ammonia, forming a ternary compound, or a submuriate of mercury and ammonia. In the London process, the muriate of ammonia, and the oxymuriate of mercury, when dissolved in the water, combine together, and form a solution of a ternary compound of muriatic acid, ammonia, and oxide of mercury, or a soluble supermuriate of mercury and ammonia. By the addition of the subcarbonate of potass, a great part of the acid of the muriate of ammonia is abstracted, and the same triple insoluble compound is precipitated as in the former process; and the fluid retains in solution muriate of potass, the carbonic acid having been dissipated in the gaseous form; at least this is very probable. It is to be regretted, that the quantity ordered is inadequate to the effect intended.

*Qualities.* — This Muriate of Mercury and of Ammonia is inodorous and insipid: of a snowy whiteness, smooth, and insoluble in water, and does not become black when triturated

<sup>1</sup> This name is completely at variance with the principles on which the reformed nomenclature is founded; and the reasons which might have excused the adoption of *calomel*, and some other equally barbarous terms, cannot be advanced in justification in this instance.

<sup>2</sup> Mercurius præcipitatus albus, P. L. 1745. Calx hydrargyri alba, P. L. 1787.



with lime-water. It is decomposed by the sulphuric and nitric acids, the former of which converts it into oxymuriate of mercury, and sulphate of mercury and of ammonia, and the latter into the oxymuriate also, and nitrate of ammonia and of mercury. Muriatic acid restores it to the state of soluble supermuriate, the *sal alemroth* of the old chemists. If heated with potass, it is decomposed, and its ammonia expelled, owing to the combination of the muriatic acid with the potass.<sup>1</sup> It is sometimes adulterated with white lead; to discover which, digest one part of it in four parts of acetic acid, and add to the solution a small quantity of sulphuret of ammonia; a black precipitate insoluble in sulphuric acid indicates the presence of lead. Chalk and starch are also sometimes mixed with it; and may be detected by heating the preparation in an iron spoon; if pure, it is completely volatilized; but if adulterated with starch, a black coal is left; or if with chalk, a white powder, at the bottom of the spoon. It is said to be a compound of 81 parts of Peroxide of Mercury, 16 of Muriatic Acid and 3 of Ammonia in 100 parts.

*Medical properties and uses.* — This preparation is only used in combination with lard, as an ointment for the cure of itch, and some other cutaneous eruptions.

Officinal preparation. *Unguentum Hydrargyri Præcipitati Albi*, L. D.

HYDRARGYRUM PURIFICATUM, Lond. *Purified Mercury.*<sup>2</sup>

“Pour Mercury into an Iron retort, and heat being applied, distil the Purified Mercury.”

HYDRARGYRUS PURIFICATUS, Edin. *Purified Mercury.*

“Take of Mercury, *six parts*; Filings of Iron, *one part*. Rub them together, and distil from an iron retort.”

HYDRARGYRUM PURIFICATUM, Dub. *Purified Mercury.*

“Take of Mercury, *six pounds*. Distil off slowly four pounds.”

*Syn.* Mercure (F.), Quecksilber (G.), Mercurio (I.), Azogogue (S.).

By this mode of treating mercury it is certainly obtained more bright and mobile; but although it is generally supposed that the foreign metals with which the mercury of commerce is supposed to be alloyed, are separated by this process, yet, the necessity of it may be well questioned.

HYDRARGYRI SUBMURIAS<sup>3</sup>, Lond. *Submuriate of Mercury.*

“Take of Purified Mercury, by weight, *four pounds*; Sul-

<sup>1</sup> *Trans. of the Pharm.* p. 122.

<sup>2</sup> *Argentum vivum purificatum*, P. L. 1745.

<sup>3</sup> Old names, *Aquila alba*, *Aquila mitigata*, *Manna Metallorum*, *Panchymagogum minerale*, *Panchymagogus quercetanus*, *Sublimatum dulce*, *Mercurius dulcis sublimatus*, *Calomelas*.



phuric Acid, by weight, *thirty ounces*; Muriate of Soda, *one pound and a half*; Muriate of Ammonia, *eight ounces*. Boil two pounds of the Mercury with the Sulphuric Acid in a glass vessel, until the Sulphate of Mercury is dry; when this is cold triturate it with two pounds of Mercury in an earthen mortar, that they may be perfectly mixed. Then add the Muriate of Soda, and rub them together until all globules disappear; afterwards sublime. Reduce the sublimed matter to a very fine powder, pass it through a sieve, and mix it carefully with the Muriate of Ammonia previously dissolved in a gallon of Boiling Distilled Water. Set it aside that the powder may subside. Pour off the solution, and wash the powder repeatedly with Boiling Distilled Water, until solution of Ammonia dropped into it throw nothing down. Finally, reduce to a very fine powder, in the same manner as is directed for the prepared chalk."

SUBMURIAS HYDRARGYRI MITIS, sive CALOMELAS, Edin. *Mild Submuriate of Mercury, or Calomel.*

"Take of Muriate of Mercury, *four parts*; Purified Mercury, *three parts*. Rub the Muriate in a glass mortar with a little water, in order to prevent the acrid powder from rising, then add the Mercury, and again triturate until it be extinguished; put the dried mass into an oblong phial, one-third of which only it shall fill, and sublime it in a sand-bath. Again triturate the sublimed powder, and again sublime it, then reduce it to a fine powder, which is, lastly, to be well washed with boiling distilled water."

SUBMURIAS HYDRARGYRI SUBLIMATUM, sive CALOMELAS, Dub. *Sublimed Submuriate of Mercury, or Calomel.*

"Take of Corrosive Muriate of mercury, *a pound*; Purified Mercury, *nine ounces*. Rub them together until the globules disappear, and sublime with a sufficient degree of heat. Let the sublimed matter be rubbed to powder, and again sublimed. Pulverize it, and wash it with frequent affusions of distilled water, until the poured off solution no longer lets any sediment fall on the addition of a few drops of Carbonate of Kali. Finally, dry it."

*Syn.* Muriate de Mercure doux (*F.*), Mildes Salzsauers Quecksilber (*G.*), Mercurio dolce sublimato (*I.*).

This very important preparation is a Protochloride of Mercury.<sup>1</sup> In the process of the London College, following the

---

<sup>1</sup> It is very remarkable that all the Colleges have erred in naming this preparation, which in no point of view can be regarded as a *submuriate*, but is as much a *muriate* as the corrosive sublimate; the sole difference depending on the degree of oxidizement of the metal, which in this preparation is at a minimum. In a medical point of view, we are of opinion that the name *calomel*, however absurd, ought to have been retained, as the syllables *oxy* and *sub* are scarcely sufficient to



old doctrines, the Mercury is first formed into a persulphate, which is mixed with the common salt, and converted into corrosive sublimate, but which at the moment of its formation, is again decomposed by the Ammonia of the Muriate of Ammonia, and converted in Calomel. But supposing, as is actually the case, that the Corrosive Sublimate which is formed, is a perchloride, this is changed into Calomel, by one-half of the chlorine uniting with the additional portion of mercury and forming a protochloride. By triturating metallic Mercury, as directed by the two other Colleges, with the Corrosive Muriate, the whole mass assumes a grey colour. The sublimations render the combination of the mercury with the chloride, and its reduction to the state of a protochloride complete; but this is not the case in the first sublimation: for both metallic mercury and corrosive muriate are found unchanged in the sublimed mass; and thence the necessity of the second trituration and the subsequent sublimations. By repeating, however, the sublimations too often the product is injured, as corrosive muriate is formed in each sublimation. The final trituration and levigation are intended to separate any corrosive muriate that may have been formed, and the test of the Dublin Pharmacopœia ought always to be had recourse to for ascertaining this point. In performing the process, the addition of a little water during the trituration of the ingredients, in the first instance, is very necessary; as otherwise, the operator is apt to suffer extremely from the acrid powder of the corrosive muriate which is elevated.

*Qualities.* — Calomel is obtained by the above processes in the form of a dull white, semitransparent mass, the specific gravity of which is 7.175. It is a protochloride of Mercury as we have already stated, inodorous, insipid, and when pulverized has a light yellow or ivory colour, which deepens by long exposure to the light, owing, perhaps, to a partial decomposition. It is regarded as insoluble, one part requiring 1152 parts of water at  $212^{\circ}$  for its solution.<sup>1</sup> Nitric acid converts it into corrosive muriate, much nitrous gas being evolved; and the same change is effected by oxymuriatic acid (*chlorine*). Lime-water and the alkalies, when triturated with it, instantly render it black, a circumstance which supplies us with a test of its purity, for if it contain any corrosive muriate, a yellow tint is mingled with the black on the addition of lime-water. It is also decomposed by sulphuretted hydrogen; and the hydro sulphurets by antimony, iron, lead, copper,

---

distinguish the two salts to blundering assistants and apprentices, by whom the most dangerous mistakes may be committed.

<sup>1</sup> Rouelle.



and soap. According to Chenevix, using the terms of the old doctrine, 100 parts of it contain 11·5 of muriatic acid, and 88·5 protoxide of mercury, consisting of 79 of mercury and 9·5 of oxygen; but according to Zaboada, the proportions are, 10·6 of acid, and 89·4 of oxide, consisting of 85 of mercury and 4·4 of oxygen. Correcting these statements, and using the new terms, the true constituents of this salt appear to be,

		Chenevix.		Zaboada.
Chlorine	-	14·9	-	14·04
Mercury	-	85·1	-	85·96
		<hr/>		<hr/>
		100·0		100·00 <sup>1</sup>

According to Mr. Phillips, “considered as a chloride or protochloride, it is composed of

Chlorine	-	15·25	or 1 atom Chlorine =	36
Mercury	-	84·75	1 atom Mercury =	200

---

100·00      weight of its atom = 236”<sup>2</sup>

*Medical properties and uses.*—This is the most useful of the preparations of mercury, and is more generally employed than almost any other remedy in the whole range of the materia medica.<sup>3</sup> It is antisyphilitic, antispasmodic, alterative, deobstruent, pugnitive, and errhine. As a remedy in syphilis, it can be fully confided in, when its disposition to run off by the bowels is counteracted by opium. In the same state of combination, it is also found efficacious in several convulsive affections, as epilepsy, trismus, and tetanus; and in that species of spasmodic stricture which occurs in virulent gonorrhœa. As an alterative and deobstruent, it is employed with advantage in cutaneous eruptions, as lepra and scabies, in which cases it is combined with antimonials and guaiacum; and in hepatitis, and glandular obstructions. In dropsies it assists the action of squill and fox-glove; and as a purgative it may be employed with safety in almost every form of disease not attended with visceral inflammation, or where there is not great irritability and delicacy of habit. Calomel, however, does not act with certainty as a purgative even in large doses, and, therefore, it is generally combined with scammony, jalap, or some other active cathartic. The usual dose to affect the

<sup>1</sup> Thomson's Chemistry, 5th edit. i. p. 467.

<sup>2</sup> Trans. of the London Pharm. 1824.

<sup>3</sup> It was described in 1608, by Beguin, who calls *Draco mitigatus*, Corrosive Sublimate being called by the alchemists *Draco*, the dragon: but from Mr. Hatchett's experiments on the Calomel of Thibet, it would appear that it has been much longer known to the natives of that part of India. Brande's Manuel, p. 302.



habit and produce ptyalism is from gr. j. to grs. ij. in a pill with opium, given night and morning; and from grs. iij. to grs. viij. act in general as a purgative: but in some complaints, as yellow fever, hydrocephalus and croup for example, in which it is supposed to exert a specific effect, this dose has been repeated every two or three hours, until upwards of 100 grains have been taken in a very short space of time.

On account of its insolubility and great specific gravity, it can be given only in the form of pills and powders.

SUBMURIAS HYDRARGYRI PRÆCIPITATUS, Edin. *Precipitated Submuriate of Mercury.*

“Take of Diluted Nitrous Acid, Purified Mercury, of each *eight ounces*; Muriate of Soda, *four ounces and a half*; Boiling Water, *eight pounds*. Mix the Mercury with the Diluted Nitrous Acid, and towards the termination of the effervescence digest with a gentle heat, frequently shaking the vessel. It is requisite, however, that more Mercury be mixed with the Acid than it can dissolve, so that a completely saturated solution be obtained.

“Dissolve at the same time the Muriate of Soda in the Boiling Water; then to this add the other solution while it is yet warm, and mix them very quickly together. After the precipitate has subsided, pour off the saline fluid, and wash the Submuriate of Mercury by frequent affusions of warm water, which are to be poured off each time after the precipitate subsides, until the water comes off tasteless.”

SUBMURIAS HYDRARGYRI PRÆCIPITATUM, Dub. *Precipitated Submuriate of Mercury.*

“Take of Purified Mercury, *seven ounces*; Diluted Nitrous Acid, *five fluid ounces*. Pour the Acid upon the Mercury in a glass vessel, and when the effervescence has ceased, digest with a gentle heat for six hours, with frequent agitation. Then raise the heat, that the solution may boil a little, which is to be poured off from the residual Mercury, and quickly mixed with *ten pounds* of Boiling Water, in which *four ounces* of Muriate of Soda have been previously dissolved; wash the powder that subsides with warm distilled water, as long as the fluid poured off from it yields a precipitate on the addition of a few drops of the solution of (*sub*) Carbonate of Kali; lastly, let it be dried.”

These processes are framed on the process originally suggested by Scheele, and the error into which he was led by reasoning from a false analogy has not been corrected by the Colleges; the product of the above process being a mild muriate of mercury mixed with subnitrate of mercury which modifies its powers: a smaller proportion also of mild muriate being obtained than should follow from the quantity of mer-



cury employed. The cause of this effect is, that by dissolving mercury in nitric acid with the assistance of heat, the metal contained in the acid solution is oxidized to a maximum, and when water is added to it, a subnitrate is precipitated, while a supernitrate remains in solution. Hence, on the addition of the watery solution of muriate of soda, the water occasions the subnitrate to be precipitated, before the decomposition which takes place is effected, at the same time part of the oxide combines with the chlorine of the muriate of soda, and forms a portion of corrosive muriate, which is held in solution with the newly formed nitrate of soda, while the mild muriate is precipitated in combination with insoluble subnitrate of mercury.

To obtain, therefore, the greatest proportion of pure, mild muriate of mercury by precipitation, the nitrate must be prepared slowly and without the aid of heat, which should not be employed in any part of the process. Dr. Murray ascertained, that “the quantity of mild muriate obtained from a solution of  $\mathfrak{z}\text{j}$ . of mercury in diluted nitric acid in the cold, is a little more than  $\mathfrak{z}\text{j}$ .; whereas, from the same quantity dissolved with the application of heat, the precipitate did not exceed  $\mathfrak{z}\text{ss}$ . and the liquor held much more corrosive muriate in solution than the other.”<sup>1</sup>

*Qualities.*—Precipitated mild muriate of mercury, when properly prepared, is inodorous and insipid. It is whiter, smoother and lighter, than the sublimed preparation, but otherwise agrees with it, both in its chemical qualities and medicinal effects. As prepared, however, according to the directions of the pharmacopœias, subnitrate of mercury, which it contains, may have some effect in altering its powers in a small degree.

*Medical properties and uses.*—It is said to be more liable to run off by the bowels than common calomel; but, as its properties are essentially the same, it may be regarded as a superfluous preparation.

HYDRARGYRI SULPHURETUM NIGRUM<sup>2</sup>,  
Lond. *Black Sulphuret of Mercury.*

“Take of Purified Mercury, by weight, *one pound*; Sublimed Sulphur, *a pound*. Triturate them together until the globules disappear.”

SULPHURETUM HYDRARGYRI NIGRUM, Edin.  
Dub. *Black Sulphuret of Mercury.*

“Take of Purified Mercury, Sublimed Sulphur, of each *equal weights*. Rub them together in a glass mortar with a glass pestle, until the globules of mercury altogether dis-

<sup>1</sup> *System of Materia Medica*, &c. ii. 319.

<sup>2</sup> Hydrargyrus e Sulphure, P. L. 1787.



appear. It may be made with double the quantity also of mercury.”

*Syn.* Sulphure de Mercure noir (*F.*), Schwarzes Schwefelquecksilber (*G.*), Solfuro di Mercurio nero (*I.*).

During the trituration of the mercury with the sulphur, Fourcroy supposes that the metal is imperfectly oxidized by attracting oxygen from the atmosphere; but this opinion has been disproved by the experiments of Proust<sup>1</sup>; and although a chemical combination be effected between the mercury and the sulphur, yet the real nature of the preparation is not understood. Mr. Brande<sup>2</sup> regards it as a compound of 58 parts of bisulphuret of Mercury, and 42 of Sulphur, and this opinion accords with that of Mr. Phillips.<sup>3</sup>

*Qualities.* — Black Sulphuret of Mercury is in the form of a very black, impalpable, inodorous, insipid powder. When heated in an open vessel it emits sulphurous acid gas; becomes first of a deep violet hue, and afterwards sublimes of a brilliant red colour. It is often ill prepared, which may be known by rubbing a portion of it on gold; to which, if it be good, no whiteness will be communicated. It is, also, sometimes adulterated with ivory black, which may be detected in it by throwing a little of the suspected sulphuret on a red-hot iron; if ivory black be present, some ashes will be left after the volatilization, which will not be the case when it is good, the pure sulphuret being completely dissipated. Mr. Phillips says it is sometimes adulterated with sulphuret of antimony, which may be detected by boiling a little of the powder in strong muriatic acid, and then pouring the clear solution into water, if the sulphuret be present, submuriate of antimony will be precipitated.

*Medical properties and uses.* — This mercurial preparation is alterative and anthelmintic. It is chiefly employed against scrophulous swellings, and in cutaneous affections; and has been found useful in ascarides. But it is on the whole a very uncertain preparation, and requires to be long used to produce any sensible effects. The dose is from grs. v. to ʒss., given twice or three times a day.

HYDRAR'GYRI SULPHURE'TUM <sup>4</sup> RU'BRUM.  
Lond. *Red Sulphuret of Mercury.*

“Take of Purified Mercury, by weight, *forty ounces*; Sublimed Sulphur, *eight ounces*. Having melted the Sulphur over the fire, mix in the Mercury, and, immediately the mass swells, remove the vessel from the fire, and cover it with force

<sup>1</sup> *Journal de Physique*, liii. 92.

<sup>2</sup> *Manual of Pharmacy*, p. 303.

<sup>3</sup> *Trans. of Pharm.* p. 123.

<sup>4</sup> *Cinnabaris factitia*, P. L. 1745. *Hydrargyrus sulphuratus ruber*, P. L. 1787.



to prevent it from catching fire; then rub it into powder and sublime.”

SULPHURETUM HYDRARGYRI RUBRUM, Dub. *Red Sulphuret of Mercury.*

“Take of Purified Mercury, *forty ounces*; Sublimed Sulphur, *eight ounces*. Mix the Mercury with the melted Sulphur; and if the mixture take fire, extinguish it by covering the vessel; then rub the mass to powder, and sublime it.”

*Syn.* Sulphure de Mercure rouge (*F.*), Zinnober (*G.*), Solfuro di Mercurio rosso (*I.*), Cinabrio (*S.*), Shengerf (*H.*).

By these processes the mercury and sulphur are more intimately combined, and a more complete sulphuret produced than in the former preparation. The inflammation which is apt to happen after the mixture of the mercury with the melted sulphur, when the mass swells and explodes, as frequently occurs, is similar to the combustion during the union of sulphur by heat with some other metals, independent of the presence of air: hence, covering the vessel, without removing it from the fire, does not check the combustion, although, by excluding the air, a real inflammation of the materials may be prevented. In the second part of the process great caution is necessary to prevent the neck of the vessel in which it is sublimed from being choaked up by the sublimed sulphuret; as by the occurrence of such an accident the vessel would be burst by the confined vapours. To avoid this, a wide-necked vessel should be used.

The Cinnabar of commerce, which is chiefly used as a pigment, is manufactured in Holland, on a very extensive scale<sup>1</sup>; and the following method has been proposed by Mr. Kirchoff, for obtaining it in the humid way. First, form ethiops mineral, by trituration, in a porcelain cup with a glass pestle, 300 grains of mercury, and 68 of sulphur, moistened with a few drops of solution of potass, and then add to it 160 grains of potass, dissolved in an equal weight of water. Heat the vessel with the ingredients over the flame of a candle, continuing the trituration, and adding, as the fluid evaporates, pure water from time to time, so as to keep the ingredients covered to the depth of an inch. At the end of two hours, if the trituration have been continued, the colour of the mixture changes from black to brown and then to red; after which no more water should be added, but the trituration must be uninterruptedly continued until the mass have acquired the consistence of a jelly, and the red colour attained considerable brightness and beauty: the heat must be then imme-

---

<sup>1</sup> See a description of the method, *Annales de Chimie*, li. 196.



diately withdrawn, otherwise the red soon changes to a dirty brown. <sup>1</sup>

*Qualities.* — Red Sulphuret of Mercury sublimes in the form of a vivid, red, crystalline cake, and yields, by trituration, a powder of a very bright-red colour, which is inodorous, insipid, and insoluble in water, alcohol, and the majority of acids. It is decomposed, however, by nitro-muriatic acid, which combines with the mercury, and disengages the sulphur; but is not altered by solutions of the alkalies, even when boiled with them; although lime, potass, soda and most of the metals decompose it when melted with it. Vauquelin supposed that it contains the metal in a state of high oxidization; a supposition, which the experiments of Proust and Seguin have completely disproved. According to Proust, 100 parts of the sulphuret consists of 85 of unoxidized mercury, and 15 of sulphur; but according to the later experiments of Guibert <sup>2</sup>, the proportions are 86.1803 of mercury, and 13.8197 of sulphur. This preparation is sometimes adulterated with red-lead, dragon's blood, and chalk; the first is discovered by the same process as was described for discovering it in the red oxide; spirit of wine detects the second by extracting the colouring matter; and the last is discovered by an effervescence being excited by muriatic acid; and the production of sulphate of lime on adding sulphuric acid.

*Medical properties and uses.* — Red Sulphuret of Mercury is alterative and deobstruent. It was formerly much used in cutaneous diseases, gouty, and rheumatic affections, and in worms. It is now, however, scarcely ever prescribed. It has been recommended for fumigations in syphilis; but, on account of the sulphurous vapours, it is less fit for this purpose than the grey oxide. The dose for internal use is from grs. x. to ʒss., made into an electuary or bolus.

SUB-SULPHAS HYDRARGYRI FLAVUS. <sup>3</sup> Edin.  
*Yellow Sub-Sulphate of Mercury.*

“Take of Purified Mercury, *two parts*; Sulphuric Acid, *three parts*. Put them into a glass cucurbit, placed in a sand-bath, and boil them to dryness. Pulverize the white mass which is left at the bottom of the vessel, and throw it into boiling water. It will immediately be converted into a yellow powder, which is to be washed with frequent affusions of warm water.”

OXYDUM HYDRARGYRI SULPHURICUM, Dub. *Sulphuric Oxide of Mercury.*

<sup>1</sup> *Nicholson's Journal*, 4to. ii. 1.

<sup>2</sup> *Journ. de Pharmacie*, Aout. 1816. p. 371.

<sup>3</sup> *Hydrargyrus vitriolatus*. P. L. 1787.



“ Take of Purified Mercury, *a pound*; Sulphuric Acid, *a pound and a half*. Dissolve in a glass vessel, with a sufficiently strong heat, and gradually raise the fire until the mass be completely dried. This, by the affusion of a large quantity of hot water, will immediately become yellow and fall into powder, which is to be well triturated with the water in an earthenware mortar.

“ After pouring off the supernatant fluid, let the powder be washed with repeated affusions of hot, distilled water, as long as any precipitate is produced in the decanted liquor on the addition of a few drops of Water of Subcarbonate of Kali; and, lastly, dry it.”

*Syn.* Subsulphate de Mercure (*F.*), Gelbes Schwefelsaures Quecksilberoxyd (*G.*), Turpeto Minerale Mercuriale (*I.*).

Sulphuric acid scarcely acts on mercury unless aided by a high temperature. When it is boiled on it, as directed in these processes, the acid is partially decomposed by the metal which is oxidized, while sulphurous gas is evolved; and the oxide thus formed uniting with the remaining acid, the whole becomes a supersulphate of mercury. By continuing the application of heat, at a higher temperature, a considerable portion of the acid is expelled, and partly decomposed, by which the metal is still more highly oxidized, and the resulting dry mass is a subsulphate of mercury. When boiling water is poured on this salt, the fluid, acting by its powerful affinity for sulphuric acid, decomposes it, abstracts the acid, and precipitates the oxide; but as the acid still holds combined with it a small portion of oxide, and the precipitated oxide retains some acid, the result of this part of the process is a supersulphate of mercury held in solution by the water, and a subsulphate precipitated in the form of a yellow powder. To obtain this effect completely, the saline mass must be made entirely dry before pouring over it the hot water; for if the vessel be sooner taken from the fire, the precipitation is partial only, the greater part of the salt being dissolved without being decomposed. Perhaps the best mode is to continue the exsiccation until a little of the white mass dissolved in cold water does not redden litmus paper. The proportions for obtaining the largest quantity of product are two parts of acid, and one of mercury: hence, while the quantity ordered by the Dublin College is rather too small, the proportions of the Edinburgh formula are productive of a very unnecessary waste of acid.

*Qualities.* — Subsulphate of Mercury is inodorous, and acrid to the taste. It is obtained in the form of a beautiful bright-yellow powder, of a specific gravity of 6.444, and nearly insoluble in water, requiring 2000 parts at 60°, and 600 at



212°, for its solution, which is colourless. By trituration with mercury it is changed into the black oxide; and at a red heat is decomposed, the oxygen being given out and the metal reduced. According to the analysis of Braumcamp and Segueira, its constituents are 84·7 parts of oxide of mercury, 15 of sulphuric acid, and three of water<sup>1</sup>; while Fourcroy makes them 87 of oxide, 10 of acid, and 3 of water.

*Medical properties and uses.* — This preparation is emetic, discutient, alterative, and errhine; but from the violence of its action it is seldom administered as an internal remedy. As an errhine, however, it has been found extremely useful in chronic ophthalmia, and diseases of the head; but even for this purpose its acrimony requires to be sheathed with some bland powder, as starch, or liquorice root powder, in the proportion of grs. v. to gr. j. of the subsulphate. In doses of gr. v. it operates as a very powerful emetic.

---

In concluding the account of the preparations of mercury, it may not be improper to observe that the exhibition of any of them in certain states of the habit, when, at the same time, the body is under exposure to cold, is apt to excite an erythematic eruption of the skin, accompanied with much fever. This disease does not at all depend on the use of any particular preparation of the remedy; but, as far as I have been able to observe, it is liable to show itself in such an irritable state of the habit as produces hysteria in females, when the body is very suddenly exposed to a current of cold air, or to a cold, moist atmosphere, while under the influence of mercury. When it occurs, the mercurials must be immediately discontinued, bark, opium, and purgatives internally administered, and the affected surface sprinkled with dry flour, or covered with the *linimentum aquæ calcis* of the Edinburgh and Dublin Pharmacopœias; while at the same time the warm bath is to be used at least twice a day. Under this treatment the disease generally disappears, and the use of the mercurial may be renewed; but sometimes the morbid symptoms increase under every mode of treatment, and a fatal termination of the disease ensues.

---

<sup>1</sup> *Annales de Chimie*, liv. 123.



## PRÆPARATA E STANNO.

## PREPARATIONS OF TIN.

PULVIS STANNI, Dub. *Powder of Tin.*

“Take of Tin *any quantity*. Melt it over the fire in an iron mortar, and stir it while it is cooling, until it become a powder, which, when cold, is to be passed through a sieve.”

*Syn.* Poudre d’Etain (*F.*), Zinn (*G.*), Stagno in polvere (*I.*).

By this process Tin is reduced to the form of a fine granular powder, and, perhaps, by the constant stirring, it is also very slightly oxidized, for the powder has less brilliancy than the entire metal.

*Medical properties and uses.*—Powder of Tin is a mechanical anthelmintic. It has been chiefly given to expel the tape-worm; and is supposed to operate by the grittiness of its particles irritating the worm, and dislodging it from the mucus in which it is imbedded. It is given in doses of ʒj. or ʒij. mixed in treacle, for two or three successive mornings, and a brisk cathartic afterwards exhibited. But it is likely to be, henceforth, seldom used, oil of turpentine being a much superior remedy for the expulsion of tape-worm.

## PRÆPARATA E PLUMBO.

## PREPARATIONS OF LEAD.

PLUMBI ACETAS, Lond. *Acetate of Lead.*<sup>1</sup>

“Take of Subcarbonate of Lead, *a pound*; Strong Acetic Acid, *one pint*; Boiling Distilled Water, *one pint and a half*. Mix the Acid with the Water; and the Subcarbonate of Lead gradually; then filter the solution through paper, and, having evaporated it until a pellicle appears on its surface, set it apart that crystals may form. Pour off the fluid, and dry the crystals upon bibulous paper.”

ACETAS PLUMBI, Edin. *Acetate of Lead.*

“Take of White Oxide of Lead, *any quantity*; Weaker Acetic Acid, *a sufficient quantity*. Put the Oxide into a cucurbit, and pour over it ten times its weight of the acid. Let the mixture stand upon a warm sand-bath until the acid become sweet; then let this be poured off, and add fresh por-

<sup>1</sup> Saccharum saturni, P. L. 1720—45. Cerussa acetata, P. L. 1787.



tions of acid successively, until no more sweetness is communicated. Evaporate all the fluid, freed from impurities, in a glass vessel to the consistence of thin honey, and set it aside in a cold place that crystals may form, which are to be dried in the shade. Evaporate again the residuary liquor, that new crystals may be obtained; and repeat the evaporation until no more are formed."

ACETAS PLUMBI, Dub. *Acetate of Lead.*

"Take of Subacetate of Lead, called CERUSSA, *any quantity*, Distilled Vinegar, *ten times its weight*. Digest them in a glass vessel until the Vinegar become sweet; and having poured this off, add more, until it cease to become sweet. Filter the solution, and crystallize by alternate slow evaporation and cooling. Dry the crystals in the shade."

*Syn.* Acetate de Plomb cristallisé (*F.*), Essigsaures Blei (*G.*), Zuccher o di Saturno (*I.*).

In the London process the acetic acid unites with the sub-carbonate<sup>1</sup>, or more correctly carbonate of lead, expelling the carbonic acid; and by the subsequent evaporations the salt crystallizes in the form of an acetate. But on account of the smallness of the quantity of product, the trouble and expense of the process, and the difficulty of obtaining the white lead perfectly free from whitening (carbonate of lime), with which it is generally adulterated, the preparation of this salt is seldom undertaken by the apothecary; so that the acetate (*sugar of lead*), usually found in the shops is the salt which is manufactured on a large scale, for the use of the calico printers, purified. It is chiefly prepared in Holland, in the following manner: Sheets of lead, coiled up, are put into pots, in which they are half immersed in distilled vinegar, and digested a sufficient time. Before long, the upper half, or that which is not immersed, is covered with an efflorescence of cerusse, after which it is immersed in the vinegar, and the part which was before immersed is now brought up to be converted into cerusse as before, when the plate is again turned; and this is repeated two or three times a day, until the vinegar become milky. This solution is next boiled in tinned vessels down to about one-third of the original quantity, then strained, and the salt crystallized by slow cooling. The crystals obtained by a second evaporation of the mother-water are browner and deliquescent<sup>2</sup>: and the whole requires to be again dissolved in rain or distilled water, and re-crystallized.

*Qualities.*—This salt, when pure, is inodorous, has a sweet,

<sup>1</sup> Carbonate of Lead consists of Protoxide of Lead 85.6, and Carbonic Acid 14.4 in 100 parts.

<sup>2</sup> *Aikin's Dictionary of Chemistry*, ii. 26.



astrigent taste, and crystallizes in white, glossy, right oblique-angled<sup>1</sup> prisms, terminated by dihedral summits, which are generally aggregated into irregular masses that have the appearance of lumps of sugar. Its specific gravity is 2.35.<sup>2</sup> Acetate of lead slightly effloresces, is soluble in 25 parts of distilled water either hot or cold; but after standing for some time, a slight decomposition takes place, and a small portion of white powder is deposited, which is an insoluble carbonate. It is also soluble in alcohol. In pump or hard water, which always contain carbonic acid, it is instantly decomposed, forming a milky solution, and a copious precipitate falls: and for the same reason it is decomposed by *Liquor Ammoniae Acetatis*. It is decomposed by the alkalies and their carbonates, most of the acids, and neutral salts, lime, magnesia, and all the sulphurets; but it is not affected by a solution of gum. The constituents of 100 parts are 59.45 of oxide of lead, 26.25 of acid, and 14.30 of water.<sup>3</sup>

*Medical properties and uses.*—Taken internally, Acetate of Lead is a very powerful astringent and sedative. It requires to be exhibited, however, with caution, and is admissible only in cases of very urgent danger, as in pulmonary, uterine, and intestinal hæmorrhages; in restraining which it has a very powerful influence. Combining it with opium prevents the deleterious effects which salts of lead are apt to produce when taken into the stomach; but, even when so combined, the smallest dose, in certain habits, is productive of very serious mischief. Some years ago, Dr. Hildebrand, of Lemberg, tried this salt in combination with opium with seeming advantage in phthisis; and it has been since occasionally used in this country; but from the effects of it in that disease, as far as I have observed, it is not likely to be generally employed by British practitioners. Dissolved in a large proportion of water, with a small quantity of distilled vinegar to prevent decomposition, it forms an excellent collyrium in ophthalmia; and somewhat less diluted, its solution is in common use as an external application in superficial inflammation. Objections have, nevertheless, been raised to the long-continued, external use of the preparations of lead; but the daily extensive employment of them in this form, without any bad effects, is a sufficient proof that, if they occasionally have produced mischief, it is rather to be attributed to some peculiar idiosyncrasy, than to the nature of the remedy.

The dose of Acetate of Lead, when internally exhibited, should not exceed gr. ss. given every six or eight hours. It

<sup>1</sup> Phillips' *Trans. of the Pharm.* 1824.

<sup>2</sup> Hassenfratz.

<sup>3</sup> Phillips' *Trans. of the Pharm.* 1824.



may be made into a pill with crumb of bread, and a proportion of opium, according to the circumstances of the case. As a collyrium or lotion, the proportions may be from gr. x. to ℥j. of the salt in fʒviii. of distilled water. The addition of a small quantity of distilled vinegar is necessary to prevent decomposition, when distilled water is not employed. Many practitioners object to even the external application of Salts of Lead; but I have used them extensively, and have never found any bad effects to result from them.

Officinal preparation. *Ceratum Plumbi Acetatis*, L.

LIQUOR PLUMBI SUBACETATIS, Lond. *Solution of Subacetate of Lead.*<sup>1</sup>

“Take of Semi-vitrified Oxide of Lead, *two pounds*; Acetic Acid (distilled vinegar), *a gallon*. Mix them, and boil down to six pints, assiduously stirring; then set the solution aside, that the impurities may subside, and strain it.”

LIQUOR SUBACETATIS LITHARGYRI, Dub. *Solution of Subacetate of Lead.*

“Take of Litharge, *a pound*; Distilled Vinegar, *eight pints*. Put them into a glass vessel, and boil to six pints, assiduously stirring; then set the solution aside, and strain it after the fæces have subsided.”

*Syn.* Acetate de Plomb liquide (*F.*), Blaiwasser (*G.*), Aceto di Saturno (*I.*).

In these processes, the acetic acid, which the distilled vinegar contains in a highly diluted state, attracts a portion of the litharge which is a protoxide of lead, and forms a subacetate, which remains dissolved in the water. The proportion of litharge ordered in both formulæ is too large, a gallon of distilled vinegar of the specific gravity 1.007 being capable of dissolving ten ounces only of the oxide: but the preparation according to strength of the vinegar.

*Qualities.*—This Solution of Subacetate of Lead, when properly prepared, is of a greenish straw-colour, has a slight acetous odour, and an austere, somewhat sweetish, taste. It is partially decomposed when largely diluted with distilled water; and with pump water, a heavy precipitate instantly takes place: it is also precipitated in the form of a white subsalt by the alkalies and their carbonates; and a black precipitate is produced by the alkaline sulphurets. It is, indeed, the best test for detecting sulphuretted hydrogen in any compound. The quantity of sulphur is always  $\frac{2}{15}$  of the sulphuret of lead, to which, if we add  $\frac{1}{15}$ , we obtain the weight of the sulphuretted hydrogen. This solution is also incompatible with solutions of mucilage, the gum of which it coagulates; and, indeed, it is the most delicate test for mucilage with which we are acquainted. According to the experiments of



Dr. Bostock<sup>1</sup>, the constituents of 100 parts of the saturated solution are 23·1 of oxide of lead, 5 of acetic acid, and 71·9 of water, which agree with the statement of Thenard<sup>2</sup>, who found that the salt, when crystallized, consists of 17 parts of acid, 78 of oxide of lead, and 5 of water, in 100 parts.<sup>3</sup>

*Medical properties and uses.*—This solution is used only externally, and when diluted with *distilled* water forms a very useful cooling, discutient application to phlegmonous inflammations and burns. It was introduced into practice by M. Goulard, a surgeon of Montpellier; and thence its appellation of Goulard's Extract.

LIQUOR PLUM'BI SUBACETA'TIS DILU'TUS, Lond. *Diluted Solution of Subacetate of Lead.*<sup>4</sup>

“Take Solution of Subacetate of Lead, *a fluid drachm*; Distilled Water, *a pint*; Proof Spirit, *a fluid drachm*. Mix.”

LIQUOR SUBACETATIS LITHARGYRI COMPOSITUS, Dub. *Compound Solution of Subacetate of Litharge.*

The same as the London formula, with double the quantity of each of the ingredients.

This preparation, as an article in the pharmacopœia, is superfluous, every surgeon being in the habit of ordering lotions with different proportions of the solution of subacetate of lead, according to the circumstances of the case.

---

## PRÆPARATA E ZINCO.

### PREPARATIONS OF ZINC.

CALAMINA PRÆPARATA, Lond. *Prepared Calamine.*

“Calcine the Calamine, and beat it to powder; then bring

---

<sup>1</sup> *Nicholson's Journal*, xi. 75.

<sup>2</sup> *Ibid.* vi. 223.

<sup>3</sup> The nature of the salt in this solution was first pointed out by Scheele, who changed a solution of the acetate of lead into Goulard's extract, by keeping it in a plate of lead for the space of a day; but this experiment was overlooked until Dr. Bostock's analysis of the preparation. An excellent mode of preparing it is employed in the French hospitals. Three parts of acetate of lead is dissolved in a sufficient quantity of hot distilled water, and to the solution one part of semi-vitreous oxide of lead is added, in fine powder. The whole is then evaporated until it marks 28° of Beaumé's areometer; and when cold, is filtered. Vide *Journ. de Pharm.* Dec. 1816. p. 565.

<sup>4</sup> *Aqua lithargyri acetati composita*, P. L. 1787.



it into the state of a very fine powder, in the manner directed for the preparation of Chalk.”

CARBONAS ZINCI IMPURUS PRÆPARATUS, E. *Prepared Impure Carbonate of Zinc.*

“Impure Carbonate of Zinc, roasted by those who make brass, being rubbed to powder in an iron mortar, and levigated with a little water on a porphyry, is to be put into a large vessel, and water poured over it, which, after frequently agitating the vessel, is to be poured off loaded with the powder. The fine powder which subsides after the water has remained at rest, is then to be dried. The coarse, which the water cannot suspend, is to be again levigated, and treated as before.”

LAPIS CALAMINARIS PRÆPARATUS, Dub. *Prepared Calamine Stone.*

“Reduce Calcined Calamine-stone to powder, and separate the very fine parts in manner directed for the preparation of chalk.”

*Syn.* Calamine préparé (F.), Galmei (G.), Kalmei (Dutch), Galmija (Russ.), Calamina (I.), Calamina (S.).

The nature of this ore of Zinc has been already stated. (Part ii.) As it is frequently used in the form of a dry powder to excoriations, ichorous ulcers, and superficial inflammations, dusted on the part, it requires to be rendered extremely fine.

Officinal preparations. *Ceratum Calaminæ*, L. E. *Unguentum Calaminaris*, D.

OXIDUM ZINCI IMPURUM PRÆPARATUM, Edin. *Prepared Impure Oxide of Zinc.*

“It is prepared in the same manner as the Impure Carbonate of Zinc.”

This substance, the nature of which has been already stated (Part ii.), is used for the same purposes as the former article.

ZINCI OXYDUM, Lond. *Oxide of Zinc.*<sup>1</sup>

“Take of Sulphate of Zinc, *one pound*; Solution of Ammonia, *one pint*, or as much as may be required; Distilled Water, *one pint*. Dissolve the Sulphate of Zinc in the Distilled Water, and add as much Solution of Ammonia as may be requisite to precipitate the Oxide of Zinc entirely. Pour off the Solution, wash the powder repeatedly with distilled water, and dry it upon a sand-bath.

OXIDUM ZINCI, Edin. *Oxide of Zinc.*

“Let a large crucible be placed in a furnace filled with burning coals, in such a manner as to be somewhat inclined

<sup>1</sup> The ancients, who were acquainted with it, called it Pompholyx; and by the early chemists, it was named *Nihil album*, *Lana philosophica*, and *Floras Zinci*. *Zincum calcinatum*, P. L. 1787. In Holland it was prepared as a secret remedy, and sold under the names *Arcanum Ludemanni* and *Luna fixata*, until Gaubius made public its composition.



to its mouth, and when the bottom of it is heated to a moderate degree of redness, throw into it a piece of Zinc about the weight of one drachm. The Zinc is soon inflamed, and converted into white flocculi, which are occasionally to be removed from the surface of the metal by means of an iron spatula, that the combustion may be more complete; and when the inflammation is over, remove the Oxide of Zinc from the crucible. Throw in then another piece, and let the operation be repeated as often as is necessary. Finally, let the Oxide of Zinc be prepared in the same manner as the Impure Carbonate of Zinc."

*Dublin.*

"Take of Zinc broken into small pieces, *any quantity*. Throw these, at intervals, into a sufficiently large crucible heated to whiteness, and placed with its mouth inclined towards the mouth of the furnace. After each piece of Zinc is thrown in, invert over the crucible another crucible, but loosely so as not to exclude the air. Preserve the light, very white, sublimed powder for use."

*Syn.* Oxide de Zinc (*F.*), Weisser Zinkoxyd (*G.*), Per Ossido di Zinco; fiori di Zinco (*I.*).

In the process of the London College, too small a quantity of water is ordered for the solution of the Sulphate of Zinc. It is requisite not to add too much Ammonia, as it redissolves the precipitates. Mr. Brande properly remarks, that the quantity of water ordered is too small, and that the word *Congium* should be substituted for *Octarium*.<sup>1</sup> In the two other processes the crucible must be heated above 700° of Fahrenheit, which is the point of ignition of zinc. At this temperature the metal inflames, burning with a dazzling white and green flame; and by attracting the oxygen of the air is converted into a white oxide, which is partly volatilized in the form of very light flocculi. The elevation of these flocculi, however, is owing to the current of air excited by the force of the combustion; for the oxide itself is not volatile, but accumulates in the crucible so rapidly, that it must be withdrawn to allow the access of the air for keeping up the combustion. If the crucible be sufficiently capacious, there is no necessity for covering it with another, by which the operation is always impeded.<sup>2</sup>

*Qualities.*—Oxide of zinc thus prepared is inodorous, insipid, of a pure white colour, infusible in the fire, insoluble in

<sup>1</sup> *Manual of Pharmacy*, p. 313.

<sup>2</sup> This oxide may also be readily prepared by dissolving zinc in diluted sulphuric or nitric acid, and precipitating by potass, a process proposed by Marabelli, Professor of Pharmacy at Pavia, in 1798. The washed precipitate is oxide of zinc, containing, according to Vauquelin, 0.21 of oxygen.



water and alcohol, but entirely soluble in acids, and is not altered by exposure to the air. According to Proust, 100 parts of it consist of 80 of zinc, and 20 of oxygen; or 100 zinc + 25 oxygen. It often contains small portions of carbonic acid.<sup>1</sup> It is often adulterated with chalk, and sometimes contains white lead. By pouring sulphuric acid on the specimen, the first is discovered by the effervescence which is excited, the second by an insoluble sulphate of lead being formed.

*Medical properties and uses.* — Oxide of Zinc is tonic and antispasmodic; and has been advantageously used in chorea, epilepsy<sup>2</sup>, and some other spasmodic affections. It has been employed in whooping-cough on the continent; and Lœffler recommends it to be used externally as well as internally in that disease. He employs a liniment composed of linseed oil, and oxide of zinc. It is chiefly used as an external application. (See *Ung. Zinci*.)

The dose, as an internal remedy, may be from gr. j. to grs. vj. given twice a day.

Officinal preparation. *Unguentum Zinci*, L. E. D.

ZINCI SULPHAS<sup>3</sup>, Lond. *Sulphate of Zinc*.

“Take of Zinc broken into small pieces, *four ounces*; Sulphuric Acid, *by weight six ounces*; Water, *four pints*. Mix them in a glass vessel, and the effervescence being over, filter the solution through paper; then boil it until a pellicle begin to form on the surface, and set it aside to crystallize.”

*Edinburgh.*

“Take of Zinc cut into small pieces, *three parts*; Sulphuric Acid, *five parts*; Water, *twenty parts*. Mix them, and the effervescence being finished, digest for a short time on hot sand. Then filter the decanted solution through paper, and after due evaporation, set it apart that crystals may be formed.”

*Dublin.*

“Take of Zinc reduced to powder in the same manner as Tin, *three ounces*; Sulphuric Acid, *five ounces*; Water, *a pint*. Pour the Acid previously diluted with the Water upon the Zinc, put into a glass vessel; digest for a short time after the effervescence ceases; then evaporate the strained solution, to a proper point, and set it aside to crystallize.”

*Syn.* Sulphate de Zinc (*F.*), Schwefelsaures Zink (*G.*), Solfato di Zinco (*I.*), Vitriolo bianco (*S.*).

The directions of the Dublin College for granulating the

<sup>1</sup> *Annales de Chimie*, xxxv. 51. The more recent experiments of Dr. Thomson make the proportions to be metal 100 + 24.16 oxygen; those of Berzelius, metal 100 + 24.4 oxygen.

<sup>2</sup> *Duncan's Med. Comment.* iii. 216.

<sup>3</sup> *Zincum vitriolatum*, P. L. 1787.



Zinc are to be adopted in preference to those of the other Colleges for dividing it. In these processes, the acid enables the Zinc to decompose the water, and the metal is oxidized by attracting its oxygen, while its hydrogen is disengaged with effervescence. The Oxide thus formed combines with the acid, forming Sulphate of Zinc, which is obtained in crystals by the subsequent evaporation. The greater part, however, of the sulphate of zinc of the shops is prepared on a large scale, and purified in the manner that shall be immediately noticed. It is denominated *White Vitriol* in the language of commerce, and is manufactured largely both in Germany<sup>1</sup> and England. In Germany it is prepared by exposing roasted Blende to the air and humidity; by which means the metal is gradually oxidized, and combined with the sulphuric acid also formed from the sulphur contained in the Blende. The sulphate thus produced is separated from the earthy parts of the Blende by lixiviation, and after being boiled down is crystallized, or rather concreted, into hard granular masses resembling loaf-sugar, which generally contain sulphate of iron, of lead, and sometimes of copper. In England it is prepared generally by the direct combination of its constituents; but although purer than the foreign salt, yet the English White Vitriol, almost always, contains iron. Both kinds are purified by solution in water, and then allowing the solution to evaporate very slowly in an open vessel containing some granulated zinc; the sulphate of lead will subside, and the other foreign salts be decomposed by the metallic zinc. The purified Sulphate of Zinc may be then crystallized by lixiviation and evaporation.<sup>2</sup>

*Qualities.* — Pure Sulphate of Zinc, or rather *Supersulphate*, for it contains an excess of acid, and reddens the vegetable blues, is inodorous, and has a slightly acidulous, styptic, metallic taste. It crystallizes in transparent, colourless, right rhombic prisms<sup>3</sup>, terminated by quadrangular pyramids; effloresces slightly in the air; is soluble in 2.5 times its weight of water at 60°, and in less than its own weight of boiling water. It is decomposed by the alkalies, earths, and hydrosulphurets; and throws down a dirty-looking precipitate from astringent vegetable infusions, with which, therefore, it is incompatible in prescriptions. According to the analysis of Dr. Wollaston, the constituents of 100 parts of the pure crystallized salt are 28.4 of oxide of zinc, 27.3 of acid, and 44.3

---

<sup>1</sup> Beckman, in his *History of Inventions*, says, it was first made at Raimelsberg, in Germany, about the middle of the 16th century. He ascribes the invention to Julius, Duke of Brunswick.

<sup>2</sup> *Aikin's Dictionary of Chemistry*.

<sup>3</sup> *Phillips' Trans. of Pharm.* 1824.



of water. Mr. Phillips makes the proportions to be 30·9 of Oxide of Zinc, 29·4 of acid, and 30·7 of water.

*Medical properties and uses.* — Sulphate of Zinc is tonic and astringent, and in large doses emetic. As a tonic, it is less heating and stimulant than Sulphate of Iron, and thence is preferable in phthisis and other diseases attended with great irritability and general weakness. It is also useful in dyspepsia, fluor albus, and some convulsive affections, as pertussis, chorea, and epilepsy; in which diseases it is generally combined with myrrh, bitter extracts, opium, extract of hemlock, or digitalis, according to the circumstances of the case. As an emetic it operates almost instantaneously, and, therefore, is often employed to empty the stomach at the commencement of the paroxysm of intermittent fever, and in other cases in which quick vomiting is required. As an external application, this salt dissolved in rose-water, in the proportion of grs. iſs. to f℥j. of rose-water, forms an excellent collyrium in the latter stage of ophthalmia, after the inflammatory action has subsided; it is a good injection in a similar stage of gonorrhœa; and a lotion in some kinds of superficial inflammations. In double the strength, this solution is the best application that can be used to scrophulous tumours, after they have suppurated, and the abscess has been discharged.

The dose to produce vomiting is from grs. x. to ʒss. and as a tonic from gr. j. to grs. ij. may be given twice a day.

Officinal preparations. *Solutio Sulphatis Zinci*, E. *Liquor Aluminis compositus*, L. *Solutio Acetatis Zinci*, E. D.

**SOLUTIO SULPHATIS ZINCI**, Edin. *Solution of Sulphate of Zinc.*

“Take of Sulphate of Zinc, *sixteen grains*; Water, *eight ounces*; Diluted Sulphuric Acid, *sixteen drops*. Dissolve the Sulphate of Zinc in the Water, and having added the Acid, filter the solution through paper.”

This formula is given under the idea of the common Sulphate of Zinc, (which often contains some excess of oxide, and some oxide of iron,) being employed. The superabundant oxide, if present, is dissolved by the acid, so that a solution of an uniform strength is always obtained. It is rather too strong for the purposes of a collyrium in chronic ophthalmia; and the addition of the acid renders it less fit to be used as an injection in gonorrhœa.

**SOLUTIO ACETATIS ZINCI**, Edin. *Solution of Acetate of Zinc.*

“Take of Sulphate of Zinc, *one drachm*; Acetate of Lead, *four scruples*; Distilled Water, *twenty ounces*. Dissolve. Mix the salts separately in ten ounces of the Water; then mix the solutions, and after the precipitate subsides, filter.”



*Syn.* Dissolution d'Acetate de Zinc (*F.*), Liquore de l'Acetato di Zinco (*I.*).

In this process a double decomposition takes place: the sulphuric acid of the Sulphate of Zinc unites with the oxide of lead of the acetate of lead, whilst its acid combines with the disengaged oxide of zinc. The former salt being insoluble, is precipitated in the form of a heavy white powder, but the Acetate of Zinc remains dissolved; and thus its solution, which is colourless and limpid, is easily separated by filtration.

*Medical properties and uses.* — This solution is astringent; and was long employed before it was introduced into the pharmacopœia, and even before its nature was clearly understood. It is an useful collyrium in chronic ophthalmia, and in the acute variety of this disease after the inflamed vessels are unloaded, and the inflammatory action subdued. It is also an useful injection in the advanced stage of gonorrhœa.

TINCTURA ACETATIS ZINCI, Dub. *Tincture of Acetate of Zinc.*

“Take of Sulphate of Zinc, Acetate of Kali, of each *an ounce*. Rub them together, and add of Rectified Spirit of Wine, *one pint*. Macerate for a week, with occasional agitation, and filter through paper.”

In this process, a double decomposition also takes place, Acetate of Zinc, and Sulphate of Potass being produced; the former of which is dissolved in the spirit, while the latter remains undissolved, and, therefore, is easily separated. It is a tedious process, and possesses no advantages over the former to recommend it.

*Medical properties and uses.* — This tincture is astringent; but requires to be diluted with water, before it can be used either as a collyrium or an injection. It might be advantageously employed as an internal remedy in dyspepsia, and other debilities of the stomach.

---

## PREPARATIONS OF SULPHUR.

PURE SULPHUR is generally regarded a simple substance; but, according to Sir H. Davy's experiments, it is a triple compound of oxygen, hydrogen, and a peculiar, unknown base. It unites readily with metals, some oxides, earths, and the fixed alkalies, forming compounds which have been denominated *Sulphurets*. These are formed by the fusion of the substances in a dry state; and the compounds require to be carefully preserved from the atmosphere, as they attract moisture from it, deliquesce, and are decomposed. When, however, the union of sulphur and alkaline or earthy bases is



effected by means of water, the products are not simple *Sulphurets*, but *Sulphurets* combined with *Sulphuretted Hydrogen*, and have been named *Hydroguretted Sulphurets*. They are equally susceptible of decomposition by exposure to the air as the *Sulphurets*.

OLEUM SULPHURA'TUM<sup>1</sup>, Lond. *Sulphuretted Oil*.

“Take of Washed Sulphur, *two ounces*; Olive Oil, *a pint*. Add the Sulphur gradually to the oil heated in a very large iron pot, and stir the mixture after each addition till they have united.”

*Edinburgh.*

“Take of Olive Oil, *eight parts*; Sublimed Sulphur, *one part*. Boil them with a gentle heat in a large iron vessel, stirring constantly, until they unite.”

*Syn.* Huile sulphur (*F.*), Oleo sulfurato (*I.*).

Sulphur is soluble in many fixed oils, but particularly so in linseed oil. This combination was formerly termed a *Balsam*; and was sometimes united with other substances; as, for instance, the oil of Aniseed, in forming the *Anisated Balsam of Sulphur*. Great attention is required in these processes to prevent the mixture from boiling over, or its vapour from catching fire. If either of these accidents occur, the combustion may be stopped, by instantly covering the pot with a close lid. The iron pot should be sufficient to contain thrice the bulk of the ingredients.

*Qualities.* — The odour of this solution of Sulphur is extremely foetid, and the taste acrid. It is of a reddish-brown colour; has a thick viscid consistence; and, when heated, emits sulphuretted hydrogen. When it is much concentrated, the sulphur crystallizes in octahedrons.

*Medical properties and uses.* — Sulphuretted oil is stimulant, and externally detergent. It was formerly, when regarded as Balsamic, recommended in catarrh, asthma, and phthisical affections: but its internal use is now properly exploded. It is sometimes still externally applied for cleansing foul ulcers. The dose was from *℥v.* to *℥xxx.* taken in water.

POTA'SSÆ SULPHURETUM<sup>2</sup>, Lond. *Sulphuret of Potass.*

“Take of Washed Sulphur, *an ounce*; Subcarbonate of Potass, *two ounces*. Rub them together, and place the mixture over the fire in a covered crucible until they unite.”

SULPHURETUM POTASSÆ, Edin.

“Take of Subcarbonate of Potass, *two parts*; Sublimed

<sup>1</sup> Balsamum Sulphuris simplex, P. L. 1745.

<sup>2</sup> Kali Sulphuratum, P. L. 1787.



Sulphur, *one part*. Rub them together, and put them into a large covered crucible, to which, having adapted a cover, apply the fire cautiously, until they melt. Preserve the mass in a well-closed vessel."

SULPHURETUM KALI, Dub. *Sulphuret of Kali*.

"Take of Subcarbonate of Kali, Sublimed Sulphur, each *two ounces*. Having mixed them together, put them into a crucible, and having adapted to it a cover, expose it to a fire gradually raised until they unite."

*Syn.* Sulphure de Potasse (*F.*), Schweflichtes Kali (*G.*), Solfuro di Potassa (*I.*).

This Sulphuret cannot be properly formed by following the directions of any of the Colleges; for, to render the combination complete, it is necessary to expose the Subcarbonate in a crucible to a red heat, previously to its being rubbed with the sulphur: the water of the Subcarbonate is thus dissipated, and at the same time a portion of the Carbonic Acid is expelled, both of which, when not driven off, alter the product. When the fusion is effected, the mixture is to be poured upon a marble slab, and, as soon as it concretes, the mass must be broken in pieces and preserved in a closely-stopped bottle. In this process the Potass, after the Carbonic Acid is totally expelled, is partly decomposed, and the oxygen of that portion combining with a portion of the sulphur forms sulphuric acid, which combines with the undecomposed portion of the Potass and forms Sulphate of Potass. The remainder of the sulphur combines with the potassium or metallic base of the Potass, and forms a Sulphuret; thus making the result of the process a compound of Sulphate of Potass and Sulphuret of Potassium<sup>1</sup>, with perhaps an excess of sulphur; varying in different proportions.

*Qualities.* — Well prepared Sulphuret of Potass, as this preparation is erroneously termed, is inodorous while dry; but when moistened or dissolved in water, a partial decomposition of both the water and the sulphuret is effected, and sulphuretted hydrogen is formed. It has an acrid, bitter taste; changes the vegetable blues to green; is hard, brittle, breaking with a glassy fracture, of a liver-brown colour, and stains the skin brown.<sup>2</sup> By exposure to the air it attracts moisture; its colour changes to a pale green, the foetid odour noticed above is emitted, and it is gradually converted into hydroguretted sulphuret of potass, combined with a small

---

<sup>1</sup> A simple or monosulphuret is formed by decomposing sulphate of potass by means of charcoal at a high temperature. The oxygen of both the Potass and the Sulphuric Acid is carried off by the carbon, whilst the Sulphur and the Potassium remain in combination as a Sulphuret.

Hence its old name, *Hepar Sulphuris*.



portion of sulphate of potass. It is also decomposed by all the acids which combine with the potassium, and the sulphur is precipitated. In a violent heat the sulphur sublimes, leaving the potass.

*Medical properties and uses.* — Sulphuret of Potass is expectorant and diaphoretic. It has been frequently given in chronic asthma and chronic catarrh, without much benefit; but has been found useful in arthritic, rheumatic, and herpetic affections; and in combination with Conium as a palliative in cancerous cases.<sup>1</sup> It has also been employed in France, for the cure of scabies, in the form of an ointment made with one part of the sulphuret, sixteen of soap, and thirty-two of oil. From a theory founded on its chemical action on metallic salts out of the body, it has, also, been strongly recommended as an antidote against arsenical, saturnine, and mercurial preparations, when these have been taken in doses sufficient to produce deleterious effects; but it has, hitherto, been too seldom employed to ascertain its real value in these cases.

The usual dose is grs. iij. or grs. iv. combined with soap, in the form of pills, for the first-mentioned cases; or from grs. v. to grs. x. as an adjunct to Conium in cancer, given several times a day.

SULPHUR LOTUM, Lond. *Washed Sulphur.*<sup>2</sup>

“Take of Sublimed Sulphur, *a pound*. Pour upon it Boiling Water, that the Acid, if there be any, may be entirely washed away; then dry it.”

SULPHUR SUBLIMATUM LOTUM, Edin. *Washed sublimed Sulphur.*

“Take of Sublimed Sulphur, *one part*; Water, *four parts*. Boil the Sulphur for a short time in the Water; then pour off this Water, and by repeated affusions of cold Water, wash away all the Acid: lastly, dry the Sulphur.”

*Dublin.*

“Let Warm Water be poured upon Sublimed Sulphur, and the washing be repeated as long as the water employed shall appear acid. This is known by means of litmus. Dry the Sulphur on bibulous paper.”

*Syn.* Soufre lavé (*F.*), Geewaschner Schwefel (*G.*), Zolfo lavato (*I.*).

In subliming sulphur, a small portion of it is apt to be acidified, by attracting the oxygen of the heated air of the vessels, or the chamber in which the process is conducted. The quantity is however very minute, and is completely removed by the above processes. The sulphur does not afterwards undergo

<sup>1</sup> Pearson's *Practical Synopsis*, &c. i. 283.

<sup>2</sup> Flores Sulphuris loti, P. L. 1787.



any change from exposure to the air at the ordinary temperature of the atmosphere.

**SULPHUR PRÆCIPITATUM**, Lond. *Precipitated Sulphur.*<sup>1</sup>

“Take of Sublimed Sulphur, *a pound*; Fresh Burnt Lime, *two pounds*; Water, *four gallons*. Boil the Sulphur and the Lime together in the Water; then filter the liquor through paper, and drop into it as much Muriatic Acid as may be sufficient to precipitate the Sulphur. Finally, wash this with repeated affusions of Water, until it become tasteless.”

*Syn.* Soufre précipité (*F.*), Schwefel niederschalg (*G.*), Zolfo Precipitato (*I.*).

In the first part of this process, a hydroguretted sulphuret of lime is produced, by the combination of the lime and sulphur occasioning a decomposition of a part of the water, the hydrogen of which unites with a portion of the sulphur, and forms a hydrosulphuret; while the oxygen with another portion forms sulphuric acid that combines with part of the lime: and thus the solution contains a small portion of sulphate of lime, and a sulphuret of lime or rather the base of lime, *Calcium*, combined with sulphuretted hydrogen. This hydroguretted sulphuret is then decomposed by the muriatic acid, which unites with the lime, and forms a soluble muriate, while the sulphur is precipitated, and sulphuretted hydrogen gas disengaged.

*Qualities.*—Precipitated sulphur is white, with a very slight greenish tinge. When heated in a retort in a low heat, it acquires the colour of common sulphur, and water is deposited in the beak of the retort; from which circumstance, and the same degree of whiteness being produced when sulphur is sublimed into a vessel filled with steam, there is reason for supposing that precipitated sulphur owes its whiteness to the presence of a little water. It differs in no other respect from sublimed sulphur, and is an unnecessary refinement for the sake of appearance in the composition of ointments.

**AQUA SULPHURETI KALI**, Dub. *Water of Sulphuret of Kali.*

“Take of Sublimed Sulphur, *half an ounce*; Water of Caustic Kali, *nine fluid ounces*. Boil them together for ten minutes, and filter through paper. Preserve the preparation in well-stopped phials. The specific gravity of this liquor is, to that of distilled water, as 1120 to 1000.”

*Syn.* Liquore di solfuro di Potassa (*I.*).

The name given to this preparation conveys an erroneous idea of its nature. When an alkaline sulphuret is dissolved in water, changes exactly similar to those we have mentioned (*Sulphur præcipitatum*) as taking place during the solution of

<sup>1</sup> Lac Sulphuris, P. L. 1720. Sulphur præcipitatum, P. L. 1787.



an earthy sulphuret occur, altering the character of the product; and as the same happen by the direct combination of sulphur with a liquid alkali, this preparation is not a simple aqueous solution of sulphuret of potass, but, in fact, a solution of *hydroguretted sulphuret of potass*, or sulphuret of potass combined with sulphuretted hydrogen.<sup>1</sup>

*Qualities.* — This Solution has a slightly foetid odour, and a nauseous, acrid, bitter taste. Its colour is reddish-yellow, approaching to deep orange; its feel soapy; and it stains the cuticle a greenish-black. Acids decompose it, precipitating the sulphur, and disengaging a portion of sulphuretted hydrogen gas; and it is also decomposed by exposure to the air, the oxygen of which being absorbed by the sulphur forms sulphuric acid, which produces a sulphate with the potass; so that in process of time the whole is changed into a solution of sulphate of potass. Hence the necessity of preserving it in well-stopped phials.

*Medical properties and uses.* — This solution does not differ in its medicinal properties from the solid sulphuret of potass. It is, however, chiefly employed as an external application; and as such has been found very beneficial in porrigo, scabies, and herpetic eruptions. When given internally, the dose is from *mxx.* to *fziss.* twice a day.

**HYDRO-SULPHURETUM AMMONIÆ**, Edin. *Hydro-Sulphuret of Ammonia.*

“Take of Water of Ammonia, Sulphuret of Iron, of each *four ounces*; Muriatic Acid, *eight ounces*; Water, *two pounds and a half*. Pour the Acid, previously mixed with the Water, on the Sulphuret, and transmit the gas evolved, through the Water of Ammonia. Preserve the solution in well-stopped phials.”

**SULPHURETUM FERRI**, Dub. *Sulphuret of Iron.*

“Take of Filings of Iron, *six ounces*; Sublimed Sulphur, *two ounces*. Mix them, and expose them in a covered crucible to a gentle heat until they unite.”

**HYDRO-SULPHURETUM AMMONIÆ**, Dub. *Hydro-Sulphuret of Ammonia.*

“Take of Sulphuret of Iron in coarse powder, *four ounces*; Muriatic Acid, *seven fluid ounces*; Water, *two pints*; Water of Caustic Ammonia, *four ounces*. Put the Sulphuret of Iron into a matrass, and gradually pour over it the acid diluted with water; and in a proper apparatus transmit the gas evolved from it through Water of Ammonia. Toward the conclusion of the operation apply a moderate heat to the mattress.”

The proportions of the ingredients ordered in the Dublin

---

<sup>1</sup> It was formerly denominated *Liquid Hepar*, or *Liquid Liver of Sulphur*.



formula for producing the Sulphuret of Iron enable them to unite at a low heat; and the combination is attended with a brilliant ignition, which takes place without the presence of air. The compound is of a grey colour intermixed with yellow, and resembles bronze; has a metallic lustre, and a crystalline texture, with a considerable degree of brittleness, and when pulverized yields a black powder. According to Proust, 100 parts of it consist of 62·5 of Iron, and 37·5 of Sulphur.<sup>1</sup> The addition of the diluted muriatic acid, by oxidizing the iron, enables it to decompose the water, the hydrogen of which dissolving, a part of the sulphur escapes, in the form of sulphureted hydrogen gas, which combines at a low temperature with the ammonia of the solution through which it is made to pass. Mr. Cruikshank<sup>2</sup> advises the Sulphuret of Iron to be prepared “by raising a piece of iron in a smith’s forge to a white heat, and then rubbing it against the end of a roll of sulphur; the iron at this temperature, immediately combines with the sulphur, and forms globules of pyrites (*sulphuret*), which should be received into a vessel filled with water; these globules are to be reduced to powder, and introduced into the proof, to which a sufficient quantity of the muriatic acid is to be added.” Various other modes have been also recommended for the preparation of the sulphuret; but the process directed by the Dublin College is easy and perfectly adequate for the purpose.

*Qualities.* — Hydro-sulphuret of Ammonia is of a dark green colour; has a very foetid odour, and an acrid, disagreeable taste. It is decomposed by the acids.

*Medical properties and uses.* — This preparation is a powerful sedative, lessening the action of the stomach, and of the arterial system in a remarkable degree; and even in moderate doses producing sickness, vomiting, and vertigo. It was first proposed as a remedy by Mr. Cruikshank, with the view of diminishing the morbid appetite and powerful action of the digestive organs, which attend those labouring under diabetes mellitus; and its subsequent use has been confined to the treatment of that disease. The dose to an adult should not at first exceed  $\mathfrak{m}\mathfrak{v}$ . or  $\mathfrak{m}\mathfrak{vj}$ . given in a large tumbler of water, three or four times a day; and the number of drops should be gradually increased, until a slight degree of giddiness takes place, when any further increase must be stopped.

AQUA SULPHURETI AMMONIÆ, Dub. *Water of Sulphuret of Ammonia.*

“Take of Lime recently prepared, Muriate of Ammonia in powder, of each *four ounces*; Sublimed Sulphur, Hot

<sup>1</sup> *Journal de Physique*, liii. 89.

<sup>2</sup> *Rollo on Diabetes and Lues Venerea*.



Water, of each *two fluid ounces*. Sprinkle the Water on the Lime, in an earthen vessel, and cover it until the Lime fall to powder; mix this when cold by trituration with the Sulphur and Muriate of Ammonia, avoiding the vapours; then put the mixture into a retort, and distil with a strong heat suddenly raised. Preserve the liquor thus obtained in a phial closely stopped with a glass stopper."

In this process the Lime decomposes the Muriate of Ammonia, attracting its acid, and forming a Muriate of Lime, while the disengaged Ammonia unites with the sulphur, one part of which, however, is converted into a hydro-sulphuret, by hydrogen arising from partial decomposition of the water, which, combining with the sulphuret of ammonia, thus produces a hydroguretted sulphuret of ammonia. It was formerly known by the name *Fuming Liquor of Boyle*, having been first prepared by Mr. Boyle.

*Qualities.* — This liquid is of a deep orange-colour, has a strong, ammoniacal, foetid odour, and emits white fumes, owing, as Berthollet ascertained, to an excess of Ammonia. The addition of an acid precipitates sulphur, and occasions the disengagement of sulphuretted hydrogen gas. Water of Sulphuret of Ammonia consists of hydro-sulphuret of ammonia holding an excess of sulphur, which it gradually deposits. As it loses the property of fuming, it becomes nearly pure Hydro-sulphuret of Ammonia.<sup>1</sup>

We are ignorant of any medicinal use to which this preparation has been applied.

---

## VEGETABILIA.

### VEGETABLES.

THE collection of vegetables cannot be attended to by the apothecary, and, consequently, the directions necessary for that purpose are of less importance to him than a knowledge of the botanical characters of plants, and the appearances they assume when they are collected under proper circumstances and well dried: for inert plants are often introduced by the collectors among those which possess the most active properties; and from a careless or an improper mode of drying plants, the medicinal virtues of the majority of them are altogether destroyed. When, however, opportunities permit the

---

<sup>1</sup> Thomson's *Chemistry*, 4th edit. iii. 380.



apothecary to be his own collector, these should not be neglected; and the collection and drying of some plants, particularly of foxglove and hemlock, should never be left to the common collector. The following general directions are, therefore, given in the London Pharmacopœia for collecting vegetable substances.

“VEGETABLES are to be gathered from the soil and situations where they spontaneously grow, in a dry season, and when no dew is upon them: they are to be collected every year, and any which shall have been longer kept are to be thrown away.” This is a direction of the utmost importance as far as regards the more active vegetable remedies, such as digitalis; for the skill of the most experienced physician will little avail the patient, if the drug which is ordered have lost its medicinal properties from being too long kept.

“ROOTS, for the most part, are to be dug up before their stems or leaves shoot forth.” This direction may be followed when the roots are cultivated: but if the prior directions be attended to, it is not easy to conceive by what means the roots are to be discovered before the stems and leaves are put forth. The object of the order is the obtaining the roots with their active principles in the most concentrated state; and this may be effected by digging them up late in autumn, or early in winter, after the sap is completely detruded to the root, and the stem is withered, but yet attached to the root, by which its situation is pointed out. If any change in the composition of the juices takes place during the cessation of vegetation in winter, it is probable that the same will happen, if the root, after being dug up, be preserved in sand.

“BARKS are to be collected at that season in which they are more easily separated from the wood.” Spring is the season here alluded to; as at this time, after the sap begins to ascend, the bark is in general very easily separated. But a more important reason may be given for preferring this period, as in spring the active principles deposited in the proper cells of the bark are most abundant: thus, oak bark collected in spring contains four times more tannin than that which is collected in winter.<sup>1</sup>

“LEAVES are to be gathered after the flowers have expanded, and before the seeds are mature.” These should be in the most perfect state, free from disease, and full grown.

“FLOWERS are to be gathered when just opened.” There is, however, one exception to this rule as far as it refers to the red rose, which must be gathered before the buds are expanded.

---

<sup>1</sup> Vide Biggin's Table, *Phil. Trans.* 1799.



“SEEDS are to be collected when they are ripe, and before they drop from the plant. They ought to be preserved in their seed vessels.”

*VEGETABILIIUM PRÆPARATIO, Lond.*

*Preparation of Vegetables.*

“VEGETABLES, soon after they are gathered, except those which are to be used in the recent state, are to be lightly spread out, and dried as quickly as possible, with a heat so gentle that their colour will not be altered; and then preserved in proper situations or vessels, where the light and moisture are excluded.” When plants cannot be dried immediately on being gathered, they should be revived by immersing their stalks in water for twelve hours. When the leaves are the parts intended to be employed, these are then to be carefully freed from the stalks, and laid in thin layers in baskets of willow stripped of its bark, in a drying room kept quite dark. They should then be exposed to a temperature of 140° Fahr. for six or eight hours. When the leaves begin to shrivel, they should be turned, and the same temperature continued, until they crumble readily in the hand. When the process has been well conducted, the leaves should retain their green colour and their medicinal properties. The vessels best adapted for preserving them are oil jars made perfectly clean and dry; closely covered and kept in a dry warm situation. It is better to preserve those leaves, the virtues of which are particularly connected with their colour, as hemlock and fox-glove, in this state, than in the form of powder, a small portion only being occasionally powdered for current use.<sup>1</sup>

“ROOTS, which are required to be preserved fresh, should be buried in dry sand. The SQUILL ROOT (*bulb*), before it is dried, is to be denuded of the arid coats, and cut transversely into thin slices. The bulb of *Colchicum*, dug up in July or August, should be cut into thin transverse slices, dried without heat, or in a very gentle heat, and preserved in well-stopped bottles.

“PULPY FRUITS, if they be unripe, or ripe and dried, are to be placed in a damp situation, until they become soft: then the pulp is to be pressed out through a hair sieve; afterwards boiled with a gentle heat, frequently stirring; and finally, the water evaporated in a water-bath, until the pulp acquire a proper consistence.

---

<sup>1</sup> The above is the method adopted by Mr. Battley, of Fore Street, whose attempts to improve Pharmacy deserve the thanks of the profession.



“ Over the bruised pods of CASSIA, pour boiling water, so as to wash out the pulp, which is to be first pressed through a sieve with large holes, and afterwards through a hair sieve; then dissipate the water in a water-bath, until the pulp acquire a proper consistence.

“ Press through a sieve the pulp or juice of ripe and fresh fruits without boiling them.”

### VEGETABILIMUM EXSICCATIO, *Edin.*

#### *The Drying of Herbs and Flowers.*

“ HERBS and flowers are to be dried by the gentle heat of a stove or a common fire, in such a quantity at once as will admit of the operation being very quickly finished: for by this means their powers are better preserved; the indication of which is the perfect preservation of their natural colour.

“ The leaves of HEMLOCK (*Conium maculatum*), and of other plants containing a subtile, volatile matter, are, when dried, to be immediately reduced to powder, and preserved in well-stopped glass vessels.”

“ The root (*bulb*) of the SEA-SQUILL (*Scilla maritima*), freed from its external coat, is to be cut transversely into thin slices. The indication of its being properly dried is the retention of its bitterness and acrimony after it has become friable.” The directions of the Dublin College are similar. (See *Pulvis Scillæ* among the Powders.) After the squill has been properly dried, in which operation it loses seven-eighths of its weight, it must be kept in a dry place, as it is apt to retain its moisture in some degree, and become mouldy. It cannot, however, be long preserved in the state of powder without becoming almost inert.

### HERBARUM EXSICCATIO, *Dub.*

#### *The Drying of Herbs.*

“ PUT the fresh leaves of the herb, gathered when it is in flower, into paper bags, and expose them to a low heat for an hour; then strew them lightly upon a sieve, and dry them as quickly as possible, taking care that their green colour be not injured by too much heat; but if the herbs are to be used under the form of powder, let them be immediately powdered, and the powder preserved in well-closed, opaque phials.

“ Herbs and flowers from which oils and distilled waters are to be obtained, should be dried as soon as they are collected.”



*PULPARUM EXTRACTIO, Edin.**Extraction of Pulps.*

“FRUITS which afford a pulp, if unripe, or if ripe and dry, are to be boiled in a small portion of water, till they become soft; then the pulp is to be pressed through a hair sieve, and afterwards boiled in an earthen vessel with a gentle heat, stirring frequently to prevent it from burning, until it acquires the consistence of honey.

“In like manner the pulp of *CASSIA FISTULA* is to be boiled out from the bruised pod, and then brought to a proper consistence by evaporating the water.

“The pulps of recent and ripe fruits are to be pressed through a sieve without being previously boiled.”

*Dublin.*

“If the Fruits, the pulps of which are to be extracted, be unripe, or ripe and dry, they are to be boiled in a small portion of water until they become soft; and then the pulps, pressed through a hair sieve, are to be reduced by slow evaporation to a proper thickness.”

*SUCCI SPISSATI, Edin.**Inspissated Juices.*

“BEAT the fresh substance, and press it strongly through a canvass bag, in order to obtain the juice; which being put into a wide shallow vessel, and heated by means of boiling water saturated with sea-salt, is to be reduced to the consistence of honey. The mass when cold is to be put into glazed earthen vessels, and moistened with strong alcohol.”

The juices of fresh vegetables obtained by expression contain, besides the sap of which they chiefly consist, mucilage, fecula, extractive matter, and the other proper juices of the plant. When newly expressed, these matters are mixed together, and form a viscid, heterogeneous fluid, which gradually separates, by rest, into two parts; the one formed of a deposit of all the insoluble components of the juice generally involved in mucilaginous matter; the other a clear liquor, consisting of water, holding some mucilage in solution, with the acids and salts, if any, and other soluble principles of the juice. As the clear liquor is that which is wished to be obtained for medical use, it is separated by first decanting it from the deposit, then filtering it repeatedly through a linen cloth, and adding about one-fortieth part of its weight of



alcohol; after which it is allowed to remain at rest for some time, and again filtered previous to being put into the bottles in which it is intended to be preserved. The bottles should be kept in a cool cellar, and sunk up to the neck in sand.

Various other methods, also, are employed for depurating vegetable juices; but as these preparations are now almost obsolete, we do not think it necessary to detail them. By whatever means they are prepared, vegetable juices undergo chemical changes, and spontaneous decompositions from keeping, which must necessarily affect their virtues as medicines. They are therefore properly rejected from all the Pharmacopœias.

The articles given in the Edinburgh Pharmacopœia under the title *Succi spissati* being associated by the London College with the extracts, and the difference between these preparations being scarcely sufficient to constitute a generic distinction, we have thought it proper not to alter the London arrangement in this respect, and have therefore placed the whole under the title *Extracts*.

---

## GUM RESINS.

### GUMMI-RESINÆ, Lond.

“Separate OPIUM very carefully from all extraneous matters, particularly those adhering to its outside. Let it be kept in a SOFT state fit for forming pills; and in a HARD state, such as can be produced by drying it in the heat of a water-bath, capable of being rubbed into powder.

“Those GUM-RESINS are to be preferred which can be selected in such a state of purity as to require no purification. If, however, they appear to be less pure, boil them in water until they soften, and squeeze them by a press through a hempen bag: then set them aside, that the resinous part may subside. Pour off the supernatant fluid, and evaporate it by the heat of a water-bath, adding the resinous part towards the end of the operation, and mixing it intimately with the gummy part so as to form one mass.

“Those GUM-RESINS which easily liquify are to be purified by putting them into an ox bladder, and holding them in boiling water, until they become soft enough to be freed from their impurities by pressing them through a hempen bag.

“Dissolve the BALSAM OF STORAX in rectified spirit, and



strain; then distil off the spirit by a gentle heat, until the Balsam acquire a proper consistence."

Gum-resins which require to be treated in the above manner are unfit for internal use, and should be kept chiefly for forming plasters and for other external purposes. The degree of heat, although not more than sufficient for the liquefaction of the substances, is nevertheless enough to dissipate many of their odorous and volatile principles, and occasion some changes of composition. The directions for the treatment of opium are sufficient for freeing it from all the grosser impurities; and as the remedy remains unaltered, its strength is preserved unimpaired; whereas when spirit is employed, as was formerly ordered in the London Pharmacopœia, and is still ordered by the Dublin College, it always suffers in efficacy. The direction to preserve it in the soft state is injudicious; for the uniformity of strength of the preparation can rarely be determined, and must vary according as accidental circumstances evaporate more or less that fluid. Pills are easily made by beating up the powdered opium with mucilage.

#### OPIUM PURIFICATUM, Dub. *Purified Opium.*

*Syn.* Opium pur (*F.*), Mohnsaft (*G.*), Oppio puro (*I.*), Oppio puro (*S.*).

"Take of Opium cut into small pieces, *one pound*; Proof Spirit, *twelve pints*. Digest with a gentle heat and frequent agitation, until the Opium be dissolved: then filter the solution through paper, and distil it from a retort to separate the Spirit; pour out the residuary liquor, and evaporate it until the extract be of a proper thickness. Purified Opium must be kept in two states; one *soft*, proper for forming pills, and one *hard*, capable of being reduced to powder."

#### STYRAX PURIFICATA. Dub. *Purified Storax.*

*Syn.* Styrax in Kërnern (*G.*), Storace depurato (*I.*).

"Digest the Storax in tepid water until it softens; then press it in a press between iron plates heated with boiling water; and, finally, separate it from the water."

In this process a considerable part of the Benzoic Acid of the Storax is dissipated by the heat of the iron plates, and the efficacy of the remedy consequently diminished. Hence the directions of the London College for purifying this substance are to be preferred,

---

## OLEA EXPRESSA.

### EXPRESSED OILS.

VEGETABLES yield two distinct species of oil, one of which is volatile at a high temperature, but the other cannot be vola-



tilized without suffering decomposition. The first of these is termed *Volatile Oils*, the second, *Fixed Oils*. The latter name is properly adopted by the Edinburgh College, and is more suitable than *Expressed Oils*, the epithet given to this class of substances by the London and the Dublin Colleges.

FIXED OILS are obtained from fruits and seeds either by expression or decoction with water. The dicotyledons, or seeds with two seed-lobes, yield the greatest proportion of oil. When the first process is employed, the fruit or seed is put into a strong hempen or hair bag, and subjected to the press; during the action of which, the oil is forced out, generally combined with some other of the vegetable principles, which are afterwards separated by subsidence. The process is facilitated, and the quantity of oil increased, by heating the plates of the press, or previously roasting the seeds; but the oil thus obtained is more liable to become rancid, and hence the cold-drawn oils are always preferred for medicinal purposes. When the oil is to be obtained by decoction, the fruits or seeds are to be bruised previously to being boiled; and the oil which is separated is to be skimmed off from the surface of the water on which it swims.

Fixed Oils have different degrees of consistence; they are, 1. Fluid at the ordinary temperature of the atmosphere, congealing in a temperature a little higher than the freezing point of water. 2. Concrete at the ordinary temperature of the atmosphere, and require a higher degree for their liquefaction. The first are denominated *Fluid Oils*; the second, *Vegetable Butters*.

1. FLUID FIXED OILS are generally inodorous and nearly insipid, or have a mild taste. They are transparent, viscid so as to run in streaks upon the sides of glass vessels, and have generally a slight tinge of colour, which may be removed by digestion with charcoal. They are generally lighter than water, but differ from each other in specific gravity. At about 600° of Fahrenheit they boil, and are then volatilized, but in a state of partial decomposition; the vapour readily catches fire, and burns with a yellow flame. When exposed to the atmosphere at a high natural temperature, such as exists in summer, or in heated rooms, the fixed oils expressed without heat become thick, lose much of their transparency, acquire a sharp taste and a disagreeable odour, and are then said to be rancid; but when heat has been used in their expression, they only become thick, and acquire resinous properties. In both cases the changes are produced by the absorption of oxygen; but in the first case, owing to the combination of the oxygen with some of the vegetable mucilage present in the cold-drawn



oil, sebacic acid is formed; and by its diffusion through the oil, the change in its properties is produced.

Fixed oils are insoluble in water; but they may be mingled through water, and kept suspended in it by means of mucilage or yolk of egg. They are, with one or two exceptions, nearly insoluble in alcohol and ether; but unite readily with each other, with volatile oils, and with resinous substances. They dissolve sulphur, and form what was formerly regarded a Balsam with it. With the alkalies they combine, and form soaps; but with the acids undergo decomposition: and when boiled with some of the metallic oxides, tough solid compounds or plasters are produced.

2. CONCRETE fixed oils possess nearly the same properties as the fluid fixed oils. They are, however, more soluble in alcohol and ether, but are not capable of entering so readily into combinations with the alkalies. The ultimate constituents of fixed oil are carbon, hydrogen, and oxygen.<sup>1</sup>

For medicinal purposes, these oils are required to be free from rancidity; consequently, they must be preserved in closed vessels, and carefully excluded from the air.

OLEUM AMYGDALARUM, Lond. *Oil of Almonds.*

“Macerate Almonds, either *bitter* or *sweet*, in cold Water, for twelve hours, and bruise them; afterwards express the Oil without heat.”

OLEUM AMYGDALI COMMUNIS, Edin. *Oil of the Almond.*

“Take fresh Almonds, and bruise them in a stone mortar, then put them into a hempen sack, and express the Oil by a press, without heat.”

OLEUM AMYGDALARUM, Dub. *Oil of Almonds.*

“Bruise fresh Almonds in a mortar; and then express the Oil by a press, without heat.”

<sup>1</sup> When oil is distilled, as soon as it begins to boil a large quantity of gas is disengaged, consisting of a mixture of Carbonated Hydrogen and Oxide of Carbon. The *first* product of the distillation is a yellow coloured, very odorous substance, of a moderate consistence, very soluble in alcohol, reddening strongly the tincture of litmus, and forming a true soap with dilute liquor of Potass. It may be regarded as a mixture of acetic, sebacic, oleic and margaric acids, empyreumatic oil, a volatile oil slightly odorous, and a peculiar volatile matter very foetid, not acid, soluble in water; and which excites coughing and tears in the operator when the receiver is opened. The *second* product is a light-green fluid, which becomes brown on exposure to the air, has a slight empyreumatic odour, but does not excite coughing or tears. It does not redden tincture of litmus; alcohol even when aided by heat dissolves a very small portion only of it, and it is not affected by Potass. It burns like an essential oil; and is volatilized without change in close vessels. The *third* product is an orange-red, solid, transparent substance, inodorous and insipid: breaking with a waxy fracture, melting at a heat above 100° Centig.; and dissolving in hot alcohol from which it is precipitated unaltered on cooling. Ether is its proper solvent. *Annales de Chimie et de Phys.* tom. xxx. p. 1—20.



*Syn.* Huile d'Amandes (F.), Mandelnöl (G.), Olio di Mandarleh (I.), Vādomcottay unnay (Tam.).

The Oil obtained from both the Sweet and the Bitter Almond is equally free from bitterness, if heat be not employed. Sixteen ounces of Almonds yield about five ounces<sup>1</sup> of a bland inodorous oil, of a very slightly sweetish taste, which is at first a little turbid, but soon becomes clear. Its colour is a very pale greenish-yellow, and its specific gravity  $\cdot 932$ .<sup>2</sup> The oil from the bitter almond, it is said, keeps longer without growing rancid than that from the sweet almond. It is soluble in ether, in the proportion of f̄3x. to f̄3iv. of ether.

*Medical properties and uses.* — This oil is demulcent and emollient, and is used in coughs and other pulmonary complaints, united with water by means of mucilage or the yolk of egg and sugar. A mixture of f̄3iv. of almond oil, and ℥viij. of acetate of lead, forms a useful injection at the commencement of gonorrhœa. The dose of the oil is from f̄3iv. to f̄3j.

OLEUM LINI, Lond. Dub. OLEUM LINI USITATISSIMI, Edin. *Oil of Linseed.*

“Bruise the seeds of Common Flax, and afterwards express the Oil without heat.”

*Syn.* Huile de graine de lin (F.), Leinöl (G.), Olio di Lino (I.), Azéyte de Laxor (S.).

The proportion of Oil thus obtained is about 20 per cent. of the seed employed. It is combined with a considerable portion of mucilage<sup>3</sup>, has a strong disagreeable odour, and a nauseous taste; is not congealed except by a cold below 0° of Fahrenheit; and boils at 600° of the same scale. Its colour is a high yellow; and its specific gravity  $\cdot 932$ .<sup>4</sup> Four ounces of alcohol are required to dissolve one drachm of it: but the same quantity of ether takes up a fluid ounce and a half. One hundred weight of seeds yields from 18 to 20 lbs. of oil.<sup>5</sup>

*Medical properties and uses.* — Linseed oil is emollient, demulcent, and slightly laxative. On account of its nauseous taste, it is seldom used as an internal remedy, although it has been given with advantage in ileus when purgatives have failed. It is chiefly employed in the form of glyster in flatulent colic, attended with costiveness, and in abrasions of the rectum: and is an useful application to burns, especially when combined with lime water. The dose, when taken by the mouth, is from f̄3ss. to f̄3j.; but f̄3vj. may be given at once, per anum.

Officinal preparation. *Linimentum Aquæ Calcis*, E.

<sup>1</sup> About 3iij. more may be obtained by impregnating the marc with the steam of boiling water.

<sup>2</sup> Fabroni.

<sup>3</sup> The oil usually prepared on a great scale is more free from mucilage, the seeds being roasted before they are subjected to the press.

<sup>4</sup> Shaw's Boyle, ii. 346.

<sup>5</sup> Brande's Manuel, p. 336.



OLEUM RICINI, Lond. *Castor Oil.*

“Bruise Castor seeds, previously decorticated; and express the Oil, without heat.”

*Syn.* Huile de Ricin (*F.*), Rizinusöhl (*G.*), Olio di Ricino (*I.*), Sitt. āmamāhā unnay (*Tam.*).

The mode of obtaining this oil, with its qualities and medicinal virtues, have been already noticed. The purer it is, the more soluble it is in alcohol of spec. grav. .820. (See RICINUS, *Part ii.*) This cold drawn oil is considerably less purgative than the oil from the West Indies, which is of a deeper colour and extremely nauseous.

---

## OLEA DISTILLATA.

### DISTILLED, or VOLATILE OILS.

VOLATILE OILS, as they are properly denominated by the Edinburgh College, are vegetable products, found in almost every part of the vegetable body, except the cotyledons of the seeds, the part in which, almost always, the fixed oils are contained. In some plants, the volatile oil exists in distinct vesicles, and is obtained by simple expression, but in general it can only be obtained by distillation; whence the name *Distilled Oils*, given to this class of substances by the London College: and as the odour of plants generally depends on their volatile oils, the Dublin College, following the example of the elder chemists, who denominated them Essences, have adopted the term *Essential Oils*. The expressed volatile oils are now rejected from all the Pharmacopœias; and the whole of those used in pharmacy retained, are procured by distillation.

Volatile oil is obtained from both recent and dried plants. When fresh plants are to be employed, they require no previous treatment; but when the plants are dry, or woods or barks are to be employed, the plants must be macerated in water for some time, and the woods and barks be previously rasped. The distillation is performed in the following manner. The plants, or the parts of them containing the oil, are to be put into a tinned copper still, and closely pressed down; after which, as much water is to be poured in as will be sufficient to cover the materials. The head of the still, which should be low, is then to be luted on; the fire lighted, and so regulated as to keep the contents of the still scarcely up to the boiling point; and the distillation continued, until the condensed vapour comes over nearly insipid and inodorous. During this process the volatile oil rises with the watery vapour, from which, however, the greater part of it again separates, after it has re-



mained at rest for some hours in a cool place, and either floats on the surface of the water, or sinks to the bottom, according to its specific gravity. The complete separation of the oil is effected by an instrument called a separatory (see *Part i.*): and the water is to be again used for a second distillation of fresh materials, by which, as it is already impregnated with as much of the oil as it can dissolve, the product of oil of the second, and every subsequent distillation will be consequently greater than that of the first; but it is not till "the tenth distillation, in some cases, that the produce of the oil attains its maximum."<sup>1</sup> By the same process, volatile oils are obtained from balsams, resins, gum-resins, and turpentine. They have not their characteristic qualities in perfection immediately after their distillation, but have a disagreeable, empyreumatic odour; to dissipate which they must be allowed to stand for some days in vessels loosely covered with paper, before they be put into the bottles in which they are to be preserved, which should be opaque.

Although all volatile oils agree in their chemical properties sufficiently to constitute them members of the same class of substances, yet they differ greatly in their qualities from each other, and in the proportions in which they are obtained.

Volatile oils have a penetrating odour, and hot taste. They are completely evaporated when heated in the open air; a property which is taken advantage of as a test of their purity; for if they be adulterated with fixed oil, which is not unfrequently the case, by heating a small portion of the oil on a piece of clean paper, a greasy spot will remain, whereas if the volatile oil be pure, the paper will be left perfectly clean. In a higher temperature, volatile oils are readily ignited, and burn with a bright white flame, emitting a large quantity of black dense smoke; and with the production of a large proportion of carbonic acid and water.

Volatile oils exposed to the light are changed in colour, or become colourless; when exposed to the air, they become more viscid, less odorous, redden the tincture of turnsole, and gradually assume the form of resins. These changes, Dr. Priestley ascertained<sup>2</sup>, depend upon the absorption of oxygen; and hence the necessity of preserving volatile oils in small phials, completely full, and well corked. An oil which has become thick and scentless, may be rectified by re-distilling it, with some of the same kind of plant from which it was originally extracted, or with alcohol or sulphuric ether<sup>3</sup>; a limpid, odorous oil comes over, and resin remains in the retort.

<sup>1</sup> *Aikin's Dictionary of Chemistry*, Art. Oil.

<sup>2</sup> *Priestley on Air*, ii. 232.

<sup>3</sup> *Nicholson's Journal*, 8vo. vii. 6.



These oils are very sparingly soluble in water, and render it milky when agitated with it, communicating to it their odour: they are all soluble in alcohol, ether, and the fixed oils in various proportions. From their solubility in alcohol they are sometimes adulterated with that fluid; but the fraud may be detected by agitating some of the suspected oil with water; when if the oil contain alcohol, an increase of temperature will be indicated by the thermometer, but not if the oil be pure.<sup>1</sup> The more expensive oils are also occasionally adulterated with the cheaper, particularly with oil of turpentine, which, however, is readily discovered by its peculiar odour, if a piece of paper be dipped in the suspected oil and dried with a gentle heat. They are also, sometimes, adulterated with Castor oil; and as the mixture, when the ingredients are in equal proportions, is soluble in alcohol, the fraud cannot be detected by that test; but it is rendered obvious by the adulterated oil leaving a greasy stain upon paper, which has been touched with it, and held before the fire, whereas no stain is left by the genuine oil.

Volatile oils unite with sulphur, in a temperature sufficient to melt the sulphur, and form brown-coloured, foetid mixtures, which have been denominated *Balsams of Sulphur*. The alkalies and earths combine imperfectly with them, and constitute a class of bodies, which the French chemists have denominated *saponules*. The action of the acids is much more violent than on the fixed oils: and several of them detonate when rubbed with oxymuriate (*chlorate*) of potass.

As medical agents, volatile oils are stimulant and aromatic. They are chiefly employed to remove nausea and flatulence, to correct the griping qualities of some purgatives, and the disagreeable taste of other remedies. They may be given, triturated with water and mucilage; or dropped first on a lump of sugar, and through its medium diffused in water, forming a solution of what has been denominated *oleum saccharum*. The quantity of sugar must be more than ten times the weight of the oil; and when they are well triturated together, the oil becomes thus completely soluble in water, and may be diluted to any extent.

Some of the more stimulant of these oils are added to embrocations to be used as rubefacients in cases of numbness, pains, and paralytic affections of the joints.

The three British Colleges give the following general rules for the preparation of *volatile oils*.

---

<sup>1</sup> Marqueron, *Annales de Chimie*, xlviii. 267.



*OLEA DISTILLATA, Lond.**Distilled Oils.*

“THE seeds of Anise and Carraway, the flowers of Chamomile and Lavender, the berries of Juniper and All-spice, the tops of Rosemary, and the entire plants of the other articles, dried, are to employed.

“Put any one of these into an alembic, then pour in as much water as will cover it, and distil the Oil into a large refrigeratory.

“The water which distils over with the Oils of Carraway, Peppermint, Spearmint, All-spice, and Penny-royal, is to be preserved for use.”

*OLEA VOLATILIA, Edin.**Volatile Oils.*

“As much water only is to be employed as will prevent empyreuma during the distillation. The distillation may be immediately commenced after a proper maceration : and the oil afterwards separated from the water.

“It is also necessary to observe, in preparing these Oils and the distilled waters, that the quality of the substances, their texture, the season of the year, and similar circumstances, must occasion so many differences, that is scarcely possible to give any certain and general rules which shall strictly apply to every example. Many things, therefore, which must be regulated by the judgment of the operator, are omitted, and the more general only given.”

*OLEA ESSENTIALIA, Dub.**Essential Oils.*

“LET the Oil be extracted by distillation, from the substance previously macerated in water, as much water being added during the distillation as may be sufficient to prevent empyreuma.

“In distilling Fennel, Peppermint, Spearmint, Penny-royal, and All-spice, the watery fluid which comes over in distillation with the Oil is to be preserved for use according to the directions under the head of Distilled Waters.”

Few of the volatile oils are prepared by the apothecary. The oils of Anise, Chamomile, Juniper, Origanum, Rosemary,



and Pimento are usually imported into this country; while those of Lavender, Peppermint, Spearmint, and Penny-royal, are annually prepared on a large scale.<sup>1</sup>

OLEUM ANISI, Lond. OLEUM VOLATILE PIMPINELLÆ ANISI, Edin. OLEUM SEMINUM ANISI, Dub. *Oil of Aniseed.*

*Syn.* Huile d'Anis (*F.*), Anisöhl (*G.*), Olio di Anice (*I.*).

This oil is of a whitish or a pale straw-colour, has the odour of the plant, and a slightly pungent, bitter, sweetish taste. It crystallizes at 50° in flat tables. One hundred pounds of Anise-seeds yield about ℥ijss. of oil; but the produce varies. It is sometimes adulterated with wax, Spermaceti, or Camphor; but the fraud is easily detected, for on moderately warming the genuine oil the crystals dissolve, which is not the case with sophisticated.<sup>2</sup> The greater part of the oil of Anniseed consumed in this country is prepared in Spain. The specific gravity of that made in England is 9768 of that imported, or the foreign 9903.

*Medical properties and uses.*—This oil is used chiefly as a carminative; and as it is less pungent than many of the other volatile oils, it is better adapted for relieving flatulence in children. It is given in doses of from ℥v. to ℥xv. triturated with sugar.

OLEUM ANTHEMIDIS, Lond. OLEUM VOLATILE ANTHEMIDIS NOBILIS, Edin. *Oil of Chamomile.*

*Syn.* Huile de Camomille Romaine (*F.*), Kamillenöhl (*G.*), Olio di Camomilla Romana (*I.*), Azeyte de Manganella de Botera (*S.*).

The odour of this oil is unpleasant, and the taste pungent. When recently distilled the colour is cærulean blue, but by exposure to light it changes to yellow. Eighty-two pounds of chamomile flowers yield eighteen drachms of oil.<sup>3</sup> The spec. grav. of the English oil, from flowers only, is 9083, of the foreign it is 9289.

*Medical properties and uses.*—This oil is supposed to possess antispasmodic powers, and is therefore sometimes recommended in cramp of the stomach, and as an adjunct to purgative pills. The dose is from ℥v. to ℥x. but it is seldom used.

OLEUM CARUI, Lond. OLEUM SEMINUM CARUI, Dub. *Oil of Carraway.*

*Syn.* Huile de Carvi (*F.*), Kümmelöhl (*G.*), Olio di Carvi (*I.*), Azeyte de Alcorovea (*S.*).

Six pounds of Carraway-seeds yield four ounces and a half of oil<sup>4</sup>, which has an aromatic odour, and a sweetish pungent taste; is viscid, and of a yellow colour. Its specific gravity is .946.<sup>5</sup> W. Brande says 9310.<sup>6</sup>

<sup>1</sup> *Lond. Med. Review*, April, 1810. 156.

<sup>4</sup> Baumé.

<sup>5</sup> *Ibid.*

<sup>2</sup> Baumé.

<sup>3</sup> *Ibid.*

<sup>6</sup> *Manual*, p. 342.



*Medical properties and uses.*—Oil of carraway is stimulant and carminative. It is chiefly used as an adjunct to purgative pills, and to cover the disagreeable flavour of other substances. The dose is from ℥j. to ℥x.

**OLEUM SEMINUM FŒNICULI DULCIS, Dub.**  
*Oil of Fennel Seeds.*

*Syn.* Huile essentielle de Fenoule (*F.*), Fenchelohl (*G.*), Olio di Finoichio (*I.*), Azey de l'Eneldo hinojo (*S.*).

Seventy-five pounds of Fennel seeds yield thirty ounces of oil<sup>1</sup>, which is colourless, and congeals at 50°. It has the odour of the plant, and a hot, sweetish taste. Its specific gravity is .997.<sup>2</sup>

*Medical properties and uses.*—The same as those of the plant; the usual dose is from ℥j. to ℥xx. It is rarely used.

**OLEUM JUNIPERI, Lond. OLEUM VOLATILE JUNIPERI COMMUNIS, Edin. OLEUM BACCARUM JUNIPERI, Dub.**  
*Oil of Juniper.*

*Syn.* Huile essentielle de Genevrier (*F.*), Wachholder beeröhl (*G.*), Olio di Ginepro (*I.*).

Forty-eight pounds of bruised Juniper berries yield six ounces of oil<sup>3</sup>, of a specific gravity .611<sup>4</sup>, according to Lewis, but Mr. Brande states that of the English to be .8688, the foreign .8834.<sup>5</sup> Its odour is similar to that of turpentine, and the taste hot and acrid. It has a greenish-yellow colour, is viscid, and deposits a feculent matter when long kept. When genuine it is soluble in alcohol.

*Medical properties and uses.*—This Oil is carminative, diaphoretic, and diuretic. It is sometimes given in dropsy, and may be added to Foxglove when it is exhibited in the form of pills. The dose is from ℥j. to ℥x. combined with water by means of sugar or of mucilage.

**OLEUM LAVANDULÆ, Lond. OLEUM VOLATILE LAVANDULÆ SPICÆ, Edin. OLEUM FLORUM LAVANDULÆ, Dub.**  
*Oil of Lavender.*

*Syn.* Huille essentielle de Lavendelöhl (*G.*), Olio di Lavanda (*I.*), Azeyte del' Espliego (*S.*).

One pound nine ounces of this oil are obtained from eighty pounds of Lavender flowers. The odour is very fragrant, and the taste warm and agreeable. Its colour is a very pale, lemon-yellow; and its specific gravity, according to Lewis, .936<sup>6</sup>, but Mr. Brande says that the specific gravity of oil obtained from the flowers only is .8960, and that from the whole herb .9206.

*Medical properties and uses.*—This Oil is stimulant and cordial. It is chiefly used in hysteria and nervous headache, in doses of from ℥j. to ℥v. given on a lump of sugar.

<sup>1</sup> Dehne.

<sup>4</sup> Lewis.

<sup>2</sup> Lewis.

<sup>5</sup> Ibid.

3 G

<sup>3</sup> Dehne.

<sup>6</sup> Ibid.



OLEUM VOLATILE LAURI SASSAFRAS, Edin. OLEUM CORTICIS ET LIGNI SASSAFRAS, Dub. *Oil of Sassafras.*

Sixty pounds of bruised Sassafras yield twelve ounces<sup>1</sup> of a viscid, yellow oil, heavier than water, its specific gravity being 1·094.<sup>2</sup> Its odour is fragrant and its taste hot and acrid, excoriating the lips when incautiously tasted. The Edinburgh College orders it to be distilled from the bruised root.

*Medical properties and uses.*—This Oil is stimulant, and supposed to be also sudorific and diuretic. It has been given in chronic rheumatism, scurvy, and some cutaneous affections. The dose is from ℥ij. to ℥x. but it is scarcely ever ordered.

OLEUM MENTHÆ PIPERITÆ, Lond. OLEUM VOLATILE MENTHÆ PIPERITÆ, Edin. OLEUM HERBÆ FLORESCENTIS MENTHÆ PIPERITIDIS, Dub. *Oil of Peppermint.*

*Syn.* Huile essentielle de Menthe Poivrée (*F.*), Pfeffermünzöhl (*G.*), Olio di Menta piperitide (*I.*).

Four pounds of the recent plant yield from one drachm and a half to three drachms and a half of this oil.<sup>3</sup> Its odour is strong, and its taste very pungent, but at the same time it impresses a sensation of coldness. Its colour is brownish-yellow: but it becomes white when exposed to the light. Its specific gravity is ·9070.

*Medical properties and uses.*—Oil of Peppermint is stimulant and carminative. It is a common domestic remedy in cramp of the stomach, flatulent colic, and anorexia; and is usually rubbed up with sugar or mucilage.<sup>4</sup> The dose is from ℥j. to ℥iij.

OLEUM MENTHÆ VIRIDIS, Lond. OLEUM HERBÆ FLORESCENTIS MENTHÆ SATIVÆ, Dub. *Oil of Spearmint.*

*Syn.* Huile essentielle de Baumé verte (*F.*).

This Oil has a flavour similar to that of Peppermint, but is less grateful; its taste is warm and less pungent than that of Peppermint. According to Lewis its specific gravity is ·975<sup>5</sup>; but Mr. Brande states it to be ·9394. Its colour is greenish.

*Medical properties and uses.*—The same as those of Oil of Peppermint. The dose is from ℥ij. to ℥v. given on a lump of sugar.

Officinal preparation. *Infusum Menthæ compositum.*

OLEUM ORIGANI, Lond. OLEUM VOLATILE ORIGANI MARJORANÆ, Edin. OLEUM HERBÆ FLORESCENTIS ORIGANI. Dub. *Oil of common Marjoram.*

*Syn.* Huile essentielle d'Oriang (*F.*), Dostöhl (*G.*), Olio di Origano (*I.*), Azeyte de Origane Sylvestre (*S.*).

<sup>1</sup> Baumé.

<sup>2</sup> Ibid.

<sup>3</sup> Ibid.

<sup>4</sup> Peppermint drops or Lozenges are a mixture of Starch, Sugar, and mucilage of Tragacanth.

<sup>5</sup> Lewis.



One hundred and fifty pounds of dried leaves of common Marjoram yield fifteen ounces of oil<sup>1</sup>, of a yellow colour, having the odour of the plant, and a hot acrid taste. Its specific gravity is  $\cdot 940$ .<sup>2</sup>

*Medical properties and uses.*—On account of its acrid quality, this oil is never exhibited internally. As a local stimulant it is sometimes used to allay the pain of tooth-ach, two or three drops on a piece of cotton being put into the carious tooth.

OLEUM PIMENTÆ, Lond. OLEUM VOLATILE MYRTI PIMENTÆ, Edin. OLEUM BACCARUM PIMENTÆ, Dub. *Oil of Pimento.*

*Syn.* Huile essentielle de Poivre de Jamaïque (F.), Nelherpfefferöhl (G.), Olio di Pimento (I.).

This Oil has the agreeable odour of the Pimento, with its pungent taste in an increased degree. It is of a reddish-brown colour, and is heavier than water, its specific gravity being  $1\cdot 021$ .

*Medical properties and uses.*—It has the same properties as All-spice in a greater degree; and is given in dyspeptic affections, colic, and tympanitis, in doses of from  $\mathfrak{mij}$ , to  $\mathfrak{m}\mathfrak{v}$ . rubbed with sugar, or in any proper vehicle.

OLEUM PULEGII, Lond. OLEUM HERBÆ FLORESCENTIS PULEGII, Dub. *Oil of Pennyroyal.*

*Syn.* Huile essentielle de Menthe Peuliot (F.), Poleiöhl (G.), Olio di Puleggio (I.), Azeyte de Peleo (S.).

This Oil is of a reddish-yellow colour, and resembles in its other qualities the oil of Peppermint. Its specific gravity according to Lewis is  $\cdot 978$ ,<sup>3</sup> but according to Mr. Brande  $\cdot 9390$ .

*Medical properties and uses.*—It is stimulant and antispasmodic, but is scarcely ever used. The dose may be from  $\mathfrak{mj}$ . to  $\mathfrak{m}\mathfrak{v}$ . given on a lump of sugar.

OLEUM ROSMARINI, Lond. OLEUM VOLATILE RORISMARINI OFFICINALIS, Edin. OLEUM HERBÆ FLORESCENTIS RORISMARINI, Dub. *Oil of Rosemary.*

*Syn.* Huile essentielle de Romarin (F.), Rosmarinöhl (G.), Olio di Rosmarino (I.).

Twenty-four pounds of the plant yield one ounce of a fluid colourless oil<sup>4</sup>, the odour of which is less agreeable than that of the plant. It deposits crystals of Camphor when long kept. Its specific gravity is  $\cdot 9108$ .

*Medical properties and uses.*—It is stimulant; and frequently enters into the composition of liniments. The dose, as an internal remedy, may be from  $\mathfrak{mij}$ . to  $\mathfrak{m}\mathfrak{vj}$ . but it is scarcely ever ordered.

<sup>1</sup> Baumé.

<sup>2</sup> Ibid.

<sup>3</sup> Lewis.

<sup>4</sup> Baumé.

<sup>5</sup> Lewis.



**OLEUM HERBÆ FLORESCENTIS RUTÆ, Dub.**  
*Oil of Rue.*

Twenty-one pounds of Rue yield fifty-nine grains of<sup>1</sup> oil, which has the strong ungrateful odour and taste of the plant. When recently drawn the colour of this oil is yellow, but it deepens to a brown by age, and deposits a brownish, resinous sediment. It congeals at 40° Fahrenheit.

*Medical properties and uses.*—Oil of Rue is stimulant and antispasmodic. It is sometimes given in hysteria, and the convulsive affections of infants attendant on dentition; and is sometimes used as a rubefacient in palsy. The dose is from ℥ij. to ℥v. triturated with sugar or mucilage.

**OLEUM HERBÆ JUNIPERI SABINÆ, Edin.**  
**OLEUM FOLIORUM SABINÆ, Dub.** *Oil of Savine.*

*Syn.* Huile essentielle de Sabine (F.), Sevenbaumöhl (G.), Olio di Sabina (I.), Azeyt de Enebrio Sabina (S.).

Two pounds of Savine are said to yield five ounces of<sup>2</sup> oil. It is limpid, has the odour of the plant, and is extremely acrid to the taste. Its colour is yellow; but it becomes colourless on being kept exposed to light.

*Medical properties and uses.*—This Oil is the principle on which the virtues of Savine depend; thence it possesses the same properties, and is applicable to the same purposes as the plant. The dose may be from ℥ij. to ℥vj. triturated with sugar.

**OLEUM CORNU CERVINI RECTIFICATUM, Dub.** *Rectified Oil of Hartshorn.*

“Take of the Oil which rises in the distillation of the volatile liquor of Hartshorn, *three pounds*; Water, *six pints*.

“Distil the Oil, then remix it with the Water, and re-distil, repeating the distillations until the Oil become limpid. It ought to be preserved in a dark place, in small phials completely filled, and closely stopped.”

This empyreumatic Oil is first formed by the decomposition of animal matter by heat; and arises from a new combination of part of the Hydrogen and Carbon of the substance distilled. As first obtained it is thick, of a dark colour, and has a very offensive odour; but, by the rectification above ordered, it is rendered thinner and less offensive.

*Qualities.*—Rectified Oil of Hartshorn is nearly colourless and transparent; has a strong, slightly aromatic odour, and a penetrating taste. It is very light and volatile, strikes a green colour with syrup of violets; is partially soluble in water, and unites readily with alcohol, ether, and oils. The acids form with it a thick saponaceous compound; and with the alkalies

<sup>1</sup> Baumé.

<sup>2</sup> Murray.



it forms a true soap. Exposure to light and air destroys its transparency, and gives it a deep-brown colour.

*Medical properties and uses.* — This Oil is stimulant, antispasmodic, anodyne, and sudorific. It was formerly regarded as a remedy of much efficacy in fever, particularly when given a few hours before the accession of the paroxysm of intermittents; and was also much employed in epilepsy, hysteria, and all convulsive affections. It is now almost discarded from practice, being only occasionally used as an external application to paralytic limbs. The dose may be from ℥x. to ℥xxx. in a sufficient quantity of water.

OLEUM SUCCINI, Lond. *Oil of Amber.*

“Put the Amber into an alembic, and distil from a sand-bath, with a fire gradually raised, an Acid Liquor, the Oil, and a Salt impregnated with the Oil. Then re-distil the Oil a second and a third time.”

OLEUM SUCCINI, Edin. *Oil of Amber.*

“Take of Amber in powder and of pure sand, *equal parts*. Mix them together in a glass retort, the capacity of which the mixture only half fills; and having adapted to it a large receiver, distil in a sand-bath, with a gradually augmented heat. An aqueous fluid, tinged with a little yellow Oil, will first come over; then a yellow Oil with an acid salt, and, lastly, a black and reddish Oil. Pour the fluid from the receiver and separate the Oil from the water.”

OLEUM SUCCINI PURISSIMUM, Edin. *Pure Oil of Amber.*

“Distil the Oil of Amber mixed with six times its quantity of Water, from a glass retort, until two-thirds of the Water pass over into the receiver. Then separate this purified volatile Oil from the Water, and keep it in well-stopped phials.”

OLEUM SUCCINI RECTIFICATUM, Dub. *Rectified Oil of Amber.*

“Take of the Oil which comes over in the preparation of Succinic Acid, *a pound*; Water, *six pints*. Distil until two-thirds of the Water have passed into the receiver: then separate the Oil.”

*Syn.* Huile de Succin. (*F.*), Bernsteinöhl (*G.*), Olio di Succino Rettificato (*I.*).

The Oil of Amber, as immediately procured by the distillation of Amber, is of a dark colour, a thick consistence, and has a very foetid odour; but by successive distillations it is rendered thinner, of a lighter colour, and at length is obtained nearly limpid. It is the result of the process, and does not pre-exist in the Amber.

*Qualities.* — Rectified Oil of Amber has a strong ungrateful odour, and a hot acrid taste. It is light, volatile, and inflam-



mable, insoluble in water, and only partially soluble in alcohol.

*Medical properties and uses.* — Oil of Amber is stimulant, antispasmodic, and rubefacient. It has been found serviceable in deficient menstruation, and in hysteria, epilepsy, and some other convulsive affections; but it is now scarcely ever administered as an internal remedy. The dose may be from ℥v. to ℥xij. combined with any distilled water by means of mucilage. It is more generally employed externally as a rubefacient in rheumatism and paralysis; and a mixture of fʒj. of this oil with fʒss. of tincture of opium has been found beneficial as a friction to the affected part in tic douloureux; and in hooping-cough, rubbed upon the chest twice or three times a day.<sup>1</sup>

Officinal preparation. *Spiritus Ammoniac succinatus.*

OLEUM TEREBINTHINÆ, Dub. *Oil of Turpentine.*

“Take of common Turpentine, *five pounds*; Water, *four pints*. Distil the Oil from a copper alembic. Yellow Resin will remain in the retort after the distillation.”

OLEUM TEREBINTHINÆ RECTIFICATUM, Lond. Dub. *Rectified Oil of Turpentine.*

“Take of Oil of Turpentine, *a pint (two pints, Dub.)*; Water, *four pints*. Distil the Oil (*a pint and a half of the oil, Dub.*)

OLEUM VOLATILE PINI PURISSIMUM, Edin. *Purified Oil of Turpentine.*

“Take of Oil of Turpentine, *one part*; Water, *four parts*. Distil as long as any oil passes over.”

*Syn.* Huile essentielle de Tirébenthine (*F.*), Terbenthinöhl (*G.*), Olio di Trementina (*I.*), Azeyte de Pino (*S.*).

The chemical qualities and medicinal properties of Oil of Turpentine have been already noticed. (See PINUS, *Part ii.*) The rectification of it is a troublesome process, and on account of the great inflammability of the vapours, much caution is required to prevent them from escaping through the lutings of the vessels, and catching fire. The rectified oil is a little lighter than the common oil, and completely free from any resinous admixture; but in other respects it has no peculiar excellence to recommend it. What remains in the retort is a thick resinous matter, and is denominated balsam of turpentine.

*Medical properties and uses.* — These have been already mentioned (*Part ii.*). I have had several opportunities of ascertaining the efficacy of Oil of Turpentine as a remedy for tape-

---

<sup>1</sup> The empirical nostrum, known by the name of *Roche's Embrocation*, for hooping cough, consists of two parts of Olive oil, one part of oil of Amber, and one part oil of Cloves.



worm. In every case in which I have administered it, the worm has been expelled, and the symptoms relieved. In general, the animal has been voided of a livid hue, and evidently killed; but in one instance, in which a portion of five feet in length was passed after two fluid ounces of the oil had been taken, it was not livid, and when voided, exhibited evident signs of animation. In no instance have I perceived that the large doses of the Oil, which were taken for the above purpose, produced any particular effect on the urinary organs. The more usual sensible effects are temporary intoxication, accompanied with considerable nausea, and sometimes vomiting, which, after two or three alvine evacuations, subside, and leave a degree of languor for ten or twelve hours. The pain of the stomach and side, which is an usual concomitant of the disease, is always removed by the Oil. I have also given this oil in combination with Cinchona, with evident benefit in rheumatism; particularly in that modification of the disease, which attacks one side only of the head, and is periodical; the paroxysms generally coming once or twice in the twenty-four hours. Tincture of Capsicum, in doses of ℥iij. is a useful adjunct to the bark and turpentine in this affection. In some persons, however, Turpentine affects the kidney, producing pain and bloody urine, and in others, its administration has produced a severe erythematic eruption over the body. These effects, however, are in some degree lessened by purifying the oil of Turpentine by means of alcohol, as Dr. Nimmo suggests. To eight parts of the Oil add “one part of alcohol and agitate; on the mixture being left at rest, the oil falls to the bottom, and the spirit holding the impurities in solution floats at the top. On repeating this process three or four times, pouring off the spirit each time, the oil is left nearly tasteless, and inodorous<sup>1</sup>.” The dose in rheumatism is fʒi. repeated every four hours; but in tænia it may be given in doses of fʒj. combined with syrup of poppies, repeated every six hours until the worm is expelled.

---

## AQUÆ DISTILLATÆ.

### DISTILLED WATERS.

It has been already remarked that the volatile oil, on the presence of which the odour and the taste of plants in a con-

---

<sup>1</sup> *Journ. of Science*, vol. xiii. pp. 65, 66.



siderable degree depend, is elevated during distillation with water; and a portion of it being retained in solution, the water thus acquires the odour and taste of the vegetable with which it is distilled. The qualities, however, thus acquired by water, are scarcely, in any case, sufficient to give it much power as a remedy; and thence, the distilled waters are generally employed as elegant vehicles only for the exhibition of more active substances.

The following general directions are given by the London College for the preparation of these waters.

“WATERS are to be distilled from dried plants, unless it be otherwise ordered, because fresh plants cannot be procured at all times of the year. When fresh plants are employed, the weight of them ordered is to be doubled.

“To every gallon of these waters add five fluid ounces of proof spirit, to preserve them from spoiling.”

The Edinburgh College orders half an ounce of proof spirit, and the Dublin College half a fluid ounce of rectified spirit, to be added to each pound of the water.

Waters distilled from aromatic plants are more grateful when the plant is used in the dried state; but when delicate odorous flowers or herbs are employed, and the water acquires little more than odour by the distillation, the vegetable should always, if possible, be used in the recent state. Much care is required in conducting the process, to prevent any of the vegetable matter from being scorched, and to stop the distillation before the water is tainted by empyreuma. Notwithstanding, however, every attention which can be given, distilled waters, when newly prepared, have a very disagreeable empyreumatic odour, to dissipate which the vessels holding the waters must be left open to the air as long as any of the unpleasant odour remains; but, afterwards, it is essential for the preservation of the waters that they be preserved in closely-corked vessels.

When long kept, many of the distilled waters undergo a species of decomposition; they become slightly sour, and a ropy, viscid matter forms in them, owing to the essential oil they contain undergoing decomposition, and changing into mucilage. The addition of the spirit is intended to prevent this from taking place, but it is not adequate to the effect intended; and a much preferable mode is to re-distil the waters, after which they will keep good for several years.

Several of these waters are prepared on a great scale of a superior quality to any that the apothecary can prepare, and cheaper.

AQUA DISTILLATA, Lond. *Distilled Water.*

“Take of Water, *ten gallons.* First distil four pints,



which are to be rejected, and then distil four gallons. Preserve the Distilled Water in a glass bottle."

*Edinburgh.*

"Let Water be distilled in clean vessels, until two-thirds of the quantity employed have distilled over."

*Dublin.*

"Take of Spring Water, *twenty pounds*. Put them into a glass retort, and having rejected the first pound which comes over, let one gallon be distilled over with a gentle heat."

*Syn.* Eau distillée (*F.*), Einfaches destillirtes wasser (*G.*), Gemeenes Destillert Water (*Dutch*), Watten (*Swed.*), Acqua Distillata (*I.*), Aqua Distillada (*S.*).

Water is almost universally diffused over the surface of the earth, but it is not found perfectly pure in any place, which is owing to its great solvent powers enabling it to take up a portion of many substances with which it must come into contact, in its natural state. These impregnations, however, in spring and in river water, are not sufficient in general to give them any very sensible taste, or render them unfit for the ordinary purposes of life; but for many Pharmaceutical purposes it is necessary that the water be absolutely free from every foreign ingredient. *Rain water* is the purest kind of natural water, but it nevertheless contains a portion of carbonic acid gas, and minute quantities of carbonate of lime, and of muriate of lime; in spring water, besides these ingredients, is found a small portion of muriate of soda; *well water*, which is spring water obtained from a greater depth, holds in solution a much larger portion of carbonic acid, and several earthy salts, the principal of which are sulphate and carbonate of lime; and *river water* is impregnated with different proportions of carbonate of lime, sulphate of lime, and muriate of soda. By distillation water is freed from these ingredients, and rendered nearly pure; but it is tasteless and vapid owing to the absence of the air which it contains in its natural state. The process should be conducted slowly, with a moderate degree of heat, and not continued longer than the time specified in the formulæ, otherwise a minute portion of the saline matter contained in the natural water passes over in the distillation.

Although the necessity of distilled water for many Pharmaceutical operations is very obvious, yet, by too much refinement in this particular having been erroneously insisted upon in the former London Pharmacopœia, apothecaries have of late years almost altogether neglected its use, even in cases where it is absolutely necessary. This error the College has avoided in its present Pharmacopœia, and therefore it may be



expected that the directions for using it will be strictly attended to. But, as it is not always easy for the apothecary to prepare distilled water, *rain water*, filtered through alternate strata of well-washed sand, or powdered flints and charcoal, will answer every purpose for which distilled water is required. Soft water is a more powerful menstruum of vegetable matter than hard water; and resinous substances cannot easily be mixed with water containing calcareous matter, even when mucilage is used, whereas they readily mix with very soft or distilled water. Perhaps it should be a rule to use filtered rain water only in all Pharmaceutical operations. In extemporaneous prescriptions distilled water is often ordered when there is no necessity for its use, and often neglected to be ordered when it is absolutely necessary. It may, therefore, be useful to know that it is necessary in formulæ containing any of the following substances: *Acidum Sulphuricum, Acidum Citricum, Antimonium Tartarizatum, Argenti Nitras, Cuprum Ammoniatum, Ferrum Tartarizatum, Hydrargyri Oxymurias, Liquor Ammoniaë, Liquor Plumbi Acetatis, Liquor Potassæ, Plumbi Superacetat, Solutio Muriatis Barytæ, Vinum Ferri, Zinci Sulphas, Ferri Sulphas.*

#### AQUA ANETHI, Lond. *Dill water.*

*Syn.* Eau d'Aneth puant (F.), Acqua di Aneto Puzzolente (I.).

“Take of Dill seeds bruised, *a pound*. Pour on them so much water, that during the distillation there may be a sufficiency to prevent empyreuma. Distil one gallon.”

This water has an unpleasant odour and little pungency. It is used principally as a carminative for infants: and is a good vehicle for giving Magnesia and Rhubarb to these tender patients.

#### AQUA CARUI, Lond. *Carraway Water.*

*Syn.* Eau de Carvi (F.), Feldkumel wasser (G.), Acqua di Carvi (I.).

“Take of Carraway seeds bruised, *a pound*. Pour on them so much Water, that during the distillation there may be a sufficiency to prevent empyreuma. Distil a gallon.”

Carraway water possesses a considerable share of the aromatic flavour and pungency of the seeds, and may be used for the same purposes as Dill water.

#### AQUA CITRI AURANTII, Edin. *Water of Orange-peel.*

*Syn.* Eau d'Orange (F.), Pomeranzen wasser (G.), Acqua de Arancio (I.).

“Take of fresh Orange-peel, *two pounds*. Add so much water that when ten pounds have been drawn off by distillation, there shall remain a quantity sufficient to prevent empyreuma. After due maceration distil ten pounds, to which add five ounces of diluted alcohol.”

The water has the flavour only of the orange-peel.



**AQUA CITRI MEDICÆ, Edin.** *Water of Lemon-peel.**Syn.* Eau de Citronier (*F.*), Citronen wasser (*G.*), Acqua di Limone (*I.*).

This is prepared in the same manner as the former; ten pounds of Water being distilled from two pounds of fresh Lemon-peel. It has the flavour of the peel, but is seldom used.

**AQUA CINNAMOMI, Lond. Dub. AQUA LAURI CINNAMOMI, Edin.** *Cinnamon Water.**Syn.* Eau de Cannelle (*F.*), Zimmt wasser (*G.*), Acqua di Cinamomo (*I.*).

“Take of Cinnamon bark bruised, *a pound*; Water, *a pint*. Macerate the bark in the water for twenty-four hours; then add a sufficient quantity of water to prevent empyreuma during the distillation. Distil a gallon.”

This water is milky, and has the agreeable flavour and pungency of the Cinnamon, without its astringency; but the oil being ponderous is apt to separate, leaving the water clear and insipid. It is a gentle stimulant and aromatic, but is chiefly used to cover the nauseous taste of other medicines.

**AQUA LAURI CASSIÆ, Edin.** *Water of Cassia Bark.**Syn.* Eau de Casse (*F.*), Acqua di Cannella (*I.*).

This is prepared from one pound of bruised Cassia bark, in the same manner as the former, for which it is often substituted, being less expensive; but it is also less agreeable.

**AQUA FÆNICULI, Lond. AQUA FÆNICULI DULCIS, Dub.** *Fennel Water.**Syn.* Eau de Fenouil (*F.*), Fenchel wasser (*G.*), Acqua di Finocchio (*I.*).

“Take of Fennel seeds bruised, *a pound*. Pour over them as much water as will prevent empyreuma during the distillation. Distil a gallon.”

It is rarely if ever employed.

**AQUA MENTHÆ PIPERITÆ, Lond. Edin. AQUA MENTHÆ PIPERITIDIS, Dub.** *Peppermint Water.**Syn.* Eau de Menthe poivrée (*F.*), Pfeffermunz wasser (*G.*), Acqua di Menta Piperitide (*I.*).

“Take of Peppermint, *a pound and a half*, (*three pounds*, Edin.) Pour over it as much Water as will prevent empyreuma during the distillation. Distil a gallon, (*ten pounds*, Edin.)”

Peppermint water has the flavour and taste of the plant in a considerable degree. It is sometimes used alone as a carminative, but more generally for the purpose of covering the taste of other medicines.

**AQUA MENTHÆ VIRIDIS, Lond. AQUA MENTHÆ SATIVÆ, Dub.** *Spearmint Water.*

“Take of Spearmint, *a pound and a half*. Distil a gallon of water in the same manner as above.”

**AQUA PIMENTÆ, Lond. AQUA MYRTI PIMENTÆ, Edin. AQUA PIMENTO, Dub.** *Pimenta Water.*



*Syn.* Eau de poivrée de Jamaïque (F.), Nelherpfeffer wasser (G.), Acqua di Pimenti (I.).

“Take of Pimento Berries bruised, *half a pound*; Water, *a pint*. Macerate the berries in the water for twenty-four hours; and with a sufficient quantity of water to prevent empyreuma, distil a gallon, (ten pounds, *Edin.*)”

This water has the odour and aromatic quality of the Jamaica pepper, but is not very agreeable to the taste. It is used as a carminative in dyspepsia.

AQUA PULEGII, Lond. Dub. AQUA MENTHÆ PULEGII, Edin. *Pennyroyal Water.*

*Syn.* Eau de Menthe peuliot (F.), Poley wasser (G.), Acqua di Puleggio (I.).

“Take of Pennyroyal, *a pound and a half*, (*three pounds*, Edin.) Pour over it a sufficient quantity of Water to prevent empyreuma, and distil a gallon, (ten pounds, *Edin.*)”

Pennyroyal water has the flavour and taste of the green herb. It is used for the same purposes as Peppermint water.

AQUA ROSÆ, Lond. AQUA ROSÆ CENTIFOLIÆ, Edin. *Rose Water.*

*Syn.* Eau des Roses (F.), Rosin wasser (G.) Acqua di Rose (I.), Agua rosada (S.).

“Take of the petals of the hundred-leaved Rose, *eight pounds*, (*six pounds*, Edin.) Pour over them as much water as will prevent empyreuma during the distillation. Distil a gallon, (ten pounds, *Edin.*)”

AQUA ROSÆ, Dub. *Rose Water.*

“Take of the fresh petals of the Damask-rose freed from their claws, *six pounds*; Water, *a sufficient quantity to prevent empyreuma*. Distil a gallon.”

This water has the agreeable odour of the Rose in great perfection when properly prepared; which, however, is seldom the case, except when it is made on a large scale. It is very apt to spoil, unless it be rectified by a second distillation: but spirit of wine ought not to be added to Rose water.

As Rose water is perfectly free from any acrimony, and, except in point of odour, does not differ from simple distilled water, it is very generally employed in collyria, with acetate and superacetate of lead, and acetate and sulphate of zinc.

## INFUSA.

### INFUSIONS.

THESE are solutions of vegetable matter, obtained by maceration either in cold or in boiling water. As in the case of de-



coction, the substance must be sliced or bruised if in a recent state, or pulverized if dry, in order to expose a large surface to the action of the menstruum. The term *Infusion*, in pharmaceutical language, is confined to watery solutions.

The substances which water, without the aid of boiling, can extract from vegetable matter submitted to its action, are gum, mucus, extractive, tannin, the bitter and narcotic principles, gum-resin, volatile oil, acids, and alkalies, a range which includes most of the principles on which the medicinal properties of plants depend. These principles, also, are less liable to be altered by infusion than by decoction, and consequently, this form of preparation is to be preferred in every instance to which it is applicable. The strength and quality of the infusions are varied by the degree of temperature of the water: those made with hot water being necessarily stronger, but particularly in the case of bitters; cold infusions are more grateful.

In making infusions, when heat is required, the vessel is to be placed near the fire, so that the temperature of the water may be kept up to the necessary point for a sufficient length of time to produce the effect intended. Perhaps it might be an advantage, were the external surface of infusion-pots covered with a metallic coating and polished; by which as the heat would be much more slowly radiated than from the vessels usually employed, the effect of it would be more uniform and certain in promoting the solvent powers of the water.

Infusions, like decoctions, are liable to undergo spontaneous decomposition, if kept even for a few days; and therefore the London College has properly directed half a pint only to be made at one time, thus regarding them as extemporaneous preparations.

INFUSUM ANTHEMIDIS, Lond. INFUSUM ANTHEMIDIS NOBILIS, Edin. *Infusion of Chamomile.*

“Take of Chamomile flowers, *two drachms*; Boiling Water, *half a pint*. Macerate for ten minutes (twenty-four hours, Edin.) in a lightly-covered vessel, and strain.”

This infusion is clear, of a pale yellow colour, and has the odour and taste of the flowers. It precipitates solution of Isinglass, whitish; infusion of Yellow Cinchona bark, white; solution of Sulphate of Iron and of tincture of Muriate of Iron, black; solution of Nitrate of Silver, white; of oxymuriate of Mercury, pale brown; and of Acetate and Superacetate of Lead, yellowish white. These substances, therefore, are incompatible in prescriptions with this infusion.

*Medical properties and uses.*—This infusion is a good stomachic and tonic; and may be given in dyspepsia and



other complaints attended with debility of the stomach, in doses of from fʒj. to fʒij. two or three times a day. When exhibited warm it excites nausea, and is occasionally employed to assist the operation of emetics.

**INFUSUM ARMORACIÆ COMPOSITUM**, Lond.  
*Compound Infusion of Horse-radish.*

“Take of fresh Horse-radish root, sliced, Mustard Seed, bruised, of each, *an ounce*; Boiling Water, *a pint*. Macerate for two hours in a lightly-covered vessel, and strain; then add of Compound Spirit of Horse-radish, *one fluid ounce*.”

This infusion, after it is strained, deposits by rest a whitish feculent matter, which should be separated. The supernatant, clear part is of a sulphur-yellow colour, and holds dissolved in every fluid ounce rather more than grs. x. of solid matter. It has a very pungent odour, and a hot, biting taste; precipitates infusion of Galls, yellowish, and infusion of yellow Cinchona bark, white. The solutions of the pure alkalies do not affect it; but with their carbonates whitish precipitates are produced, as is also the case with solution of Oxymuriate of Mercury; while Nitrate of Silver produces a precipitate of a brown colour. Hence all those substances, except the pure alkalies, are incompatible in formulæ with this infusion. This infusion soon spoils in hot weather, and emits an offensive odour.

*Medical properties and uses.* — This is not an unusual form of giving Horse-radish, the stimulant property of which is aided by that of the Mustard. It is particularly serviceable in paralysis, and in dropsies occurring after intermittents. The dose is from fʒj. to fʒij. given three or four times a day.

**INFUSUM AURANTII COMPOSITUM**, Lond.  
*Compound Infusion of Orange-peel.*

“Take of dried Orange-peel, *two drachms*; fresh Lemon-peel, *one drachm*; Cloves bruised, *half a drachm*; Boiling Water, *half a pint*. Macerate for fifteen minutes in a lightly-covered vessel, and strain.”

This infusion has the agreeable compound odour and taste of the ingredients from which it is made. It is clear, and has the brown hue of deep-coloured sherry wine. It precipitates Sulphate of Iron, black; and also produces precipitates with Superacetate of Lead, infusion of yellow Cinchona bark, and lime-water.

*Medical properties and uses.* — It is an excellent and grateful stomachic. The dose may be from fʒj. to fʒv. given twice or thrice a day.

**INFUSUM CALUMBÆ**, Lond. **INFUSUM COLOMBÆ**, Edin. *Infusion of Calumba.*

“Take of Calumba Root, sliced, *one drachm*; Boiling



Water, *half a pint*. Macerate for two hours in a lightly-covered vessel, and strain."

The active matter of Calumba is not all extracted by water. The infusion is inodorous, and tastes bitter. It is clear and of a pale brown colour: affords precipitates with infusion of Yellow Cinchona bark, Lime-water, and solution of Oxymuriate of Mercury, which, therefore, ought not to be ordered in conjunction with it. This infusion soon spoils.

*Medical properties and uses.* — Infusion of Calumba is a good stomachic bitter in dyspeptic cases, and for restraining the nausea and vomiting which occur in pregnancy. It is also useful in the severe diarrhoea and vomiting which often attend dentition. The dose may be from  $\text{f}\overline{\text{z}}\text{jss.}$  to  $\text{f}\overline{\text{z}}\text{ijj.}$  given several times a day.

INFUSUM CARYOPHYLLORUM, Lond. *Infusion of Cloves.*

"Take of Bruised Cloves, *a drachm*; Boiling Water, *half a pint*. Macerate for two hours in a lightly-covered vessel, and strain."

This infusion contains all the active matter of the Cloves; one fluid ounce holding nearly grs.  $\text{vj.}$  in solution. It is of a deep, clear, brown colour, has an aromatic odour, and a bitterish, aromatic taste, affording precipitates with infusion of Yellow Cinchona bark, the strong acids, and Lime-water. Solution of Sulphate of Iron occasions a copious black precipitate; Sulphate of Zinc, Superacetate of Lead, and Nitrate of Silver, brown precipitates. It also decomposes Tartarized Antimony.

*Medical properties and uses.* — It is a warm and grateful stomachic; and may be advantageously used in dyspepsia, particularly when it arises from the abuse of ardent spirits, accompanied with a sensation of coldness at the stomach; in chronic gout, and flatulent colic. The dose is from  $\text{f}\overline{\text{z}}\text{jss.}$  to  $\text{f}\overline{\text{z}}\text{ij.}$  given three or four times a day.

INFUSUM CASCARILLÆ, Lond. *Infusion of Cascarrilla.*

"Take of Cascarrilla bark, bruised, *half an ounce*; Boiling Water, *half a pint*. Macerate for two hours in a lightly-covered vessel, and strain."

This is a clear, pale reddish-brown infusion, having the aromatic odour of the bark, and a bitterish, aromatic taste. It is incompatible in formulæ with the following substances, which it precipitates; Lime-water, infusion of Galls, infusion of Yellow Cinchona bark, solutions of Nitrate of Silver, Acetate and Superacetate of Lead, Sulphate of Zinc, and Sulphate of Iron, which is slowly thrown down, of a pale olive colour.

*Medical properties and uses.* — It is a light stimulant, and



tonic; and is advantageously given in some alvine fluxes, particularly such as occur after measles; and in the aptha gangrenosa of infants. In combination with Carbonate of Soda, it forms an excellent tonic in those affections of children which are dependent on a weak state of the digestive organs, and accompanied with acidity. The dose may be from  $\text{f}\text{ʒ}\text{jss}$  to  $\text{f}\text{ʒ}\text{ij}$ .

**INFUSUM CATECHU COMPOSITUM, Lond.**  
*Compound Infusion of Catechu.*

“Take of extract of Catechu, *two drachms and a half*; Cinnamon bark, bruised, *half a drachm*; Boiling Water, *half a pint*. Macerate for an hour in a lightly-covered vessel, and strain.”

**INFUSUM ACACIÆ CATECHU, Edin.** *Infusion of Catechu.*

“Take of pulverized Extract of Catechu, *two drachms and a half*; Cinnamon Bark, bruised, *half a drachm*; Boiling Water, *seven ounces*; Simple Syrup, *one ounce*. Macerate the Extract and Bark with the Water for two hours, in a covered vessel; then strain, and add the Syrup.”

*Syn.* Infusion de Cachou (*F.*), Katechu infusum (*G.*), Infuso di Cato (*I.*).

In these formulæ it is intended that the whole of the soluble matter of the Catechu taken up by the boiling water should remain dissolved after the infusion cools; but we find that a considerable portion is deposited. When the extract is triturated with water at  $212^{\circ}$ , as much of it is dissolved as the water can hold in solution, so that a preparation similar to this infusion may be immediately made by simply triturating the materials together. The addition of the syrup ordered by the Edinburgh College prevents the preparation from keeping longer than two or three days, although without the syrup it will keep good for months.

*Qualities.* — This infusion is inodorous, and has a slightly bitter austere taste, leaving, even when it contains no syrup, an agreeable sweetness in the mouth. The colour, when the pale Catechu is used, is a light brown or ale colour; but when the dark Catechu is employed, a deep red-brown. The following substances precipitate its tannin, or otherwise alter its properties, and therefore ought not to be ordered in formulæ with it; solution of Isinglass, infusion of Yellow Cinchona bark, the strong Acids, Sulphate of Iron, Sulphate of Zinc, Oxymuriate of Mercury, Tartarized Antimony, and Supracetate of Lead. The Alkalies only deepen the colour.

*Medical properties and uses* — This infusion, which is a powerful, agreeable astringent, is the best form under which Catechu can be prescribed; and is very useful in long-continued diarrhœa, and other fluxes, proceeding from a weakened state of the intestines. The dose is from  $\text{f}\text{ʒ}\text{j}$ .



to fʒiij. given after every liquid dejection, or every four hours.

INFU'SUM CINCHO'NÆ, Lond. *Infusion of Cinchona Bark.*

*Syn.* Infusion de Quinquina (*F.*), Chinainfusum (*G.*), Infuso di China (*I.*).

“Take of Lance-leaved Cinchona Bark<sup>1</sup>, bruised, *half an ounce*; Boiling Water, *half a pint*. Macerate for two hours in a lightly-covered vessel, and strain.”

This infusion contains a portion of Quinia, the febrifuge principle of the bark; but this is considerably augmented when fʒj. of diluted sulphuric acid is added to the boiling water before it is poured on the powdered bark. An infusion made with this addition contains a greater quantity of Quinia than the ordinary infusion, in the proportion of 7 to 6; and the salt, being a sulphate, is more active in its effects. It is slightly turbid, has a pale, pinkish-yellow colour, more of the aromatic odour of the bark than the decoction possesses, and an equal degree of bitterness and astringency. It ferments spontaneously in the course of a few days during summer. It affords precipitates with the following substances; the strong acids, the alkaline carbonates, lime-water, solutions of Sulphate of Iron, Sulphate of Zinc, Nitrate of Silver, Oxymuriate of Mercury, Oxide of Arsenic, Subcarbonate of Potass, and Tartarized Antimony; the aqueous infusions and decoctions of Chamomile flowers, Calumba, Cascarilla, Horse-radish, Cloves, Catechu, Orange-peel, Foxglove, Senna, Rhubarb, Valerian, Simaruba, and Elm-bark.

Any considerable portion of the tinctures also produces precipitates in this infusion. Some of these precipitates take place immediately, others not till after several hours have elapsed: the febrifuge virtue is perhaps not destroyed by them, but the mixtures are certainly rendered inelegant. Sulphuric acid destroys the bitterness of the infusion, but not its astringency; and, as has been already stated, adds considerably to its efficacy.

*Medical properties and uses.*—The Cinchona in this form agrees better with most stomachs than when in powder; but its powers are necessarily diminished. It is chiefly serviceable in dyspepsia, and convalescencies, particularly after the

<sup>1</sup> The other species of Cinchona may be used in the same manner and proportions.

<sup>2</sup> Since writing the article Cinchona (see Part I.) I have seen a statement in a French work, (*“Moyens propres a reconnaitre la purété de diverses substances médicamenteuse;”* par M. J. Fontenelle,) that Sulphate of Quinia is adulterated with Sulphate of Lime, and Carbonate of Magnesia. To detect these frauds put the suspected Sulphate of Quinia into rectified spirit of wine, which will dissolve the whole of the Sulphate and leave the Salts of Lime and Magnesia. These salts when washed with distilled water are insipid.



maturation of the pustules in *Ecthyma vulgare*. The dose is from fʒj. to fʒij. three or four times a day.

INFUSUM CINCHONÆ LANCIFOLIÆ, Edin. *Infusion of Cinchona*. INFUSUM CINCHONÆ SINE CALORE, Dub. *Cold Infusion of Cinchona*.

“Take of Cinchona Bark, bruised, *one ounce*; Water, *one pound* (*twelve ounces by measure*, Dub.) Macerate for twenty-four hours, agitating frequently, and strain. (Triturate the bark with a little of the water, and whilst triturating add the remainder: then macerate for twenty-four hours, occasionally agitating, and decant the clear liquor. *Dub.*”)

The directions of the Dublin College for making this infusion are preferable to those of the Edinburgh College. It is nearly clear, but deposits by rest a small quantity of a brick-red sediment. It is affected by the same substances, and its properties and use are the same as those of the former preparation, from which it differs chiefly in strength. The residuum may be used for some purposes, as its active principle is not nearly exhausted.

INFUSUM CUSPARIÆ, Lond. *Infusion of Cusparia*.

“Take of Cusparia Bark, bruised, *two drachms*; Boiling Water, *half a pint*. Macerate for two hours in a lightly-covered vessel, and strain.”

This infusion is slightly turbid, and of a brownish colour; has a somewhat aromatic odour, and a bitter taste. The solution of Sulphate of Iron throws down a greenish-yellow precipitate, and Sulphate of Zinc a yellowish one; Nitrate of Silver, Oxymuriate of Mercury, Superacetate of Lead, infusions of Galls, and of Catechu also, produce precipitates in it. Tartarized Antimony is slowly decomposed. These substances, therefore, cannot properly be ordered in formulæ with this infusion.

*Medical properties and uses.*—This infusion possesses the stimulant and tonic properties of the bark, and is an useful form of giving it in typhoid fevers, obstinate bilious diarrhœa, and in dysentery, after proper evacuations. The tincture of Cinnamon both covers its taste, and makes it sit lighter on the stomach. The dose is from fʒj. to fʒij. given every three or four hours.

INFUSUM DIGITALIS, Lond. *Infusion of Foxglove*.

“Take of dried Foxglove Leaves, *a drachm*; Boiling Water, *half a pint*. Macerate for four hours in a lightly-covered vessel, and strain; then add of Spirit of Cinnamon, *half a fluid ounce*.”

INFUSUM DIGITALIS PURPUREÆ, Edin. *Infusion of Foxglove*.



“Take of dried Foxglove Leaves, *one drachm*; Boiling Water, *eight ounces*; Spirit of Cinnamon, *one ounce*. Macerate the Leaves with the Water for four hours in a lightly-covered vessel; then, having added the Spirit, strain.”

*Syn.* Infusion de Digitale purpurine (F.), Fingerhut aufgus (G.), Infuso di Digitale porporina (I.).

The faint odour and nauseous bitter taste of the Foxglove are covered by the Spirit of Cinnamon in these infusions, which are clear, and of a brownish-yellow colour. The solution of Sulphate of Iron slowly throws down in them a pale-olive precipitate; Superacetate of Lead and infusion or decoction of Yellow Cinchona Bark produce instantaneous and copious precipitates.

*Medical properties and uses.*—These infusions do not differ materially from that made by the formula of Withering, and are well calculated to obtain speedily the diuretic effects of the remedy. The dose is from  $f\text{ʒss.}$  to  $f\text{ʒj.}$  given twice a day; or every eight hours, if the patient be strong, and the symptoms very urgent. For the necessary cautions to be observed in administering them, see the article *Digitalis*, Part ii.

INFUSUM GENTIANÆ COMPOSITUM, London.  
*Compound Infusion of Gentian.*

“Take of Gentian Root, sliced, Orange-peel, dried, of each *a drachm*; fresh Lemon-peel, *two drachms*; Boiling Water, *twelve fluid ounces*. Macerate for an hour in a lightly-covered vessel, and strain.”

*Edinburgh.*

“Take of Gentian Root, sliced, *half an ounce*; Dried Orange-peel, bruised, Coriander Seeds, bruised, of each *a drachm*; Diluted Alcohol, *four ounces*; Water, *one pound*. First, pour on the Alcohol, and, after three hours, the Water; then macerate without heat for twelve hours, and strain.”

*Dublin.*

“Take of Gentian Root, bruised, *two drachms*; Fresh Lemon-peel, *half an ounce*; Dried Orange-peel, *a drachm and a half*; Proof Spirit, *four ounces by measure*; Boiling Water *twelve ounces by measure*. First, pour on the Spirit, and, three hours afterwards, the Water; then macerate for the space of two days, and strain.”

The Spirit ordered by the Edinburgh and Dublin Colleges, is intended to aid the solvent power of the water, and to preserve the infusion, which in summer very soon becomes ropy, and spoils; but as infusions can always easily be prepared, and boiling water takes up the greater part of the active matter of the ingredients, the spirituous addition, and the length of time ordered for the maceration, are certainly ob-



jectionable. The formula of the London College is free from both these objections, and produces a clear infusion, of a yellowish colour, with the agreeable odour of the Orange-peel, and a pleasant bitter taste. The solution of Acetate of Lead throws down a copious precipitate in this infusion; and Sulphate of Iron strikes a brown colour, but no precipitate takes place for twelve hours.

*Medical properties and uses.*—These are very commonly used, and are elegant tonic and stomachic infusions. They are given in dyspepsia and chlorosis, united with chalybeates, or with alkalies; in atonic gout and diarrhœa, with absorbents and aromatics; in jaundice, with Rhubarb and saline purgatives; and in dropsies, with Squills and neutral salts. From fʒj. to fʒij. may be given for a dose, three or four times a day.

INFUSUM LINI COMPOSITUM, Lond. *Compound Infusion of Linseed.*

“Take of Linseed, bruised, *an ounce*; Liquorice Root, sliced, *half an ounce*; Boiling Water, *two pints*. Macerate for four hours near the fire, in a covered vessel, and strain.”

INFUSUM LINI USITATISSIMI, Edin. *Infusion of Linseed.*

“Take of Linseed, *an ounce*; Liquorice Root, bruised, *two drachms*; Boiling Water, *two pounds*. Digest for four hours in a lightly-covered vessel, and strain.”

*Syn.* Infusion de Semence de Lin (*F.*), Leinsamen aufgus (*G.*), Infuso d. Semi di Lino (*I.*).

This infusion is a solution of mucus nearly in its pure state. It is clear, colourless, inodorous, and nearly insipid. Alcohol precipitates the mucus in white flocculi; and precipitates are also produced by Subacetate and Acetate of Lead, and the tincture of muriated Iron; hence these substances are incompatible in formulæ with this infusion.

*Medical properties and uses.*—Infusion of Linseed is a cheap and very useful demulcent, in the various cases in which this class of remedies is indicated, and during the internal exhibition of corrosive muriate of mercury. The dose is fʒij. frequently repeated.

INFUSION MENTHÆ COMPOSITUM, Dub. *Compound Infusion of Mint.*

“Take of the Leaves of Spearmint, dried, *two drachms*; Boiling Water, *a sufficient quantity to afford six ounces by measure when strained*. Digest for half an hour in a covered vessel, and strain the liquor when cold; then add of Refined Sugar, *two drachms*, Oil of Spearmint, *three drops*, dissolved in *half an ounce* (fluid?) of Compound Tincture of Cardamoms. Let them be mixed.”

*Medical properties and uses.*—This is a grateful stomachic,



and is also slightly diaphoretic. It may prove serviceable in anorexia and nausea, and as a vehicle to cover the disagreeable taste of other medicines. The dose may be from  $\text{f}\mathring{\text{z}}\text{j.}$  to  $\text{f}\mathring{\text{z}}\text{iiij.}$  or *ad libitum*.

INFUSUM QUASSIÆ, Lond. *Infusion of Quassia.*

“Take of Quassia Wood, chipped, *a scruple*; Boiling Water, *half a pint*. Macerate for two hours in a lightly-covered vessel, and strain.”

INFUSUM QUASSIÆ EXCELSÆ, Edin. *Infusion of Quassia.*

“Take of Quassia Wood, rasped, *half a drachm*: Boiling Water, *eight ounces*. Macerate for two hours in a lightly-covered vessel, and strain.”

The active matter of Quassia taken up by water appears to be a pure, simple bitter. It is not altered by any of the substances usually employed as adjuncts to bitters; and by two only of the metallic salts. Nitrate of Silver slowly throws down soft, yellow flakes; and Acetate of Lead, a white precipitate.

*Medical properties and uses.* — This infusion is a light tonic, very efficacious in dyspepsia, and other cases in which tonics are indicated. In hysteria it may be combined with purgatives and Tincture of Valerian; in atonic gout, with aromatics; and in dyspeptic affections with chalybeates, Sulphate of Zinc, or mineral acids. The dose is from  $\text{f}\mathring{\text{z}}\text{j.}$  to  $\text{f}\mathring{\text{z}}\text{iiij.}$  given twice or thrice a day.

INFUSUM RHEI, Lond. *Infusion of Rhubarb.*

“Take of Rhubarb Root, sliced, *a drachm*; Boiling Water, *half a pint*. Macerate for two hours in a lightly-covered vessel, and strain.”

*Edinburgh.*

“Take of Rhubarb Root, bruised, *half an ounce*; Boiling Water, *eight ounces*; Spirit of Cinamon, *one ounce*. Macerate the Root with the Water in a covered vessel for twelve hours; then add the Spirit, and strain.”

*Syn.* Infusion de Rhubarbe (*F.*), Rhabarber aufguss (*G.*), Infuso de Rabarbaro (*I.*).

These infusions differ chiefly in point of strength; and the Edinburgh is rendered pleasanter by the spirituous addition. Neither of them is quite clear; and both have a reddish-brown colour, which is very much deepened by the addition of alkalies. The following substances either occasion precipitates in these infusions, or otherwise alter their properties, and are therefore incompatible in formulæ with them; the strong acids, and Lime-water, solutions of Sulphate of Iron, Sulphate of Zinc, Nitrate of Silver, Oxymuriate of Mercury, Acetate of Lead, and Tartarized Antimony; Infusions of Catechu, Cinchona, and Cusparia.



*Medical properties and uses.* — These infusions are good forms for exhibiting rhubarb, when it is intended to act on the bowels: but they are considerably less active than the powder. The dose of the London infusion may be from fʒj. to fʒiv. and of the Edinburgh half the quantity, united with neutral salts or aromatics, as circumstances may direct.

INFUSUM ROSÆ COMPOSITUM, Lond. Dub.  
*Compound Infusion of Roses.*

“Take of the dried petals of the Red Rose, *half an ounce*; Boiling Water, *two pints and a half* (three pounds by measure, Dub.); Diluted Sulphuric Acid, *three fluid drachms* (three drachms by weight, Dub.); Double-refined Sugar, *an ounce and a half*. Pour the water on the Rose-petals in a covered glass vessel; then drop in the Acid, and macerate for half an hour. Finally, strain the liquor, and add the Sugar to it.”

INFUSUM ROSÆ GALLICÆ, Edin. *Infusion of Red Roses.*

“Take of the dried petals of the Red Rose, *one ounce*; Boiling Water, *two pounds and a half*; Sulphuric Acid diluted, *half an ounce*; Refined Sugar, *one ounce*. Macerate the petals with the water in an earthen vessel, which has not been glazed with lead, for four hours; then pour in the Acid, strain the liquor, and add the Sugar.”

*Syn.* Infusion de Roses (*F.*), Rosen aufguss (*G.*) Infuso di Rose (*I.*).

This infusion is clear, of a beautiful red colour, and has an acid, pleasantly austere taste.<sup>1</sup> The addition of the sugar prevents it from keeping so long as it might otherwise be kept. The incompatible substances are those which are decomposed by the sulphuric acid. The Sulphates of Iron and of Zinc, although they do not immediately alter it, yet slowly produce dark-coloured precipitates after some hours.

*Medical properties and uses.* — Infusion of Roses is indebted for any astringency it possesses chiefly to the acid it contains. It is used alone in the colliquative sweats of phthisis; and as a gargle in cynanche tonsillaris; but it is chiefly employed as an elegant vehicle for more active remedies, particularly Sulphate of Magnesia, the nauseous taste of which it completely covers. The dose is from fʒij. to fʒiv.

INFUSUM SENNÆ COMPOSITUM, Lond. *Compound Infusion of Senna.*

“Take of Senna Leaves, *an ounce and a half*; Ginger Root, sliced, *a drachm*; Boiling Water, *a pint*. Macerate for an hour in a lightly-covered vessel, and strain the liquor.”

INFUSUM CASSIÆ SENNÆ, Edin.

“Take of Senna Leaves, *six drachms*; Ginger Root,

<sup>1</sup> Dr. Clarke, of Cambridge, supposed that he had detected iron in the petals of the Rose.



bruised, *a scruple* ; Boiling Water, *nine ounces*. Macerate for an hour in a lightly-covered vessel, and strain."

INFUSUM SENNÆ, Dub. *Infusion of Senna.*

"Take of Senna Leaves, *three drachms* ; Lesser Cardamom Seeds, freed from the capsules and bruised, *half a drachm* ; Boiling Water, *as much as will yield, when strained, six ounces by measure*. Digest for an hour, and when the liquor is cold, strain it."

*Syn.* Infusion de Senè (F.), Senna aufguss (G.), Infuso di Senna (I.).

These infusions should be clear, and have a deep red-brown, nearly black colour; with a slightly bitter, mawkish taste, which is scarcely corrected by the aromatic. In warm weather they spoil in forty-eight hours; and by simple exposure to the air attract oxygen, which occasions a yellowish precipitate of oxidized extractive; that is not purgative, but gripes violently. On this account they should be preserved in a well-closed vessel, or made only when wanted. Dr. Paris (*Pharmacologia*) observes, that the nauseous taste of these infusions is completely covered by the addition of *Bohea tea*. Decoction of Guaiac is, also, said to increase their powers, and to render them milder. They are also precipitated by the strong acids, the alkaline carbonates, Lime-water, solutions of Nitrate of Silver, Oxymuriate of Mercury, Superacetate of Lead, Tartarized Antimony, and infusion of Yellow Cinchona bark, which are consequently incompatible in formulæ with these infusions.

*Medical properties and uses.* — Both these infusions contain all the purgative principles of the plant, whilst the aromatics correct its griping properties: but there is perhaps a waste of Senna in the London formula. They are given alone, or more generally combined with neutral salts and manna. The dose of the simple infusions may be from fʒij. to fʒiv.; but with the addition of ʒj. of the Tartrate of Potass, or ʒij. of the Sulphate of Magnesia, which are the usual adjuncts, fʒij. are sufficient.

INFUSUM SENNÆ COMPOSITUM, Edin. *Infusion of Tamarinds and Senna.*

"Take of Preserved Tamarinds, *one ounce* ; Senna Leaves, *one drachm* ; Coriander Seeds, bruised, *half a drachm* ; Raw Sugar, *half an ounce* ; Boiling Water, *eight ounces*. Macerate in a covered earthen vessel, which is not glazed with lead, shaking frequently, and after four hours strain.

"It may be made also with double or triple the proportion of Senna."

INFUSUM SENNÆ CUM TAMARINDIS, Dub. *Infusion of Senna and Tamarinds.*

Made in the same manner as the infusion of Senna, ex-



cept that ℥j. of Tamarinds is added before straining the liquor.

These infusions are pleasanter than the simple infusions, the nauseous taste being well covered by the sugar and the acid of the Tamarinds: in other respects they agree both in their properties, and in the effects of the incompatible substances; to which, however, must be added all salts having potass for their base.

INFUSUM SIMAROUBÆ, Lond. *Infusion of Simaruba.*

“Take of Simaruba bark, bruised, *half a drachm*; Boiling Water, *half a pint*. Macerate for two hours in a lightly-covered vessel, and strain.”

This infusion is inodorous; has a slightly bitter taste, is clear, and of a greenish straw-colour. The alkaline carbonates and lime-water render it milky, and the following substances occasion precipitates: Nitrate of Silver, Oxymuriate of Mercury, Superacetate of Lead; infusions of Galls, Catechu, and Yellow Cinchona bark.

*Medical properties and uses.* — Simaruba infusion possesses the same properties as the bark, and is the best form of exhibiting the remedy, but it is not much used in this country. The dose is f℥ij. combined with Tincture of Opium, or an aromatic.

INFUSUM TABACI, Lond. *Infusion of Tobacco.*

“Take of Tobacco Leaves, *a drachm*; Boiling Water, *a pint*. Macerate for an hour in a lightly-covered vessel, and strain.”

This infusion is clear, of a reddish-brown colour; has the odour of the plant in a slight degree, and a hot, very acrid taste.

*Medical properties and uses.* — Tobacco infusion is chiefly intended to be given under the form of enema; for although it has been occasionally employed as an emetic, it cannot be recommended. As an enema, it has been found useful in ileus, colica pictonum, incarcerated hernia and dysury; the practice of employing it in cases of suspended animation is now justly condemned.

INFUSUM VALERIANÆ, Dub. *Infusion of Valerian.*

“Take of Valerian Root, coarsely powdered, *two drachms*; Boiling Water, *seven ounces by measure*. Digest for an hour, and when the liquor is cold, strain.”

Valerian infusion is clear, of a pale-brown colour; with the odour of the Valerian, and a bitterish pungent taste. Solutions of Nitrate of Silver, Sulphate of Iron, and infusion of Yellow Cinchona, afford precipitates with this infusion; and are therefore incompatible in formulæ with it.



*Medical properties and uses.*—This is an useful form of giving Valerian in hysterical and nervous affections, in which the stomach will not always bear the powder. The dose may be from fʒjss. to fʒij. twice or thrice a day.

AQUA CALCIS COMPOSITA, Dub. *Compound Lime-water.*

“Take of Raspings of Guaiacum Wood, *half a pound*; Liquorice Root, sliced and bruised, *an ounce*; Sassafras Bark, bruised, *half an ounce*; Coriander Seeds, *three drachms*; Lime-water, *six pints*. Macerate without heat for two days, and strain.”

This is a very inert preparation; and unless great care be taken to exclude the air completely from the vessel in which it is made, the lime-water will be decomposed.

AQUA PICIS LIQUIDÆ, Dub. *Tar Water.*

“Take of Tar, *two pints*; Water, *a gallon*. Mix, stirring with a wooden rod for a quarter of an hour; then, after the Tar shall have subsided, let the liquor be strained, and preserve it in well-corked bottles.”

Water readily dissolves a portion of Tar; and is impregnated with empyreumatic oil, a small portion of resinous matter, and acetic acid, the components of the Tar. The solution has the colour of Madeira wine, and a sharp empyreumatic taste.

*Medical properties and uses.* — Tar water is stimulant and diuretic; but to produce the latter effect, its operation requires to be aided by bodily exercise. It may prove useful in scurvy, and some cutaneous diseases; but the reputation which it obtained on the faith of the judgment of the worthy Bishop of Cloyne<sup>1</sup> has long since been lost, and it is now scarcely ever employed. From ʒj. to ʒij. may be taken in the course of a day.

---

## MUCILAGINES.

### MUCILAGES.

MUCILAGES, correctly speaking, are simple solutions of gum or mucus in water; but the term Mucilage, in pharmaceutical language, implies also any solution of a thick and adhesive nature, resembling in its appearance the solutions of gum.

---

<sup>1</sup> Berkley's *Siris* — passim.



MUCILAGO ACACIÆ, Lond.<sup>1</sup> *Mucilage of Acacia.*

“Take of Acacia Gum, in powder, *four ounces*; Boiling Water, *half a pint*. Rub the Gum with the Water, gradually added, until it form a Mucilage.”

MUCILAGO ACACIÆ ARABICÆ, Edin. *Mucilage of Gum Arabic.*

“Take of Gum Arabic, in powder, *one part*; Boiling Water, *two parts*. Digest with occasional agitation, until the gum be dissolved; then strain the Mucilage through linen.”

MUCILAGO GUMMI ARABICI, Dub. *Mucilage of Gum Arabic.*

“Take of Gum Arabic, in coarse powder, *four ounces*; Boiling Water, *eight ounces*. Digest with frequent agitation, until the Gum be dissolved; then strain the Mucilage through linen.”

*Syn.* Mucilage de Gomme Arabique (*F.*), Schlieim de Arabiche Gummi (*G.*), Mucilagine de Gomma Arabica (*I.*).

The straining through linen is very necessary, as the gum is often mixed with small pieces of wood and other impurities. The mucilage thus obtained is viscid, thick, and adhesive; semipellucid, and nearly colourless, if the gum be good. It has a faint, peculiar odour, is insipid, and may be kept without altering for a considerable time; but at length it becomes sour, and acetic acid is formed. The strong acids act on it as they do on gum; but when diluted, they do not alter mucilage. Alcohol converts it into a white curd; but proof spirit produces scarcely any alteration; no change is produced by Spirit of Nitric Ether; but Sulphuric Ether and Compound Spirit of Ether precipitate a thick, white, tenacious curd. Tincture of Muriate of Iron, even when diluted, converts mucilage into a brownish or orange-coloured, insoluble jelly; and Subacetate of Lead gives a copious, dense, flaky precipitate, which is a compound of gum and oxide of lead; while no change is produced by the solutions of the following metallic substances: Superacetate of Lead, green Sulphate of Iron, Sulphate of Zinc, Oxymuriate of Mercury, and Tartarized Antimony; nor by the alkalies or the neutral salts. Mucilage, like gum, serves to combine resins, oils, and balsams with water, for which purpose, and to give tenacity to pills, it is much employed in pharmacy.

*Medical properties and uses.* — The properties of mucilage are the same as those of gum. (See *Part ii.*) It is the usual basis of demulcent mixtures for allaying the tickling which

---

<sup>1</sup> This appellation is certainly exceptionable. It is a mucilage of gum of the *Acacia vera*. The Edinburgh name is liable to nearly the same objection, except that the specific name of the plant is used, whereas *Acacia* is the name of the genus.



excites cough in catarrh and phthisis<sup>1</sup>; and, combined with Opium and other narcotics, it is useful in diarrhœa, dysentery, calculous affections, and ardor urinæ. The dose of mucilage may be from fʒss. to fʒj. frequently repeated.

Officinal preparations. *Mistura Guaiaci*, L. *Potio Carbonatis Calcis*, E.

MUCILAGO ASTRAGALI TRAGACANTHÆ, Edin.  
*Mucilage of Tragacanth.*

“Take of Gum Tragacanth, in powder, *two drachms*; Boiling Water, *eight ounces*. Macerate for twenty-four hours, and triturate the gum carefully, that it may be dissolved; then strain the mucilage through linen.”

MUCILAGO GUMMI TRAGACANTHÆ, Dub. *Mucilage of Gum Tragacanth.*

“Take of Gum Tragacanth, in powder, *two drachms*; Water, *eight fluid ounces*. Macerate in a covered vessel until the gum be dissolved; then strain the mucilage through linen.”

*Syn.* Mucilage de Gomme Adraganthe (*F.*), Schlieim de Traganth (*G.*), Mucilage di Gomma Adragante (*I.*).

Tragacanth treated in this manner forms a thick, soft, very viscid mucilage, but the diffusion in the water is not uniform; nor does it become so even when boiled. The water separates from the Tragacanth, on standing; and this separation is increased, if mucilage of gum be mixed with the Tragacanth. It may be used in the same cases as mucilage of Gum Arabic; and has been recommended by M. Blaire, a French surgeon, as a remedy in burns. He directs linen rags, or bibulous paper, soaked in the mucilage, to be applied over the affected part, which must be also kept moist with the mucilage for some days.<sup>2</sup> It is chiefly employed for making pills and troches.

MUCILAGO AMYLI, Lond., Edin., Dub. *Mucilage of Starch.*

“Take of Starch, *three drachms*; Water, *a pint*. Rub the Starch, gradually adding the water to it; then boil till a mucilage be produced.”

Starch thus treated forms a strong, insipid, inodorous, opaline-coloured, gelatinous mucilage. In cases of phthisis, hectic fever, and abrasions of the stomach, it is given as a demulcent by the mouth; but it is more generally, and more advantageously exhibited in the form of enema in diarrhœa, dysentery, and abrasions of the rectum. It is the common vehicle for exhibiting Opium in the form of enema.

<sup>1</sup> R Mucilaginis Acaciæ fʒj. Olei Amygdalarum, Syrupi Papaveris Albi, ā ā fʒss., Aquæ fʒiv., Acidi Citrici q. s. ad gratam acidulat. M. ut fiat mistura cujus sumat æger cochl. med. unum subindè.

<sup>2</sup> Vide *London Med. Repository*, vol. iii. p. 257.

401

*ad gratam acidulat*  
*subindè = 7*  
*sumat æger*



*DECOCTA.*

## DECOCTIONS.

THESE are aqueous solutions of the active principles of vegetables obtained by boiling. They are intended to afford more powerful remedies than can be obtained by the simple infusion of the same substances in cold or even in boiling water; but, although, by the operation of boiling, the solvent power of the water is increased, and a greater quantity of the soluble parts of any vegetable body is consequently taken up by it, yet, it does not always follow, that the medicinal virtues of decoctions are greater than those of infusions. On the contrary, if the active principles of a plant be volatile, or if they consist chiefly of extractive matter, this form of preparation often renders the remedy altogether inert, either by dissipating the volatile matters, or by favouring the oxidizement of the extractive, which, in a continued temperature of  $212^{\circ}$ , attracts the oxygen of the atmosphere so rapidly, that it is soon converted into a soluble, insipid, inert matter, and precipitated in the fluid. This is the case with some substances, which are nevertheless ordered to be prepared in this form by the Colleges, and which we shall particularly notice in treating of the individual decoctions.

For making decoctions, the substances employed must be divided, if in the dry state, by pulverization, or, if fresh, by slicing, so as to expose an extended surface to the action of the water; which is thus enabled to take up their soluble principles in a shorter space of time, a circumstance, for the reasons already stated, of much importance in the preparation of decoctions. By covering the vessel in which they are made, the action of the air is prevented from affecting the ingredients; but there is reason for believing, that by long coction in water, even in covered vessels, the constituents of some vegetable bodies re-act upon one another, and produce entirely new compounds, possessed of properties altogether different from those which they previously constituted. On this account, decoctions should be quickly made; and when aromatic or volatile ingredients are to enter into them, these should not be boiled with the more fixed substances, but the decoction, after it is made, should be poured over them, and allowed to remain covered up until it be nearly cold, before it is strained. In general, however, it is better to strain decoctions while they



are hot through a sieve; for as boiling water dissolves a larger proportion of vegetable matter than it can retain in solution at a lower temperature, a deposit almost always takes place as the decoction cools; and if this be of active matter, it is lost by deferring the straining; whereas by straining the decoction while hot, the deposit can be mingled, by being shaken, with the clear fluid, when it enters into extemporaneous compositions, or when the dose of it is taken.

Decoctions, from the nature of their constituents, very soon ferment and spoil: consequently, they should be prepared in small quantities only, and never used, particularly in summer, forty-eight hours after they have been made.

**DECOCTUM ALOES COMPOSITUM, Lond.** *Compound Decoction of Aloes.*

“Take of Extract of Liquorice, *half an ounce*; Subcarbonate of Potass, *two scruples*; Extract of Spiked Aloes, powdered, Myrrh, powdered, Saffron, of each, *a drachm*; Water, *a pint*. Boil down to twelve fluid ounces, and strain: then add of Compound Tincture of Cardamoms, *four fluid ounces*.”

By the addition of the alkali in this preparation, the water is enabled to hold in solution a greater portion of the Aloes than it could otherwise hold, while another portion is suspended by the mucilage of the Liquorice and the Myrrh. The addition of the tincture prevents any spontaneous decomposition from taking place. The taste of the decoction is extremely nauseous, notwithstanding the bitter of the Aloes is in some degree covered by the Liquorice: but the tastes of even children are soon reconciled to it; and in some instances we have seen it preferred to all other purgatives simply on account of its flavour. It is decomposed, and a flaky precipitate thrown down in it by all the strong acids; Corrosive Muriate of Mercury produces a pale brown precipitate, while Tartarized Antimony, Sulphate of Zinc, and Superacetate of Lead, produce white curdy precipitates: thence these substances are incompatible in formulæ with this decoction. This decoction may be kept for a much longer time than any other without spoiling.

*Medical properties and uses.* — It is gently cathartic and emmenagogue; and is introduced as being analogous to the well-known *Beaume de Vie*; which, however, contained no alkali. It may be given with advantage in habitual costiveness, dyspepsia, hypochondriasis, jaundice, and chlorosis, in the dose of from  $\text{f}\overline{\text{z}}\text{ss}$ . to  $\text{f}\overline{\text{z}}\text{ij}$ . taken in the morning.

**DECOCTUM ALTHÆÆ OFFICINALIS, Edin.** *Decoction of Marsh Mallows.*



“Take of Marsh Mallow Root, dried and bruised, *four ounces*; Raisins stoned, *two ounces*; Water, *seven pounds*. Boil down to five pounds; set aside the strained liquor until the dregs have subsided, and then decant it.”

Marsh Mallow roots contain a considerable quantity of mucus, which is thus extracted unaltered by water. The simple decoction of the roots is viscid, of a pale-yellow colour, sweetish, and has a peculiar odour resembling that of boiled turnips. In the above preparation, the raisins increase its sweetness, and render it more palatable.

*Medical properties and uses.* — This decoction is a useful demulcent in visceral inflammations, calculous affections, gonorrhœa, strangury, and other diseases of the urinary organs; and is supposed to be particularly indicated in inflammation of the kidneys after the abscess bursts. The dose is a cupful frequently taken; but in inflammation of the urinary organs, and in similar cases, it may be drunk *ad libitum*, as common beverage.

DECOCTUM ANTHEMIDIS NOBILIS, Edin. *Decoction of Chamomile.*

“Take of Chamomile Flowers, dried, *one ounce*; Carraway Seeds, bruised, *half an ounce*; Water, *five pounds*. Boil for a quarter of an hour, and strain.”

DECOCTUM CHAMÆMELI COMPOSITUM, Dub. *Compound Decoction of Chamomile.*

“Take of Chamomile Flowers, dried, *half an ounce*; Fennel Seeds, *two drachms*; Water, *a pint*. Boil a little, and strain.”

These decoctions contain in solution bitter extractive, and a small portion of essential oil. Were their mode of preparation a matter of any consequence, we would recommend the aromatic seeds not to be added till towards the conclusion of the boiling; but for the purposes of fomentation and glyster, for which they are intended, as much benefit is probably derived from the warm water, as the principles it holds in solution.

DECOCTUM CINCHONÆ, Lond. *Decoction of Cinchona.*

“Take of Lance-leaved Cinchona Bark bruised, *an ounce*; Water, *a pint*. Boil for ten minutes in a lightly-covered vessel, and strain the liquor while it is hot.”

DECOCTUM CINCHONÆ LANCIFOLIÆ, Edin. *Decoction of Lance-leaved Cinchona.*

“Take of Cinchona Bark, in powder, *one ounce*; Water, *one pound and a half*. Boil for ten minutes in a covered vessel, and strain the liquor while it is hot.”



DECOCTUM CORTICIS CINCHONÆ, Dub. *Decoction of Cinchona Bark.*

“Take of Cinchona Bark in coarse powder, *an ounce*; Water, *a pint*. Boil for ten minutes in a vessel almost covered, and strain the liquor through a linen cloth while it is hot.”

*Syn.* Decoction de Quinquina (*F.*), Chinadekokte (*G.*), Decotto di China (*I.*).

Cinchona bark is one of those substances which suffers by long coction with water; and therefore, the Colleges have properly limited the time of boiling to ten minutes, and ordered the vessel to be covered, and the liquor to be strained while it is hot. As the strained decoction cools, it becomes turbid, and lets fall a reddish or yellowish powder, according to the kind of bark used: this, however, must not be rejected, but diffused through the clear decoction when it is about to be used in compounding extemporaneous mixtures, or when the dose is to be taken. When the decoction is made with the lance-leaved Cinchona it contains a Kinate or rather Cinchonate of Cinchonia; and on this its peculiar properties depend. By long boiling, says Pelletier, the tannin and starch form a compound insoluble in cold water, and which is therefore in conjunction with colouring matter, cinchonate of lime &c., precipitated as the decoction cools, and carries down also a portion of the Cinchonia; to remedy which Pelletier recommends a larger quantity of water to be used, and the decoction to be filtered and evaporated. This, however, is a tedious process, and, as Sulphate of Cinchonia is very soluble, we are of opinion that the addition of fʒj. of Diluted Sulphuric Acid to the water for making the decoction will answer the same intention.

This decoction is more bitter, but less aromatic than the infusion. It is affected by the same re-agents, and used in the same cases, and in similar doses, as the infusion. (See *Infusum Cinchonæ*.)

DECOCTUM CYDONIÆ<sup>1</sup>, Lond. *Decoction of Quince Seeds.*

“Take of Quince Seeds, *two drachms*; Water, *a pint*. Boil them over a gentle fire for ten minutes, then strain.”

Quince Seeds abound with mucus, which is extracted by boiling water. It is considerably viscid, transparent, nearly colourless, insipid and inodorous. It is coagulated by Alcohol, acids, and most of the metallic salts, which, therefore, are incompatible in formulæ with it; and it must be used as

---

<sup>1</sup> This title would lead to the inference, that the preparation is a decoction of the quince, and not of the seeds. It should have been *Cydoniæ Seminum*.



soon as it is made, for it soon spoils, owing perhaps to its containing some of the other constituents of the seeds.

*Medical properties and uses.* — This is often preferred to the other mucilages as a local demulcent in tenesmus, and in aphthous affections and excoriations of the mouth. A diluted solution of it injected beneath the eye-lids is useful for obtunding the acrimony of the discharge in violent inflammations of the eye.

**DECOCTUM DAPHNES MEZERII**, Edin. *Decoction of Mezereon.*

“Take of the Bark of Mezereon Root, *two drachms*; Liquorice Root bruised, *half an ounce*; Water, *three pounds*. Boil with a gentle fire down to two pounds, and strain.”

*Syn.* Décoction de Daphne Mezereon (*F.*), Scioblbastrinde-dekokte (*G.*), Decotto di Daphne Mezereon (*I.*).

This Decoction is slightly mucilaginous and of a yellowish-brown colour; has the sweet taste of the liquorice root with a slight degree of bitterness; and leaves in the mouth a sensation of heat and pungency, which, however, is scarcely felt until a few minutes after the dose has been swallowed.

*Medical properties and uses.* — This decoction was first made public by Dr. Alexander Russel<sup>1</sup> as an appropriate remedy for venereal nodes, arising from a thickening of the periosteum; and for removing those nocturnal pains with which venereal patients are afflicted. This opinion, however has not been supported by experience; and Mr. Pearson<sup>2</sup> asserts, that it “has not the power of curing the venereal disease in any one stage, or any one form;” and adds, “except in an instance or two of lepra, in which the decoction conferred a temporary benefit, I have very seldom found it possessed of medicinal virtue, either in syphilis, or in the sequelæ of that disease, in scrophula, or in cutaneous affections.” It has been given with seeming benefit in chronic rheumatism. The dose is from fʒiv. to fʒvi. three or four times a day.

**DECOCTUM DIGITALIS**, Dub. *Decoction of Foxglove.*

“Take of Foxglove Leaves dried, *a drachm*; Water, *as much as will afford of strained liquor eight fluid ounces*. Place the vessel over a gentle fire, and as soon as the liquor boils, remove it; then digest for a quarter of an hour, and strain.”

This decoction is almost inodorous, and has a bitter nauseous taste. It is affected by the same re-agents as the infusion, and used in dropsies with the same intention. (See *Infusum Digitalis*.)<sup>3</sup>

<sup>1</sup> *Medical Observations and Inquiries*, vol. iii.

<sup>2</sup> *Pearson on the Remedies for Lues Venerea*, p. 47.

<sup>3</sup> For cases of the successful use of this decoction, in making which, however, the fresh leaves were employed instead of the dried, see *Darwin's Zoonomia*, vol. i. p. 326. 331.



DECOCTUM DULCAMARÆ, Lond. *Decoction of Woody Nightshade.*

“Take of the Stalks of Woody Nightshade, sliced, *one ounce*; Water, *a pint and a half*. Boil down to a pint, and strain.”

This decoction appears to have been introduced into the Pharmacopœia merely to fix the proportions of the ingredients. It has a strong unpleasant odour, and a bitter, nauseous taste followed by a degree of sweetness.

*Medical properties and uses.* — It is possessed of diuretic and narcotic properties; and has been found useful in humoral asthma, dropsy, lepra vulgaris and alphas, and ptyriasis. The dose is from fʒiv. to fʒj. combined with any aromatic tincture, given three times a day.

DECOCTUM GEOFFRÆÆ INERMIS, Edin. *Decoction of Cabbage-tree Bark.*

“Take of Cabbage-tree Bark in powder, *one ounce*; Water, *two pounds*. Boil with a gentle heat down to one pound, and strain.”

This decoction has the colour of Madeira wine, a disagreeable odour, and a bitter, mucilaginous taste. It is given to children in doses of fʒij. and to adults to the amount of fʒij. An overdose, or the drinking cold water during its use, produces vomiting, fever, and delirium; effects which are to be remedied by castor oil, warm water and acids. It is seldom employed in this country.

DECOCTUM GUIAIACI COMPOSITUM, Edin. *Compound Decoction of Guaiacum.*

“Take of Guaiacum Wood rasped, *three ounces*; Raisins, *two ounces*; Sassafras Root sliced, Liquorice Root bruised, of each, *one ounce*; Water, *ten pounds*. Boil the Guaiacum Wood and the Raisins in the Water over a gentle fire down to five pounds, adding the roots towards the end of the boiling; then strain.”

*Syn.* Decoction de Guajac composée (*F.*), Guajack-dekokte (*G.*), Decotto di Guajaco composito (*I.*).

This decoction derives less of its efficacy from the Guaiacum than is generally imagined, a small portion of extractive matter only being taken up by the water. It is, however, supposed to be useful in chronic rheumatism, some cutaneous diseases, and in syphilis during a mercurial course; but, probably, at best it is only serviceable as a demulcent. It may be taken in doses of a quarter of a pint, to the amount of Oj. or Oij. in the day.

DECOCTUM HORDEI, Lond. Dub. DECOCTUM HORDEI DISTICHI, Edin. *Decoction of Barley.*

“Take of Pearl Barley, *two ounces*; Water, *four pints and a half*, (*five pounds*, Edin.) First wash away any extraneous



substances that may adhere to the Barley ; then, having poured on it half a pint of Water, boil for a few minutes. This Water being thrown away, let the remainder be added boiling ; then boil down to two pints, and strain."

DECOCTUM HORDEI COMPOSITUM, Lond. *Compound Decoction of Barley.*

"Take of Decoction of Barley, *two pints* ; Figs sliced, *two ounces* ; Liquorice Root sliced and bruised, *half an ounce* ; Raisins stoned, *two ounces* ; Water, *a pint*. Boil down to two pints, and strain."

*Dublin.*

"Take of Decoction of Barley, *four pints* ; Raisins stoned, Figs sliced, of each, *two ounces* ; Liquorice Root sliced and bruised, *half an ounce*. During the boiling add first the Raisins, then the Figs, and lastly, the Liquorice Root a short time before it is finished ; when it is completed, the strained liquor ought to measure two pints."

*Syn.* Decoction d'Orge (*F.*), Gerstedekokte (*G.*), Decotto d'Orzo (*I.*).

The preparation of these decoctions is generally intrusted to nurses and the attendants of the sick-room ; but a practitioner ought not to be ignorant of the best manner of making them, as his directions may be occasionally necessary. They are elegant and useful demulcents in cases of fever, phthisis, gonorrhœa, and strangury, and indeed in all acute diseases, given *ad libitum*. A few drops of Tincture of Opium may be added to the compound decoction, to obviate its laxative effect, where this might prove hurtful. Equal parts of this decoction, and of decoction of Bark, form a useful gargle in cynanche maligna. The simple decoction mixed with an equal quantity of good milk and a small portion of sugar, is an excellent substitute for the breast milk, in those cases, in which infants are so unfortunate as to require being brought up with the spoon.

DECOCTUM LICHENIS,<sup>1</sup> Lond. DECOCTUM LICHENIS ISLANDICI, Edin. *Decoction of Liverwort.*

"Take of Liverwort, *an ounce* ; Water, *a pint and a half*, (*two pounds*, Edin.) Boil down to a pint, and strain."

DECOCTUM LICHENIS ISLANDICI, Dub. *Decoction of Iceland Liverwort.*

"Take of Iceland Liverwort, *half an ounce* ; Boiling Water, *a pint*. Digest for two hours ; then boil for a quarter of an hour, and strain the liquor while it is hot."

In these decoctions the bitter principle of the lichen is united

---

<sup>1</sup> The impropriety of using the generic name only of the plant is here very obvious, particularly as another species of this extensive family, the *Lichen roccella*, is now introduced into the Dublin Pharmacopœia.



with its fecula, which is thus rendered extremely nauseous; and although its operation in the stomach may be thus augmented, yet few patients will be persuaded to take it in this form. The dose is from  $\text{f}\text{ʒ}\text{iv.}$  to  $\text{f}\text{ʒ}\text{ij.}$  three times a day. We have already stated its use as a demulcent, when freed from the bitter, and the mode of preparing it. (See *Lichen*, Part ii.)

DECOCTUM MALVÆ COMPOSITUM, Lond. *Compound Decoction of Mallows.*<sup>1</sup>

“Take of Mallows dried, *an ounce*; Chamomile Flowers dried, *half an ounce*; Water, *a pint*. Boil for a quarter of an hour, and strain.”

This decoction is intended for fomentations and enemas, for which purposes it answers sufficiently well.

DECOCTUM PAPAVERIS, Lond. *Decoction of Poppy.*<sup>2</sup>

“Take of the Capsules of the White Poppy bruised, *four ounces*; Water, *four pints*. Boil for a quarter of an hour, and strain.”

In making this decoction, the seeds should not be rejected, as they contain a considerable portion of bland oil, which, added to the mucilage and narcotic principle of the capsule, increases the emollient quality of the decoction. It is a very useful fomentation in painful swellings, and in the excoriations produced by the thin acrid discharge of ulcers, and those common to infants.

DECOCTUM QUERCUS, Lond. DECOCTUM QUERCUS ROBORIS, Edin. *Decoction of Oak Bark.*

“Take of Oak Bark, *an ounce*; Water, *two pints*, (*two pounds and a half*, Edin.) Boil down to a pint, and strain.”

From Oak Bark thus treated the greater part of its astringent matter is extracted. The decoction is nearly inodorous, has a brown colour, and the austere taste of the bark. It reddens tincture of Litmus, and is precipitated by solutions of Isinglass, infusion of Yellow Cinchona bark, the carbonates of the alkalis, the aromatic spirit of Ammonia, Lime-water, and solutions of Sulphate of Iron, Acetate of Lead, Oxymuriate of Mercury, and Sulphate of Zinc, which are, therefore, incompatible in formulæ with it. The precipitates produced by the two last salts do not take place for a considerable time.

*Medical properties and uses.*—This is the usual form under which oak bark is exhibited. We have already noticed its internal use. (See *Quercus*, Part ii.) As a local astringent, it is used as a gargle in cynanche and relaxation of the uvula; as an injection, in passive uterine hæmorrhages, in leucorrhœa,

<sup>1</sup> Decoctum pro enemate, P. L. 1787.

<sup>2</sup> Decoctum pro fomento, P. L. 1787.



and the gleet discharge which often remains after miscarriages. It is also a useful wash in piles and procedentia recti.

**DECOCTUM SARSAPARILLÆ**, Lond.<sup>1</sup> *Decoction of Sarsaparilla.*

“Take of Sarsaparilla Root sliced, *four ounces*; Boiling Water, *four pints*. Macerate for four hours in a vessel lightly covered, and placed near the fire; then take out the Sarsaparilla, and bruise it. Return it again to the liquor, and macerate in a similar manner for two hours more; then boil it down to two pints, and strain.”

**DECOCTUM SMILACIS SARSAPARILLÆ**, Edin. *Decoction of Sarsaparilla.*

“Take of Sarsaparilla sliced, *six ounces*; Water, *eight pounds*. Digest for two hours in a temperature of about 195°, then take out the Root and bruise it: in this state put it again into the liquor, and boil it with a gentle fire down to four pounds; then express it, and strain.”

**DECOCTUM SARSAPARILLÆ**, Dub. *Decoction of Sarsaparilla.*

“Take of Sarsaparilla Root sliced, *an ounce and a half*; Boiling Water, *two pints*. Digest for two hours in a moderate heat; then take out the Sarsaparilla, and bruise it; return it to the liquor, and again digest for two hours; then boil down to one half, express, and strain the liquor through a linen cloth.”

*Syn.* Decoction de Sarsaparille (F.), Sarsaparille-dekokte (S.), Decotto di Sarsaparilla (I.).

We have already stated the claims which Sarsaparilla has to the attention of the practitioner as a remedy in syphilis. All the above formulæ display a great deficiency of enquiry, in those who introduced them into the Pharmacopœias: for, as the whole of the active matter of the root resides in the cortical part, and can be extracted from this by moderate coction, there is no necessity for the various macerations and boilings ordered by the Colleges; which, in fact, injure the remedy. The entire root, merely bruised, and macerated in water at 180° Fahr., will yield up all its medicinal properties.<sup>2</sup> This decoction may be regarded as useful during the exhibition of Mercury; and is found to be so in dysuria, and incon-

<sup>1</sup> The Lisbon Sarsaparilla has long been esteemed the best; but, from the experiments of the late Mr. Pope, (*See Trans. of the Medico Chirurg. Soc.*) it appears that the Red Sarsaparilla, which is an uncultivated species collected on the Spanish main, is superior both in the quantity of Extract which it yields and the medicinal virtues of that extract.

<sup>2</sup> The profession is indebted to Mr. Battley, of Fore Street, for his remarks on this subject. See *London Med. Repos.* vol. xi. 130: but more particularly to the late Mr. Pope, of Oxford Street. See *Trans. of the Medico-Chirurg. Soc. of London*, vol. xii. p. 344.



tinence of urine arising from a morbid irritability of the bladder. It affords precipitates with Lime-water, solution of Muriate of Barytes and of Acetate of Lead, which are therefore incompatible in formulæ with it.

**DECOCTUM SARSAPARILLÆ COMPOSITUM**, Lond. *Compound Decoction of Sarsaparilla.*

“Take of Decoction of Sarsaparilla boiling, *four pints*; Sassafras Root sliced, Guaiacum Root rasped, Liquorice Root bruised, of each, *an ounce*; Bark of Mezereon Root, *three drachms*. Boil for a quarter of an hour, and strain.”

*Dublin.*

“Take of Sarsaparilla Root sliced and bruised, *one ounce and a half*; Raspings of Guaiacum Wood, Bark of Sassafras Root, Liquorice Root bruised, of each, *two drachms*; Bark of Mezereon Root, *a drachm*; Boiling Water, *three pints*. Digest the Sarsaparilla, the Guaiacum, and the Sassafras, in the Water, with a moderate heat for six hours; then boil down to one-half, adding, towards the end of the coction, the Liquorice and the Mezereon; finally, strain.”

This decoction is an imitation of the once celebrated *Lisbon Diet-drink*. Its efficacy depends, chiefly, on the Mezereon root bark, the quantity of which, therefore, ordered by the Dublin College is undoubtedly too small. It operates as a diaphoretic and alterative, and is found to be useful in the treatment of secondary syphilis, chronic rheumatism, and in lepra, and some other cutaneous affections. The dose is from  $f\text{℥}iv.$  to  $f\text{℥}vj.$  taken three or four times a day.

**DECOCTUM SENEGÆ**, Lond. **DECOCTUM POLYGALÆ** SENEGÆ, Edin. *Decoction of Seneka.*

“Take of Seneka root, *an ounce*; Water, *two pints*. Boil down to a pint, and strain.”

*Syn.* Decoction de Polygale Senega (*F.*), Senegawurzel-dekokte (*G.*), Decotto di Poligala Senega (*I.*).

This decoction is of a brownish-olive colour, inodorous, and has a hot, pungent taste. Its virtues have been already discussed under the account of the root. (*Part ii.*) The dose is from  $f\text{℥}jss.$  to  $f\text{℥}iij.$  taken three or four times a day.

**DECOCTUM ULMI**, Lond. Dub. **DECOCTUM ULMI CAMPESTRIS**, Edin. *Decoction of Elm Bark.*

“Take of Fresh Elm Bark bruised, *four ounces*; Water, *four pints*, (*five pints*, Edin.) Boil to two pints, and strain.”

This decoction is thick, slightly mucilaginous, and of a brown colour; has a faint odour and a bitterish taste. Alcohol added to it produces a precipitate of light brown flakes; tinctures, therefore, in any considerable quantity, are inadmissible in formulæ with it. Its medicinal properties have



been already noticed. (See *Ulmus*, Part ii.) The dose is from fʒiv. to fʒvj. taken twice or three times a day.

DECOCTUM VERATRI, Lond. *Decoction of White Hellebore.*

“Take of white Hellebore Root bruised, *an ounce*; Water, *two pints*; Rectified Spirit, *two fluid ounces*. Boil the Hellebore Root with the water down to a pint, and strain; then, when the decoction is cold, add the spirit.”

This decoction is stimulant, acrid, and cathartic; but its operation is too violent for internal use. As a lotion it often proves beneficial in scabies, tinea capitis, and other cutaneous eruptions; but it requires to be used with caution even as an external remedy.

## EXTRACTA.

### EXTRACTS.<sup>1</sup>

THESE are preparations obtained by evaporating aqueous and alcoholic solutions of vegetable substances, until a mass of a somewhat firm, tenacious consistence remains. When water has been employed for making the solution, the Extract may consist of Gum or Mucilage, Albumen, Extractive, Tannin, Cinchonin, Saccharine matter, and the salts which the vegetable contained, and is termed a *Watery Extract*; but if alcohol have been the menstruum, Resin, Extractive, and all the above matters, except the gum, may be the ingredients, and the Extract is denominated a *Spirituos Extract*. The latter appellation also is used if proof spirit be employed. The proper menstruum, therefore, for the preparation of any extract must be that fluid which most readily dissolves the peculiar principles on which the medicinal efficacy of the vegetable is supposed to depend.

When water is employed, the substance to be subjected to its action should be in the dried state, and coarsely powdered; and the solution, whether made by decoction or infusion, should be evaporated immediately after it is strained, and whilst it is yet hot; for, as we observed in treating of Decoctions, water at the temperature of 212° takes up much more of the active matter of vegetables than it can hold in

<sup>1</sup> Chemists are much divided as to the nature of the substances to which the term *Extract* can be chemically applied; the term, therefore, as used in this work and in the Pharmacopœias, is confined solely to preparations obtained in the manner described under the title, without reference to their chemical properties.



solution at a lower temperature; therefore, by allowing them to cool, with the view of defæcation, and evaporating the clear fluid only, a considerable portion of the active matter does not enter into the extract, and is necessarily lost. In performing the evaporation, a higher temperature than that of boiling water must not be employed; but it must, nevertheless, be conducted as quickly as possible; and, therefore, the evaporating vessel should be broad and shallow, and set in boiling water; or the water-bath recommended by Dr. Powell<sup>1</sup> should be employed. (See *Instruments*, Part i.) A method of preparing extracts in vacuo has lately been introduced by Mr. Barry; and certainly, if the presence of air is likely to alter the properties of extracts, considerable advantages will accrue from Mr. Barry's mode of conducting the evaporation.<sup>2</sup>

Alcohol is used only in cases where the active ingredient of the vegetable is chiefly resin, or too volatile to bear the heat which is necessary for evaporating the water without being dissipated, or without suffering some decomposition, which would materially alter its properties. A tincture of the substance is first obtained, which is then evaporated by a very gentle heat in a water-bath; but the alcohol need not be allowed to evaporate in the air, as by employing a distilling apparatus, the greater part of it may be again obtained, either altogether free from any vegetable principle, or containing a small portion only of the more volatile; which renders it fitter for being again employed for the preparation of the same kind of extract.

Whether water, proof spirit, or pure alcohol be employed, the medicinal properties of the extract are always in some degree injured, the volatile parts are dissipated, and some of the fixed decomposed by the degree of heat required for the evaporation, particularly if water be the menstruum; or the proper extractive is oxidized, and consequently rendered inert. These are strong objections to this form of preparation; and, as Dr. Murray has properly observed, "with the exception of some of the pure bitters, as Gentian; or some of the saccharine vegetables, as Liquorice; there is no medicine perhaps but what may be given with more advantage under some other form."<sup>3</sup>

Extracts require to be kept in a hard and a soft state. A hard extract should be in such a state as to admit of its being easily pulverized, and the soft extract should be such as to

---

<sup>1</sup> *Translation of the London Pharmacopœia*, p. 201.

<sup>2</sup> For a description of the apparatus Mr. Barry employs, see *Journal of Science and the Arts*, vol. viii. p. 360.

<sup>3</sup> *System of Materia Medica*, &c. ii. 119.



retain the round form of a pill, without the addition of any powder. Both kinds should be preserved in pots which are so close that all the external air can be excluded; and these should be kept in a dry place. The soft should be wrapped in oiled bladder, and kept also in close-covered pots.

The London College does not arrange the Extracts under the titles *Watery* and *Spirituos*, which is the arrangement of the Edinburgh College, nor *Simple* and *Resinous*, which is that of the Dublin; it also classes the inspissated juices, which require no menstruum, as extracts: and the London Pharmacopœia being our text book, we have followed its arrangement. The following general directions are given by the LONDON COLLEGE for the preparation of Extracts.

“ In preparing all kinds of Extracts, evaporate the fluid as quickly as possible in a broad, shallow dish placed in a water-bath, until the extract acquire a consistence proper for forming pills, and, towards the end of the operation, stir assiduously with a spatula.

“ Sprinkle a small quantity of rectified spirit upon all the softer extracts.”

The EDINBURGH COLLEGE gives its general directions for the preparation of the *Extracts by Water*, under the Extract of Gentian; and for the *Extracts by Water and Alcohol*, under the Extract of Bark.<sup>1</sup>

---

<sup>1</sup> The following method of preparing narcotic extracts, invented by Mr. Battley, Chemist, Fore Street, London, has been found to answer every purpose in preserving the green colour and the medicinal properties of the plants.

“ Plants which from circumstances cannot be operated upon immediately after they are collected, must be revived by immersing their stalks in water for twelve or eighteen hours. Such as perfectly recover by this means, which will be known by the leaves becoming as fresh as when growing, are to be bruised and pressed; and the juice from them passed through a fine hair-sieve, and immediately placed on the fire. Some time before it is raised to the boiling temperature, a quantity of green-coloured matter begins to float on the surface of the fluid: in the juice of some plants this matter is very considerable: it is to be carefully removed by means of a thin perforated tin dish. By the time the liquor boils, or soon afterwards, the green matter ceases to appear. The boiling is to be continued until rather more than half the fluid have been evaporated, when the decoction is to be put into a conical pan, and suffered to stand in it until cold. A large precipitation of dark green-coloured feculent matter will then be found to have taken place; from which the supernatant fluid is to be poured off, and again exposed to evaporation, until half of it be consumed; when it is to stand for precipitation.

“ The precipitated matter from the second coction is by no means so green as the first. The remaining fluid is now to be suffered to boil till it acquire the consistence of syrup; when the matter which had been collected at the commencement by filtration and precipitation is to be mixed with it, and placed in a metallic pan in a water-bath, and further evaporated till of the consistence of an extract. In this latter part of the process, the operator must give his constant attention until it be quite completed. It is not necessary that the matter should be kept constantly stirred; but the operator ought never to suffer it to stick, or become hard on the sides of the pan; for, if it be allowed to harden, the extract loses its green colour; and in proportion to such loss is the deterioration of its medicinal virtues.”—*London Medical Repository*, vol. iv. p. 198.



EXTRACTA SIMPLICIORA, Dub. *Simple Extracts.*

“ All Simple Extracts, unless otherwise ordered, are to be prepared according to the following rule.

“ The vegetable matter is to be boiled in eight times its weight of Water, which is to be reduced by boiling to one-half; the liquor is then to be expressed, and after the fæces have subsided, to be filtered, and evaporated by the heat of boiling water, until it begin to thicken; and is to be, finally, inspissated by a medium heat, frequently stirring, until it acquire a consistence proper for forming pills.

“ All Extracts, when they begin to thicken, ought to be frequently stirred with a clean, iron spatula. They may be reduced to a proper degree of thickness by means of a stove heated for the purpose.

“ They ought to be preserved as much as possible from the contact of the air; and the softer are to be sprinkled with Rectified Spirit.” A direction which is useless, and consequently unnecessary.

EXTRACTUM CACUMINUM ABSYNTII, Dub.  
*Extract of Wormwood.*

*Syn.* Extrait d'Absynthe (*F.*), Wermuth-extrakt (*G.*), Estratto d'Assenzo (*I.*).

This is ordered to be prepared according to the above directions. It is nearly a simple bitter, the volatile oil being dissipated during the evaporation. It may be used in those cases for which bitters are commonly prescribed; but it is scarcely ever used. The dose is from gr. x. to ℥j. taken three times a-day.

EXTRACTUM ACONITI, Lond. *Extract of Aconite or Wolfsbane.*

“ Take of the Fresh Leaves of Aconite, *a pound*. Bruise them in a stone mortar, sprinkling over them a little water; then express the juice, and without any depuration, evaporate it to a proper consistence.”

SUCCUS SPISSATUS ACONITI NAPELLI, Edin. *Inspissated Juice of Aconite.*

“ Let the Fresh Leaves of Aconite be bruised; inclose them in a hempen bag, and press them strongly, until they yield their juice; which is to be evaporated in flat vessels, heated with boiling water saturated with Muriate of Soda (*common salt*), and immediately reduced to the consistence of thick honey.

“ After the mass is cold, let it be put into glazed earthen vessels, and moistened with alcohol.”

*Syn.* Extrait d'Aconit (*F.*), Eisenhütlein-extrakt (*G.*), Estratto d'Aconito Napello (*I.*).

Mr. Brande informs us that 1 cwt. of fresh Aconite yields about 5 lbs. of extract.<sup>1</sup> The activity of the plant is very un-



certain, and depends on soil and the nature of the seasons. Its active principle is an alkaloid, which has been named *Aconita*.

This extract, or inspissated juice, is the form under which Stoerk introduced Wolfsbane into practice. It has an obscure, brownish-red colour, a disagreeable odour, and an acrid, slightly styptic taste. Its medicinal properties are the same as those of the plant, but it is very seldom used. (See *Part ii.*) The dose at first should be gr.  $\frac{1}{2}$  only, and gradually increased to grs. vj. taken night and morning.

**EXTRACTUM ALOES PURIFICATUM**, Lond. *Extract of Aloes*.

“Take of Extract of spiked Aloes in powder, *a pound*; Boiling Water, *a gallon*. Macerate for three days in a gentle heat, then strain the solution, and set it aside that the dregs may subside. Pour off the clear liquor, and evaporate it to a proper consistence.”

*Syn.* Extrait d'Aloes (*F.*), Aloeextrakt (*G.*), Estratto d'Aloe (*I.*).

This Extract consists chiefly of the mucous and extractive matter of the Aloes; but, during the inspissation the extractive is partially oxidized and rendered less soluble, and, consequently, the extract is not completely soluble in water. It is employed in the same cases as the aloes, and is said to be less stimulant and griping; but it is at best an unnecessary preparation. The dose is from grs. x. to grs. xv. given in the form of pills.

**EXTRACTUM ANTHEMIDIS**<sup>1</sup>, Lond. **EXTRACTUM ANTHEMIDIS NOBILIS**, Edin. **EXTRACTUM FLORUM CHAMÆMELI**, Dub. *Extract of Chamomile Flowers*.

“Take of Chamomile Flowers dried, *a pound*; Water, *a gallon*. Boil down to four pints, and strain the liquor while it is hot; then evaporate it to a proper consistence.”

*Syn.* Extrait de Camomille romaine (*F.*), Kamillenextrakt (*G.*), Estratto di fiori di Camomillo (*I.*).

The Edinburgh Extract is to be prepared in the same manner as the Extract of Gentian of that College; the Dublin, after the manner directed for the preparation of the simple extracts.

In these processes the volatile oil is dissipated, and a simple bitter extract remains, possessing scarcely any of the properties of the plant, 1 cwt. of the flowers yield 48 lbs. of extract. It is of a deep-brown colour, and has a grateful bitter taste, but scarcely any odour. It has scarcely any efficacy when used alone; but is a useful adjunct to rhubarb, and sulphate of zinc as a stomachic pill. The dose may be from grs. x. to ℥j. given twice or thrice a-day.

<sup>1</sup> Extractum Chamæmeli, P. L. 1787.



EXTRACTUM BELLADONNÆ, Lond. SUCCUS SPIS-SATUS ATROPÆ BELLADONNÆ, Edin. *Extract of Belladonna.*

“Take of Fresh Leaves of Belladonna, *a pound*. Bruise them in a stone mortar, sprinkling a little water over them; then express the juice, and, without any separation of the sediment, evaporate it to a proper consistence.”

*Syn.* Extrait de Belladonne (*F.*), Belladonnaextrakt (*G.*), Estratto de l'erba di Belladonna (*I.*).

The inspissated juice of the Edinburgh Pharmacopœia is to be prepared in the same manner as the inspissated juice of Aconite.

This extract is inodorous, and has a bitterish taste. Its medicinal properties are the same as those of the plant, but weaker. They depend on an alkaloid, which has been named *Atropia*. We have witnessed great benefit from extract of Belladonna in whooping-cough, in combination with Subcarbonate of Soda; but it requires to be exhibited with great caution. A plaster composed of equal parts of the extract and carbonate of ammonia, or of soap plaster, is efficacious in relieving local pains. An ointment also made with equal proportions of the extract, and simple ointment is useful in allaying the pain of chordee, and the extract rubbed on the affected part has been found a useful means of procuring temporary relief from the pain of neuralgia. The dose is from gr. j. gradually increased to grs. v. given in the form of pills. This extract, when taken in very large doses, produces all the symptoms of a powerful narcotic poison. The most striking are dilatation and immobility of the pupils, confused vision or insensibility to the impression of external objects; the vessels of the conjunctiva appearing as if injected with blue blood; dryness of the lips, tongue, palate, and fauces; sometimes difficult deglutition, nausea, syncope, frequent flexion of the body forward, continual movements of the hands and fingers, lively delirium, with an idiotic smile, difficulty of articulation, and a kind of tenesmus. If these effects subside the healthy functions are restored, but every recollection of the previous state is lost. Sometimes, the patient sinks into a state of fatuity, livid spots appear on different parts of the body, the pulse sinks, the intestines become paralyzed, profuse colliquative sweats break forth, and death supervenes.

Dissections display signs of inflammation of the mucous coat throughout the digestive canal, sometimes proceeding to ulceration; the liver and lungs are more or less inflamed, and the latter gorged with black blood. The body rapidly passes into the putrefactive decomposition.

The remedy in poisoning by Belladonna or its extract, is, after removing the poison by the stomach syringe, affusions of



cold water on the head and body, acidulated drinks, and vomiting and purging.

EXTRACTUM CASCARILLÆ RESINOSUM, Dub.  
*Resinous Extract of Cascarilla.*

“Take of Cascarilla Bark in coarse powder, *a pound*; Rectified Spirit of Wine, *four pints*. Digest for four days, then pour off the coloured spirit, and filter. Boil what remains of the Cascarilla in ten pints of Water down to two pints; then evaporate the strained decoction, and at the same time distil the tincture from a retort, until both begin to thicken; then mix them together, and evaporate the mixture to a consistence proper for making pills. Lastly, mix the Extracts intimately together.”

*Syn.* Extrait de Cascarille (F.), Cascarillenextrakt (G.).

This preparation is expensive, and does not appear to possess any peculiar advantages to recommend it. The dose is from grs. x. to ℥j. given twice or thrice a-day, in the form of pills.

EXTRACTUM CINCHONÆ, Lond.<sup>1</sup> *Extract of Bark.*

“Take of Lance-leaved Cinchona Bark bruised, *a pound*; Water, *a gallon*. Boil down to six pints, and strain the liquor while it is warm. In the same manner boil it down again four successive times, in an equal quantity of Water, and strain. Finally, mix the solutions together, and evaporate the mixture to a proper consistence.

“This Extract ought to be kept in a *soft* state fit for making pills, and in a *hard* state that it may be reduced to powder.”

*Dublin.*

“Take of Cinchona Bark in coarse powder, *a pound*; Water, *six pounds*. Boil for a quarter of an hour in a vessel nearly covered; then filter the decoction while it is yet hot, and set it aside. Boil the residue again in the same quantity of Water, and filter it in the same manner: repeat this a third time; and finally, mix all the liquors, and evaporate the mixture to a proper consistence.

“This extract should be kept in two states: one *soft*, fit for making pills; and the other *hard*, or in a state proper to be reduced to powder.”

*Syn.* Extrait de Quinquina (F.), Wäss rigtes Chinaextrakt (G.), Estratto di China aquosa (I.).

The operation of the same causes as those which we stated to be unfavourable to decoction, as a form of preparation for the exhibition of Cinchona, are still more hurtful to its efficacy in the form of extract; and, according to Sir John Pringle,

<sup>1</sup> Extractum Corticis Peruvianæ, P. L. 1754.



the extract is less efficacious, even in equal quantities, than the simple powder. This extract, however, certainly contains Cinchonia, and is not devoid of utility: it often sits very lightly on the stomach when the powder is rejected. It is usually ordered in doses of from grs. x. to ʒss. dissolved in any distilled water; but it is necessary to observe, that owing to the oxidizement of the extractive matter, the solubility of the extract is diminished during its formation: scarcely more than one-half is soluble in water. It has a very bitter taste, but is less austere than the bark.

EXTRACTUM CINCHONÆ RESINOSUM, Lond. *Resinous Extract of Bark.*

“Take of Lance-leaved Cinchona bark, bruised, *a pound*; Rectified Spirit, *four pints*. Macerate for four days, and strain. Distil the tincture in a water-bath, until the extract has acquired a due consistence.”

EXTRACTUM CINCHONÆ LANCIFOLIÆ, Edin. *Extract of officinal Cinchona Bark.*

“Take of Lance-leaved Cinchona bark in powder, *one pound*; Alcohol, *four pounds*. Digest for four days, and pour off the tincture. Boil the residue in five pounds of Distilled Water for fifteen minutes, and strain the decoction while it is hot through a linen cloth. Repeat this coction with an equal quantity of Distilled Water, strain again, and evaporate the liquor to the consistence of thin honey. Distil the Alcohol from the tincture until it be reduced to a similar consistence. Then mix the inspissated liquors, and evaporate them to a proper consistence in a bath of boiling water, saturated with Muriate of Soda.”

EXTRACTUM CINCHONÆ RUBRÆ RESINOSUM, Dub. *Resinous Extract of Red Cinchona Bark.*

This is ordered to be prepared in the same manner as the resinous extract of Cascarella.

One cwt. of fine Bark, yields 25 lbs. of extract. <sup>1</sup>

*Syn.* Extrait resineux de Quinquina (*F.*), Estratto di China collaresina (*I.*).

The Extract prepared by these processes has the bitter, austere taste of the bark, which it nearly equals in efficacy, and is more grateful to the stomach. It is altogether a preferable preparation to the watery extract; for, by the separate action of the spirit and the water, all the soluble and active principles of the drug are taken up: less heat is required to evaporate the menstruum; and, owing to the presence of the Alcohol, the extractive matter absorbs less oxygen: indeed, the expense of the spirit, of which there is always some waste, is

<sup>1</sup> *Brande's Manual*, p. 391.



the only objection to its general use. The dose is from grs. x. to grs. xxx. formed into pills.

EXTRACTUM COLOCYNTHIDIS, Lond. *Extract of Colocynth.*

“Take of the Pulp of Colocynth, *a pound*; Water, *a gallon*. Boil down to four pints, and strain the liquor while it is hot; then evaporate it to a proper consistence.”

*Syn.* Koloquinten-extrakt (G.).

This Extract is a milder but less powerful cathartic than the pulp from which it is prepared, and with the addition of Calomel forms an excellent purgative pill, which operates without griping. It has, however, the disadvantage of becoming extremely tough, when kept. From grs. v. to ʒss. is the usual dose.

EXTRACTUM COLOCYNTHIDIS COMPOSITUM<sup>1</sup>, Lond. *Compound Extract of Colocynth.*

“Take of Colocynth Pulp, sliced, *six ounces*; Extract of the Spiked Aloe, powdered, *twelve ounces*; Scammony, powdered, *four ounces*; Cardamom Seeds, powdered, *one ounce*; Hard Soap, *three ounces*; Proof Spirit, *one gallon*. Macerate the Colocynth Pulp in the spirit, with a gentle heat, for four days. Strain the liquor, and add to it the Aloes, the Scammony, and the Soap; then evaporate it to a proper consistence, and towards the end of the inspissation mix in the Cardamom seeds.”

*Dublin.*

“Take of the Pulp of Colocynth, cut small, *six drachms*; Hepatic Aloes, *an ounce and a half*; Scammony, *half an ounce*; Lesser Cardamom Seeds, husked, *a drachm*; Castile-soap softened with water, so as to have a gelatinous appearance, *three drachms*; Hot Water, *a pint*. Digest the Colocynth in the Water in a covered vessel, with a medium heat, for four days; express and strain the liquor, and add to it the Aloes and Scammony, first separately reduced to powder; then evaporate the mixture with a medium heat to a proper consistence for making pills, and towards the end of the inspissation add the gelatinized Soap and the powdered Seeds, and with frequent stirring mix the whole intimately together.”

By this combination of powerful cathartic substances a purgative mass is obtained, more manageable and less irritating than any of its components separately taken. It forms a very useful pill for relieving the habitual costiveness of leucophlegmatic habits; and in obstinate visceral obstructions when combined with Calomel, which is not decomposed, as might

<sup>1</sup> Extractum catharticum, P. L. 1745.



*à priori* be supposed. The dose is from gr. vj. to ʒss. repeated every eight hours until it operate.<sup>1</sup>

EXTRACTUM CONII, Lond. SUCCUS SPISSATUS CONII MACULATI, Edin. *Extract of Hemlock.*

“Take of Fresh Hemlock, *a pound*. Bruise it in a stone mortar, sprinkling over it a little Water; then express the juice, and without separating the sediment, evaporate it to a proper consistence.”

The Edinburgh preparation is to be made according to the directions ordered for the preparation of inspissated juices.

SUCCUS SPISSATUS CICUTÆ, Dub. *Inspissated Juice of Hemlock.*

“Express Hemlock Leaves, gathered when the flowers are about to appear, and allow the juice to remain six hours to deposit the fæces; then evaporate the pure juice to a proper consistence with a moderate heat.”

*Syn.* Extrait de Ciqué (*F.*), Schierlings-extrakt (*G.*), Estratto del' erba della Cicuta (*I.*).

This extract, or inspissated juice, should be of a clear olive colour, have a foetid odour, and a bitterish saline taste. One cwt. of Hemlock leaves, yield from three to five lbs. of Extract.<sup>2</sup> Although it be the form in which Stoerk introduced Hemlock into practice, yet the narcotic power of the remedy is always impaired by this mode of preparation, and it is still more weakened by keeping, being nearly lost when a saline efflorescence begins to appear on the surface of the extract. It also varies in its power according to the soil where the plant grows, and the seasons. It is used in the same cases as the powder, with which it is frequently mixed when it is to be made into pills: and is an useful adjunct to mercurials in cutaneous affections. Bergius recommends it in impotency.<sup>3</sup> The dose is gr. iij. gradually increased to ʒj. given twice or thrice a-day.

EXTRACTUM ELATERII, Lond. *Extract of Elaterium.*

“Slice ripe Wild Cucumbers, express the juice very gently, and pass it through a very fine hair-sieve into a glass vessel; then set it aside for some hours until the thicker part has subsided. Reject the thinner, supernatant part, and dry the thicker part with a gentle heat.”

*Syn.* Elaterium (*F.*), Estratto del frutto della Momordica (*I.*).

<sup>1</sup> BARCLAY'S ANTIBILIOUS PILLS are composed of *ext. of colocynth ʒij.*, *ext. of jalap ʒj.*, *almond soap ʒiss.*, *guaiaic ʒij.*, *tartarized antimony gr. viij.*, *oils of juniper, of carraway and of rosemary, of each four drops*; *syrup of buckthorn*, sufficient to form a mass, which is to be divided into sixty-four pills.

<sup>2</sup> *Brande's Manuel*, p. 394.

<sup>3</sup> “Impotentiam virilem sub usu Conii curatam observavi, in viro quodam plusquam quadragenario, qui omnem erectionem penis perdiderat, postinde tamen plures liberos procreavit.”—*Bergius Mat. Med.* i. 195.



ELATERIUM, Dub. *Elaterium*.

“Slice ripe Wild Cucumbers, and strain the juice very lightly, expressed through a fine hair-sieve into a glass vessel; then set it aside for some hours until the thicker part subside; reject the supernatant liquor, and dry the fecula, laid upon a linen cloth and covered with another, by a medium heat.”

The substance obtained by these processes is neither an extract nor an inspissated juice, but a peculiar modification of fecula combined with a very active principle which is deposited with it; and which has been named *Elatin* by Dr. Paris. It is contained in the juice which surrounds the seeds only; and subsides from this juice obtained without pressure. From Dr. Clutterbuck's experiments<sup>1</sup>, the quantity of elaterium in the fruit appears to be so small, that he obtained six grains of it only from forty cucumbers. Dr. Paris found that ten grains of the best elaterium, as it is found in the shops, contain one grain only of *Elatin*; and in general it is adulterated with starch, on which account we scarcely ever obtain two samples of it of the same strength. When good, it is of a greenish-grey colour, has a bitter taste, is light and pulverulent. The name adopted by the Dublin College is more appropriate than that imposed by the London College. It is very remarkable that the Edinburgh College has rejected so important a remedy from the last edition of its Pharmacopœia.

*Medical properties and uses.* — Elaterium is a very powerful hydragogue, and excites sickness, severe vomiting, and hypercatharsis, if it be not cautiously administered. On this account it is seldom used as a cathartic; but in ascites it often produces the entire evacuation of the fluid, when gamboge and crystals of tartar, foxglove, and every other remedy have failed. The best mode of administering it is to give it in divided doses of gr.  $\frac{1}{8}$  each, every fourth hour, until it begin to operate.

EXTRACTUM CACUMINUM GENISTÆ, Dub.  
*Extract of Broom Tops.*

This Extract is to be prepared in the same manner as the Extract of Wormwood. It is said to be diuretic, but its efficacy is doubtful, and it is scarcely ever employed. The dose is from ʒss. to ʒj. or more.

EXTRACTUM GENTIANÆ, Lond. EXTRACTUM RADICIS GENTIANÆ, Dub. *Extract of Gentian*.

“Take of Gentian root, sliced, *a pound*; Boiling Water, *a gallon*. Macerate for twenty-four hours; then boil down to four pints, strain the liquor while it is hot, and evaporate it to a proper consistence.”

EXTRACTUM GENTIANÆ LUTEÆ, Edin. *Extract of Gentian*.

<sup>1</sup> *London Med. Repos.* vol. xii.



“Take of Gentian Root, *any quantity*. Having sliced and bruised it, pour upon it eight times its weight of Boiling Water. Boil down to one half, express the liquor strongly, and strain it. Evaporate the decoction immediately to the consistence of thick honey, in a bath of boiling water saturated with Muriate of Soda.”

*Syn.* Extrait de Gentiane (F.), Enzian-extrakt (G.), Estratto di Gentiana (I.).

The bitter principle of Gentian Root is not injured by this form of preparation. The extract is inodorous, very bitter, black, shining, and tenacious. It is chiefly used as a vehicle for the exhibition of the metallic oxides. The dose is from grs. x. to ʒss. given twice or thrice a day.

EXTRACTUM GLYCYRRHIZÆ, Lond. Dub. *Extract of Liquorice.*

“Take of Liquorice Root, sliced, *a pound*; Boiling Water, *a gallon*. Macerate for twenty-four hours; then boil down to four pints; strain the hot solution, and evaporate it to a proper consistence.”

There is scarcely any of this extract prepared by the apothecary; the pure extract of Liquorice sold in the shops under the name of *Refined Liquorice* being prepared from the impure extract of commerce, by dissolving it in water, straining and inspissating it in the usual manner. It is an useful demulcent for allaying tickling cough, as from its tenacity it hangs about and sheathes the fauces.

Officinal preparations. *Pilulæ opiatæ*, E. *Pilulæ scilliticæ*, E. *Trochisci Glycyrrhizæ glabræ*, E. *Trochisci Glycyrrhizæ cum Opio*, E.

EXTRACTUM HÆMATOXYLI<sup>1</sup>, Lond. EXTRACTUM HÆMATOXYLI CAMPECHIANI, Edin. EXTRACTUM SCOBIS HÆMATOXYLI, Dub. *Extract of Logwood.*

“Take of Logwood rasped, *a pound*; Boiling Water, *a gallon*. Macerate for twenty-four hours; then boil down to four pints; strain the hot liquor, and evaporate to a proper consistence.

*Syn.* Campecheholz-extrakt (G.).

One cwt. of the wood yields 20 lbs. of the extract.<sup>2</sup> This extract is almost inodorous, has a sweet austere taste, and a deep ruby colour. It becomes extremely brittle when kept.

It is regarded as useful in the protracted stage of diarrhœa and dysentery. The dose is from grs. x. to ʒss. dissolved in cinnamon water or peppermint water.

EXTRACTUM RADICIS HELLEBORI NIGRI, Edin. Dub. *Extract of Black Hellebore Root.*

<sup>1</sup> Extractum Ligni Campachensis, P. L. 1745.

<sup>2</sup> Brande's Manuel, p. 397.



*Syn.* Extrait d'Ellebore (*F.*), Schwarz Niesenwurz-extrakt (*G.*).

This is to be prepared from the bruised root, after the manner directed for the extract of Gentian by the Edinburgh College, and the extract of Wormwood by the Dublin College. The effects of this extract are those of a drastic purgative, and the emmenagogue power which has been attributed to it, seems to depend on its action as a purgative. It has been advantageously given in chronic diseases of the skin.

**EXTRACTUM HU'MULI**, Lond. *Extract of Hops.*

“Take of the Strobiles of the Hop, *four ounces*; Water, *a gallon*. Boil down to four pints; strain the hot liquor; and evaporate it to a proper consistence.”

One cwt. of the strobules yields about 40 lbs. of the extract.<sup>1</sup> This extract is inodorous; and has the bitter taste peculiar to the Hop. Its virtues depend on Lupulin. We have found it an useful anodyne in gout, acute rheumatism, and cases which do not admit of the use of opium. The dose is from grs. v. to ℥j. given in the form of pills, or dissolved in any aqueous vehicle.

**EXTRACTUM HYOSCY'AMI**, Lond. **SUCCUS SPIS-SATUS HYOSCYAMI NIGRI**, Edin. **SUCCUS SPISSATUS HYOSCYAMI**, Dub. *Extract of Henbane.*

“Take of fresh Leaves of Henbane, *a pound*. Bruise them in a stone-mortar, sprinkling on them a little Water; then press out the juice, and without separating the sediment, evaporate it to a proper consistence.”

*Syn.* Extrait de Jusquiame (*F.*), Hyoszyamus-extrakt (*G.*), Estratta di Giusquiama nera (*I.*).

This Extract has a disagreeable slightly foetid odour, and a nauseous, bitterish, subsaline taste. It is possessed of considerable narcotic powers, and is used as a substitute for Opium in nervous affections, mania, gout, rheumatism, and all painful complaints, in which it is wished to avoid the costiveness which Opium is apt to induce. A solution of it in water, in the proportion of one drachm to the ounce, dropped into the eye, dilates very much the pupil; and has been used, on the recommendation of Professor Himly, for facilitating the operation for cataract; and also in contracted pupils not accompanied by adhesion of the iris to the capsule.<sup>2</sup> The dose is from grs. iij. to ℥j. given in the form of pills.

**EXTRACTUM JALA'PÆ**<sup>3</sup>, Lond. *Extract of Jalap.*

“Take of Jalap Root powdered, *a pound*; Rectified Spirit, *four pints*; Water, *one gallon*. Macerate the Jalap root in the

<sup>1</sup> Brande's Manuel, p. 398.

<sup>2</sup> Edin. Medical and Surg. Journal, vol. ix. p. 6. 11.

<sup>3</sup> Extractum Jalapii, P. L. 1745.



spirit for four days, and decant the tincture. Boil the residue in the water down to two pints. Then strain separately the tincture and the decoction; distil the former, and evaporate the latter, until both begin to thicken. Lastly, mix the extract with the resin, and evaporate the mixture to a proper consistence."

"This extract should be kept in a *soft* state, fit for forming pills, and in a *hard* state, so that it may be reduced to powder."

EXTRACTUM CONVULVULÆ JALAPÆ, Edin. *Extract of Jalap.*

This is ordered to be prepared from the root, in the same manner as the extract of cinchona bark. (*Edin.*)

EXTRACTUM JALAPÆ, Dub. *Extract of Jalap.*

"Let it be prepared in the same manner as the resinous extract of Cascarilla."

*Syn.* Extrait de Jalap (*F.*).

One cwt. of Jalap yields about 50 lbs. of watery extract, and fifteen of resinous extract.<sup>1</sup> These extracts contain all the active principles of the Jalap root. They are, however, apt to gripe during their operation: hence, particularly when given to children, they should be triturated with sugar and almonds, or mucilage, so as to form an emulsion, in which state they operate freely and without griping. The dose to an adult is from grs. x. to ℥j.

EXTRACTUM RADICIS JALAPÆ, Dub. *Extract of Jalap Root.*

This is to be prepared with water alone, after the manner directed for the preparation of the simple extracts. (*Dub.*) It contains chiefly the gummy part of the Jalap, very little of the resin being taken up by the water. It is milder in its operation than the root, and may be given to infants, in doses of from grs. vi. to grs. xij. triturated with sugar or testaceous powders.

EXTRACTUM LACTUCÆ, Lond. *Extract of Lettuce.*

"Take of Fresh Lettuce leaves, *one pound*; bruise them in a stone-mortar, sprinkling a little water over them; then express the juice, and evaporate it unstrained, until it acquire a proper consistence."

SUCCUS SPISSATUS LACTUCÆ SATIVÆ, Edin.

*Inspissated Juice of Garden Lettuce.*

*Syn.* Extrait de Laitue (*F.*).

This is to be prepared from the fresh leaves, according to the general directions for preparing inspissated juices.

SUCCUS SPISSATUS LACTUCÆ VIROSÆ, Edin.

*Inspissated Juice of the Wild Lettuce.*

*Syn.* Extrait de Laitue vireuse (*F.*), Estratto dell' erba della Lactuca (*I.*).

<sup>1</sup> Brande's Manuel, p. 400.



To be prepared from the fresh herb, in the same manner as the other inspissated juices.

One cwt. of lettuce yields between 3 and 4lbs. of the extract.<sup>1</sup>

The extracts of both the above species of *Lactuca* are exhibited as substitutes for Opium, in cases in which the intention is rather to allay irritation than to produce the full effect of a narcotic. The dose is grs. vj. gradually increased.

EXTRACTUM OPII, Lond. *Extract of Opium.*

“Take of Opium sliced, *sixteen ounces*; Water, *one gallon*. Pour a small portion of the water upon the Opium, and macerate for twelve hours, that it may become soft; then adding gradually the remaining water, rub them together till they be well mixed, and set the mixture apart that the fæculencies may subside. Lastly, strain the liquor, and evaporate it to a proper consistence.”

EXTRACTUM OPII AQUOSUM, Dub. *Watery Extract of Opium.*

“Take of Opium, *two ounces*; Boiling Water, *a pint*. Rub the Opium in the water for ten minutes, and after a little pour off the solution; rub the residuary opium in an equal quantity of boiling water for the same space of time, pouring off also this solution: and repeat the operation a third time. Mix together the decanted solutions, and expose the mixture in a broad open vessel to the air for two days. Lastly, strain it through linen, and by slow evaporation form it into an extract.”

*Syn.* Extrait d'Opium (*F.*), Opiums extrakt (*G.*), Estratto d'Oppio (*I.*).

Water takes up a certain proportion of all the constituents of crude Opium, but less of the resinous than of the gummy part; and the watery solution contains more of morphia, on which depends the remedial quality of Opium.<sup>2</sup> In the Dublin preparation the quantity of active matter must necessarily be greater, owing to the employment of boiling water for the second and third triturations. This extract, therefore, differs very little from Opium; but as the inspissation cannot always be conducted exactly in the same manner, its strength must consequently vary.

*Qualities.*—This extract is inodorus, has a bitter taste,

<sup>1</sup> Brande's Manuel.

<sup>2</sup> This extract, however, contains some of Derosnes salt also, or *Narcotine*, as it has been lately termed; and this is supposed to produce that excitement, which even the aqueous extract occasions previous to its sedative effect. M. Robiquet (*Journ. de Pharm.* May 1821,) proposes to free it of this principle, by agitating the extract as soon as it acquires the consistence of syrup with ether; and repeating this agitation with fresh portions of ether, as long as the ether on distillation deposits any crystals of *Narcotine*. The extract thus prepared contains only morphia, gum, and extractive.



and is of a very deep brown colour. It is not altogether soluble in water, but is not precipitated from its solution by alcohol. It, however, affords precipitates with the following substances, which ought not, therefore, to enter into prescriptions with its solution; viz. solutions of astringent vegetables, the alkaline carbonates, corrosive Muriate of Mercury, Sulphate of Copper, Sulphate of Zinc, Acetate of Lead, and Nitrate of Silver.

*Medical properties and uses.* — This extract is supposed to produce the effects of Opium, but with less subsequent derangement of the nervous system. It is therefore supposed to be well adapted for the diseases of children and very irritable habits. The dose is from gr. j. to grs. vj. for an adult.

Officinal preparation. *Syrupus Opii*, D.

EXTRACTUM PAPAVERIS, Lond. EXTRACTUM PAPAVERIS SOMNIFERI, Edin. *Extract of Poppies.*

“Take of the Capsules of the Poppy, freed from the seeds, and bruised, *a pound*; Boiling Water, *a gallon*. Macerate for twenty-four hours; then boil down to four pints; strain the hot liquor, and evaporate it to a proper consistence.”

*Syn.* Extrait de Pavot (F.).

This Extract possesses nearly the same medicinal properties as Opium, but in a much weaker degree; and is less apt to occasion the nausea, head-ach, and delirium, which Opium not unfrequently produces. It is, therefore, to be preferred for procuring sleep in diseases in which the head is much affected. The dose is from grs. ij. to ʒj. given in the form of pills.

EXTRACTUM CORTICIS QUERCUS, Dub. *Extract of Oak Bark.*

*Syn.* Estratto della Quercia (I.).

This Extract consists principally of tannin, which is, therefore, not liable to be injured by this form of preparation; but it possesses no peculiar advantages to recommend it.

EXTRACTUM RHEI, Lond. *Extract of Rhubarb.*

“Take of Rhubarb Root in powder, *a pound*; Proof Spirit, *a pint*; Water, *seven pints*. Macerate for four days in a gentle heat, then strain the solution, and set it apart that the fæculencies may subside. Pour off the clear liquor, and evaporate it to a proper consistence.”

*Syn.* Extrait de Rhubarbe (F.), Rhabarber-extrakt (G.).

Although the purgative properties of the Rhubarb be obtained to a certain degree in this extract, yet its virtues are certainly impaired during the inspissation; and the simple infusion is in every respect a preferable form of preparation. The dose is from grs. x. to ʒss. given in the form of pills.



EXTRACTUM RUTÆ GRAVEOLENTIS, Edin.  
EXTRACTUM FOLIORUM RUTÆ, Dub. *Extract of Rue.*

The Edinburgh extract is to be prepared in the same manner as the extract of Gentian; the Dublin after the manner of the simple extracts. Prepared by either process, this extract is inodorous, and has a bitter, acrid taste. Its medicinal properties are different from those of the plant, the stimulant and narcotic powers of which depend on the volatile oil it contains, which is dissipated during the inspissation of the extract. The dose is from grs. x. to ℥j. in pills.

EXTRACTUM FOLIORUM SABINÆ, Dub. *Extract of Savine.*

To be prepared in the same manner as the simple extracts. It is a simple bitter of little efficacy, for the acrid volatile oil on which the efficacy of savine depends is dissipated by the heat employed during the inspissation. The dose is from grs. x. to 3ss. in pills.

SUCCUS SPISSATUS SAMBUCI NIGRÆ, Edin.  
*The inspissated Juice of the Black Elder.*

“Take of the ripe berries of the Black Elder, *five parts*; purified Sugar, *one part*. Boil with a gentle heat to the consistence of thick honey.

*Syn.* Eingedicker Fliedersaft (G.).

We are perfectly unacquainted with the use to which this extract can be put, as a remedial agent.

EXTRACTUM SARSAPARILLÆ, Lond. *Extract of Sarsaparilla.*

“Take of Sarsaparilla Root, sliced, *a pound*; Boiling Water, *a gallon*. Macerate for twenty-four hours, then boil down to four pints; strain the solution while it is hot, and evaporate it to a proper consistence.”

The experiments of the late Mr. Pope, Mr. Battley, and M. Robinet have demonstrated that the Red, or, as it is termed, Jamaica Sarsaparilla yields a much larger quantity of Extract than any of the other species of Sarsaparilla. It, also, appears from Mr. Pope's experiments, “that by submitting the root, cut transversely, to the action of steam or of distilled water, at a temperature somewhat below boiling, an elegant, soluble Extract may be obtained, containing all the virtues of the plant, not liable to decomposition, and applicable to the various purposes of extemporaneous prescription;” whilst by the method, ordered in the above formula of the London College, an insoluble, inefficacious extract only is obtained.



When properly prepared, without much heat, this extract possesses all the medicinal virtues of the root; but prepared according to the above formula, it has nothing to recommend it to practice. The dose is from grs. ix. to ʒj. dissolved in the decoction, or given in the form of pills.

**EXTRACTUM STRAMONII**, Lond. *Extract of Thorn Apple.*

“Take of Thorn Apple seeds, *one pound*; Boiling Water, *one gallon*. Macerate for four hours in a covered vessel near the fire; then take out the seeds; bruise them in a stone-mortar, and put them again into the liquor. Finally evaporate it until it acquire a proper consistence.”

The medicinal powers of this extract are less to be depended on than those of a tincture prepared with ʒij. of the herb, and fʒxvj. of proof spirit. The dose of the extract is ʒss. to ʒij. in the form of pills.

**EXTRACTUM TARAXACI**, Lond. **EXTRACTUM HERBÆ ET RADICIS TARAXACI**, Dub. *Extract of Dandelion.*

“Take of fresh Dandelion Root, bruised, *a pound*; Boiling Water, *a gallon*. Macerate for twenty-four hours; then boil down to four pints, strain the hot liquor and evaporate it to a proper consistence.”

*Syn.* Lowenzahn-extrakt (G), Estratto di Tarassaco (I.).

The medicinal powers of Dandelion have been already noticed. (See *Part ii.*) Dr. Pemberton affirms, that he has seen great advantage result from the use of this extract in doses of ʒss. in chronic inflammation and incipient scirrhus of the liver, and in chronic derangement of the stomach.<sup>1</sup> The usual dose is from grs. x. to ʒj. united with Sulphate of Potass.

**EXTRACTUM VALERIANÆ**, Dub. *Extract of Valerian.*

“Take of Valerian Root in coarse powder, *six ounces*; Boiling Water, *six pints*. Mix and digest for twenty-four hours with a moderate heat in a covered vessel; then express the liquor, and reduce it by evaporation to a proper consistence.”

*Syn.* Baldrian-extrakt (G.).

The odour of the plant is almost entirely dissipated in preparing this extract; and if the efficacy of the remedy be connected with that quality, which is extremely probable, it must be much inferior to the infusion, or the tincture. The usual dose is from grs. x. to ʒj. given in the form of pills.

<sup>1</sup> *On Diseases of the Abdominal Viscera*, p. 43.



## MISTURÆ.

## MIXTURES.

THE term Mixture in pharmaceutical language denotes a mingled compound, in which different ingredients are held suspended in a fluid medium by means of mucilaginous or of saccharine matter. The London College has placed under this title those medicines, also, which consist of the fixed oil of seeds, diffused through water by means of the mucilage, fecula, or saccharine matter of the seeds, and which are denominated *Emulsions*. Both these kinds of preparations should always be extemporaneous; and in prescribing them attention is required not to bring together incompatible substances, nor to order in mixtures insoluble matters of a specific gravity too great to be suspended, in the fluid vehicle, by the ordinary means.

---

MISTURA AMMONIACI, Lond. *Mixture of Ammoniacum.*

“Take of Ammoniacum, *two drachms*; Water, *half a pint*. Triturate the Ammoniacum, gradually adding the water until they be thoroughly mixed.”

LAC AMMONIACI, Dub. *Milk of Ammoniacum.*

“Take of Gum Ammoniacum, *a drachm*; Pennyroyal Water, *eight fluid ounces*. Triturate the Gum, gradually adding the Pennyroyal Water, until the mixture acquire the appearance of milk, which is to be strained through linen.”

The resinous part of the Ammoniacum is suspended in the water by means of the gummy part; but after a little time the greater portion of the resin subsides. It is coagulated by distilled vinegar, the Oxydels, Ether, Spirit of Nitric Ether, Supertartrate of Potass, and Oxymuriate of Mercury, which are therefore incompatible in prescriptions with mixture of Ammoniacum. It is advantageously employed as an expectorant in doses of from fʒss. to fʒj. combined with an equal quantity of Almond mixture.

MISTURA AMYGDALARUM, Lond. *Almond Mixture*

“Take of Almond Confection, *two ounces*; Distilled Water, *a pint*. Add the Water gradually to the Almond Confection whilst triturating, and then strain.”

EMULSIO AMYGDALI COMMUNIS, Edin. *Almond Emulsion.*

“Take of Sweet Almonds, *an ounce*; Refined Sugar, *half an ounce*; Water, *two pounds and a half*. Beat diligently the



Blanched Almonds in a stone-mortar, adding the Water gradually; then strain."

LAC AMYGDALÆ, Dub. *Almond Milk.*

"Take of Sweet Almonds, blanched, *an ounce and a half*; Purified Sugar, *half an ounce*; Water, *two pints and a half*. Rub the Almonds with the sugar, adding the Water gradually; then strain."

EMULSIO ACACIÆ ARABICÆ, Edin. *Emulsion of Gum Arabic.*

"Take of Mucilage of Gum Arabic, *two ounces*; Almonds, *an ounce*; Refined Sugar, *half an ounce*; Water, *two pounds and a half*. Blanch the Almonds, and then beat them in a stone-mortar with the Sugar and the Mucilage, gradually adding the Water; then strain through linen."

EMULSIO ARABICA, Dub. *Arabic Emulsion.*

"Take of Gum Arabic in powder, *two drachms*; Sweet Almonds blanched, Purified Sugar, of each *half an ounce*; Decoction of Barley, *a pint*. Dissolve the Gum in the warm decoction, and when it is almost cold, pour it gradually upon the Almonds previously beaten to a paste with the sugar, triturating at the same time so as to form a milky mixture; then strain."

*Syn.* Emulsion d'Amandes (*F.*), Mandelmilch (*G.*), Latte di Mandorle (*I.*).

In these preparations the oil of Almonds is diffused through the water, and suspended in it by the mucilage and fecula which the Almonds contain; the gum in the two latter preparations contributing nothing to this effect. The confection ordered by the London College affords an expeditious mode of making the mixture, but does not prevent the necessity of straining. The use of distilled water is an unnecessary refinement. Bitter Almonds are sometimes employed instead of sweet; and as the Prussic Acid is contained in the bitter Almond, and this is a direct sedative, the Emulsion of the bitter Almond is certainly a better vehicle for squills and other expectorant remedies than that of the sweet Almond.

*Qualities.* — These emulsions are inodorous, bland, milky fluids. The oil, after some time, rises like a thick cream to the surface; and in forty-eight hours the acetous fermentation commences, and the mixtures become sour. They are decomposed by Acids, Oxymel and Syrup of Squills, Spirits, and Tinctures, (unless these be in small quantity,) Tartrate and Supertartrate of Potass, Supersulphate of Potass, Nitrate of Potass, Oxymuriate of Mercury, Acetate of Lead, and Spirit of Nitric Ether, which are therefore incompatible in prescriptions with Almond emulsions.

*Medical properties and uses.* — These mixtures are in common use as diluents and demulcents in inflammatory fevers,



strangury, dysury and other affections of the urinary organs; but they are chiefly useful as pleasant vehicles for the exhibition of more active remedies. The dose is from  $f\text{℥}ij.$  to  $0ss.$  frequently repeated.

**MISTURA ASSAFŒTIDÆ, Lond.** *Mixture of Assafoetida.*

“Take of Assafoetida, *two drachms*; Water, *half a pint*. Triturate the Assafoetida, gradually adding the Water to it, until they be thoroughly mixed.”

**LAC ASSAFŒTIDÆ, Dub.** *Milk of Assafoetida.*

“Take of Assafoetida, *a drachm*; Pennyroyal Water, *eight fluid ounces*. Triturate the Assafoetida, gradually adding the Water until it form an emulsion.”

Owing to the disagreeable flavour of Assafoetida, it is seldom given by the mouth in this form. The milk is chiefly employed as an enema in flatulent colic, worms, and the convulsions of infants arising from irritations of the bowels during dentition. When given by the mouth, the dose may be from  $f\text{℥}ss.$  to  $f\text{℥}jss.$  frequently repeated.

**MISTURA CAMPHORÆ, Lond.** *Mixture of Camphor.*

“Take of Camphor, *half a drachm*; Rectified Spirit, *ten minims*; Water, *a pint*. Rub the Camphor first with the spirit, then add the water gradually, and strain.”

**MISTURA CAMPHORATA, Dub.** *Camphorated Mixture.*

“Take of Camphor, *a scruple*; Rectified Spirit of Wine, *ten drops*; Refined Sugar, *half an ounce*; Water, *a pint*. Rub the Camphor first with the spirit, and then with the sugar; add the water during the trituration, and strain the mixture through linen.”

*Syn.* Mixture Camphré (*F.*), Kämpfermixtur (*G.*), Mistura Canforata (*I.*).

A pint of water takes up scarcely more than one-half the quantity of Camphor ordered by the London College; but the camphor communicates both odour and taste to the water in a considerable degree. Solution of Pure Potass separates the Camphor. It is an elegant vehicle for more active remedies in low fevers and nervous affections: but cannot be regarded as acting on the habit from the quantity of Camphor it contains. The dose is from  $f\text{℥}j.$  to  $f\text{℥}ij.$  given every three or four hours.

**EMULSIO CAMPHORÆ, Edin.** *Camphor Emulsion.*

“Take of Camphor, *a scruple*; Sweet Almonds blanchéd, Refined Sugar, each, *half an ounce*; Water, *a pint and a half*. It is to be made in the same manner as the common Almond Emulsion.”

In this preparation the whole of the Camphor is diffused through the mixture; the medicinal powers of which are consequently more considerable than those of the foregoing pre-



paration. It is less apt to excite nausea and uneasiness at the stomach than Camphor taken in the solid state, and is given with advantage, in typhus and nervous cases, in doses of  $\text{f}\text{ʒ}\text{ij}$ . every three or four hours. Its preparation should always be extemporaneous, as the Camphor separates and swims on the surface of the mixture after a few days.

**MISTURA CORNU USTI**, Lond. **DECOCTUM CORNU CERVINI**, Dub. *Mixture of burnt Hartshorn.*

“Take of Burnt Hartshorn, *two ounces*; Acacia Gum, in powder, *an ounce (three drachms, Dub.)*; Water, *three pints*. Boil down to two pints, constantly stirring, and strain.”

This is the most unchemical, injudicious, and useless of any of the preparations in the pharmacopœias which have admitted it; being a simple diffusion of insoluble Phosphate of Lime in a thin mucilage.

**MISTURA CRETÆ**, Lond. Dub. *Mixture of Chalk.*

“Take of Prepared Chalk, *half an ounce*; Refined Sugar, *three drachms*; Acacia Gum, in powder, *half an ounce*; (*an ounce, Dub.*); Water, *a pint*. Mix, by trituration.”

**POTIO CARBONATIS CALCIS**, Edin. *Chalk Potion.*

“Take of Prepared Carbonate of Lime (chalk), *one ounce*; Refined Sugar, *half an ounce*; Mucilage of Gum Arabic, *two ounces*. Rub them together, and then gradually add of Water, *two pounds and a half*; Spirit of Cinnamon, *two ounces*. Mix them.”

These are common and useful forms of giving chalk in acidity of the primæ viæ; and combined with Opium or Catechu in diarrhœa. The dose is from  $\text{f}\text{ʒ}\text{j}$ . to  $\text{f}\text{ʒ}\text{ij}$ . given every three or four hours; or after every liquid evacuation.

**MISTURA FERRI COMPOSITA**, Lond.<sup>1</sup> *Compound Mixture of Iron.*

“Take of Myrrh in powder, *a drachm*; Subcarbonate of Potass, *twenty-five grains*; Rose-water, *seven fluid ounces and a half*; Sulphate of Iron, in powder, *a scruple*; Spirit of Nutmeg, *half a fluid ounce*; Refined Sugar, *a drachm*. Rub together the Myrrh, the Subcarbonate of Potass, and the Sugar, and while triturating, add first the Rose-water and the Spirit of Nutmeg, and afterwards the Sulphate of Iron. Put the mixture immediately into a proper glass vessel, and keep it closely stopped.”

In this mixture the Sulphate of Iron is decomposed by the Subcarbonate of Potass forming, by the change of constituents which takes place, Sulphate of Potass and Subcarbonate of Iron; the former of which is dissolved, while the latter is

<sup>1</sup> This name is certainly improper; but it is not easy to invent one which would be descriptive of the compound, and yet be sufficiently concise: *Mistura Subcarbonatis Ferri cum Myrrha* would have been less objectionable.



diffused through the mixture, and kept suspended by the Myrrh, which forms a saponaceous compound with the excess of alkali. The Iron is in the state of a protocarbonate; and as it rapidly attracts oxygen in this state, and is converted into the red oxide, it is necessary to keep the mixture very well excluded from the air: or rather it should not be kept prepared.

*Medical properties and uses.*—This mixture, which is nearly the same as the celebrated antihectic mixture of Dr. Griffith, it an useful tonic in all cases in which preparations of Iron are indicated, particularly in hysteria and chlorosis, and in phthisis, when no active inflammatory diathesis subsists. The dose is from  $\text{f}\frac{3}{4}\text{j}$ . to  $\text{f}\frac{3}{4}\text{ij}$ . given two or three times in a day.

MISTURA GUAIACI, Lond. *Mixture of Guaiacum.*

“Take of Guaiacum, *a drachm and a half*; Refined Sugar, *two drachms*; Mucilage of Acacia Gum, *two fluid drachms*; Cinnamon Water, *eight fluid ounces*. Rub the Guaiacum with the Sugar, then with the Mucilage, and during the trituration add gradually the Cinnamon Water.”

This is a convenient mode of exhibiting Guaiacum. It is given in doses of from  $\text{f}\frac{3}{4}\text{ss}$ . to  $\text{f}\frac{3}{4}\text{ij}$ . two or three times a-day; diluting freely with tepid barley-water or gruel to assist its operation.

MISTURA MOSCHI, Lond. *Mixture of Musk.*

“Take of Musk, Acacia gum, in powder, Refined Sugar, of each *a drachm*; Rose-water, *six fluid ounces*. Rub the Musk with the Sugar, then with the Gum, and add gradually the Rose-water.”

*Syn.* Mixture avec le musc (*F.*), Mixtura Muschiato (*L.*).

The quantity of Gum ordered is scarcely sufficient to retain the Musk suspended in the mixture. It is a convenient form of exhibiting the remedy, and may be given to the extent of  $\text{f}\frac{3}{4}\text{ij}$ . every three or four hours in spasmodic affections, and the sinking state of typhus. The late Mr. White of Manchester, found this mixture combined with Ammonia  $\frac{3}{4}\text{ss}$ . Spirit of Lavender  $\text{f}\frac{3}{4}\text{j}$ . and Spirit of Juniper  $\text{f}\frac{3}{4}\text{j}$ . of great utility in sloughing phagedenic ulcers of a syphilitic and strumous nature.

ENEMA CATHARTICUM, Dub. *Purging Clyster.*

“Take of Manna, *an ounce*; dissolve it in Compound Decoction of Chamomile, *ten fluid ounces*; then add of Olive Oil, *an ounce*; Sulphate of Magnesia, *half an ounce*. Mix them.”

ENEMA FÆTIDUM, Dub. *Fætid Clyster.*

“It is to be prepared by adding to the purging clyster *two drachms* of Assafoetida.”



*SPIRITUS.*

## SPIRITS.

UNDER this title are placed Alcohol and spirituous solutions of vegetable matters, formed by simple mixture, by maceration, and by distillation. They are uniform, transparent, unchanging solutions, containing, in general a large proportion of volatile oil; and when well prepared, are free from empyreuma. Pure Alcohol is more volatile than many of the volatile oils, which do not therefore rise in distillation with it: and, consequently, proof or distilled spirit is employed. As medicinal agents the spirits are stimulant and cordial; but sometimes bad habits are acquired from their continued use. They are employed to cover the taste and flavour of disagreeable medicines; and to make some, which are apt to produce nausea, sit light upon the stomach.

ALCOHOL, Lond. *Alcohol.*

“Take of Rectified Spirit, *a gallon*; Subcarbonate of Potass, *three pounds*. Add a pound of the Subcarbonate previously heated to 300 degrees to the spirit, and macerate for twenty-four hours, frequently shaking the mixture: then pour off the spirit, and add the remainder of the subcarbonate heated to the same degree: lastly, distil the Alcohol from a water-bath, and preserve it in a well-closed vessel. The specific gravity of Alcohol is to that of distilled water, as .815 to 1.000.”

ALCOHOL, Dub. *Alcohol.*

“Take of Rectified Spirit of Wine, *a gallon*; Pearl-ashes dried, at a heat of 300°, and still hot, *a pound*; Caustic Kali, in powder, *an ounce*; Muriate of Lime, dried, *half a pound*. Mix the Spirit and the Kali; add the Pearl-ashes, previously reduced to powder, and digest the mixture for three days in a closed vessel, frequently shaking it; then pour off the spirit; mix with it the Muriate of Lime; and, lastly, distil with a moderate heat until the residue begin to thicken. The specific gravity of this spirit is to that of distilled water, as .815 to 1.000.

“The Muriate of Lime may be conveniently obtained from the residue of the distillation of water of Ammonia.”

*Syn.* Alcohol (*F.*), Höchst rektifizirter Weingiest (*G.*), Alcoole (*I.*).

Rectified spirit, of the specific gravity of 835°, contains about fifteen per cent. of water; and to free it from this is the intention of the above processes. The affinity of the alkali and the muriate of lime for water is much greater than that of spirit; the water, therefore, which the Rectified Spirit con-



tains, is attracted by these substances, and prevented from rising with the spirit during the distillation, by which means the alcohol comes over in a very highly concentrated state. The process of the Dublin College is to be preferred, Muriate of Lime being a much more powerful agent for separating the Water than Subcarbonate of Potass. By its means, Dr. Black obtained alcohol of the specific gravity of  $800^{\circ}$ ; and Richter procured it so low as  $0.792$ , in the temperature of  $68^{\circ}$  Fahrenheit<sup>1</sup>, at which degree of concentration it may be regarded almost as pure Alcohol, or Alcohol perfectly free from water. The Alcohol of the pharmacopœias, therefore, is not free from water, but contains about seven per cent. of water. It is, however, sufficiently concentrated for all the purposes of Pharmacy.

*Qualities.* — Alcohol chemically combines with water: the bulk of the resulting mixture is less and its specific gravity greater than the mean of the two liquids before admixture; and much caloric is evolved. It is highly inflammable, burning with a pale blue flame: and during its combustion, water and carbonic acid are formed, the quantity of the water exceeding that of the alcohol consumed. Alcohol boils at  $174^{\circ}$ , and as its boiling point is higher the more water it contains, its strength may be known by the degree at which it boils; allowing for the atmospheric pressure under which it is tried. In vacuo, it boils at  $56^{\circ}$ . According to Saussure's experiments, it is a compound of Carbon  $52.17$ , Oxygen  $34.79$ , and Hydrogen  $13.04$  in 100 parts of water.<sup>2</sup> It cannot be frozen by any known degree of cold. It dissolves about 60 parts of sulphur, when both the alcohol and the sulphur are in a state of vapour. It dissolves, also, the carbonic and boracic acids, pure Ammonia, Soda, and Potass; volatile oils, resins, and gum-resins; soaps, camphor, sugar, extractive, and the vegetable alkaloids. As a pharmaceutical agent, alcohol, both in its pure and diluted state, is of the utmost importance.

---

<sup>1</sup> Crell's *Annals*, 1796, ii. 211.

<sup>2</sup> *Trans. of the Pharm.* p. 193.



THE following Table drawn up by Lowitz, with an additional column by Dr. Thomson, shows the Specific Gravity of different Mixtures of pure Alcohol of a specific gravity .791, and Distilled Water, at the temperature of 60° and 68° of Fahrenheit.

100 parts by weight.		Sp. Gravity		100 parts by weight.		Sp. Gravity		100 parts by weight.		Sp. Gravity	
Alco.	Wat.	at 68°	at 60°	Alco.	Wat.	at 68°	at 60°	Alco.	Wat.	at 68°	at 60°
100	—	791	796	66	34	877	880	32	68	952	955
99	1	794	798	65	35	880	883	31	69	954	957
98	2	797	801	64	36	882	886	30	70	956	958
97	3	800	804	63	37	885	889	29	71	957	960
96	4	803	807	62	38	887	891	28	72	959	962
95	5	805	809	61	39	889	893	27	73	961	963
94	6	808	812	60	40	892	896	26	74	963	965
93	7	811	*815	59	41	894	898	25	75	965	967
92	8	813	817	58	42	896	900	24	76	966	968
91	9	816	820	57	43	899	903	23	77	968	970
90	10	818	822	56	44	901	904	22	78	970	972
89	11	821	825	55	45	903	906	21	79	971	973
88	12	823	827	54	46	905	908	20	80	973	974
87	13	826	830	53	47	907	910	19	81	974	975
86	14	828	832	52	48	909	912	18	82	976	
85	15	831	†835	51	49	912	915	17	83	977	
84	16	834	838	50	50	914	917	16	84	978	
83	17	836	840	49	51	917	920	15	85	980	
82	18	839	843	48	52	919	922	14	86	981	
81	19	842	846	47	53	921	924	13	87	983	
80	20	844	848	46	54	923	926	12	88	985	
79	21	847	851	45	55	925	928	11	89	986	
78	22	849	853	44	56	927	†930	10	90	987	
77	23	851	855	43	57	930	933	9	91	988	
76	24	853	857	42	58	932	§935	8	92	989	
75	25	856	860	41	59	934	937	7	93	991	
74	26	859	863	40	60	936	939	6	94	992	
73	27	861	865	39	61	938	941	5	95	994	
72	28	863	867	38	62	940	943	4	96	995	
71	29	866	870	37	63	942	945	3	97	997	
70	30	868	871	36	64	944	947	2	98	998	
69	31	870	874	35	65	946	949	1	99	999	
68	32	872	875	34	66	948	951	—	100	1000	
67	33	875	879	33	67	950	953				

\* Alcohol of the Lond. and the Dub. Pharm.

† Ditto, (*Edin.*) rectified spirit, (*Lond.*)

‡ Proof spirit, (*Lond. Dub.*)

§ Proof spirit, (*Edin.*)



TABLE, extracted from the Tables of Mr. Gilpin, showing the Real Specific Gravity of different mixtures of Spirit and Water at every 5° of temperature from 50° to 70°.¹ The standard spirit employed was of the specific gravity 0·825; or contained 89 pure alcohol, and 11 water, in 100 parts.

Proportions by weight of		REAL SPECIFIC GRAVITY.				
Spirit.	Water.	at 50°.	at 55°.	at 60°.	at 65°.	at 70°.
100	—	·82977	·82736	·82500	·82262	·82023
100	5	·84076	·83834	*·83599	·83362	·83124
100	10	·85042	·84802	·84568	·84334	·84092
100	15	·85902	·85664	·85430	·85193	·84951
100	20	·86676	·86441	·86208	·85975	·85736
100	25	·87384	·87150	·86918	·86680	·86415
100	30	·88030	·87796	·87569	·87337	·87105
100	35	·88626	·88393	·88169	·87938	·87705
190	40	·89174	·88945	·88720	·88490	·88254
100	45	·89684	·89458	·89232	·89006	·88773
100	50	·90160	·89933	·89707	·89479	·89252
100	55	·90596	·90367	·90144	·89920	·89695
100	60	·90997	·90768	·90549	·90328	·90104
100	65	·91370	·91144	·90927	·90707	·90484
100	70	·91723	·91502	·91227	·91066	·90847
100	75	·92051	·91837	·91622	·91400	·91181
100	80	·92358	·92145	·91933	·91715	·91493
100	85	·92647	·92436	·92215	·92010	·91793
100	90	·92919	·92707	·92499	·92283	·92069
100	95	·93177	·92960	·92758	·92546	·92333
100	100	·93419	·93208	†·93002	·92794	·92580
100	95	·93658	·93462	·83247	·93040	·92828
109	90	·93897	·93696	‡·93493	·93285	·93076
100	85	·94149	·93948	·93749	·93546	·93337
100	80	·94414	·94213	·94018	·93822	·93616
105	75	·94683	·94486	·94296	·94099	·93898
100	70	·94958	·94767	·94579	·94388	·94193
100	65	·95243	·95087	·94876	·94689	·94500
100	60	·95534	·95467	·95181	·95000	·94813
100	55	·95831	·95662	·95493	·95318	·95139
100	50	·96126	·95966	·95804	·95635	·95469
100	45	·96420	·96262	·96122	·95962	·95802
100	40	·96708	·96595	·96437	·96288	·96143
100	35	·96995	·96277	·96752	·96620	·96484
100	30	·97284	·97181	·97074	·96959	·96836
100	25	·97589	·97800	·97410	·87309	·97203
100	20	·97920	·97887	·97771	·97688	·97596
100	15	·98293	·98289	·98176	·98106	·98028
100	10	·98745	·98702	·98654	·98594	·98527
100	5	·99316	·99284	·99244	·99194	·99134

¹ *Phil. Trans.* for 1794, p. 320—370.  
spirit, (*Lond.*)

† Proof spirit, (*Lond. Dub.*)

\* Alcohol, (*Edin.*) Rectified  
‡ Proof spirit, (*Edin.*)



**SPIRITUS AMMONIÆ**<sup>1</sup>, Lond. *Spirit of Ammonia.*

“Take of Rectified Spirit, *three pints*; Muriate of Ammonia, *four ounces*; Subcarbonate of Potass, *six ounces*. Mix, and, by a gentle heat, distil over a pint and a half of Spirit of Ammonia into a receiver which is kept cold.”

**ALCOHOL AMMONIATUM**, Edin. *Ammoniated Alcohol.*

“Take of Alcohol (835), *thirty-two ounces*; Lime, recently burnt, *twelve ounces*; Muriate of Ammonia, *eight ounces*; Water, *six ounces*. From these Ammoniated Alcohol is prepared exactly in the same manner as Water of Ammonia.”

**SPIRITUS AMMONIÆ**, Dub. *Spirit of Ammonia.*

“Take of Proof Spirit, *three pints*; Muriate of Ammonia, *four ounces*; Potass, *six ounces*. Mix them, and distil with a moderate heat two pints.”

*Syn.* Alcohol Ammoniacale (F.), Gristiger Ammonium liquor (G.), Alcoolé Ammoniato (I.).

In the London process, the Muriate of Ammonia is decomposed by the Subcarbonate of Potass, the alkali of which attracts the muriatic acid, while the ammonia is extricated in the state of a carbonate, is volatilized, and readily combines with the alcohol. Muriate of Potass, or rather Chloride of Potassium, remains in the retort. The present formulæ of the London and of the Dublin Colleges are not essentially the same; the latter containing a solution of pure ammonia in the spirit, while the former contains also a portion of carbonate of ammonia in water. A portion of subcarbonate of ammonia also sublimes, and remains undissolved in the distilled product.

This Spirit properly prepared has the pungent odour and acrid taste of Ammonia, with which it coincides in its medicinal properties. It is used chiefly for Pharmaceutical purposes.

Officinal preparations. *Spiritus Ammoniac compositus*, L. E. D. *Spiritus Ammoniac foetidus*, L. E. D. *Tinctura Castorei composita*, E. *Tinctura Guaiaci composita*, E. *Tinctura Opii ammoniata*, E.

**SPIRITUS AMMONIÆ AROMATICUS**<sup>2</sup>, Lond. *Aromatic Spirit of Ammonia.*

“Take of Cinnamon Bark bruised, Cloves bruised, of each, *two drachms*; Lemon-peel, *four ounces*; Subcarbonate of Potass, *half a pound*; Muriate of Ammonia, *five ounces*; Rectified Spirit, *four pints*; Water, *a gallon*. Mix, and distil over six pints.”

**ALCOHOL AMMONIATUM AROMATICUM**, Edin. *Aromatic ammoniated Tincture.*

<sup>1</sup> Spiritus Salis ammoniaci dulcis, P. L. 1745.

<sup>2</sup> Spiritus volatilis Aromaticus, P. L. 1748. Spiritus salis volatilis oleosus P. L. 1720. Spiritus Ammoniac compositus, P. L. 1787.



“ Take of Ammoniated Alcohol, *eight ounces* ; Volatile Oil of Rosemary, *a drachm and a half* ; Volatile Oil of Lemons, *a drachm*. Mix them so as to dissolve the oils.”

SPIRITUS AMMONIÆ AROMATICUS, Dub. *Aromatic Spirit of Ammonia.*

“ Take of Spirit of Ammonia, *two pints* ; Essential Oil of Lemons, *two drachms* ; Nutmegs, bruised, *half an ounce*. Digest in a covered vessel for three days, frequently shaking the vessel ; then distil *a pound and a half*.”

For these latter preparations it is necessary that the oils be pure ; for if they contain fixed oil, as is often the case with the volatile oils imported into this country, the mixture is rendered turbid and coloured. It is turbid also when prepared with pure oils, if the spirit of ammonia contain any carbonate of ammonia, as must be the case in the Dublin preparation ; in which case it is necessary to distil the mixtures.

*Medical properties and uses.* — This Spirit is an useful stimulant in languors, and flatulant colic ; and the oils render it more grateful to the stomach than the simple spirit of ammonia. The dose is from fʒss. to fʒj. in any convenient vehicle.

Officinal preparations. *Tinctura Guaiaci ammoniata*, L. D. *Tinctura Valerianæ ammoniata*, L. D.

SPIRITUS AMMONIÆ FŒTIDUS<sup>1</sup>, Lond. Dub. *Fœtid Spirit of Ammonia.*

“ Take of Spirit of Ammonia, *two pints* ; Assafoetida, *two ounces (one ounce and a quarter, Dub.)*. Macerate for twelve hours (for three days, in a covered vessel, with frequent agitation, *Dub.*) ; then, by a gentle fire, distil one pint and a half into a cold receiver.”

TINCTURA ASSAFOETIDÆ AMMONIATA, Edin. *Ammoniated Tincture of Ammonia.*

“ Take of Ammoniated Alcohol, *eight ounces* ; Assafoetida, *half an ounce*. Digest them in a close vessel for twelve hours ; then distil eight ounces by the heat of Boiling Water.”

In these processes the fœtid volatile oil of the gum resin is dissolved in the spirit of ammonia, and its odour and flavour communicated to it ; but very little else is taken up. Its medicinal properties are not different from those of the preceding spirit ; and its dose is the same. It acquires colour from age.

SPIRITUS AMMONIÆ SUCCINATUS, Lond. *Succinated Spirit of Ammonia.*

“ Take of Mastich, *three drachms* ; Alcohol, *nine fluid drachms* ; Oil of Lavender, *fourteen minims* ; Oil of Amber, *four minims* ; Solution of Ammonia, *ten fluid ounces*. Macerate

<sup>1</sup> Spiritus volatilis fœtidus, P. L. 1745.



the Mastich in the Alcohol, that it may be dissolved, and pour off the clear tincture; then add the other ingredients, and mix them by agitation."

*Syn.* Eau de Luce (*F.*), Ammoniaca Succinata (*I.*).

The preparation of this name in the Pharmacopœia of 1787 did not preserve the milky appearance characteristic of the *Eau-de-luce*, for which it was intended to be a substitute: and therefore the present formula is given as furnishing a compound capable of preserving its milkiness for a very considerable time. It is stimulant and antispasmodic: and has been successfully used in India against the bite of the rattle-snake. The dose is from  $\mathfrak{m}x.$  to  $\mathfrak{f}3ss.$  given in any convenient vehicle.

**SPIRITUS ANISI**, Lond. *Spirit of Aniseed.*

"Take of Aniseeds bruised, *half a pound*; Proof Spirit, *a gallon*; Water, *a sufficient quantity to prevent empyreuma*. Macerate for twenty-four hours; then distil by a gentle fire."

**SPIRITUS ANISI COMPOSITUS**, Dub. *Compound Spirit of Aniseed.*

"Take of Aniseeds bruised, Angelica Seeds bruised, of each, *half a pound*; Proof Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Distil one gallon."

*Syn.* Alcohol d'Anis (*F.*), Alcoole Anisato (*I.*).

These are pleasant carminatives, useful in flatulent colic and similar affections. The dose is from  $\mathfrak{f}3ss.$  to  $\mathfrak{f}3iv.$  in water.

**SPIRITUS ARMORACIÆ COMPOSITUS**, Lond. *Compound Spirit of Horse-Radish.*

"Take of Fresh Horse-radish, sliced, Orange-peel dried, of each, *a pound*; Nutmegs bruised, *half an ounce*; Proof Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours; and distil a gallon by a gentle fire."

**SPIRITUS RAPHANI COMPOSITUS**, Dub. *Compound Spirit of Horse-Radish.*

"Take of Fresh Horse-radish dried, Peel of Seville Oranges, of each, *two pounds*; Fresh Garden Scurvy-grass, *four pounds*; Nutmegs bruised, *an ounce*; Proof Spirit, *two gallons*; Water, *sufficient to prevent empyreuma*. Distil two gallons."

These spirits were formerly used as antiscorbutics, but they possess little value as such; and are, now, chiefly used in dropsies attended with much debility. The dose is from  $\mathfrak{f}3j.$  to  $\mathfrak{f}3iv.$  combined with infusion of foxglove or of juniper berries.

**SPIRITUS CAMPHORÆ**, Lond. **SPIRITUS CAMPHORATUS**, Dub. *Spirit of Camphor.*

"Take of Camphor, *four ounces*; Rectified Spirit, *two pints*. Mix, that the Camphor may be dissolved."



TINCTURA CAMPHORÆ, Edin. *Tincture of Camphor.*

“Take of Camphor, *an ounce*; Alcohol (sp. grav. 835), *a pound*. Mix, that the Camphor may be dissolved. It may also be made with double or triple the quantity of Camphor.”

*Syn.* Alcohol Camphré (F.), Kampfer Spiritus (G.), Alcoole Canforato (I.)

The strength of the Spirit renders this preparation unfit to be given internally; and the addition of water separates the camphor. It is an useful external application to chilblains, and in chronic rheumatism, paralytic numbness, and gangrene.

SPIRITUS CARUI, Lond. Dub. *Spirit of Carraway.*

“Take of Carraway Seeds bruised, *a pound and a half*, (*half a pound*, Dub.); Proof Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours, then distil a gallon by a gentle fire.”

SPIRITUS CARI CARUI, Edin. *Spirit of Carraway.*

“Take of Carraway Seeds bruised, *half a pound*; Proof Spirit, *nine pounds*. Macerate for two days in a close vessel; then add a sufficient quantity of Water to prevent empyreuma, and distil nine pounds.”

*Syn.* Alcoole con Carvi (I.).

An useful carminative, and adjunct to griping purgatives.

SPIRITUS CINNAMOMI, Lond. Dub. *Spirit of Cinnamon.*

“Take of Oil of Cinnamon, *by weight, five scruples*; Rectified Spirit, *four pints and a half*. Add the Spirit to the Oil, with the addition of as much Water as will be sufficient to prevent empyreuma; then distil a gallon by a slow fire.”

SPIRITUS LAURI CINNAMOMI, Edin. *Spirit of Cinnamon.*

“To be prepared with a pound of Cinnamon Bark, in the same manner as the Spirit of Carraway.”

This spirit is an agreeable cordial in languor and debility. The dose is from f ʒj. to f ʒiv. in any convenient vehicle.

Officinal preparation. *Infusum Digitalis*, L.

SPIRITUS COLCHICI AMMONIATUS, Lond. *Ammoniated Spirit of Colchicum.*

“Take of Seeds of Colchicum bruised, *two ounces*; Aromatic Spirit of Ammonia, *a pint*. Macerate for fourteen days and strain.”

The least useful of the preparations of Meadow Saffron. The dose is from f ʒss. to f ʒj. in water or any mild vehicle.

SPIRITUS JUNIPERI COMPOSITUS, Lond. Dub. Edin. *Compound Spirit of Juniper.*

“Take of Juniper Berries bruised, *a pound*; Carraway Seeds bruised, Fennel Seeds bruised, of each, *an ounce and a half*; Proof Spirit, *a gallon* (*nine pounds*, Edin.); Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours



(two days, *Edin. Dub.*); then distil a gallon (nine pounds, *Edin.*) by a gentle heat."

This Spirit is a grateful and useful addition to infusions of foxglove, and other diuretics, in dropsy.

**SPIRITUS LAVANDULÆ**, Lond. *Spirit of Lavender.*

"Take of Fresh Lavender flowers, *two pounds*; Rectified Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours; then distil a gallon by a gentle heat."

*Dublin.*

"Take of Fresh Flowers of Lavender, *a pound and a half*; Proof Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Distil five pints by a moderate fire."

**SPIRITUS LAVANDULÆ SPICÆ**, Edin. *Spirit of Lavender.*

"Take of Fresh Flowers of Lavender, *two pounds*; Alcohol, *eight pounds*. Distil, with the heat of a water-bath, seven pounds."

*Syn.* Teinture alcoolique de Lavande (*F.*), Lavandel-spiritus (*G.*).

The Oil of Lavender is sufficiently volatile to be brought over with rectified spirit, which is also required to extract all the oil from the flowers; for this reason, the Dublin process produces a spirit less highly impregnated with the oil. Spirit of Lavender is chiefly used as a perfume.<sup>1</sup>

Officinal preparations. *Spiritus Lavandulæ compositus*, L. E. D. *Linimentum Camphoræ compositum*, L.

**SPIRITUS LAVANDULÆ COMPOSITUS**, Lond. Dub. *Compound Spirit of Lavender.*

"Take of Spirit of Lavender, *three pints*; Spirit of Rosemary, *a pint*; Cinnamon Bark bruised, Nutmegs bruised, of each, *half an ounce*; (Cloves, *two drachms*, Dub.); Red Saunders-wood chipped, *an ounce*. Macerate for fourteen days, (ten days, *Dub.*) and strain."

*Edinburgh.*

"Take of Spirit of Lavender, *three pounds*; Spirit of Rosemary, *one pound*; Cinnamon Bark bruised, *an ounce*; Nutmegs bruised, *two drachms*; Red Saunders-wood rasped, *three drachms*. Macerate seven days, and strain."

The addition of these aromatics to the Spirit of Lavender renders it a grateful cordial and stimulant, useful in languors and faintings, and as an adjunct to tonic and stomachic infusions. Its dose is from ℥xxx. to fʒij.

<sup>1</sup> Lavender Water is seldom, as Mr. Brande remarks, a distilled spirit, and each manufacturer has his own recipe. The following is said to be the most approved:—Take of Rectified Spirit of Wine, *five gallons*; Essential Oil of Lavender, *twenty ounces*; Essential Oil of Bergamotte, *five ounces*; Essence of Ambergris, *half an ounce*. Mix.



SPIRITUS MENTHÆ PIPERITÆ, Lond. Edin.  
*Spirit of Peppermint.*

“Take of Peppermint dried, *a pound and a half*; Proof Spirit, *a gallon (nine pounds, Edin.)*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours; then distil a *a gallon (nine pounds, Edin.)* by a gentle heat.”

*Syn.* Teinture alcoolique de Menthe Poivrée (*F.*), Alcoole con Menta piperitide (*I.*).

An useful carminative in nausea and flatulence, and as an adjunct to purgative remedies.

SPIRITUS MENTHÆ VIRIDIS, Lond. *Spirit of Spearmint.*

“Take of Spearmint dried, *a pound and a half*; Proof Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours; then distil a gallon by a gentle heat.”

In the same cases as the former.

SPIRITUS MYRISTICÆ, Lond. SPIRITUS MYRISTICÆ MOSCHATÆ, Edin. SPIRITUS NUCIS MOSCHATÆ, *Spirit of Nutmeg.*

“Take of Nutmegs bruised, *two ounces*; Proof Spirit, *a gallon (nine pounds, Edin.)*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours; then distil a gallon (*nine pounds, Edin.*) by a gentle heat.”

SPIRITUS PIMENTÆ, Lond. SPIRITUS PIMENTO, Dub. *Spirit of Pimenta.*

“Take of Pimento Berries bruised, *two ounces (three ounces, Dub.)*; Proof Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours; then distil a gallon by a gentle heat.”

SPIRITUS MYRTI PIMENTÆ, Edin. *Spirit of Pimenta.*

“It is to be prepared with *half a pound* of bruised Pimenta Berries, in the same manner as Spirit of Carraway.”

An useful carminative in flatulent colic, atonic gout, and dyspepsia.

SPIRITUS PULEGII, Lond. *Spirit of Pennyroyal.*

“Take of Pennyroyal dried, *a pound and a half*; Proof Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours; then distil a gallon by a gentle fire.”

Similar to Spirit of Spearmint in its qualities and medicinal properties.

SPIRITUS ROSMARINI, Lond. *Spirit of Rosemary.*

“Take of Fresh Rosemary-tops, *two pounds*; Proof Spirit, *a gallon*; Water, *sufficient to prevent empyreuma*. Macerate for twenty-four hours; then distil a gallon in a gentle heat.”

SPIRITUS ROSMARINI OFFICINALIS, Edin. *Spirit of Rosemary.*



“Take of Fresh Rosemary-tops, *two pounds* ; Alcohol (sp. grav. 835.), *eight pounds*. Draw off seven pounds by distillation in a water-bath.”

SPIRITUS RORISMARINI, Dub. *Spirit of Rosemary.*

“Take of Fresh Rosemary-tops, *a pound and a half* ; Proof Spirit, *a gallon*. Distil five pints by a moderate fire.”

Syn. Esprit de Rosmarin (F.), Rosmarien-spiritus (G.), Alcoole Rosmarinato (I.).

Oil of Rosemary is sufficiently volatile to rise in distillation with rectified spirit, which the Edinburgh College has, therefore, ordered to be used. It is a fragrant perfume, and is chiefly used in the undermentioned preparations.<sup>1</sup>

Officinal preparations. *Linimentum Saponis compositum*, L. E. D. *Spiritus Lavandulæ compositus*, L. E. D.

TABLE shewing which of the officinal Spirits are prepared with rectified Spirit of wine, and which with proof Spirit.

Rectified Spirits is employed in preparing	Proof Spirits is employed in preparing
Spiritus Ammoniaë aromaticus,	Spiritus Ammoniaë,
——— Ammoniaë succinatus,	——— Anisi,
——— Camphoræ,	——— Armoraciæ compositus,
——— Cinnamomi,	——— Carui,
——— Lavandulæ,	——— Juniperi compositus,
——— Rosmarini, E.	——— Menthæ piperitæ,
	——— Menthæ viridis,
	——— Myristicæ,
	——— Pimentæ,
	——— Pulegii,
	——— Rosmarini, L. D.

TINCTURÆ.

TINCTURES.<sup>2</sup>

THESE are Spirituous Solutions of such of the proximate principles of vegetables and animals, as are soluble in pure

<sup>1</sup> The following are the best recipes for *Hungary Water* and *Eau de Cologne*.

*Hungary Water.*

Take of Fresh Rosemary in blossom, *four pounds* ; Fresh Sage, in blossom, *eight ounces* ; Ginger Root, in blossom, *two ounces*. Cut, bruise, and pour upon them twelve pints of rectified spirit, and two pints of water. Distil, with a slow fire, eleven pints.

*Eau de Cologne.*

Take of Alcohol, *one pint* ; Oil of Bergamotte, Oil of Orange-peel, Oil of Rosemary, of each *one drachm* ; bruised Cardamom seeds, *one drachm* ; Orange flower Water, *one pint*. Distil, from a water-bath, *one pint*.

<sup>2</sup> Arnold de Villa Nova, who was professor of medicine at Montpellier, invented tinctures about the end of the 13th century.



Alcohol or in Proof Spirit. From vegetable matter submitted to its action, Alcohol takes up *Sugar, Resin, Extractive, Tannin, Cinchona, Quinia, Camphor, Volatile Oils, several Acids, Veratrine, Scillatin, Elatin, and Morphia*; Proof Spirit also takes up the whole of these partially, and is besides the proper menstruum for gum-resins; so that Alcohol, either in a concentrated or diluted form, is capable of separating the greater part of the active principles of vegetables from the ligneous inert fibres. The Tinctures obtained from animal substances are very few in number, and the principles taken up by the Spirit are analogous to those enumerated above, belonging to the vegetable kingdom. Pure Alcohol is required in a very few instances only for the formation of Tinctures, Proof Spirit being adequate for almost every purpose. The dilution of the Spirit, however, must be varied according to the known principles of the substance to be submitted to its action: when resin predominates, it must necessarily be more concentrated; when gum-resin or extractive are the most abundant constituents, Proof Spirit then must be employed. In consequence of the great affinity of water for Alcohol, the addition of it to Alcoholic Tinctures separate the resin, camphor, and volatile oils they contain; but water is generally miscible with tinctures made with proof spirit, without producing any decomposition. Tinctures are not liable to suffer spontaneous decomposition, as is the case with infusions and decoctions; and, independent of the loss which takes place from the evaporation of the Spirit and the Volatile Oils, if the bottles containing tinctures be closely corked, they may be kept for an indefinite length of time, and their virtues remain unimpaired.

Tinctures are prepared by macerating the ingredients in the Spirit in a temperature not exceeding  $80^{\circ}$ , at which degree, by allowing the menstruum to remain on the ingredients for a sufficient length of time, all the principles that can prove useful in the tincture are extracted, and the solvent saturated. The ingredients must be dried and reduced to a coarse powder, and the maceration made in close vessels, and assisted by frequent agitation. When completely made, Tinctures should not be allowed to remain upon the ingredients, but be filtered through bibulous paper, and kept in this state in well-corked bottles. Parmentier<sup>1</sup> has proposed that one-half only of the spirituous menstruum be added to the ingredients at first; and, after digesting for six days, this part is to be poured off, and the remainder added. In six more the whole is to be

---

<sup>1</sup> *Annales de Chimie*, lxii. 40.



strongly expressed, and the two portions of tincture mixed together. By this method he imagines more of the active principles of the ingredients are extracted, and the Tinctures obtained of a more uniform strength.

Tinctures are not of very extensive use as remedies, except in cases where stimulants are indicated; the solvent, even in doses of a few fluid drachms, often acting more powerfully on the living system than the principles it holds in solution. In ordinary cases, this action, when continued for some time, produces the same deleterious effects as the habitual use of ardent spirits; and often lays the foundation of the pernicious custom of dram-drinking. When the action of a substance is the reverse of stimulant, it cannot with propriety be exhibited in this form, unless the dose be so small that the operation of the spirit cannot be taken into account, as in tincture of foxglove. The chief use of this class of preparations, therefore, is to enable infusions and decoctions, to which they are added, to sit lighter on the stomach, or to add to them some active principle which water is incapable of extracting.

The general rule given in the London Pharmacopœia for the preparation of tincture, is, “to prepare them in closed vessels, and to shake them frequently during the maceration.”

TINCTURA ALOES, Lond. *Tincture of Aloës.*

“Take of Extract of Spiked Aloës powdered, *half an ounce*; Extract of Liquorice, *an ounce and a half*; Water, *a pint*; Rectified Spirit, *four fluid ounces*. Macerate in a sand-bath until the extracts are dissolved; then strain.”

*Dublin.*

“Take of Socotorine Aloës in powder, *half an ounce*; Extract of Liquorice, dissolved in eight ounces of Boiling Water, *an ounce and a half*; Proof Spirit, *eight fluid ounces*. Digest for seven days; then strain.”

TINCTURA ALÖES SOCOTORINÆ, Edin. *Tincture of Socotorine Aloës.*

“Take of Socotorine Aloës in powder, *half an ounce*; Extract of Liquorice, *one ounce and a half*; Alcohol, *four ounces*; Water, *a pound*. Digest for seven days, with a gentle heat, in a close vessel, which is to be frequently shaken (a circumstance to be attended to in preparing all the tinctures); then pour off the clear tincture.”

*Syn.* Teinture d'Aloës (F.), Tintura d'Aloe (I.).

This may be regarded rather an aqueous solution than a tincture, the quantity of spirit being too small to serve any other purpose than that of preventing decomposition. It may be used in the same cases as the extract of aloës; but notwithstanding the presence of the liquorice, the bitterness of the



alöes is so intense and disagreeable, as to prevent it from being often prescribed. Its dose is from  $f\text{ʒss.}$  to  $f\text{ʒjss.}$

TINCTURA ALOES ÆTHEREA, Edin. *Ethereal Tincture of Alöes.*

“Take of Socotorine Alöes, Myrrh, of each, in powder, *an ounce and a half*; English Saffron cut, *an ounce*; Sulphuric Ether with Alcohol, *a pound*. Digest the Myrrh with the Ether for four days in a closed bottle; then add the Saffron and the Alöes. Digest again for four days, and when the dregs have subsided, pour off the Tincture.”

The Spirit of Sulphuric Ether is supposed to afford a more grateful tincture than Spirit of Wine; and in cases attended with spasm, as in hysteria connected with obstructed menstruation, this solvent may prove serviceable, independent of the matter it holds in solution. It is a warm stomachic purgative, and is advantageously given in dyspeptic affections, jaundice, gout, chlorosis, and other cases in which alöetics are indicated. In doses of  $f\text{ʒj.}$  or  $f\text{ʒij.}$  it acts chiefly as a stomachic; but purges briskly in large doses.

TINCTURA ALOES COMPOSITA, Lond. Dub. *Compound Tincture of Alöes.*<sup>1</sup>

“Take of Extract of Spiked Alöes powdered, Saffron of each, *three ounces*; Tincture of Myrrh, *two pints*. Macerate for fourteen days (seven days, *Dub.*), and strain.”

TINCTURA ALOES ET MYRRHÆ, Edin. *Tincture of Alöes and of Myrrh.*

“Take of Myrrh in powder, *two ounces*; Alcohol, *a pound and a half*; Water, *half a pound*. Mix the Alcohol with the Water; then add the Myrrh; digest for four days: and, lastly, add of Socotorine Alöes in powder, *one ounce and a half*; English Saffron cut in pieces, *one ounce*. Digest again for three days, and pour off the clear tincture.”

*Syn.* Alcohol avec l'Aloë et la Myrrhe (*F.*), Alcoole Aloë Mirrato (*I.*).

This tincture, which differs in little, except the solvent, from the former, may be used in the same cases. It resembles the *Elixir Proprietatis* of the old pharmacopœias. It is occasionally used as a local stimulant to foul ulcers. The dose is from  $f\text{ʒj.}$  to  $f\text{ʒij.}$

TINCTURA ASSAFŒTIDÆ, Lond. *Tincture of Assafœtida.*

“Take of Assafœtida, *four ounces*; Rectified Spirit, *two pints*. Macerate for fourteen days, and strain.”

*Dublin.*

“Take of Assafœtida, *four ounces*; Rectified Spirit of Wine, *two pints*; Water, *eight fluid ounces*. Add the Spirit

<sup>1</sup> Elixir Proprietatis, P. L. 1720.



to the Assafoetida, previously triturated with Water; then digest for seven days, and strain."

TINCTURA FERULÆ ASSAFOETIDÆ, Edin. *Tincture of Assafoetida.*

"Take of Assafoetida, *four ounces*; Alcohol, *two pounds and a half*. Digest for seven days, and filter through paper."

Syn. Teinture d'Assafoetide (F.), Ascande tinktur (G.), Alcoole asfetidato (I.).

When this Tincture is added to water or aqueous infusions, it renders them of a milky hue, owing to the separation of the resin. It is given in the same cases as crude Assafoetida, in doses of fʒj. or more.

Officinal preparation. *Enema foetidum*, D.

TINCTURA AURANTII, Lond. Dub. *Tincture of Orange-peel.*

"Take of Fresh Orange-peel, *three ounces*; Proof Spirit, *two pints*. Macerate for fourteen days, (three days, *Dub.*) and filter."

Syn. Pomeranzan schalen tinktur (G.).

This tincture is not decomposed by water, and may therefore be added to infusions and decoctions, to which it is an useful adjunct in dyspepsia, besides communicating its agreeable flavour.

TINCTURA BENZOINI COMPOSITA, Lond. TINCTURA BENZOES COMPOSITA, Dub. *Compound Tincture of Benzoin.*

"Take of Benzoin, *three ounces*; Storax Balsam strained, *two ounces*; Balsam of Tolu, *an ounce*; Extract of Spiked Aloës, *half an ounce*; Rectified Spirit, *two pints*. Macerate for fourteen days, (seven days, *Dub.*) and filter."

Syn. Teinture de Benzoin composé (F.), Zusammengesetzte Benzoe tinktur (G.).

TINCTURA BENZOINI COMPOSITA, Edin. *Compound Tincture of Benzoin.*

"Take of Benzoin in powder, *three ounces*; Balsam of Peru, *two ounces*; Hepatic Aloes in powder, *half an ounce*; Alcohol, *two pounds*. Digest for seven days, and filter through paper."

Syn. Tintura del Commendatore; Alcoole Benzoato composito (I.).

This Tincture is a stimulating expectorant, and as such is sometimes prescribed in chronic catarrh and old asthmatic cases; but it is chiefly employed as an external application to wounds and languid ulcers, which it gently stimulates, and shields from the action of the air.<sup>1</sup> It is decomposed by water, and therefore, when given internally, must be triturated with yolk of egg or with mucilage, to suspend it in aqueous fluids. Its dose is from fʒss. to fʒij. or more.

TINCTURA BONPLANDIÆ TRIFOLIATA, Edin. *Tincture of Bonplandia, or Augustura.*

<sup>1</sup> It is an improved form of Wade's Balsam, or Friar's Balsam.



“Take of the Bark of Trifoliate Bonplandia bruised, *two ounces*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper.”

TINCTURA ANGUSTURÆ, Dub. *Tincture of Angustura.*

“Take of Angustura Bark in coarse powder, *two ounces*; Proof Spirit, *two pints*. Digest for seven days; then filter.”

*Syn.* Teinture d'Angusture (F.).

This tincture, which contains the active principles of the angustura, is given in doses of fʒj. or fʒij. in the same cases as the bark. (See *Cusparia*, Part ii.)

TINCTURA CALUMBÆ, Lond. *Tincture of Calumba.*

“Take of Calumba root sliced, *two ounces and a half*; Proof Spirit, *two pints*. Macerate for fourteen days, and filter.

TINCTURA COLOMBÆ, Edin. TINCTURA COLOMBO, Dub. *Tincture of Calumba.*

“Take of Colomba root in powder, *two ounces*; Proof Spirit, *two pounds*. Digest for seven days, and filter through paper.”

An useful addition to stomachic infusions and decoctions.

TINCTURA CAMPHORÆ COMPOSITA<sup>1</sup>, Lond. *Compound Tincture of Camphor.*

“Take of Camphor, *two scruples*; Hard Opium powdered, Acid of Benzoin, of each, *one drachm*; Proof Spirit, *two pints*. Macerate for fourteen days, and filter.

TINCTURA OPII CAMPHORATA, sive ELIXIR PAREGORICUM ANGLORUM, Edin. Dub. *Camphorated Tincture of Opium, or Paregoric Elixir.*

“Take of Camphor, *two scruples*; Hard Purified Opium in powder, Benzoic Acid, of each, *a drachm*; Proof Spirit, *two pints*. Digest for seven (ten, *Dub.*) days, then filter.”

*Syn.* Teinture Camphrée d'Opium (F.), Opiumstinktur mit Benzoe (G.).

Half a fluid ounce of this tincture contains nearly a grain of opium. The present tincture differs from that which has been long known under the titles *Paregoric Elixir* and *Asthmatic Elixir*. It is an useful anodyne in chronic asthma, hooping-cough, and catarrh, after the inflammatory symptoms have abated; in which it contributes to allay the tickling which induces the frequent cough. The dose is from fʒj. to fʒij. occasionally in the above cases, using after it the inhaler; and fʒij. in cases where quiet, rather than sleep, is required.

TINCTURA CANTHARIDIS, Lond. *Tincture of Blistering Fly.*<sup>2</sup>

<sup>1</sup> The change of name here imposed, appears to us to be more likely to produce errors than the old name; and it is directly opposed to the principles of nomenclature adopted by the College.

<sup>2</sup> Tinct, Cantharidis, P. L. 1787.



“Take of Blistering Flies bruised, *three drachms*; Proof Spirit, *two pints*. Macerate for fourteen days, and filter.”

TINCTURA CANTHARIDIS VESICATORIÆ, Edin. *Tincture of Blistering Fly.*

“Take of Blistering Flies bruised, *a drachm*; Proof Spirit, *a pound*. Digest for seven days, and filter through paper.”

TINCTURA CANTHARIDIS, Dub. *Tincture of Blistering Fly.*

“Take of Blistering Flies in powder, *two drachms*; Cochineal in powder, *half a drachm*; Proof Spirit, *a pound and a half*. Digest for seven days, then strain.”

Syn. Teinture de Cantharides (F.), Cantharidentinktur (G.), Tintura di Cantaridi (I.).

Proof Spirit extracts the active matter of Blistering Flies, and is a more convenient form for exhibiting it internally than in substance.

This tincture is useful in gleet, fluor albus, incontinence of urine, and in some cutaneous eruptions. The dose is from  $\mathfrak{m}\mathfrak{x}$ . to  $\mathfrak{f}\mathfrak{z}\mathfrak{j}$ . given in some demulcent infusion. As an external application it is efficaciously employed, in conjunction with *soap or camphor liniment*, as an embrocation against rheumatic pains; and I have found that a rag moistened with it is an useful application in that peculiar species of mortification of the extremities which sometimes happens without any apparent cause; and to frost-bitten parts.

TINCTURA CAPSICI, Lond. *Tincture of Capsicum.*

“Take of Capsicum Berries, *an ounce*; Proof Spirit, *two pints*. Macerate for fourteen days, and filter.”

This is a convenient form for exhibiting Capsicum in tympanitis, cynanche maligna, the low stage of typhus, and in similar cases. The dose is from  $\mathfrak{m}\mathfrak{x}\mathfrak{i}\mathfrak{i}$ . to  $\mathfrak{f}\mathfrak{z}\mathfrak{ss}$ .; and a mixture of  $\mathfrak{f}\mathfrak{z}\mathfrak{i}\mathfrak{j}$ . with half a pint of water will answer all the purposes of the capsicum gargle.

TINCTURA CARDAMOMI, Lond. Dub. *Tincture of Cardamoms.*

“Take of Cardamom Seeds husked and bruised, *three ounces*; Proof Spirit, *two pints*. Macerate for fourteen days (seven days, *Dub.*), and filter.”

TINCTURA AMOMI REPENTIS, Edin. *Tincture of Cardamoms.*

“Take of Lesser Cardamom seeds bruised, *four ounces*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper.”

TINCTURA CARDAMOMI COMPOSITA, Lond. Dub. *Compound Tincture of Cardamoms.*

“Take of Cardamom Seeds (husked, *Dub.*), Carraway Seeds, Cochineal, of each, in powder, *two drachms*; Cinnamon Bark bruised, *half an ounce*; Raisins stoned, *four ounces*;



Proof Spirit, *two pints*. Macerate for fourteen days and filter."

The raisins are properly omitted in the Dublin formula. Both the simple and the compound tinctures of cardamoms are agreeable cordials, and form elegant adjuncts to stomachic infusions.

TINCTURA CASCARILLÆ, Lond. Dub. *Tincture of Cascarilla*.

"Take of Cascarilla Bark powdered, *four ounces*; Proof Spirit, *two pints*. Macerate for fourteen days (seven days, *Dub.*), and filter."

TINCTURA CROTONIS ELEUTHERIÆ, Edin. *Tincture of Croton Eleutheria, or Cascarilla*.

"Take of Croton Eleutheria bruised, *four ounces*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper."

*Syn.* Cascarillentinktur (*G.*).

This tincture may be regarded as superfluous, as it is scarcely ever ordered.

TINCTURA CASTOREI, Lond. TINCTURA CASTOREI (ROSSICI, CANADENSIS), Dub. *Tincture of Castor*.

"Take of Castor powdered, *two ounces*; Rectified Spirit, (Proof Spirit, *Dub.*) *two pints*. Macerate for seven days, and filter."

*Edinburgh.*

"Take of Castor powdered, *an ounce and a half*; Alcohol, *a pound*. Macerate for seven days, and filter through paper."

*Syn.* Teinture de Castor (*F.*), Castoreumstinktur (*G.*), Tintura di Castoro (*I.*),

Rectified Spirit is the preferable solvent for the active parts of castor, which are resin and a volatile oil; and it also affords a more grateful tincture than that made with proof spirit. The tincture is supposed to possess the medical properties of the castor, and is used in similar cases. The dose is from ℥xx. to fʒij.

The Dublin College orders a "*Tincture of Russian Castor*," and a "*Tincture of Canadian Castor*," which is an unnecessary refinement. The Russian Castor is the best, and should always be preferred when it can be obtained. A few drops of the Tincture of Castor, dropped into distilled water, produces a milky mixture, which is again cleared and completely decolorized by the addition of Ammonia, if the tincture have been made with Russian Castor; but no such effect is produced if the Tincture be of Canadian Castor.

TINCTURA CASTOREI COMPOSITA, Edin. *Compound Tincture of Castor*.

"Take of Russian Castor powdered, *an ounce*; Assafoetida, *half an ounce*; Ammoniated Alcohol, *one pound*. Digest for seven days, and filter through paper."



This is a more active preparation than the former, and is advantageously given in hysteria, cramp of the stomach, and flatulent colic, to the extent of fʒij. for a dose.

TINCTURA CATECHU<sup>1</sup>, Lond. Dub. *Tincture of Catechu.*

“Take of Extract of Catechu, *three ounces*; Cinnamon Bark bruised, *two ounces*; Proof Spirit, *two pints*. Macerate for fourteen days (seven days, *Dub.*), and filter.”

TINCTURA CATECHU ACACIÆ, Edin. *Tincture of Catechu.*

“Take of Extract of Catechu in powder, *three ounces*; Cinnamon bark bruised, *two ounces*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper.”

*Syn.* Teinture de Cachou (*F.*), Katechutinktur (*G.*).

Proof Spirit dissolves all the soluble parts of catechu, except the mucilage, which in ʒij. of Bengal catechu amounts to 94 grains; besides which 72 grains of impurities remain undissolved. The tincture is a solution of tannin, extractive matter, and the volatile oil of Cinnamon. It is a grateful warm astringent, useful in all cases in which astringents are indicated. The dose is from fʒj. to fʒij. taken in water or in wine, or in the cretaceous mixture of the pharmacopœias.

TINCTURA CINCHONÆ<sup>2</sup>, Lond. *Tincture of Cinchona.*

“Take of Lance-leaved Cinchona bark in powder, *seven ounces*; Proof Spirit, *two pints*. Macerate for fourteen days, and filter.”

TINCTURA CINCHONÆ, Edin. TINCTURA LANCIFOLIÆ CINCHONÆ, Dub. *Tincture of Lance-leaved Cinchona.*

“Take of Cinchona Bark in powder, *four ounces*; Proof Spirit, *two pounds and a half* (*two pints*, *Dub.*). Digest for seven days, and filter through paper.”

*Syn.* Teinture de Quinquina (*F.*), Chinatinktur (*G.*), Tintura di China (*I.*).

Although this tincture contains the active principles of Cinchona bark in considerable quantity, yet, from the nature of the vehicle, it cannot be given in sufficiently large doses to produce the beneficial effects of the bark in substance; it is therefore used chiefly as an adjunct to the infusion or decoction. The dose is from fʒj. to fʒiv.

TINCTURA CINCHONÆ AMMONIATA, Lond. *Ammoniated Tincture of Bark.*

“Take of Lance-leaved Cinchona Bark powdered, *four ounces*; Aromatic Spirit of Ammonia, *two pints*. Macerate for ten days, and filter.”

This is the formula of 1787, which was rejected without

<sup>1</sup> Tinctura Japonica, P. L. 1745.

<sup>2</sup> Tinctura corticis Peruvianæ simplex, P. L. 1745.



any proper reason in 1809. It is extremely useful in dyspeptic complaints attended with much acidity.

**TINCTU'RA CINCHO'NÆ COMPOS'ITA**, Lond. Edin. Dub. *Compound Tincture of Cinchona.*

“Take of Lance-leaved Cinchona Bark powdered, *two ounces*; Dried Orange-peel, *one ounce and a half*, (*half an ounce* Dub.); Virginian Snake-root bruised, *three drachms*; Saffron, *a drachm*; Cochineal in powder, *two scruples*; Proof Spirit, *twenty fluid ounces*. Macerate for fourteen (*seven*, Edin.) days, and filter.”

*Syn.* Zusammengesetzte Chinatinktur (G.).

This Tincture is more grateful than the former; and although it contains less Cinchona, yet the addition of the other ingredients renders it more useful both as a stomachic and a febrifuge. It is the same as the celebrated *Tincture of Huxham*<sup>1</sup>, who generally gave it in intermittents, and low nervous fevers, in diluted wine or any proper vehicle, with ten or fifteen drops of elixir of vitriol, (aromatic sulphuric acid, Edin.) The dose is from fʒj. to fʒiij. or more in intermittents.

**TINCTU'RA CINNAMO'MI**, Lond. Dub.<sup>2</sup> **TINCTURA LAURI CINNAMOMI**, Edin. *Tincture of Cinnamon.*

“Take of Cinnamon Bark bruised, *three ounces*; (*three ounces and a half*, Dub.) Proof Spirit, *two pints*. Macerate for fourteen days (*seven days*, Dub.), and filter.”

*Syn.* Teinture de Cannelle (F.), Zimmtinktur (G.), Tintura di Cinnamonomo (I.).

This Tincture contains the active principles of the Bark, and is an elegant and useful adjunct to the chalk mixture and astringent infusions. The dose is from fʒj. to fʒiij.

**TINCTU'RA CINNAMO'MI COMPOS'ITA**<sup>3</sup>, Lond. Dub. *Compound Tincture of Cinnamon.*

“Take of Cinnamon Bark bruised, *six drachms*; Cardamom Seeds bruised, *three drachms*; Long Pepper powdered, Ginger-root sliced, of each, *two drachms*; Proof Spirit, *two pints*. Macerate for fourteen days (*seven days*, Dub.), and filter.”

**TINCTURA CINNAMOMI COMPOSITA**, Edin. *Compound Tincture of Cinnamon.*

“Take of Cinnamon Bark bruised, lesser Cardamom Seeds bruised, of each, *one ounce*; Long Pepper in powder, *two drachms*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper.”

*Syn.* Teinture de Saffron (F.).

This is a much warmer aromatic than the simple tincture; and is frequently advantageously used in flatulencies, atonic gout, languors, and debility, in doses of fʒj or fʒij. properly diluted.

<sup>1</sup> *Essay on Fever*, 122.

<sup>2</sup> *Aqua Cinnamomi fortis*, P. L. 1720.

<sup>3</sup> *Tinctura aromatica*, P. L. 1745.



Official preparation. *Æther sulphuricus cum Alcohole aromaticus*, E.

TINCTURA CONII MACULATI, Edin. *Tincture of Hemlock.*

“Take of Dried Leaves of Hemlock, *two ounces*; Cardamom Seeds, bruised, *half an ounce*; Proof Spirit, *sixteen ounces*. Digest for seven days, and filter through paper.”

This is an elegant form of administering Conium; and admits of its being added to mixtures. It possesses all the active properties of the plant.

TINCTURA CROCI SATIVI, Edin. TINCTURA CROCI, Dub. *Tincture of Saffron.*

“Take of English Saffron cut in shreds, *one ounce*; Proof Spirit, *fifteen ounces* (*a pint*, Dub.). Digest for seven days, and filter through paper.”

This Tincture contains almost pure extractive, and is supposed to be stimulant and diaphoretic, but its chief value perhaps arises from its colour.

TINCTURA DIGITALIS, Lond. *Tincture of Foxglove.*

“Take of Foxglove Leaves dried, *four ounces*; Proof Spirit, *two pints*. Macerate for fourteen days, and filter.”

Syn. Teinture de Digitale (F.), Fingerhautinktur (G.), Tintura di Digitale porporina (I.).

*Dublin.*

“Take of Foxglove Leaves (the larger ones being rejected) dried and reduced to coarse powder, *two ounces*; Proof Spirit, *a pint*. Digest for seven days, and filter.”

TINCTURA DIGITALIS PURPUREÆ, Edin. *Tincture of Foxglove.*

“Take of Foxglove Leaves dried, *one ounce*; Proof Spirit, *eight ounces*. Digest for seven days, and filter through paper.”

This is a convenient form for exhibiting Foxglove. It contains Digitalia, the alkaloid principle on which the virtues of the plant depend, and has the advantage of preserving these virtues unimpaired for any length of time. The dose should be *m̄x.* at first, and gradually increased, the same cautions being necessary as in the exhibition of the plant in substance.

TINCTURA GALBANI, Dub. *Tincture of Galbanum.*

“Take of Galbanum cut into small pieces, *two ounces*; Proof Spirit, *two pints*. Digest for seven days, then filter.”

It is used in the same cases as tincture of Assafœtida; but, if less nauseous, it is also less powerful.

TINCTURA GALLARUM, Edin. Dub. *Tincture of Galls.*



“Take of Galls in powder, *two ounces (four ounces, Dub.)*; Proof Spirit, *sixteen ounces (two pints, Dub.)*. Macerate for seven days, then filter through paper.”

Proof Spirit dissolves tannin; consequently this tincture contains all the astringency of the galls, and may be employed in the same cases. The dose is from fʒj. to fʒiij.

TINCTURA GENTIANÆ COMPOSITA<sup>1</sup>, Lond. Dub. *Compound Tincture of Gentian.*

“Take of Gentian Root cut, *two ounces*; Orange-peel dried, *an ounce*; Cardamom Seeds bruised, *half an ounce*; Proof Spirit, *two pints*. Macerate for fourteen days (seven days, *Dub.*), and filter.”

TINCTURA GENTIANÆ COMPOSITA, Edin. *Compound Tincture of Gentian, commonly called Stomachic Tincture.*

“Take of Yellow Gentian Root sliced and bruised, *two ounces*; Orange-peel dried and bruised, *one ounce*; Canella Alba bruised, *half an ounce*; Cochineal in powder, *half a drachm*; Proof Spirit, *two pints and a half*. Digest for seven days, and filter through paper.”

Syn. Teinture de Gentiane composée (*F.*), Enziantinktur (*G.*), Tintura di Genziana (*I.*).

This is an elegant stomachic bitter and cordial; but in dyspepsia, in which it is more particularly indicated, the infusion is preferable.

TINCTURA GUAIACI, Lond. Dub. *Tincture of Guaiacum.*

“Take of Guaiacum powdered, *half a pound (four ounces, Dub.)*; Proof Spirit, *two pints*. Macerate for fourteen days (seven days, *Dub.*), and filter.”

TINCTURA GUAIACI OFFICINALIS, Edin. *Tincture of Guaiacum.*

“Take of Guaiacum, in powder, *six ounces*; Alcohol, *two pounds and a half*. Digest for seven days, and filter through paper.”

Syn. Teinture de Guajac (*F.*), Guajaktinktur (*G.*), Tintura di Guajac (*I.*).

The difference in the proportion of the Guaiacum in these formulæ is much to be regretted. It is separated from the alcohol by the addition of water; and, therefore, when this tincture is to be given in the form of draught, it must be triturated with yolk of egg, or with mucilage, to enable it to combine with water. The dose is from fʒj. to fʒiij. in any convenient vehicle.

TINCTURA GUAIACI AMMONIATA<sup>2</sup>, Lond. Dub. Edin. *Ammoniated Tincture of Guaiacum.*

“Take of Guaiacum in powder, *four ounces*; Aromatic

<sup>1</sup> Tinctura amara, P. L. 1745.

<sup>2</sup> Tinct. Guaiacina volatilis, P. L. 1745.



Spirit of Ammonia, *one pint and a half*. Macerate for fourteen days (seven days, *Edin., Dub.*), and filter."

*Syn.* Teinture Ammoniacal de Guajac (*F.*), Ammonium guajaktinktur (*G.*), Alcoole Ammoniato con Guajaco; Tintura Guajachina volatile (*I.*).

As the ammonia coincides with the operation of Guaiacum more than spirit, this tincture is more efficacious as a stimulating sudorific than the former preparation. Water decomposes it, separating the Guaiacum in dark curdy flakes. Chlorine, nitrous acid, and the spirit of nitric ether separate the Guaiacum into curdy coagulated masses, and impart to the whole an intense blueish-green colour: but sulphuric and muriatic acids produce no change. Dr. Paris, from whose Pharmacologia the above remark is quoted, adds, "if equal parts of quicklime and powder of Guaiacum be rubbed together, and a quantity of water be poured over them, and the mixture allowed to stand until it become fine, we shall obtain a solution of this substance, which will mix, in any proportion, with aqueous vehicles without decomposition, and to which the aromatic spirit of ammonia may be subsequently added with effect." The dose is from  $\text{f3j.}$  to  $\text{f3ij.}$ , triturated with any mucous or viscid matter.

TINCTU'RA HELEBO'RI NI'GRI', Lond. *Tincture of Black Hellebore.*

"Take of the Root of Black Hellebore sliced, *four ounces*; Proof Spirit, *two pints*. Macerate for fourteen days, and filter."

*Edinburgh. Dublin.*

"Take of Black Hellebore-root bruised (powdered, *Dub.*), *two ounces*; Cochineal in powder *fifteen grains* (*two scruples*, *Dub.*); Proof Spirit, *fifteen ounces* (*two pints*, *Dub.*). Digest for seven days, then filter through paper."

*Syn.* Teinture d'Ellebore noir (*F.*), Tintura d'Elleboro (*I.*).

The smallness of the fibres of the Root of Black Hellebore, which are the parts medicinally employed, renders it almost impossible to follow the direction of the London formula; and it is better to powder it coarsely, as ordered by the Dublin College. This tincture was regarded by Dr. Mead as a powerful emmenagogue, and is still ordered in uterine obstructions, and in some cutaneous affections. The dose is from  $\text{mxxx.}$  to  $\text{f3j.}$ , in any appropriate vehicle.

TINCTU'RA HU'MILI, Lond. Edin. *Tincture of Hops.*

"Take of Hops, *five ounces*; Proof Spirit, *two pints* (*two pounds and a half*, *Edin.*). Macerate for fourteen days, and strain (seven days; express the tincture, and filter through paper, *Edin.*)."



The lightness and bulk of the Hops render it difficult to make the quantity of spirit here ordered act equally on the ingredients, therefore their surface should be several times changed by stirring, during the maceration, and the tincture expressed as ordered by the Edinburgh College. The tincture is supposed to possess the tonic and narcotic properties of the plant, and may be regarded as an alcoholic solution of Lupulin. It has been recommended as a substitute for tincture of opium in gout and rheumatism<sup>1</sup>; but, from the experiments of Dr. Bigsby, its efficacy is very<sup>2</sup> problematical. The dose is from fʒss. to fʒij. or more.

TINCTURA HYOSCYAMI, Lond. *Tincture of Henbane.*

“Take of the Dried Leaves of Henbane, *four ounces*; Proof Spirit, *two pints*. Macerate for fourteen days, and filter.”

*Dublin.*

“Take of the Dried Leaves of Black Henbane in coarse powder, *two ounces and a quarter*; Proof Spirit, *a pint*. Digest for seven days; then strain.”

TINCTURA HYOSCYAMI NIGRI, Edin. *Tincture of Black Henbane.*

“Take of the Dried Leaves of Black Henbane, *an ounce*; Proof Spirit, *eight ounces*. Digest for seven days, and filter through paper.”

I have found this a more useful substitute for Tincture of Opium, than the Tincture of Hops. In a dose of fʒj. it seldom fails of procuring sleep and quiet; and does not affect the head, or produce costiveness. In cases of diarrhœa, when this tincture is given, it will be necessary to add a few drops of tincture of opium to counteract the tendency it has to run off by the bowels.

TINCTURA JALAPÆ, Lond.<sup>3</sup> *Tincture of Jalap.*

“Take of Jalap-root, powdered, *eight ounces*; Proof Spirit, *two pints*. Digest for fourteen days, then filter.”

*Dublin.*

“Take of Jalap-root in coarse powder, *five ounces*; Proof Spirit, *two pints*. Digest for seven days, then filter.”

TINCTURA CONVULVULI JALAPÆ, Edin. *Tincture of Jalap.*

“Take of Jalap-root in powder, *three ounces*; Proof Spirit, *fifteen ounces*. Digest for seven days, and filter through paper.”

<sup>1</sup> *Freak's Observations on the Humulus Lupulus*, 9. et passim.

<sup>2</sup> *London Medical Repository*.

<sup>3</sup> *Tinctura Jalapæ*, P. L. 1745, 1787.



Both water and alcohol separately extract part of the active principles of jalap, and proof spirit extracts the whole of them; the combination of the gum, extractive, and resin of the root being requisite for the production of its full cathartic effect. The great difference in point of strength of these tinctures is much to be regretted.

TINCTU'RA KINO, Lond. *Tincture of Kino.*

"Take of Kino, in powder, *three ounces*; Proof Spirit, *two pints*. Macerate for fourteen days, and strain."

*Edinburgh. Dublin.*

"Take of Kino, *two ounces (three ounces, Dub.)*; Proof Spirit, *a pint and a half*. Digest for seven days, and filter through paper."

*Syn.* Teinture de Kino (F.).

The matter in solution in this tincture is chiefly tannin. It is administered in chronic diarrhœa, the latter stage of dysentery, fluor albus, and in all cases in which astringents are indicated; but it is less certain in its operation than the tincture of catechu. The dose is from fʒj. to fʒij.

TINCTURA MOSCHI, Dub. *Tincture of Musk.*

"Take of Musk in powder, *two drachms*; Rectified Spirit, *a pint*. Digest for seven days, then strain."

*Syn.* Teinture de Musc (F.), Tintura di Muschio (I.).

The only effectual form in which Musk can be exhibited is in powder; much larger doses of it being requisite to do any good than can be given in a spirituous vehicle.

TINCTU'RA MYRRHÆ, Lond.<sup>1</sup> *Tincture of Myrrh.*

"Take of Myrrh, bruised, *four ounces*; Rectified Spirit, *three pints*. Macerate for fourteen days, and filter."

*Edinburgh.*

"Take of Myrrh, in powder, *three ounces*; Alcohol, *twenty ounces*; Water, *ten ounces*. Digest for seven days, and filter through paper."

*Dublin.*

"Take of Myrrh, bruised, *three ounces*; Proof Spirit, *a pint and a half*; Rectified Spirit, *half a pint*. Digest for seven days, then strain."

*Syn.* Teinture de Myrrhe (F.), Myrrhentinktur (G.), Tintura de Mirra (I.).

A transparent tincture of a golden yellow colour may be prepared by treating myrrh with alcohol or rectified spirit alone, as now ordered by the London College; but, when water is used, a rather turbid tincture is obtained; a circumstance which, *à priori*, might have been suspected. This tincture is tonic and deobstruent; but it is more generally used in gargles, combined with infusion of roses and acids;

<sup>1</sup> Tinct. Myrrhæ simplex, P. L. 1720.



or as an application to foul ulcers, and exfoliating bones; or diluted with water, as a wash for the mouth when the gums are spongy. The dose is from  $\text{f}\text{ʒss.}$  to  $\text{f}\text{ʒj.}$

TINCTURA OPII, Lond. *Tincture of Opium.*

“Take of hard Opium, powdered, *two ounces and a half*; Proof Spirit, *two pints*. Macerate for fourteen days, and strain.”

TINCTURA OPII, sive THEBAICA; vulgo, LAUDANUM LIQUIDUM, Edin. *Tincture of Opium, or Thebaic Tincture, commonly, Liquid Laudanum.*

“Take of Opium, *two ounces*; Proof Spirit, *two pounds*. Macerate for seven days, and filter through paper.”

TINCTURA OPII, sive TINCTURA THEBAICA, Dub. *Tincture of Opium, or Thebaic Tincture.*

“Take of Purified Hard Opium in coarse powder, *ten drachms*; Proof Spirit, *a pint*. Digest for seven days, then strain.”

*Syn.* Teinture d'Opium (F.), Opiums-tinktur (G.), Laudano liquido; Alcoleo opiato (I.).

Owing to crude opium being now ordered by the London College instead of hard purified opium, the strength of the tincture formerly prepared is to that of the present tincture as 3 to 2; or  $\text{m}\text{xiv.}$  of the old tincture contained one grain of opium, and were equal to  $\text{m}\text{xix.}$  of the present tincture. The Edinburgh tincture is of the same strength as the present London tincture; but the Dublin is stronger,  $\text{m}\text{xiv.}$  of it containing one grain of opium. Mr. Brande says the whole of the Morphia is not taken up, for it is found in no inconsiderable quantity in the matter which remains on the filter. This tincture is used in all cases in which opium is indicated, and is a very convenient and elegant form of giving the remedy.<sup>1</sup> The usual dose is from  $\text{m}\text{x.}$  to  $\text{m}\text{lx.}$ ; but in some morbid states of the habit very large doses can be borne, and are even necessary. In colica pictonum,  $\text{f}\text{ʒj.}$  given before using purges, facilitates their action, and renders the relief more speedy; and in tetanus  $\text{f}\text{ʒvss.}$  have been given in divided doses, with advantage in twenty-six hours.<sup>2</sup> As an external application, the tincture rubbed upon the skin produces its anodyne effects in a smaller degree, allays local pains, and assists in relaxing the spasm in lock-jaw and similar affections. Its powers as an external remedy are very much increased by combining it with vinegar; an acetate of morphia being thus produced: whilst it is decomposed by

<sup>1</sup> It ought to be kept in opaque bottles; as light, according to the experiments of Vogel, has the power of decomposing it. *Vide Journ. Pharm. Maii. 1815. p. 199.*

<sup>2</sup> *Currie's Report on Cold Water, i. 138.*



ammonia, potass, soda, and their subcarbonates, and its Morphia precipitated.

**TINCTURA OPII AMMONIATA**; olim, **ELIXIR PAREGORICUM**, Edin. *Ammoniated Tincture of Opium*; formerly *Paregoric Elixir*.

“Take of Opium, *two drachms*; Benzoic Acid, Saffron, cut in shreds, of each, *three drachms*; Volatile Oil of Aniseed, *half a drachm*; Ammoniated Alcohol, *sixteen ounces*. Digest for seven days, and filter through paper.”

This tincture is useful in whooping-cough and spasmodic asthma. Each fʒj. contains gr. j. of opium.

**TINCTURA QUASSIÆ EXCELSÆ**, Edin. *Tincture of Quassia*.

“Take of Quassia Wood rasped, *one ounce*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper.”

**TINCTURA QUASSIÆ**, Dub. *Tincture of Quassia*.

“Take of Chips of Quassia Wood, *an ounce*; Proof Spirit, *two pints*. Digest for seven days, then strain.”

*Syn.* Teinture de Quassia (F.).

This tincture contains the bitter of the Wood in perfection, and may be used in the same cases as the infusion.

**TINCTURA RHEI**<sup>1</sup>, Lond. *Tincture of Rhubarb*.

“Take of Rhubarb Root sliced, *two ounces*; Cardamom Seeds bruised,† *an ounce and a half*; Saffron, *two drachms*; Proof Spirit, *two pints*. Macerate for fourteen days in a gentle heat, and filter.”

*Dublin.*

“Take of Rhubarb Root sliced, *two ounces*; Lesser Cardamom Seeds husked and bruised, Liquorice bruised, of each, *half an ounce*; Saffron, *two drachms*; Proof Spirit, *two pints*. Digest for seven days, then filter.”

**TINCTURA RHEI**, Edin. *Tincture of Rhubarb*.

“Take of Rhubarb Root sliced, *three ounces*; Lesser Cardamom Seeds bruised, *half an ounce*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper.

*Syn.* Teinture de Rhubarbe (F.), Rhabarbertinktur (G.), Tinctura de Rhabarbaro (I.).

**TINCTURA RHEI COMPOSITA**<sup>2</sup>, Lond. *Compound Tincture of Rhubarb*.

“Take of Rhubarb Root sliced, *two ounces*; Liquorice Root bruised, *half an ounce*; Ginger Root sliced, Saffron, of each, *two drachms*; Proof Spirit, *a pint*; Water, *twelve*

<sup>1</sup> Tinctura Rhabarbari, P. L. 1720. 1787.

<sup>2</sup> Tinctura Rhabarbari composita, P. L. 1787.

*† half an ounce only*



*fluid ounces.* Macerate for fourteen days in a gentle heat and filter."

**TINCTURA RHEI ET ALOES**, Edin. *Tincture of Rhubarb and Aloes*; formerly, *Sacred Elixir*.

"Take of Rhubarb Root sliced, *ten drachms*; Socotorine Aloes powdered, *six drachms*; Lesser Cardamom Seeds bruised, *half an ounce*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper."

*Syn.* Alcohol avec Aloé et Rhubarbe (*F.*), Alcoole Aloe-Rabarbarato (*I.*).

**TINCTURA RHEI ET GENTIANÆ**, Edin. *Tincture of Rhubarb and Gentian*

"Take of Rhubarb Root sliced, *two ounces*; Gentian Root, sliced, *half an ounce*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper."

All these Tinctures of Rhubarb are purgative and stomachic; but the strength of the menstruum is too great to permit of their general use for the first intention, and they are more usually employed as adjuncts to saline purgatives, to give them warmth, or to stomachic infusions in dyspepsia, flatulent colic, diarrhœa, the costiveness of old people and of those of cold phlegmatic habits. The dose to operate as a purgative is fʒvj. and from fʒj. to fʒiij. to produce stomachic effects.

**TINCTURA SCILLÆ**, Lond. Edin. Dub. *Tincture of Squills*.

"Take of recent Squill Root (bulb) dried, *four ounces two ounces*, Edin.); Proof Spirit, *two pints* (*sixteen ounces*, Edin.). Macerate for fourteen days, and filter." (Digest for seven days, then set it aside until the dregs are subsided, and pour off the clear liquor, *Dub.*)

Proof spirit takes up the active principles of the squill, and affords a convenient form of exhibiting it in all the cases in which it is indicated. The dose is from m̄x. to m̄xxx. given in almond mixture, ammoniac mixture, or mucilage.

**TINCTURA SENNÆ**, Lond. *Tincture of Senna*.<sup>1</sup>

"Take of Senna Leaves, *three ounces*; Carraway Seeds bruised, *three drachms*; Cardamom Seeds bruised, *a drachm*; Raisins stoned, *four ounces*; Proof Spirit, *two pints*. Macerate for fourteen days in a gentle heat, and filter."

*Dublin.*

"Take of Senna Leaves, *a pound*; Carraway Seeds bruised, *one ounce and a half*; Lesser Cardamom Seeds husked and bruised, *half an ounce*; Proof Spirit, *a gallon*. Digest for fourteen days, then filter."

<sup>1</sup> Elixir Salutis, P. L. 1720.



**TINCTURA SENNÆ COMPOSITA**, Edin. *Compound Tincture of Senna.*

“Take of the Leaves of Senna, *two ounces*; Jalap Root bruised, *one ounce*; Coriander Seeds bruised, *half an ounce*; Proof Spirit, *three pounds and a half*. Digest for seven days, and to the filtered tincture add of Refined Sugar, *four ounces*.”

These tinctures are stomachic and purgative. They are very efficacious in flatulent colic, atonic gout<sup>1</sup>, and as an opening medicine for those whose bowels have been weakened by intemperance. The dose is from fʒij. to fʒj. in any appropriate vehicle.

**TINCTURA SERPENTARIÆ**, Lond. Dub. *Tincture of Snake Root.*

“Take of Snake Root (cut and bruised, *Dub.*), *three ounces*; Proof Spirit, *two pints*. Macerate for fourteen days (seven days, *Dub.*), and filter.”

**TINCTURA ARISTOLOCHIÆ SERPENTARIÆ**, Edin. *Tincture of Snake Root.*

“Take of Snake Root bruised, *two ounces*; Cochineal in powder, *a drachm*; Proof Spirit, *two pounds and a half*. Digest for seven days, and filter through paper.”

This tincture is a useful addition to infusion of Cinchona bark, in typhoid and putrid fevers, gout, and periodic head-ach. The dose is from fʒss. to fʒij; or, when taken in water, as much as can be taken without the operation of the spirit proving hurtful.

**TINCTURA TOLUIFERÆ BALSAMI**, Edin. *Tincture of Balsam of Tolu.*

“Take of Balsam of Tolu, *an ounce and a half*; Alcohol, *a pound*. Digest until the balsam is dissolved, and filter through paper.”

Tincture of Balsam of Tolu is scarcely ever used except on account of its agreeable flavour. As it is decomposed by water, it is necessary to triturate it with mucilage, in order to mix it with any aqueous fluid. It is chiefly used for making the syrup.

Officinal preparation. *Syrupus Toluiferæ Balsami*, E.

**TINCTURA VALERIANÆ**, Lond. Dub. *Tincture of Valerian.*

“Take of Valerian Root in powder, *four ounces*; Proof Spirit, *two pints*. Macerate for fourteen days (seven days, *Dub.*), and filter.”

*Syn.* Teinture de Valeriane (*F.*), Baldrian Tinktur (*G.*), Tintura di Valeriana (*I.*).

<sup>1</sup> The preparation called *Gout Cordial* is a mixture of Tincture of Senna and Tincture of Rhubarb.



Proof-spirit extracts the active matter of the Valerian, but the tincture cannot be given in doses sufficiently large to prove very efficacious.

**TINCTURA VALERIANÆ AMMONIATA**, Lond.  
*Ammoniated Tincture of Valerian.*<sup>1</sup>

“Take of Valerian Root, *four ounces*; Aromatic Spirit of Ammonia, *two pints*. Macerate for fourteen days, and filter.”

*Dublin.*

“Take of Valerian Root in powder, *two ounces*; Spirit of Ammonia, *a pint*. Digest for seven days, then filter.”

*Syn.* Baldriantinktur mit Ammonium liquor (G.).

As the Ammonia corresponds in virtue with the Valerian, this tincture is more powerful than the foregoing. It is advantageously employed in hysteria and other nervous affections, in doses of fʒj. or fʒij.; given in milk, or some other bland fluid.

**TINCTURA VERATRI ALBI**, Edin. *Tincture of White Hellebore.*

“Take of White Hellebore Root bruised, *eight ounces*; Proof Spirit, *one pound and a half*. Digest for seven days, and filter through paper.”

This tincture is sometimes employed to excite vomiting in maniacal and apoplectic cases; and as an alterative in cutaneous eruptions. It owes its efficacy to an alkaloid, which has been named *Veratria*, the same principle to which Colchicum owes its virtues. It is given in doses of ℥v. to ℥x.; but it is a very unmanageable remedy, producing sometimes the most violent effects.

**TINCTURA ZINGIBERIS**, Lond. Dub. **TINCTURA AMOMI ZINGIBERIS**, Edin. *Tincture of Ginger.*

“Take of Ginger Root sliced, *two ounces*; Rectified Spirit, *two pints*. Macerate for fourteen days (seven days, *Dub. Edin.*), and filter.” The substitution of rectified spirit in this preparation is highly judicious, as the spirit takes up the acrid principles of the Ginger without acting on the mucilage which, when proof spirit was used, rendered the tincture turbid.

This tincture possesses all the pungency of the ginger, and is useful as a stimulant and carminative, in atonic gout, when it attacks the stomach, in flatulent colic, and as a corrigent to griping purgatives.

---

<sup>1</sup> Tinctura Valerianæ volatilis, P. L. 1745.



## TABLE OF TINCTURES.

Tinctures prepared with rectified spirit of the Spec. grav. 835.	Tinctures prepared with proof spirit of the Spec. grav. 950.
<p>Tinctura Alöes,</p> <p>———— Alöes et Myrrhæ,</p> <p>———— Assafoetidæ,</p> <p>———— Benzōini comp.</p> <p>———— Castorei,</p> <p>———— Ferri Muriatis,</p> <p>———— Guaiaci,</p> <p>———— Myrrhæ,</p> <p>———— Moschi,</p> <p>———— Toluiferæ Balsami,</p> <p>———— Zingiberis.</p>	<p>Tinctura Aurantii,</p> <p>———— Bonplandiæ trifoliatæ,</p> <p>———— Calumbæ,</p> <p>———— Camphoræ comp.</p> <p>———— Capsici,</p> <p>———— Cardamomi,</p> <p>———— Cardamomi comp.</p> <p>———— Cascarillæ,</p> <p>———— Catechu,</p> <p>———— Cinchonæ,</p> <p>———— Cinchonæ comp.</p> <p>———— Cinnamomi,</p> <p>———— Cinnamomi comp.</p> <p>———— Conii maculati,</p> <p>———— Croci sativi,</p> <p>———— Digitalis,</p> <p>———— Ferri Ammoniat,</p> <p>———— Galbani,</p> <p>———— Gallarum,</p> <p>———— Gentianæ comp.</p> <p>———— Hellebori nigri,</p> <p>———— Humuli,</p> <p>———— Hyosciami,</p> <p>———— Jalapæ,</p> <p>———— Kino,</p> <p>———— Lyttæ,</p> <p>———— Opii,</p> <p>———— Quassiæ excelsæ,</p> <p>———— Rhei,</p> <p>———— Rhei composita,</p> <p>———— Rhei et Alöes,</p> <p>———— Rhei et Gentianæ,</p> <p>———— Scillæ,</p> <p>———— Sennæ,</p> <p>———— Serpentariæ,</p> <p>———— Valerianæ,</p> <p>———— Veratri albi.</p>



*ÆTHEREA.*

## PREPARATIONS OF ETHER.

THE action of the strong acids on alcohol produces an order of compounds which possess both important chemical properties and medicinal virtues. These are named ETHERS, and agree in certain general properties, but vary in some of their qualities according as they are produced from different acids. They are all extremely volatile, and require to be preserved in closely-stopped phials.<sup>1</sup> The following are medicinally used.

ÆTHER SULPHURICUS, Lond. *Sulphuric Ether.*

“Take of Rectified Spirit, Sulphuric Acid, of each, *one pound and a half*. Pour the Spirit into a glass retort, and add the Acid gradually to it, shaking it frequently, and taking care that the temperature of the mixture do not exceed 120°, until it be completed. Then cautiously place the retort in a sand-bath, previously heated to 200°, that the liquor may boil as quickly as possible, and the ether pass into a tubulated receiver to which another receiver is adapted, which is to be cooled by ice or water. Distil until a heavier fluid begin to pass over, which is seen in the bottom of the receiver below the ether. Pour on the liquor which remains in the retort twelve fluid ounces more of Rectified Spirit, and distil another portion of Æther in a similar manner.”

*Edinburgh.*

“Take of Sulphuric Acid, Alcohol, of each, *thirty-two ounces*. Pour the Alcohol into a glass retort, capable of sustaining a sudden heat; then pour the Acid on it in an uninterrupted stream. Mix them gradually by frequent and gentle agitation; then immediately distil from a sand-bath, previously heated for the purpose, into a receiver kept cold with water or snow. Let the fire be so regulated that the fluid may boil as soon as possible, and continue to boil until sixteen ounces shall have distilled over; then let the retort be removed from the sand-bath.

“To the distilled liquor, add two drachms of Potass; then distil again from a high retort, with a very gentle heat, into a receiver kept cold, until ten ounces have passed over.”

---

<sup>1</sup> The phial proposed by Dr. Dewar is the best for this purpose. It consists of a stopped phial, having a circular rim round its shoulder, not rising quite so high as the mouth of the bottle, and a glass cup with a heavy bottom, which, when inverted over the mouth of the phial, and mercury poured into the rim, hermetically closes it. *Annals of Phil.* vol. x. p. 20.



“ If sixteen ounces of Alcohol be added to the residuary acid after the first distillation, and the distillation repeated, ether will be reproduced. And this may be often repeated.”

ÆTHER RECTIFICATUS, Lond. *Rectified Ether*.<sup>1</sup>

“ Take of Sulphuric Ether, *fourteen fluid ounces*; Fused Potass, *half an ounce*; Distilled Water, *eleven fluid ounces*. First dissolve the Potass in two fluid ounces of the Water, and add the Æther to it, shaking them well together, until they be mixed; then, at a temperature of about 120°, distil from a large retort into a cooled receiver, twelve fluid ounces of ether: next shake the distilled fluid with nine fluid ounces of Water, and set it aside that the water may subside. Lastly, pour off the supernatant rectified Æther, and keep it in a well-stopped vessel.”

ÆTHER SULPHURICUS, Dub. *Sulphuric Ether*.

“ Take of Sulphuric Ethereal liquor, *twenty fluid ounces*; Subcarbonate of Kali dried and in powder, *two drachms*. Mix them, and distil from a high retort by means of a very gentle heat, into a receiver kept cold, twelve fluid ounces. The specific gravity of this fluid is to that of distilled water, as 765 to 1000.”

*Syn.* Ether (*F.*), Schwefelätther (*G.*), Etere (*I.*).

The admixture of alcohol and sulphuric acid produces an almost instantaneous formation of ether, which is made sensible by the odour of the mixture; while by the mutual action of the spirit and the acid on each other a considerable evolution of caloric takes place, and the temperature of the mixture is raised to 180°. Whatever can encourage these effects in the first instance is to be avoided, as by the sudden rise of temperature, and the disengagement of ethereal vapours before the apparatus be adjusted, not only is the retort in danger of being broken, but a considerable waste of product also takes place. The proper mode for forming the mixture is undoubtedly that of the London College, if the retort be cooled after the addition of each portion of the acid. The retort should be thin, and the sand-bath previously heated to 200°, so that the liquor may boil immediately; for the ether is formed, and distils over at this temperature: whereas, by gradually raising the heat to this point, part of the alcohol comes over unaltered. The ether, as it distils, is condensed in the cool receivers, in the form of a colourless, limpid, transparent fluid; but towards the end of the operation, a white vapour also comes over, on the appearance of which the distillation should be stopped. The receivers ought to be ample, and kept cool with ice or snow, or cold water,

---

<sup>1</sup> Ether vitriolicus, P. L. 1787.



which we have found to be preferable to ice or snow. The best mode of applying it is to lay narrow shreds of woollen cloth over the receivers, with one end of each immersed in a vessel of cold water placed higher than the receivers, by which means the water is made to trickle constantly over them; and, by the evaporation which it suffers, the receivers are kept in a sufficiently low temperature, and at the same time, the nature of their contents is distinctly seen, which cannot be conveniently done when they are immersed in snow, or ice, or even water. The luting which answers best in this operation is common paste, spread on slips of cloth, which are to be first applied, and when dry surrounded with pieces of wet bladder.

The product of the first distillation is Sulphuric Æther combined with water, some alcohol, and a small portion of sulphurous acid, forming an impure Æther of the specific gravity  $\cdot 768$ ; and that of the second distillation, or after the addition of a new portion of alcohol, is a similar Æther of the specific gravity  $\cdot 807$ : on mixing these, a fluid of the specific gravity  $\cdot 788$  is obtained, which is the unrectified Æther of the present London Pharmacopœia.<sup>1</sup> By the rectification of this Æther according to the directions of the British Colleges, it is deprived of the sulphurous acid and nearly all the water, and its specific gravity reduced to  $\cdot 732$ , or when highly rectified to  $\cdot 725$ ; but it still contains some water and alcohol, as Æther of a specific gravity so low as  $\cdot 632$  in the temperature of  $60^{\circ}$  is said to have been obtained.<sup>2</sup> The use of the alkali in the rectification is to separate and detain the acid and the water by its affinity for these substances; and this is still more completely accomplished by the addition of a portion of black oxide of manganese, which, by affording oxygen to the sulphurous acid, converts it into sulphuric acid, and thus renders it perfectly fixed at the temperature employed.

The theory of the formation of ether is still unsettled. It has been contended, that the balance of affinities between the constituents of the alcohol is broken by the acid, the oxygen of which attracting a portion of the hydrogen of the alcohol forms water; while a portion of its carbon, at the same time set free, forms the residuary black matter found in the retort;

---

<sup>1</sup> *London Medical Review*, April 1810, p. 163.

<sup>2</sup> Lowitz procured ether of this gravity by the following process. To ether reduced to  $\cdot 746$  specific gravity by means of subcarbonate of potass in the usual method, he added as much dry powdered muriate of lime as it would dissolve. On standing, the mixture separated into two parts; the alcohol holding the salt in solution sunk to the bottom; the ether swam on the surface. When separated from the inferior liquor, its specific gravity was now only  $\cdot 632$  in the temperature of  $60^{\circ}$ . *Thomson's Chemistry*, 4th ed. ii. 443.



and, by a new combination of the remaining hydrogen, carbon, and oxygen, the ether is produced. This explanation, however, which supposes a partial decomposition of the acid, has been denied by Fourcroy and Vauquelin, who, from a series of very ingenious experiments<sup>1</sup>, concluded, that the acid suffers no decomposition, except towards the end of the process, which is to be attributed to the carbonaceous matter collected in the retort; but that it produces the decomposition of the alcohol without being itself decomposed, by the exertion alone of a disposing affinity. The ether, according to them, is the result of the new combination of the components of the alcohol, part of its oxygen and hydrogen first combining to form water, and a large portion of its carbon being separated without entering into any new combination; so that ether differs from alcohol only in containing a greater proportion of hydrogen and oxygen, and a smaller proportion of carbon. Thus, alcohol, according to the analysis of Saussure, is a compound of carbon 52·17, oxygen 34·79, and hydrogen 13·04 in 100 parts; whilst æther, according to the same chemist, consists of carbon 67·98, oxygen 17·62, and hydrogen 14·40: or, according to the atomic theory, alcohol consists of 2 atoms = 12, ether of 4 atoms = 24 of carbon; alcohol of 1 atom = 8, ether of 1 atom = 8 of oxygen; alcohol of 3 atoms = 3; ether of 5 atoms = 5 of hydrogen; making the weight of an atom of the former = 23, and that of an atom of the latter = 37. Several objections have been raised to this theory; but, in a work of this nature it is not necessary to enter minutely into theoretical discussions. The sulphurous and carbonic acids, and the charry residue formed in the process, are produced by the decomposition of a portion of the alcohol by the acid. Mr. Phillips says, that in using the proportions directed by the London College, he procured  $f\frac{3}{4}xj$  of product, of which  $f\frac{3}{4}xj\frac{1}{4}$  were Sulphuric Ether of specific gravity 0·768. On adding the second portion of Spirit, the ætherial product was equal to the former, but of specific gravity 0·807, showing that the power of the acid to produce ether was much diminished by the first operation.<sup>2</sup>

*Qualities.* — Æther has a fragrant, penetrating odour, and a hot, pungent taste. It is colourless and perfectly limpid; and is the most volatile of liquids, drying immediately if poured on the hand, and producing a great degree of cold by its evaporation. It boils in the open air at 96°, and in vacuo at 20°, a temperature 12 degrees below the common freezing point; and, but for the pressure of the atmosphere, would

<sup>1</sup> *Annales de Chimie*, xxiii. 205.

<sup>2</sup> *Trans. of the Pharmacopæia*, p. 224.



always be in a gaseous form. When cooled down to  $-46^{\circ}$ , ether congeals in brilliant transparent plates. It is extremely inflammable, taking fire on the approach of any ignited body, a circumstance which requires to be attended to in pouring it from one phial to another by candle light. During its combustion, carbonic acid is formed, and traces of charcoal are left behind. It unites with alcohol in every proportion, and also readily mixes with ammonia; but ten parts of water take up or dissolve one only of ether. It dissolves balsams, wax, volatile oils, bitumens, camphor, extractive, gum-resins, resins, and sulphur in small proportions. It is decomposed by sulphuric acid, which converts it into sweet oil of wine.<sup>1</sup> Ether is sometimes adulterated; if it contain sulphuric acid from imperfect rectification, this may be detected by a precipitate being formed on the addition of solution of Barytes: when *alcohol* is present, a milky solution is formed with phosphorus, which is not the case when it is pure.

*Medical properties and uses.*—Sulphuric ether is stimulant, narcotic and antispasmodic. In its operation it resembles alcohol, but is more diffusible, and its effects are less permanent. It is beneficially employed as a cordial in typhoid and low fevers, particularly when nausea, subsultus tendinum, and other spasmodic symptoms are present. As an antispasmodic, it relieves the paroxysm of spasmodic asthma, whether it be taken into the stomach, or its vapour only be inhaled into the lungs; in which latter form it is also useful in simple dyspnœa and in catarrh. Much caution, however, is required in inhaling the vapour of ether, as the imprudent inspiration of it has produced lethargic and apoplectic symptoms. It is employed with advantage in hysteria, tetanus, cramp of the stomach, hiccough, and in cholera morbus, to check the vomiting; and also allays the violence of sea-sickness. The usual dose of sulphuric ether is from fʒss. to fʒij.; but it has been given in much larger doses with the most beneficial effects; and in all cases the dose must be repeated at short intervals to produce the full effect of the remedy. As an external application, ether acts either as a stimulant or a refrigerant, according to the mode in which it is applied. The first takes place when it is prevented from evaporating, by being confined over the spot to which it is applied; in which case it often proves useful in relieving headach and other muscular pains: and from its refrigerent effect produced by its rapid evaporation, it is applied to burns, and to assist in the reduc-

---

<sup>1</sup> When *sulphuric* and *muriatic ethers* are mixed together in equal proportions, the evaporation is very rapid, and a degree of cold considerably below 0 of Fahrenheit is produced.



tion of strangulated hernia. We have seen it produce almost immediate relief in ear-ach, when dropped into the external meatus.

Officinal preparations. *Spiritus Ætheris sulphurici*, L.

**OLEUM ÆTHEREUM**, Lond. *Ethereal Oil*.<sup>1</sup>

“After the distillation of Sulphuric Ether, distil again the remaining liquor with a gentle heat, until a black froth swells up; then immediately remove the retort from the fire. To the liquor in the retort, add water sufficient that the oily part may float upon it. Let this be skimmed off, and as much Lime-water be added to it as will neutralize any acid it may contain; and shake them together. Lastly take off the Ethereal Oil after it has separated.”

**LIQUOR ETHEREUS OLEOSUS**, Dub. *Oily Ethereal Liquor*.

“Take what remains in the retort after the distillation of Sulphuric Ether. Distil to one-half, by a moderate heat.”

*Syn.* Huile douce de vin (*F.*), Oleo dolce di Vino (*I.*).

The product of both of these processes is a thick oily matter, of a yellow colour, slightly viscid; of sp. gr. 1.05, less volatile than ether, soluble both in ether and alcohol; but insoluble in water. After being kept for a few months, it becomes viscid, and prismatic crystals form in it, which are soluble in Ether and Alcohol; melt with a slight heat and sublime unaltered. In sulphuric acid they dissolve, forming a pink solution, in nitric a deep red: but they are soluble in muriatic and acetic acids and in the pure alkalies. Its nature is not clearly ascertained; for, although Fourcroy and Vauquelin consider it as similar to ether, and differing from it principally in containing a larger proportion of carbon; yet other chemists maintain that it is merely a compound of ether and sulphurous acid. From experiments made to ascertain its nature by Mr. Hennell, there appears to be much probability in his opinion, that it is a compound of sulphorinic acid, and the crystallizable oil already described.<sup>2</sup> It can be obtained more directly, although less economically, by distilling ether with a portion of sulphuric acid. It is used only for the preparation of the compound spirit.

Officinal preparation. *Spiritus Ætheris Sulphurici compositus*, L. D.

**SPIRITUS ÆTHERIS AROMATICUS**, Lond. *Aromatic Spirit of Ether*.<sup>3</sup>

“Take of Cinnamon Bark bruised, *three drachms*; Cardamom Seeds powdered, *a drachm and a half*; Long Pepper powdered, Ginger-root sliced, of each, *a drachm*; Spirit of

<sup>1</sup> Oleum Vini, P. L. 1787.

<sup>2</sup> *Journal of Science*, vol. xix. p. 96.

<sup>3</sup> Elixir Vitrioli dulce, P. L. 1745.



Sulphuric Ether, *a pint*. Macerate for fourteen days in a stopped glass-bottle, and strain."

ÆTHER SULPHURICUS CUM ALCOHOLE AROMATICUS, Edin. *Aromatic Sulphuric Ether with Alcohol*.

"Take of Cinnamon Bark bruised, Cardamom Seeds bruised, each, *an ounce*; Long Pepper bruised, *two drachms*; Sulphuric Ether with Alcohol, *two pounds and a half*. Digest for seven days, and filter through paper."

These preparations do not differ in their medicinal properties from the former; the aromatics rendering them only a little more grateful.

SPIRITUS ÆTHERIS NITRICI, Lond. *Spirit of Nitric Ether*.<sup>1</sup>

"Take of Rectified Spirit, *two pints*; Nitric Acid (by weight), *three ounces*. Add the Acid gradually to the Spirit, and mix them, taking care that the temperature, during the mixture, does not exceed  $120^{\circ}$ ; then distil, by a gentle heat, twenty-four fluid ounces."

SPIRITUS ÆTHERIS NITROSI, Edin. *Spirit of Nitrous Ether*.

"Take of Alcohol, *three pounds*; Nitrous Acid, *one pound*; pour the Alcohol into a large phial placed in a vessel full of cold water, and add the acid gradually, with frequent agitation. Let the phial be slightly corked, and placed in a cool place for seven days; then distil the liquor by the heat of boiling water, into a receiver kept cool with snow or water, as long as any spirit comes over."

SPIRITUS ÆTHEREUS NITROSUS, Dub. *Nitrous Ethereal Spirit*.

"Add to the matter which remains after the distillation of Nitrous Ether, the Rectified Spirit of wine, employed in that operation for condensing the elastic vapour, and distil to dryness with the greater heat of a water-bath. Mix the distilled liquor with the alkaline liquor which remains after the separation of the Nitrous Ether, and also add as much dry Subcarbonate of Kali as shall be sufficient to saturate the predominant acid; which is to be determined by the test of litmus. Lastly, distil by the medium heat of a water-bath, as long as any fluid comes over. The specific gravity of this liquor is to that of distilled water, as 850 to 1000."

*Syn.* Alcool éthéréux par l'acide nitrique (*F.*), Atherischer saltpeter spiritus (*G.*), Spirito di nitro dolce (*I.*).

The products of the London and Edinburgh processes are in every respect the same; but the former is to be preferred on account of the length of time required by the latter. The small quantity of acid in proportion to the alcohol employed,

<sup>1</sup> Spiritus Nitri dulcis, P. L. 1745. Spir. Ætheris nitrosi, 1787.



permits the mixture to be effected without any violent action taking place, or the evolution of much heat, provided the acid be added in small quantities and at intervals, and each portion be thoroughly mixed with the alcohol before another be added. The heat employed for the distillation should not exceed  $212^{\circ}$ , and it should be stopped as soon as twenty-four fluid ounces come over; for when it is longer continued, the product becomes coloured, and contains too much free acid.<sup>1</sup> The theory of the operation, inasmuch as relates to the production of the nitric ether, which is thus obtained in combination with a large proportion of unchanged alcohol and a small proportion of nitric acid, is nearly the same as that already detailed; except that a partial decomposition of the acid takes place. The entire product has the same relation to nitric ether as spirit of sulphuric ether has to sulphuric ether.

The product obtained by the first part of the Dublin process is analogous to the above. The acid which the residue of the distillation of nitrous ether contains, and the alcohol already impregnated with a small portion of that fluid, when mixed and heated, act reciprocally on each other, and a compound of nitric ether, unchanged alcohol, and free acid, distils over; but the alkali with which it is mixed before the second distillation, removing the acid, its properties, both as a chemical compound and as a remedy, must be necessarily altered. The products of the former processes are those which have been longest known and most extensively employed.

*Qualities.* — Spirit of nitric ether, as procured by the London or the Edinburgh process, has an extremely fragrant odour, and a pungent acidulous taste. Its spec. grav. should not exceed 0.834. It is very volatile and inflammable; and soluble in water and in alcohol. It coagulates tincture of Guaiacum, giving it at the same time a deep blue colour; and it also strikes a deep olive with solution of green sulphate of iron.<sup>2</sup> Nitric Ether, according to Thenard, consists of Carbon 28.45, Oxygen 48.52, Hydrogen 8.54 and Azote 14.49 in 100 parts. The proportions of this Ether in the Spirit has not been determined.

*Medical properties and uses.* — Spirit of nitric ether is refrigerant, diuretic, and antispasmodic. It has long been employed under the title of *Sweet Spirit of Nitre*, as a grateful refrigerant, and to quench thirst in febrile affections; for which purpose the dose is from  $\mathfrak{mxx}$ . to  $\mathfrak{mxl}$ . given in a cup-

<sup>1</sup> *London Medical Review*, April 1810, p. 164.

<sup>2</sup> Dr. Paris says (*Pharmacologia*), that this ethereal spirit, when added in a small proportion to malt spirits, communicates to them a flavour resembling that of *French brandy*.



ful of water, or any other appropriate vehicle. In larger doses, it acts as a gentle stimulant to the stomach, relieving nausea and flatulence; and also determines to the kidneys, increasing the flow of urine; on which account it is advantageously prescribed as an auxiliary to other diuretics in dropsical complaints.

**SPIRITUS ÆTHERIS SULPHURICI**, Lond. *Spirit of Sulphuric Ether.*

“Take of Sulphuric Ether, *half a pint*; Rectified Spirit, *a pint*. Mix them.”

**ÆTHER SULPHURICUS CUM ALCOHOLE**, Edin. *Sulphuric Ether with Alcohol.*

“Take of Sulphuric Ether, *one part*; Alcohol, *two parts*. Mix them.”

**LIQUOR ÆTHEREUS SULPHURICUS**, Dub. *Sulphuric ethereal Liquor.*

“Take of Rectified Spirit of Wine, Sulphuric Acid, of each, *thirty-two ounces*. Let the Spirit heated to  $120^{\circ}$  be poured into a glass retort fit to bear a sudden heat, and add the Acid in an uninterrupted stream; let them be gradually mixed, and by means of a quick and sufficiently powerful heat, distil twenty ounces of the liquor into a receiver kept cool.

“If sixteen ounces of Rectified Spirit of Wine be poured on the residuary acid in the retort, more sulphuric ethereal liquor will be obtained by repeating the distillation.”

In the old method of preparing this spirit by distilling the charge for sulphuric ether by a slow and gradually increased heat, an alcoholized ether was obtained, owing to part of the alcohol first passing over unaltered before the ether was formed, the specific gravity of which was  $\cdot 768$ ; but the gravity of the above mixture is  $\cdot 816$ , showing that it contains considerably less ether in combination with the alcohol.

*Medical properties and uses.* — It may be used for the same purposes as the ether; but it is necessarily much less active. The dose is from  $\text{f}\text{ʒj.}$  to  $\text{f}\text{ʒiij.}$  An useful gargle for slight inflammation of the fauces is prepared by adding  $\text{f}\text{ʒj.}$  of this spirit to  $\text{f}\text{ʒvj.}$  of barley water, sweetened with  $\text{f}\text{ʒiv.}$  of syrup of marsh-mallows.

**SPIRITUS ÆTHERIS SULPHURICI COMPOSITUS**, Lond. *Compound Spirit of Ether.*

“Take of Spirit of Sulphuric Ether, *a pint*; Ethereal Oil, *two fluid drachms*. Mix them.”

*Syn.* Alcool éthereux par l'acide sulphurique (*F.*), Atherischer Schwefelegeist Liquor (*G.*), Anodino minerale dell' Hoffmann (*I.*).

This is intended as a substitute for the *Anodyne Liquor of Hoffmann*; and, besides being stimulant and antispasmodic, it is supposed to possess anodyne properties. It is an useful ad-



dition to tincture of opium, when given with the intention of procuring sleep; and often prevents the opium from exciting the nausea which it is apt to produce in some habits. The dose is from f3ss. to f3ij. in any appropriate vehicle.

**ÆTHER NITROSUS**, Dub. *Nitrous Ether*.

“Take of Nitrate of Kali, dried and coarsely powdered, *a pound and a half*; Sulphuric Acid, *a pound*; Rectified Spirit of Wine, *nineteen fluid ounces*. Put the Nitrate of Kali into a tubulated retort, placed in a bath of cold water, and pour upon it in small quantities, and at intervals, the Sulphuric Acid and the Spirit previously mixed together, and allow the mixture to become cold. Without any external heat, or at least a very gentle one, (such as may be communicated by the addition of a little tepid water to the bath,) an ethereal liquor will begin to distil. In a short time the heat of the retort will spontaneously increase, and a considerable ebullition take place, which must be moderated by adding some cold water to the bath. The receiver must also be kept cool with water or snow, and furnished with a proper apparatus for transmitting the very elastic vapour (arising from the mixture with great force, if the heat be too much increased) through a pound of Rectified Spirit of Wine in a phial which is to be kept cold.

“The Æthereal Liquor, thus spontaneously distilled, is to be put into a glass phial, fitted with a ground glass stopper, and as much Subcarbonate of Kali, dried, and in powder, is to be added as is necessary to neutralize the acid, closing the phial after each addition of the Kali, and determining the neutralization by the test of litmus: about a drachm of salt is generally sufficient for this purpose; and in a short time the Nitrous Ether will rise to the surface, and is to be separated by means of a funnel.

“If the Æther is required to be very pure, distil it again from a water-bath, at a temperature of  $140^{\circ}$ , to one-half. Its specific gravity is to that of distilled water, as 900 to 1000.”

The action of nitric acid on alcohol is so violent, that the formation of ether which it affords, has always been regarded as a process of great difficulty, to obviate which, many ingenious plans have been suggested. The operation which has been just described, is admirably adapted for procuring it with facility and safety. It was contrived by Wolfe, and was found by Pelletier to succeed better than any other. The sulphuric acid and the spirit must be mixed with the same degree of caution as is necessary in preparing sulphuric ether; and the receiver must be larger, and kept perfectly cool, with the apparatus, described in the formulæ, attached to it, which



should be kept cool by a mixture of snow or ice and muriate of lime.

In the above process, the nitrate of potass is first decomposed, and nitric acid formed, which acts upon the alcohol as it evolves. The theory of this action is very obscure: but from a number of well-contrived experiments, Thenard was led to draw the following conclusions. Both the acid and the alcohol are decomposed; the oxygen of the former combines with a large proportion of the hydrogen, and a small quantity of the carbon of the alcohol, and thence result, "1st, Much water and nitrous oxide, and small quantities of carbonic acid, nitrous acid, and nitric oxide. 2dly, The separation of a small quantity of nitrogen, and the formation of much nitric ether by the combination in large quantity of the two elements of the nitric acid, with the alcohol from which the large proportions of hydrogen and small proportion of carbon have been abstracted. 3dly, The formation of acetic acid, and of a matter disposed to pass to the state of charcoal, by the combination, in certain proportions, of the hydrogen and carbon of the alcohol with the oxygen of the nitric acid."<sup>1</sup>

*Qualities.* — Nitrous, or rather Nitric Æther has a strong ethereal odour, but is less fragrant than sulphuric ether. Its taste is strong and peculiar; and its colour slightly yellow, probably arising from the presence of a small portion of nitric oxide. When highly rectified, its specific gravity is 0·866<sup>2</sup>. It is more volatile than sulphuric ether, boiling at a temperature of 70°, and consequently producing a greater degree of cold by its evaporation; and is very inflammable. It requires 48 parts of water for its solution, but combines with alcohol in every proportion; and readily absorbs nitrous and acetic acids, both of which acids are formed in it when it is kept for some time. According to the analysis of Thenard, the constituents of 100 parts of nitric ether are 48·52 of oxygen, 28·45 of carbon, 14·49 of azote, and 8·54 of hydrogen; but this analysis is liable to some exceptions.

*Medical properties and uses.* — Nitric Æther, although introduced into the Dublin Pharmacopœia, has not yet been generally used in practice; but it is probable that its properties are the same as those of sulphuric ether, and consequently it is applicable to the same cases.

---

<sup>1</sup> Murray's Chemistry, 2d ed. iv. 447.

<sup>2</sup> Duncan, New Edinburgh Dispensatory, 5th ed. 567.



## VINA.

## WINES.

WINE acts upon vegetable substances in nearly the same manner as diluted spirit, dissolving such of their proximate principles as can be taken up by water and alcohol when combined: hence it has been long used as a menstruum for extracting the active parts of medicinal vegetables; and the solutions thus formed have been denominated *Medicated Wines*. As a solvent, however, it is liable to the objection of inequality of strength; and owing to the spontaneous decomposition which it undergoes from exposure to the air, it is still more objectionable, this change being likely to take place sooner when it is imbued with principles all of which tend to hasten the fermentative process. To remedy these disadvantages in this class of preparations, Parmentier has proposed<sup>1</sup>, that instead of preparing medicated wines in the usual method, the alcoholic tinctures well prepared should be added to wine in given quantities: by which means, he contends that the preparations are less nauseous, and, what is a still greater advantage, are always of a determinate strength. Two of the British Colleges still order medicated wines to be prepared after the old method; and when the vegetable products which are to be taken up are of an alkaline nature, as morphia, veratria, &c. it is still the best method. They should be kept in very well corked bottles, and in a cool situation. By the general term Wines, however, the London College, now designates preparations, the menstrua of which are dilute spirits of different degrees of strength. Admitting that this were a judicious change, we cannot avoid criticising the absurdity of the title *Vina*, when applied to preparations which do not contain one drop of wine; but, as will be seen in the sequel, the change is not so generally advantageous as might be expected, from its adoption by the College.

TABLE of the Quantity of Proof Spirit and Water ordered by the London College, for the Preparation of the Wines.<sup>2</sup>

Names of the Wines.	Proof Spirit.	Water.
Vinum Alöes	1 part	1 part.
—— Colchici	1 part	2 parts.
—— Ferri	1 part	1½ parts.
—— Ipecacuanhæ	1 part	1⅔ parts.
—— Opii	1 part	1⅔ parts.
—— Veratri	1 part	1½ parts.

<sup>1</sup> *Annales de Chimie*, lii. 46.

<sup>2</sup> Vide Phillips' *Trans. of the Pharm.* 1824.



VINUM ALOES, Lond. *Wine of Aloes.*

“Take of Extract of Spiked Aloes, *eight ounces*; Canella Bark, *two ounces*; Proof Spirit, Distilled Water, of each, *four pints*. Rub the extract to powder with white sand previously freed from impurities; rub the Canella Bark also into powder, and on these, mixed together, pour the Water and Spirits. Macerate for fourteen days, frequently shaking the vessel containing the mixture, and afterwards strain.”

*Dublin.*

“Take of Socotorine Aloes, *four ounces*; Canella Alba, *an ounce*; Spanish White Wine, *three pints*; Proof Spirit, *a pound*. Let the Aloes and the Canella, separately reduced to powder, be mixed together, and pour on them the wine mixed with the spirit; then digest for fourteen days with frequent agitation; and lastly, strain the solution.”

VINUM ALOES SOCOTORINÆ, Edin. *Wine of Socotorine Aloes.*

“Take of Socotorine Aloes in powder, *one ounce*; Lesser Cardamom Seeds bruised, Ginger Root bruised, of each, *a drachm*; Spanish White Wine, *two pounds*. Digest for seven days, shaking the mixture frequently, and strain.”

*Syn.* Vin d'Aloé (*F.*), Vino Aloetico (*I.*).

Wine is an excellent solvent of Aloes, and, therefore, the two latter formulæ contain all the virtues of the remedy in a more concentrated state than the formula of the London College. The sand ordered by the London College is intended to facilitate the pulverization of the aloes; but, although it does not in the least affect the solution, yet, it is seldom used; for by cutting the aloes into small pieces, and exposing them to the air, they become sufficiently pulverulent. It is preferable, when the formula of the London College is ordered, to use the water boiling hot; and, after the solution is cold, to add the spirits.

*Medical properties and uses.* — Wine of aloes is an excellent warm purgative and stomachic. It has long been employed with benefit in cold phlegmatic habits, paralysis, gout, dyspepsia, and chlorosis. The dose is from  $\text{f}\text{ʒj.}$  to  $\text{f}\text{ʒij.}$  to act as a stomachic, and from  $\text{f}\text{ʒss.}$  to  $\text{f}\text{ʒij.}$  to produce purging.

Aloes are advantageously combined with alkalies; and in this state I have long been in the habit of employing a wine containing aloes and myrrh, in dyspepsia and chlorosis; and, also, in that affection of the mesenteric glands in children which produces a tumid and tense abdomen. The following is the formula I employ, which was copied, with some modification, from a very old pharmacopœia, that accidentally fell into my hands; but of the date of which, I have unfortunately preserved no memorandum.



R Sodæ subcarbonatis ℥iij.  
 Ammoniae carbonatis ℥ivss.  
 Myrrhæ ʒvj.  
 Aloes extracti ʒvj.  
 Vini Albi (*Sherry, Anglice*) f℥xxiv.  
 Macera per dies septem et cola.

The dose is from one fluid drachm to half a fluid ounce.

VINUM COLCHICI, Lond. *Wine of Colchicum.*

“Take of the Fresh Root (*bulb*) of Colchicum bruised, *two ounces*; Proof Spirit, *twelve fluid ounces*; Distilled Water, *twenty fluid ounces*. Macerate for fourteen days, and filter.”

The medicinal power of the bulb of Colchicum depends on an alkaline principle which, as it is also found in white hellebore, *Veratrum album*, has been named Veratria. It is combined, in the bulbs of Colchicum, with a large quantity of starch, some mucilage and gluten, and much water, the proportion of the last of which differs considerably in different bulbs, according to the moist or dry nature of the soil in which the bulbs have grown or been reared, and the season of the year, in which they are dug up for use. On these considerations we must object to the above formula, inasmuch as the strength of the preparation must always be uncertain when the recent bulb is employed. The quantity of water, also, which the bulb contains in its fresh state, tends to dilute the spirit to a greater degree than is intended in the formula; and, consequently, to weaken its solvent power over the veratria, which is very sparingly soluble in water. When, on the contrary, the dried bulb, taken up at a proper season, is employed, as the proportions of starch, gum, and gluten are nearly always the same, the strength of the preparation is more likely to be uniform. But, independent of the state of the bulb, the London formula is injudicious in substituting diluted spirit for sherry wine, which is a much better solvent of the veratria, owing to the tartaric acid of the tartrate of potass, and a small quantity of acetic acid which it contains. The following is the formula which we recommend for the preparation of the Wine of *Colchicum*. Take of the Bulbs of Colchicum (dug up in July or August) sliced transversely, and dried without heat, or at a temperature not exceeding 110°, *one ounce and a half*; pulverise them and pour upon the powder put into a glass-bottle, *twelve ounces* of good Sherry Wine. Agitate the mixture twice a day for seven days, and then filter for use.

*Medical properties and uses.*—Wine of Colchicum, whether prepared according to the formula of the London College, or with the dried bulb and with wine, is a powerful sedative and purgative. It is administered with great advantage in inflam-



matory and painful nervous affections, connected with a disordered state of the liver; as for example, gout and acute rheumatism, diminishing the force and frequency of the pulse, allaying the pain, and cutting short the paroxysm: but, it rarely produces a permanently favourable result without stimulating the duodenum and biliary ducts, and producing a copious discharge of bilious stools. It is apt to nauseate the stomach, a property which is greatly subdued by combining it with magnesia: but in some habits even this does not correct its nauseating qualities, and its use is followed by great faintness and depression of nervous power. The dose of wine of Colchicum is from  $\mathfrak{mxxx.}$  to  $\mathfrak{f3jss.}$  given in conjunction with Magnesia in the effervescing draught, or with infusion of Cinchona bark, Gentian root, or any other bitter.

The seeds of Colchicum possess the same medicinal properties as the bulb; and the wine may be made by digesting *two ounces* of the unbruised seeds in *two pints* of sherry wine, for eight days.

VINUM GENTIANÆ COMPOSITUM, Edin. *Compound of Wine of Gentian.*

“Take of Gentian Root, *half an ounce*; Cinchona Bark, *one ounce*; Orange Peel dried, *two drachms*; Canella Alba, *one drachm*; Proof Spirit, *four ounces*; Spanish White Wine, *two pounds and a half*. First pour the Proof Spirit on the root and the bark sliced and bruised; and after twenty-four hours add the Wine; then macerate for seven days, and strain.”

*Syn.* Vin de Gentiane composé (*F.*), Vino di Genziana composto (*I.*).

This wine, when newly prepared, is stomachic and tonic; but, by keeping, it is very apt to become acescent. The dose is from  $\mathfrak{f3iv.}$  to  $\mathfrak{f3vj.}$  given two or three times a day.

VINUM IPECACUANHÆ, Lond. *Wine of Ipecacuanha.*

“Take of Ipecacuanha Root bruised, *two ounces*; Proof Spirit, *twelve fluid ounces*; Distilled Water, *twenty fluid ounces*. Macerate for fourteen days, and filter.”

*Edinburgh.*

“Take of the Root of Ipecacuanha bruised, *one part*; Spanish White Wine, *fifteen parts*. Macerate for seven days, and filter through paper.”

*Dublin.*

“Take of the Root of Ipecacuanha bruised, *two ounces*; Spanish White Wine, *two pints*. Digest for seven days, then filter.”

*Syn.* Vin d’Ipecacuanha (*F.*), Vino con Ipecacuana (*I.*).

From my trials, I find that a pint of sherry wine takes up 100 grains of ipecacuanha, which is the larger proportion of the soluble matter contained in an ounce of the root; and as



the active part of the root, or emetin<sup>1</sup>, is more soluble in acetic acid than in any other menstruum, the acescency of the wine is no objection. Dr. Irvine says, that 30 grains of the root administered in fʒij. of vinegar produced only some loose stools; but, if acetate of emetin produces vomiting, what is the cause of the inertness of this mixture? Does the solution of the starch, and the gum in the acid form a substance, which obtunds the action of the acetate of emetin? As an emetic, Ipecacuanha wine is equally efficacious with antimonial wine, and at the same time is milder in its operation, and is, therefore, better adapted for infants. For this purpose a tea-spoonful, or fʒss. is given for a dose, and repeated every ten minutes till it operate. In smaller doses, from m̄x. to m̄xx., it answers the same purposes as the powder, and is given in coughs, diarrhœa, dysentery, and other complaints in which a determination to the skin is indicated.

VINUM NICOTIANÆ TABACI, Edin. *Wine of Tobacco.*

“Take of Tobacco Leaves, *one part*; Spanish White Wine, *twelve parts*. Macerate for seven days, and filter through paper.”

This is the only form in which tobacco can be conveniently exhibited as an internal remedy. It is given to produce diuretic and antispasmodic effect in dropsies, colica pictonum and ileus. The dose is from m̄x. to m̄xxx. in any proper vehicle.

VINUM OPII, Lond. Edin. *Wine of Opium.*<sup>2</sup>

“Take of Extract of Opium, *an ounce*; Cinnamon Bark bruised, Cloves bruised, of each, *a drachm*; Proof Spirit, *six fluid ounces*; Distilled Water, *ten fluid ounces*. Macerate for eight days (seven days, *Edin.*), and filter.”

*Syn.* Vin d'Opium aromatique (*F.*), Vino aramo opiato (*I.*).

The aromatics which this preparation contains are supposed to modify the action of the opium, and prevent the disturbance of the brain and nervous system, which the simple tincture is apt to induce in nervous habits, and where the head is much affected. It is intended to supply the place of the liquid laudanum of Sydenham; but that preparation contained double the quantity of opium, and ʒj. of Saffron, which is altogether omitted in the formulæ of the colleges.<sup>3</sup> Mr. Ware

<sup>1</sup> Emetin was discovered by M M. Dumas and Pelletier; and according to their analysis it consists of Carbon 64·57, Azote 4·30, Hydrogen 7·77, and Oxygen 22·95, in 100 parts.

<sup>2</sup> Laudanum liquidum Sydenhami, P. L. 1020.

<sup>3</sup> The following is Sydenham's formula. R Vini Hispanici lb. j. Opii ʒij. Croci ʒj. pulv. Cinnamomi et Caryophyllarum ā. ā. ʒj. Infundantur simul in B. M. per duas vel tres dies, donec liquor debitam consistentiam acquirat: Colitura servetur pro usu. *Sydenhami opera omnia*, Lond. 1705, p. 147.



introduced the use of this preparation made with wine as a local application in the second stage of ophthalmia; when the inflammatory symptoms have subsided, and the vessels of the conjunctiva remain turgid with red blood. Two or three drops are poured into the eye every morning, until the redness be removed.

VINUM RHEI, Edin. *Wine of Rhubarb.*

“Take of Rhubarb Root sliced, *two ounces*; Canella Bark bruised, *a drachm*; Proof Spirit, *two ounces*; Spanish White Wine, *fifteen ounces*. Macerate for seven days, and filter through paper.”

This wine, when newly prepared, has the same properties, and may be applied to the same uses as the tincture, but it is liable to undergo decomposition. The dose is from fʒss. to fʒj. or more.

VINUM VERATRI, Lond. *Wine of White Hellebore.*

“Take of White Hellebore Root bruised, *eight ounces*; Proof Spirit, *a pint*; Distilled Water, *a pint and a half*. Macerate for fourteen days, and filter.”

A solution of White Hellebore in Wine is supposed to form one of the ingredients of the Eau Medicinale; and this opinion, although it is unfounded, inasmuch as White Hellebore is not an ingredient of the French nostrum, yet is correct in some degree; inasmuch as the active principle of Colchicum, the plant used in preparing Eau Medicinale, is *Veratria*, which is also the active principle of White Hellebore, in which it is combined with gallic acid.<sup>1</sup> I have no doubt that a vinous preparation of White Hellebore, exhibited with due caution, would answer every purpose of the *Wine of Colchicum*. Wine of White Hellebore is seldom employed.

The dose is ten minims gradually increased to thirty.

## ACETICA.

### PREPARATIONS OF VINEGAR.

VINEGAR is capable of dissolving all those proximate principles of plants which are soluble in water; and it is, further, found to extract, more completely, the alkaloid principles on which the efficacy of Squill and Colchicum depend. As a solvent, however, of vegetable matter, the use of Vinegar cannot be extended, for it destroys the medicinal properties of some vegetable principles, and does not accord with others in

<sup>1</sup> Veratria, like the other newly discovered vegetable alkalies, is a compound substance; and, according to the analysis of Dumas and Pelletier, consists of Carbon 66.75, Azote 5.04, Hydrogen 8.54, and Oxygen 19.60, in 100 parts.



virtue. Medicated vinegars are very apt to spoil, notwithstanding the addition of spirit which is ordered; and, therefore, they should be made in small quantities only at a time, and preserved in well-stopped, glass-bottles.

**ACIDUM ACETICUM AROMATICUM**, Edin. *Aromatic Vinegar*.

“Take of Rosemary-tops dried, Sage Leaves dried, of each, *one ounce*; Lavender Flowers dried, *half an ounce*; Cloves bruised, *half a drachm*; Distilled Vinegar, *two pounds*. Macerate for seven days, and filter the expressed liquor through paper.”

*Syn.* Vinaigre antiseptique (F.), Gewürzessig (G.), Aceto antisettico (I.).

This preparation has a pleasant, pungent, aromatic odour. It is a solution of the volatile oils of the substances employed in vinegar, and is a grateful perfume in sick-rooms; but has no right to be regarded as a neutralizer of febrile, or other contagions. It is nearly the same as the old preparation, known under the name of *Thieves' Vinegar*; with the deficiency of *Lignum Rhodii*, *Santalum album*, *Fœniculi Semina*, *Absinthium*, and *Mentha piperita*; but these add nothing to its medicinal powers. The dose is from f3ss. f3j. in any bland fluid.

**ACIDUM ACETICUM CAMPHORATUM**, Edin.  
**ACIDUM ACETICUM CAMPHORATUM**, Dub. *Camphorated Acetic Acid*.

“Take of Acetic Acid, *six ounces*; Camphor, *half an ounce*. Rub the camphor to powder with the assistance of a little Alcohol; then dissolve it in the Acid.

*Syn.* Acide acétique Camphré (F.), Aceto Canforato (I.).

The strong acetic acid readily dissolves a considerable portion of camphor; and forms a very highly pungent and stimulating perfume, which, snuffed up the nostrils, is useful in syncope and nervous languors. Owing to its extreme volatility when well prepared, it requires to be preserved in phials closely fitted with ground-glass stoppers.

**ACETUM COLCHICI**, Lond. *Vinegar of Meadow Saffron*.

“Take of Fresh Meadow Saffron Root (bulb) sliced, *an ounce*; Acetic Acid (distilled vinegar), *a pint*; Proof Spirit, *a fluid ounce*. Macerate the Meadow Saffron Root with the Vinegar in a covered glass vessel for twenty-four hours; then express, and set the liquor aside that the fæculencies may subside; lastly, add the Spirit to the clear liquor.”

The bulb of the meadow-saffron is the part intended to be ordered, not the root. When dug up in July, the bulb contains that principle for which it is employed in the greatest perfection, and of which vinegar is a good solvent. This



solution is now introduced as a better form of preserving the virtues of the remedy than the oxymel. It is given as a diuretic in ascites and hydrothorax; but is less to be depended on than the squill. Like the preparation termed *Vinum Colchici*, it should be prepared with the dried bulb, in which case the proportion should be six drachms to a pint of vinegar. It is employed in gout. The dose is from f3ss. to f3j. united with any bland fluid.

ACETUM SCILLÆ, Lond. *Vinegar of Squill.*

“Take of Fresh Squill Root (bulb) dried, *a pound*; Acetic Acid (distilled vinegar), *six pints*; Proof Spirit, *half a pint*. Macerate the Squill Root (bulb) in the Vinegar with a gentle heat, in a covered vessel, for twenty-four hours; then express the liquor, and set it aside that the fæculencies may subside; lastly, add the Spirit to the clear liquor.”

ACIDUM ACETICUM SCILLITICUM, Edin. *Vinegar of Squill.*

“Take of Squill Root (bulb) dried, *one ounce*; Distilled Vinegar, *fifteen ounces*; Alcohol, *one ounce and a half*. Macerate the Squill with the Acid for seven days; then express the liquor, and add to it the Alcohol: and when the fæculencies have subsided, pour off the clear fluid.”

ACETUM SCILLÆ, Dub. *Vinegar of Squill.*

“Take of Fresh Squill Root (bulb) dried, *half a pound*; Wine Vinegar, *three pints*; Rectified Spirit, *four fluid ounces*. Digest the Squill with the Vinegar for four days in a glass vessel, with frequent agitation; then express the Vinegar, and after the fæculencies have subsided, add to it the Spirit.”

*Syn.* Vinaigre scillitique (F.), Meerzwiebeleessig (G.).

Vinegar extracts and holds in solution the *Scillitina*, the active principle of the squill, upon which its efficacy as a remedy depends. This preparation has long been used as an expectorant and diuretic in chronic catarrh, humoral asthma, and dropsies. The dose is from f3ss. to f3ij. given in cinnamon or mint-water. In large doses it produces vomiting; and is occasionally used as an emetic in the above diseases, when the stomach is loaded.

When kept, the Vinegar of Squill deposits a precipitate, which consists of citrate of lime and tannin.<sup>1</sup>

Officinal preparations. *Oxymel Scillæ*, L. *Syrupus Scillæ*, E.

<sup>1</sup> Vogel. *Annales de Chimie*, vol. lxxxiii. p. 157.



## MELLITA.

## PREPARATIONS OF HONEY.

A MORE correct knowledge of the operation of those medicinal substances which have been named *Balsamic* or *Pectoral*, has set aside the high opinion which formerly prevailed of the efficacy of honey as a remedy in pulmonary diseases. It is, however, still employed in pharmacy, and has some advantages over syrup, particularly where it is to be employed as a local application; but, for internal purposes, its use is to a certain degree limited, owing to the unpleasant effects, which it produces on the bowels of some individuals. The Edinburgh College has altogether rejected this class of preparations, but a few of them are retained by the London and the Dublin Colleges. They are not apt to spoil, and, therefore, require less care to preserve them than the syrups.

MEL DESPUMATUM, Lond. Dub. *Clarified Honey.*

“Melt the Honey in a water-bath; then remove the scum.”

*Syn.* Miel despumé (*F.*), Geschäumter Honig (*G.*), Mele Schiumato (*I.*).

By thus liquefying honey, the wax, which it may have retained when expressed from the comb, rises to the surface; and at the same time any sand or other impurities, with which it may have been fraudulently mixed, fall to the bottom or rise with the wax, and are easily separated. The specific gravity of purified honey is 1.31. It is chiefly employed for forming the other preparations into which honey enters. It is less apt to gripe than the crude honey, owing, probably, to its being less liable to ferment. It is, undoubtedly, more agreeable to the taste and smell than crude honey.

Officinal preparations. *Confectio Rutæ*, L. *Linimentum Æruginis*, L. *Mel Boracis*, L. *Mel Rosæ*, L. D. *Oxymel Colchici*, D. *Oxymel Scillæ*, L. D. *Oxymel Simplex*, L. D.

MEL BORACIS, Lond. *Honey of Borax.*

“Take of Sub-Borate of Soda powdered, a drachm; Clarified Honey, an ounce. Mix them.”

This is a cooling, detergent, useful application to the tongue and fauces in aphthous affections. Dissolved in water it forms an excellent gargle for allaying the pain attending mercurial salivations.

MEL ROSÆ, Lond. *Rose Honey.*<sup>1</sup>

<sup>1</sup> Mel Rosarum, P. L. 1720. Mel rosaceum, P. L. 1745.



“ Take of the Petals of the Red Rose dried, *four ounces* ; Boiling Water, *three pints* ; Clarified Honey, *five pounds*. Macerate the Petals in the water for six hours ; then to the filtered liquor add the Honey, and boil it down to a proper consistence by means of a water-bath.”

*Dublin.*

“ Take of the Petals of Red Rose-buds, dried and freed from their claws, *four ounces* ; Boiling Water, *three pints* ; Honey, *five pounds*. Macerate the Petals in the Water for six hours ; then mix the Honey with the strained liquor, and boil the mixture to the consistence of a syrup, taking off the scum.”

*Syn.* Miel rosat (*F.*), Rosenhonig (*G.*), Mele rosato (*I.*).

This honey has the pleasant flavour of the Rose, and a slight degree of astringency. In making it, the clarified honey ordered by the London College is to be preferred. It is chiefly employed as an adjunct to detergent and astringent gargles.”

OXYMEL SIMPLEX, Lond. Dub. *Oxymel.*<sup>1</sup>

“ Take of Clarified Honey, *two pounds* ; Acetic Acid (distilled vinegar), *one pound*. Boil them in a glass vessel, by a gentle heat, to a proper consistence.”

*Syn.* Oxymel (*F.*), Essighonig (*G.*), Ossimele (*I.*).

Simple oxymel in doses of fʒj. or more, dissolved in barley-water, forms a pleasant and cooling beverage in fevers and inflammatory affections ; but in some individuals it has a gripping quality like honey, which prevents it from being generally used in these affections. It is often added to gargles in cyananche tonsillaris, and is a common vehicle of other remedies in catarrhal complaints. The Dublin College orders it to be prepared with unclarified honey, skimming it during the boiling ; but the London directions are to be preferred.

OXYMEL COLCHICI, Dub. *Oxymel of Meadow Saffron.*

“ Take of the Fresh Root (bulb) of Meadow Saffron cut into thin slices, *one ounce* ; Distilled Vinegar, *a pint* ; Clarified Honey, *two pints*. Digest the Colchicum with the Vinegar, in a glass vessel for two days ; then to the liquor strongly expressed from the Root, add the Honey ; and lastly boil down the mixture to the consistence of a syrup, frequently stirring it during the boiling with a wooden spoon.”

The active matter of the Colchicum is apt to be injured by the boiling ; and therefore this preparation is very uncertain in point of strength. It is given in humoral asthma, and in

<sup>1</sup> Mel acetatum, P. L. 1787.



dropsies. The dose is fʒj. gradually increased to fʒj., given in a cupful of gruel, twice a day.

OX'YMEL SCILLÆ<sup>1</sup>, Lond. Dub. *Oxymel of Squill.*

“Take of Clarified Honey, *three pounds*; Vinegar of Squill, *two pints*. Boil in a glass vessel, over a gentle fire, to a proper consistence.”

*Syn.* Meerzwiebelhonig (G.).

Oxymel of Squill is principally employed as an expectorant, and as such is very useful in humoral asthma, and chronic coughs, in doses of from fʒss. to fʒij. It is generally given in some aromatic distilled water, to prevent the nausea which it is apt to induce: in larger doses it is given to excite vomiting, and at the same time clear the chest, in hooping cough.

When kept for a considerable time, this oxymel lets fall a precipitate which has the aspect of crystallized honey. Vogel found it to consist of citrate of lime, tannin, and honey.<sup>2</sup>

---

## SYRUPI.

### SYRUPS.

THESE are saturated solutions of Sugar in Water, either simple, or united with some vegetable principle, with the view either to colour, flavour, or medicinal virtue: but for the last intention, this is perhaps the worst of all forms for obtaining the medicinal qualities of substances: and, therefore, as syrups seldom possess much activity, they are chiefly employed to render more active remedies palatable. Upon the whole, however, they are not well adapted even for this purpose, few persons thinking that sweetness renders a nauseous drug more palatable; and, with a few exceptions, they might be properly rejected from the Pharmacopœias.

In making syrups, refined sugar should always be employed, or, if coarser sugar be used, the syrup should be clarified, by beating to a froth the white of eggs with a small portion of water, and adding it to the solution of sugar and water before boiling. The albumen coagulates as the syrup boils, and, involving the impurities which the sugar contained, rises to the surface in the form of a scum, which must be carefully removed. If too much sugar be used, or if the syrup be too long boiled, the sugar soon crystallizes; and if it be in too

---

<sup>1</sup> Oxymel scilliticum, P. L. 1720, 1745.

*Annales de Chimie*, vol. lxxxiii. p. 157.



small proportion, the syrup quickly ferments, and becomes acescent. The most certain test of the proper consistence of a syrup is its specific gravity, which, when cold, should be 1.385. But, however well prepared, syrups are apt to ferment when kept in a temperature above 60°; and, therefore, the following direction relative to their preservation is given by the London College.

“Let Syrups be preserved in a place the temperature of which never exceeds 55°.”<sup>1</sup>

SYRUPUS SIMPLEX, Lond. *Syrup.*

“Take of Refined Sugar, *two pounds and a half*; Water, *a pint*. Dissolve the Sugar in the water by means of a water-bath; then set it aside for twenty-four hours; take off the scum, and, if there be any fæces, pour off the clear part from them.”

SYRUPUS SIMPLEX, Edin. *Simple Syrup.*

“Take of Purified Sugar, *fifteen parts*; Water, *eight parts*. Dissolve the Sugar in the Water by a gentle heat, and boil it a little so as to form a syrup.

SYRUPUS, Dub. *Syrups.*

“In making Syrups, for which neither the weight of the sugar nor the mode of dissolving it is specified, the following rule is to be observed:

“Take of Refined Sugar reduced to a fine powder, *twenty-nine ounces*; the liquor prescribed, *one pint*. Add the Sugar by degrees, and digest with a moderate heat, in a close vessel, until it is dissolved, frequently stirring it; set the solution aside for twenty-four hours, take off the scum, and pour off the syrup from the fæces, if there be any.”

*Syn.* Sirop (*F.*), Einfacher syrup (*G.*), Sciroppo (*I.*).

Simple Syrup, when properly prepared, should be inodorous, sweet, thickish, nearly colourless, and perfectly transparent.

SYRUPUS ACETI, Edin. *Syrup of Vinegar.*

“Take of Vinegar, *five parts*; Refined Sugar, *seven parts*. Boil, so as to form a syrup.”

*Syn.* Syrop d'acide acetique (*F.*), Sciroppo acetico (*I.*).

This syrup is very liable to undergo decomposition: it should, therefore, be made in small quantities only at a time.

It may be used for sweetening barley-water or gruels, in fevers and inflammatory diseases.

SYRUPUS ALLII, Dub. *Syrup of Garlic.*

“Take of Garlic Root (bulb) sliced, *a pound*; Boiling Water, *two pints*. Macerate the garlic in the water, in a

<sup>1</sup> Dr. Macculloch informs us, that by the addition of a small quantity of *sulphate of potass*, or of the *orymuriate of potass*, which is a tasteless salt, the fermentation of syrups may be effectually prevented. See *Essay on Wine*.



covered vessel, for twelve hours: then let the sugar be added to the strained liquor, and a syrup formed."

This is a very disagreeable syrup; but contains the virtues of the garlic in a sufficient degree to entitle it to some attention.

**SYRUPUS ALTHÆÆ**, Lond. *Syrup of Marsh Mal-*  
*lows.*

"Take of Fresh Marsh Mallow Root bruised, *half a pound* Refined Sugar, *two pounds*; Water, *four pints*. Boil down the Water with the Marsh Mallow Root to one half, and express the liquor when it is cold. Set it aside for twenty-four hours, that the fæces may subside; then decant off the clear liquor, and, having added to it the sugar, boil down to a proper consistence."

**SYRUPUS ALTHÆÆ OFFICINALIS**, Edin. *Syrup of Marsh*  
*Mallows.*

"Take of Fresh Root of Marsh Mallows sliced, *one part*; Water, *ten parts*; Refined Sugar, *four parts*. Boil the Water with the Root down to one half and, expressing it strongly, strain. Put aside the strained liquor, and, when the fæces have subsided, add to it the sugar; then boil so as to form a syrup."

*Syn.* Sirop d'Althea (*F.*), Althee syrup (*G.*), Sciroppo d'Altea (*I.*).

This is a solution of vegetable mucus and syrup, and is thence supposed to possess demulcent properties; but these are very trivial; and owing to the small proportion of sugar it contains, it very soon suffers spontaneous decomposition.

**SYRUPUS AURANTIORUM**, Lond. *Syrup of*  
*Oranges.*

"Take of Fresh Orange-peel, *two ounces*; Boiling Water, *a pint*; Refined Sugar, *three pounds*. Macerate the Peel in the Water for twelve hours in a covered vessel; then pour off the liquor, and add to it the sugar."

**SYRUPUS CITRI AURANTII**, Edin. *Syrup of Orange.*

"Take of the Fresh Peel of Seville Oranges, *three ounces*; Boiling Water, *one pound and a half*; Refined Sugar, *three pounds*. Macerate the Peel in the Water in a covered vessel for twelve hours; then add the Sugar to the strained liquor, and expose it to a gentle heat so as to form a syrup."

**SYRUPUS AURANTII**, Dub. *Syrup of Orange.*

"Take of the Fresh Peel of Seville Oranges, *eight ounces*; Boiling Water, *six pints*. Macerate for twelve hours in a covered vessel, and dissolve as much Sugar in the filtered liquor as will form a syrup."

*Syn.* Sirop d'ecorce d'orange (*F.*), Pomeranzenschaleusyrup (*G.*), Sciroppo di Corteccia di Arancio (*I.*).

The quantity of water ordered by the Edinburgh College is



too great; particularly as the application of a degree of heat sufficient to evaporate part of it would dissipate also the flavour of the Orange-peel, for which the syrup is chiefly valued. A syrup equally agreeable and efficacious may be made by adding f℥j. of tincture of orange-peel to a pint of simple syrup.

**SYRUPUS COLCHICI AUTUMNALIS**, Edin. *Syrup of Meadow Saffron.*

“Take of Fresh Meadow Saffron Root (bulb) cut into thin slices, *one ounce*; Distilled Vinegar, *sixteen ounces*; Refined Sugar, *twenty-six ounces*. Macerate the Root in the Acid for two days, shaking the vessel occasionally; then expressing gently, strain the liquor, and to it add the sugar; lastly, boil a little so as to form a syrup.”

With the substitution of Syrup for Honey, this preparation is similar to the oxymel. The dose is f℥ij. increased gradually to f℥ss. or more.

**SYRUPUS CROCI**, Lond. *Syrup of Saffron.*

“Take of Saffron, *an ounce*; Boiling Water, *a pint*; Refined Sugar, *two pounds and a half*. Macerate the Saffron in the Water for twelve hours, in a slightly covered vessel; then filter the liquor, and add to it the sugar.”

*Syn.* Sirop de Safran (*F.*), Safransyrup (*G.*).

This syrup is cordial in a small degree; but it is chiefly valued on account of its beautiful colour.

**SYRUPUS DIANTHI CARYOPHYLLI**, Edin. *Syrup of the Clove July Flower.*

“Take of recent Petals of the Clove July Flower, freed from their claws, *one part*; Boiling Water, *four parts*; Refined Sugar, *seven parts*. Macerate the Petals in the Water for twelve hours; then add the Sugar to the strained liquor, and dissolve it with a gentle heat.”

**SYRUPUS CARYOPHYLLI RUBRI**, Dub. *Syrup of Clove July Flower.*

“Take of Fresh Petals of the Clove July Flower, freed from the claws, *two pounds*; Boiling Water, *six pints*. Macerate for twelve hours in a glass vessel; and dissolve a sufficient quantity of sugar in the strained liquor to make a syrup.”

This syrup is valued for the rich colour, and the agreeable flavour of the flowers, which it possesses in perfection when well prepared. Alkalies change the colour to green, and form a test of the genuineness of the syrup; for they do not produce this effect on a counterfeit syrup, made of an infusion of cloves and coloured with cochineal, which is sometimes sold for it; but the one is as good for medical use as the other.

**SYRUPUS LIMONUM**, Lond. *Syrup of Lemon.*



“Take of strained Lemon Juice, *a pint*; Refined Sugar, *two pounds*. Dissolve the Sugar in the Lemon-juice, in the manner directed for syrup.”

SYRUPUS CITRI MEDICÆ, Edin. *Syrup of Lemons*.

“Take of Lemon-juice, strained after the fæces have subsided, *three parts*; Refined Sugar, *five parts*. Dissolve the sugar.”

SYRUPUS LIMONIS, Dub. *Syrup of Lemon*.

“Take of Expressed Lemon-juice, *two pints*. As soon as the fæces have subsided put it into a matrass, and immerse it in Boiling Water for a quarter of an hour: when cold, strain it through a sieve, and make it into a syrup.”

*Syn.* Zitronensaftsyrup (G.).

This is an agreeable syrup for acidulating barley water or other drinks in febrile diseases. It is also an useful adjunct to gargles in inflammatory sore-throat.

SYRUPUS MORI, Lond. *Syrup of Mulberry*.

“Take of Strained Mulberry-juice, *a pint*; Refined Sugar, *two pounds*. Dissolve the Sugar in the Mulberry-juice in the manner ordered for syrup.”

*Syn.* Maulbeersyrup (G.).

This syrup is used for the same purposes as the syrup of lemons, and has besides the advantage of colour.

SYRUPUS OPII, Dub. *Syrup of Opium*.

“Take of the Watery Extract of Opium, *eighteen grains*; Boiling Water, *eight ounces*. Macerate until the Opium be dissolved; then add Sugar so as to make a syrup.”

*Syn.* Sirop d'opium (F.), Opiumsyrup (G.), Sciroppo di oppio (I.).

The watery extract of opium contains the virtues of the opium, without its stimulant qualities. Each ounce of this syrup contains gr. j. of the watery extract. It is a useful anodyne for allaying the irritation which keeps up the cough in catarrh, after the inflammatory symptoms are abated; and for procuring sleep, in the diseases of children.

SYRUPUS PAPAVERIS, Lond. *Syrup of Poppy*.

“Take of the dried Capsules of the Poppy bruised and freed from the seeds, *fourteen ounces*; Refined Sugar, *two pounds*; Boiling Water, *two gallons and a half*. Macerate the capsules in the water for twelve hours; then boil down the whole in a water-bath to one gallon, and express strongly. Boil the liquor again down to two pounds, and strain it while it is hot. Set it aside twelve hours that the fæces may subside; then boil down the clear liquor to one pint, and add the sugar, in the manner ordered for making syrup.”

SYRUPUS PAPAVERIS SOMNIFERI, Edin. *Syrup of White Poppy*.

“Take of the Capsules of the White Poppy dried, and



freed from the seeds, *one part*; Boiling Water, *fifteen parts*; Refined Sugar, *two parts*. Macerate the sliced Capsules in the water for twelve hours; then boil until a third part only of the liquor remain; and, expressing strongly, strain the decoction. Boil the strained liquor to one-half, and again strain it; lastly, having added the Sugar, boil it for a short time, so as to form a syrup."

SYRUPUS PAPAVERIS ALBI, Dub. *Syrup of White Poppies.*

"Take of the Capsules of the White Poppy, gathered before they are ripe, dried and freed from the seeds, *a pound*; Boiling Water, *three pints*. Slice and bruise the Capsules; then pour over them the water, and macerate for twelve hours; express the liquor, and evaporate it by a gentle heat to a pint; strain through a thin linen cloth, and set it aside six hours that the fæces may subside; finally, add sugar to the clear liquor that it may make a syrup.

*Syn.* Sirop de Pavot blanc (*F.*), Syrup von Weissen Mohn (*G.*), Sciroppo di Papaveri bianchi (*I.*).

The narcotic principle of the Poppy is taken up by the water, but it is very probable that any variation of the degree of heat necessary to produce the evaporation, will alter in a considerable degree the nature of the extract, and must, consequently, make the syrup differ in point of strength. It ferments more readily than most other syrups, but loses its narcotic property when it becomes acescent. One fluid ounce of it contains about one grain of extract.

*Medical properties and uses.*—Syrup of Poppy is an useful anodyne for allaying the violence of the cough in catarrh, for easing pain, and procuring sleep in children's diseases, to whom, however, it should not be given when it is in a state of fermentation. The dose is from fʒj. to fʒj. according to the age of the patient.

SYRUPUS RHŒADOS, Lond. *Syrup of the Red Poppy.*

"Take of the Recent Petals of the Red Poppy, *a pound*; Boiling Water, *a pint and two fluid ounces*; Refined Sugar, *a pound and a half*. To the Water, heated in a water-bath, add gradually the Petals of the Red Poppy, stirring them occasionally; then, having removed the vessel, macerate for twelve hours; press out the liquor, and set it aside, that the impurities may subside; lastly, add the sugar in the manner directed for making syrup."

SYRUPUS PAPAVERIS ERRATICI, Dub. *Syrup of the Red Poppy.*

"Take of the Fresh Petals of the Red Poppy, *a pound*; Boiling Water, *twenty fluid ounces*. Add the Flowers gradually to the Boiling Water; then, having removed the vessel



from the fire, macerate in a lower heat or twelve hours; express the liquor, and set it aside that the fæces may subside; finally, let the sugar be added so as to make a syrup."

*Syn.* Sirop de Coquelicot (*F.*), Klapprosen-syrup (*G.*).

By attending strictly to the directions of either of the above formulæ, the petals yield their fine rich colour, for which alone the syrup is valued.

SYRUPUS RHA'MNI<sup>1</sup>, Lond. *Syrup of Buckthorn.*

"Take of the Fresh Juice of Buckthorn Berries, *four pints*; Ginger Root sliced, Pimenta Berries bruised, of each, *half an ounce*; Refined Sugar, *three pounds and a half*. Set apart the Juice for three days that the fæces may subside, then strain it. To a pint of the defecated Juice add the Ginger Root, and Pimenta Berries; then macerate by a gentle heat for four hours, and strain. Boil the remainder of the Juice down to a pint and a half; mix the liquors, and add the sugar in the manner directed for making syrup."

SYRUPUS RHAMNI CATHARTICI, Edin. *Syrup of Buckthorn.*

"Take of the Clarified Juice of Ripe Buckthorn Berries, *two parts*; Refined Sugar, *one part*. Boil so as to form a syrup."

Of these two formulæ, that of the London College is to be preferred, as the addition of the Ginger and All-spice tends to cover the unpleasant taste of the Buckthorn Juice, and prevent the violent griping which it is apt to induce. It is a brisk cathartic; but owing to the unpleasantness of its operation, and the dryness of the mouth and fauces which it occasions, it is seldom used, except as a horse medicine. The dose is from fʒss. to fʒj. drinking freely of tepid demulcent fluids during its operation.

SYRUPUS RO'SÆ<sup>2</sup>, Lond. *Syrup of the Rose.*

"Take of the Petals of the Hundred-leaved Rose dried, *seven ounces*; Refined Sugar, *six pounds*; Boiling Water, *four pints*. Macerate the Rose Petals in the Water for twelve hours, and strain. Evaporate the strained liquor in a water-bath down to *two pints and a half*; then add the sugar so as to make a syrup."

SYRUPUS ROSÆ CENTIFOLIÆ, Edin. *Syrup of Damask Roses.*

"Take of the Fresh Petals of the Damask Rose, *one part*; Boiling Water, *four parts*; Refined Sugar, *three parts*. Macerate the Petals in the Water for twelve hours; then add the sugar to the strained liquor, and boil, so as to form a syrup."

<sup>1</sup> Syr. de Spina cervina, P. L. 1720. Syrupus Spinæ cervinæ, P. L. 1787.

<sup>2</sup> Syr. e Rosis siccis, P. L. 1720. Syr. Rosarum solutivus, P. L. 1745.



This syrup has none of the agreeable odour of the Rose, but possesses a weak purgative property; on which account it is given as a laxative in very delicate habits, and to infants. The dose is from fʒij. to fʒxij. or more.

SYRUPUS ROSÆ GALLICÆ, Edin. *Syrup of Red Roses.*

“Take of the Petals of the Red Rose dried, *one part*; Boiling Water, *nine parts*; Refined Sugar, *ten parts*. Macerate the Petals in the Water for twelve hours; then boil a little, and strain. Add the sugar to the strained liquor, and again boil a little, so as to form a syrup.”

*Syn.* Sirop de Roses rouges (F.), Rosen syrup (G.), Sciroppo di Rose Rosse (I.).

This syrup is a very weak astringent; and as such is added to astringent and stomachic infusions and gargles.

SYRUPUS SARSAPARILLÆ, Lond. *Syrup of Sarsaparilla.*

“Take of Sarsaparilla Root, *one pound*; Boiling Water, *one gallon*; Refined Sugar, *one pound*. Macerate the Root in the water for twenty-four hours; then boil down to *four pints*, and strain the liquor whilst it is hot; lastly, add the sugar, and evaporate to a proper consistence.”

This is a trifling preparation, and of no use but as an adjunct to the decoction of Sarsaparilla, which can be much better and more easily supplied by rubbing up a few grains of the extract, with some simple syrup.

SYRUPUS SCILLÆ MARITIMÆ, Edin. *Syrup of Squill.*

“Take of Vinegar of Squill, *four parts*; Refined Sugar powdered, *seven parts*. Dissolve the sugar by a gentle heat, so as to make a syrup.”

*Syn.* Sirop aceteux de Scille (F.), Sciroppo di Squilla marino acetoso (I.).

This syrup has the same properties as the Oxymel of Squill. The dose is from fʒj. to fʒij. given in any aromatic distilled water.

SYRUPUS SENNÆ, Lond. *Syrup of Senna.*

“Take of Senna Leaves, *two ounces*; Fennel Seeds bruised, *an ounce*; Manna, *three ounces*; Refined Sugar, *a pound*; Boiling Water, *a pint*. Macerate the Senna Leaves and the Fennel Seeds in the Water in a gentle heat for twelve hours; strain the liquor, mix with it the Manna and the Sugar; and boil to a proper consistence.”

SYRUPUS CASSIÆ SENNÆ, Edin. *Syrup of Senna.*

“Take of Senna Leaves, *two ounces*; Boiling Water, *a pound and a half*; Burnt Syrup, *eight ounces*. Macerate the Leaves in the Water in a covered vessel for four hours, and strain; then add the syrup, and boil with a gentle heat until the whole acquire the consistence of the burnt syrup.



*Dublin.*

“Take of Manna, Refined Sugar, of each, *a pound*; Senna Leaves, *half an ounce*; Boiling Water, *a pint*. Let the Senna Leaves be macerated in the Water in a covered vessel for twelve hours; then dissolve the manna and the sugar in the strained liquor.”

This syrup contains the purgative properties of the senna, and is chiefly intended for children.

SYRUPUS TOLUTA'NUS, Lond. *Syrup of Tolu.*<sup>1</sup>

“Take of Balsam of Tolu, *an ounce*; Boiling Water, *a pint*; Refined Sugar, *two pounds*. Boil the Balsam in the Water for half an hour in a close vessel, frequently stirring it, and strain the liquor when it is cold; then add the sugar so as to make a syrup.”

SYRUPUS TOLUIFERÆ BALSAMI, Edin. *Syrup of Tolu.*

“Take of Simple Syrup, *two pounds*; Tincture of Balsam of Tolu, *one ounce*. To the Syrup immediately after it is made, and before it is quite cold, add the Tincture gradually, frequently stirring.”

*Syn.* Syrop Balsamique (*F.*).

By following the London formula a more elegant and grateful syrup is obtained than that produced by the Edinburgh method: but the syrup ordered by the Edinburgh College is sufficient for all the uses to which it can be applied. It is whitish and turbid, owing to a partial decomposition of the tincture, which deposits its resin when mixed with the syrup. It is used to give a pleasant flavour to draughts and mixtures.

SYRUPUS VIOLÆ ODORATÆ, Edin. *Syrup of Violet.*

“Take of Flowers of the Odorous Violet, *two parts*; Boiling Water, *eight parts*; Refined Sugar, *fifteen parts*. Macerate the flowers in the water for twenty-four hours, in a covered glass or glazed earthenware vessel; then strain without expression, and add the sugar.”

SYRUPUS VIOLÆ, Dub. *Syrup of Violet.*

“Take of the Fresh Petals of the Violet, *two pounds*; Boiling Water, *five pints*. Macerate for twenty-four hours; then strain the liquor through fine linen with expression; and add a sufficient quantity of sugar to make a syrup.”

*Syn.* Sirop de Violettes (*F.*), Violensyrups (*G.*), Scioppo di Viole (*I.*).

This syrup has a deep blue colour, and a very agreeable flavour. The colour, however, which constitutes its chief value, is apt to suffer by keeping; and, thence, the syrup is often counterfeited with materials, the colour of which is more permanent, and which are more easily obtained. This fraud is easily detected by adding a little acid or alkali to a portion

<sup>1</sup> Syr. balsamicus, P. L. 1720.



of the suspected syrup: if it be genuine, the acid will change the blue colour to red, and the alkali to green; but if it be counterfeit, these changes will not take place, except in the case of the juice of red cabbage being substituted for violets: but in this case the fraud is a very innocent one.

*Medical properties and uses.* — This syrup acts as a gentle laxative when given to infants; but it is chiefly used as a test of the presence of acids and alkalies.

SYRUPUS ZINGIBERIS, Lond. *Syrup of Ginger.*

“Take of Ginger Root sliced, *two ounces*; Boiling Water, *a pint*; Refined Sugar, *two pounds*. Macerate the Ginger Root in the Water for four hours, and strain; then add the Sugar so as to make a syrup.”

*Dublin.*

“Take of Ginger Root bruised, *four ounces*; Boiling Water, *three pints*. Macerate for twenty-four hours; then strain the liquor, and add sugar so as to make a syrup.”

SYRUPUS AMOMI ZINGIBERIS, Edin. *Syrup of Ginger.*

“Take of Ginger Root powdered, *six drachms*; Boiling Water, *one pound*; Refined Sugar, *twenty-two ounces*. Macerate the Root in the Water, in a covered vessel, for twenty-four hours; then add the Sugar to the strained infusion, and dissolve by a gentle heat.”

*Syn.* Sirop de Gingembre (*F.*), Sciroppo d'Amomo Zenzero (*I.*).

This syrup is moderately stimulant and carminative; and is an useful adjunct to bitter and tonic infusions.

Officinal preparations. *Electuarium Catechu*, D. *Electuarium Opiatum*, D.

## CONFECTIONES.

### CONFECTIONS.

UNDER this title the London College comprehends the *Conserves* and *Electuaries* of its former Pharmacopœia, and of the present Edinburgh and Dublin Pharmacopœias. There is, however, a distinction between Confections or Conserves and Electuaries, which prevents them in strict propriety from being classed together; and which we shall point out, although at the same time we adhere to the title of the London College.

CONFECTIONS or CONSERVES consist of fresh vegetable matters beat into an uniform mass with refined sugar. They are designed to preserve, as nearly as possible, unaltered, the virtues or properties of recent vegetables; and to prevent the decomposition to which they would otherwise be liable; and



although several delicate flowers and fruits and juicy plants can be well preserved by this means, yet, this form of preparation is not adapted for all plants; and in almost all cases the active ingredients are injured by keeping in this state. As remedies, Confections scarcely ever possess great activity; and are chiefly useful as vehicles for the exhibition of more active substances. They should be kept in closely covered jars, in order to preserve their proper degree of moisture.

ELECTUARIES<sup>1</sup> are mixtures of vegetable and light earthy powders, combined by means of honey or of syrup so as to form masses of a moderate consistence. All substances of this description may, therefore, be made into Electuaries; but as the intention of this form of preparation is to render remedies as palatable as possible, those matters only can be employed to form electuaries, the taste of which is not too ungrateful to be covered by syrup or honey. They are more active remedies than Conserves; but still the more powerful vegetable substances cannot well be exhibited in this form, on account of their taste; and the metallic salts are too ponderous to remain suspended in either syrup or honey. In making electuaries, the degree of consistence must always be regulated by the nature of the substances which enter into them.

“In Conserves,” as Mr. Murray justly observes, “the addition of the saccharine matter is in much larger proportion, and is designed to preserve the vegetable matter in Electuaries, the syrup is designed merely to communicate the required form.”

The following general rule is given by the London College for restoring the consistence of confections and electuaries, when they have become hard by keeping.

“If Confections have become hard from long keeping, they are to be moistened with water, so as to restore their proper consistence.”

CONFECTIO AMYGDALARUM, Lond. *Confection of Almonds.*

“Take of Sweet Almonds, *an ounce*; Acacia Gum in powder, *a drachm*; Refined Sugar, *half an ounce*. Macerate the Almonds in Water to free them from their cuticle, then beat all the ingredients together, until they be thoroughly incorporated.”

This preparation is introduced as affording an easy and expeditious mode of preparing the almond mixture, the extemporaneous preparation of which is tedious. A little of this paste, triturated with a sufficient portion of water, immediately forms an emulsion. It may be made with Bitter Almonds.

---

<sup>1</sup> Ἐκλεκτον Hippocratis.



CONFECTIO AROMATICA, Lond. *Aromatic Confection.*

“Take of Cinnamon Bark, Nutmegs, of each, *two ounces*; Cloves, *an ounce*; Cardamom Seeds, *half an ounce*; Saffron dried, *two ounces*; Prepared Shells, *sixteen ounces*; Refined Sugar powdered, *two pounds*; Water, *a pint*. Rub the dry substances mixed together into a very fine powder; then add the water gradually, and mix until the whole be thoroughly incorporated.”

ELECTUARIUM AROMATICUM, Edin. *Aromatic Electuary.*

“Take of the Aromatic Powder, *one part*; Syrup of Orange, *two parts*. Mix and beat them well together so as to form an Electuary.”

*Dublin.*

“Take of Cinnamon Bark, Nutmegs, of each, *half an ounce*; Refined Sugar, Saffron, of each, *an ounce*; Lesser Cardamom Seeds husked, Cloves, of each, *two drachms*; Precipitated Chalk, *two ounces*; Syrup of Orange, *a sufficient quantity*. Reduce the aromatics separately to powder, and then mix them with the syrup.”

*Syn.* Electuare Aromatique (F.), Gewürzlatwerge (G.), Elettuario Aromatico (I.).

These combinations of aromatics are stimulant, and cordial. They are given with advantage in typhoid fevers, atonic gout, and nervous languors; either alone in the form of bolus, or combined with camphor and syrup of orange-peel, in the form of mixture. The dose is from grs. x. to ʒj. or more.

CONFECTIO AURANTIORUM, Lond. *Confection of Oranges.*

“Take of the External Rind of the Fresh Orange, separated by rasping, *a pound*; Refined Sugar, *three pounds*. Beat the Rind in a stone mortar with a wooden pestle; then add the Sugar, and continue the beating until they be thoroughly incorporated.”

CONSERVA AURANTII, Dub. *Conserve of Orange.*

“To the Rind of Seville Orange rasped off, add three times its weight of Refined Sugar, while beating it.”

CONSERVA CITRI AURANTII, Edin. *Conserve of Orange.*

“Grate off the Exterior Rind of Seville Oranges, beat it into a pulp, and during the beating add gradually three times its weight of Refined Sugar.”

This confection is gently stomachic, and is a pleasant vehicle for the exhibition of tonic powders.

CONFECTIO CASSIÆ, Lond. *Confection of Cassia.*

“Take of Fresh Cassia Pulp, *half a pound*; Manna, *two ounces*; Tamarind Pulp, *an ounce*; Syrup of Roses, *half a pound*. Bruise the Manna; then dissolve it in the Syrup,



by the heat of a water-bath, and, having mixed in the Pulp, evaporate down to a proper consistence."

ELECTUARIUM CASSIÆ FISTULÆ, Edin. *Electuary of Cassia.*

"Take of Cassia Pulp, *four parts*; Tamarind Pulp, Manna, of each, *one part*; Syrup of Damask Roses, *four parts*. Bruise the Manna in a mortar, and dissolve it in the Syrup, by means of a gentle heat; then add the Pulps, and by a continued heat reduce the mixture to a proper consistence."

ELECTUARIUM CASSIÆ, Dub. *Electuary of Cassia.*

"Take of freshly extracted Cassia Pulp, *half a pound*; Manna, *two ounces*; Tamarind Pulp, *an ounce*; Syrup of Orange, *half a pound*. Bruise the Manna, then dissolve it in the Syrup by means of a moderate heat, and add the Pulp; lastly, evaporate slowly the mixture to a proper consistence."

This electuary is gently laxative, and is used to relieve habitual costiveness; as a purge for children; and as a vehicle for the exhibition of other more powerful purgatives.

CONFECTIO OPII, Lond. *Confection of Opium.*

"Take of Hard Opium powdered, *six drachms*; Long Pepper, *an ounce*; Ginger Root, *two ounces*; Carraway Seeds, *three ounces*; Syrup, *a pint*. Rub the Opium with the Syrup made hot, then add the remaining articles reduced to powder, and mix."

ELECTUARIUM OPIATUM; olim, ELECTUARIUM THEBAICUM, Edin. *Opiate Electuary*; formerly, *Thebaic Electuary*.

"Take of Aromatic Powder, *six ounces*; Virginian Snake-Root, in fine powder, *three ounces*; Opium, diffused in a sufficient quantity of Spanish White Wine, *half an ounce*; Syrup of Ginger, *a pound*. Mix, so as to make an electuary."

*Syn.* Electuaire Opiate (*F.*), Theriaklatwerge (*G.*), Elettuario Oppiato (*I.*).

The operation of the Opium, in these preparations, is modified by the aromatics. They are intended as substitutes for the *Mithridate* and *Theriaca* of the old pharmacopœias, which were too long allowed to disgrace modern pharmacy. They are stimulant narcotics; and are usefully employed in atonic gout, flatulent colic, and in diarrhœas, unattended by any inflammatory symptoms. Thirty-six grains of the London confection contain one grain of opium, and the same quantity is contained in forty-three of the Edinburgh electuary. The dose is from grs. x. to f3j.

CONFECTIO PIPERIS NIGRI, Lond. *Confection of Black Pepper.*

"Take of Black Pepper, Elecampane-root, of each, *one pound*; Fennel-seeds, *three pounds*; Honey, Refined Sugar, of each, *three pounds*. Rub together the dry ingredients to a fine powder; then add the Honey, and beat the whole into one mass."



This preparation, although a warm stimulant, yet, is intended as a substitute for Ward's Paste in hæmorrhoides. The dose is from ʒj. to ʒij.

**ELECTUARIUM CATECHU COMPOSITUM**, Edin.  
*Compound Electuary of Catechu.*

“Take of Extract of Catechu, *four ounces*; Kino, *three ounces*; Cinnamon Bark, Nutmegs, of each, *one ounce*; Opium, diffused in a sufficient quantity of Spanish White Wine, *a drachm and a half*; Syrup of Red Roses boiled to the thickness of honey, *two pounds and a quarter*. Reduce the solid ingredients to powder; then mix them with the Opium and Syrup, so as to form an electuary.”

*Dublin.*

“Take of Catechu, *four ounces*; Cinnamon Bark, *two ounces*; Kino, *three ounces*; rub them to powder, and add, of Hard Refined Opium diffused in Spanish White Wine, *a drachm and a half*; Syrup of Ginger boiled to the consistence of honey, *two pounds and a quarter*. Mix them.”

These are useful combinations of astringents and aromatics; and may be efficaciously given in diarrhœas, and the last stage of dysentery, either in the form of bolus, or diffused in some distilled water. The dose is from ʒj. to ʒij. Ten scruples contain one grain of opium.

**CONFECTIO ROSÆ CANINÆ**, Lond. *Confection of the Dog-rose.*

“Take of Pulp of the Dog-rose, *a pound*; Refined Sugar in powder, *twenty ounces*. Rub them together until they be well incorporated.”

*Edinburgh.*

“Take of the Fresh Fruit of the Dog-rose, carefully freed from the seeds and inclosed spiculæ, beat it to a pulp, and while beating add gradually three times its weight of double-refined sugar.”

**CONFECTIO ROSÆ GALLICÆ**, Lond. *Confection of the Red Rose.*

“Take of the Unblown Petals of the Red Rose, freed from the claws, *a pound*; Refined Sugar, *three pounds*. Beat the Petals in a stone mortar; then add the Sugar, and beat again until the whole be thoroughly incorporated.”

*Edinburgh.*

“Beat the Unblown Petals of the Red Rose to a pulp; and add during the beating three times their weight of refined sugar.”

**CONSERVÆ ROSÆ**, Dub. *Conserve of Roses.*

“Beat the Unblown Petals of the Red Rose, freed from



their claws; adding gradually three times their weight of refined sugar."

*Syn.* Conserve de Roses rouges (*F.*), Rosenconserve (*G.*), Conserva di Rose rosse (*I.*).

The Confection of the Red Rose possesses a small degree of astringency, and is sometimes given dissolved in new milk, as a tonic in early convalescence from acute diseases; but the chief use of the confections of both kinds of roses, is to form pleasant vehicles for more active remedies.

CONFECTIO RUTÆ, Lond. *Confection of Rue.*

"Take of Rue Leaves dried, Carraway Seeds, Laurel Berries, of each, *an ounce and a half*; Sagapenum, *half an ounce*; Black Pepper, *two drachms*; Clarified Honey, *sixteen ounces*. Rub the dry articles together to a very fine powder; then add the honey, and mix the whole together."

This electuary, we are informed, is introduced as a substitute for the old *Bay Berry Electuary*. It possesses antispasmodic virtues, and since the medicinal powers of Prussic Acid have been investigated, I am inclined to think this confection might be given with advantage in chronic, spasmodic cough, Prussic Acid being the active principle of the Laurel berries. At present it is used in the form of enema only; from ℥j. to 3j. dissolved in Oss. of gruel, being administered in the convulsive affections of infants, and flatulent colic.

CONFECTIO SCAMMONEÆ<sup>1</sup>, Lond. *Confection of Scammony.*

"Take of Scammony powdered, *an ounce and a half*; Cloves bruised, Ginger Root powdered, of each, *six drachms*; Oil of Carraway, *half a drachm*; Syrup of Roses, *a sufficient quantity*. Rub the dry substance into a very fine powder; then add gradually the Syrup, and rub them again: lastly, after adding the Oil of Carraway, mix the whole together."

ELECTUARIUM SCAMMONII, Dub. *Electuary of Scammony.*

"Take of Scammony, Ginger Root, of each reduced to powder, *an ounce*; Oil of Cloves, *a scruple*; Syrup of Orange, *a sufficient quantity*. Mix the Ginger in powder with the Syrup of Orange, then add the Scammony, and lastly the Oil."

This is a stimulating cathartic; and may be given in a dose of from fʒss. to fʒj., but it is seldom ordered."

CONFECTIO SEN'NÆ<sup>2</sup>, Lond. *Confection of Senna.*

"Take of Senna Leaves, *eight ounces*; Figs, *a pound*; Tamarind Pulp, Cassia Pulp, the Pulp of Prunes, of each, *half*

<sup>1</sup> Electuarium caryocostinum, P. L. 1720. Electuarium e scammonio, P. L. 1745. Electuarium scammoni, P. L. 1787.

<sup>2</sup> Electuarium lenitivum, P. L. 1720, Electuarium Sennæ, P. L. 1787.



*a pound; Coriander Seeds, four ounces; Liquorice Root, three ounces; Refined Sugar, two pounds and a half. Powder the Senna Leaves with the Coriander Seeds, and separate by sifting ten ounces of the mixed powder. Boil the residue with the Figs and Liquorice Root, in four pints of water, until it be reduced one half; then press out and strain the liquor. Evaporate the strained liquor in a water-bath, until a pint and a half only remain of the whole; then the sugar being added, make a syrup. Finally, mix gradually the pulps with the syrup; and, having added the sifted powder, mix the whole together."*

ELECTUARIUM SENNÆ COMPOSITUM, Edin. *Compound Electuary of Senna.*

*"Take of Senna Leaves, eight ounces; Coriander Seeds, four ounces; Liquorice Root bruised, three ounces; Figs, Pulp of Prunes, of each, a pound; Pulp of Tamarinds, half a pound; Refined Sugar, two pounds and a half; Water, four pounds. Rub the Senna with the Coriander, and separate by sifting ten ounces of the mixed powder. Boil the residue with the Figs and Liquorice Root in the Water, down to one half; then express and strain. Evaporate the strained liquor to about a pound and a half. Add the Sugar, and gradually the Pulps; and, lastly, mix in the powder."*

ELECTUARIUM SENNÆ, Dub. *Electuary of Senna.*

*"Take of Senna Leaves, in very fine powder, four ounces; Pulp of Prunes, a pound; Pulp of Tamarinds, two ounces; Molasses, one pint and a half; Essential Oil of Carraway, two drachms. Boil the Pulps with the Syrup, to the thickness of honey; then add the powder, and, when the mixture is nearly cold, the oil; finally, mix the whole thoroughly together."*

Any of these Electuaries, when properly prepared, is a mild and pleasant purgative, and well adapted for those who are afflicted with habitual costiveness; and for pregnant women. They are similar to the old *Lenitive Electuary*. The dose is from ʒj. to ʒiv. or more, taken at bed time.

---

## PULVERES.

### POWDERS.

THIS form of preparing medicines is the simplest, and perhaps the least objectionable: but it is not applicable to all the articles of the materia medica. Those remedies, which are very



unpleasant to the taste; those which deliquesce rapidly when exposed to the air, or are very volatile; and those which require to be given in large doses, or which are not diffused readily in water, cannot with propriety be administered in the form of powder. Some substances cannot be reduced to powder, unless they be very much dried; and the heat necessary for that purpose alters their properties: even the impalpable form given to powders is injurious to some resinous substances; and we cannot be surprised that a great alteration should be effected in a short time, by the action of the air, on so great an extension of surface as takes place in the operation usually adopted for reducing drugs to fine powder. *Cinchona*, *Rhubarb*, *Ipecacuanha*, and *Guaiacum*, operate much less powerfully in the state of impalpable powder, than when reduced to that degree of fineness only, which can be effected by simply beating them in a mortar, and passing them through a coarser sieve than is employed in the former case.

As powders are generally affected by the action of the air and light, all powders should be kept in opaque or green glass bottles. The effect of light on the majority of powders is rendered obvious by the labelled sides of clear bottles containing them, which are always turned to the light, becoming encrusted with the powder changed in its colour, while the other side remains clear and transparent.

In forming compound powders, it is necessary to sift the mixture after it has been well triturated. The following general rule for the formation of powders is given by the Dublin College. "Let the substances to be powdered be first dried<sup>1</sup>, and then beaten in an iron mortar; then separate the finer powder by shaking it through a hair sieve, and preserve it in close vessels."

---

<sup>1</sup> Mr. Battley, a respectable druggist in London, has proposed the following method of drying narcotic plants for powders.

Previous to the process of drying the leaves of plants, the same rules must be carefully observed in reviving them, which were recommended previous to their being pressed for extracts.

The leaves being in a high state of preservation, and entirely freed from the stalks, and as much as possible from external moisture, must be laid in thin layers in baskets of willow stripped of its bark, in a drying room, from which the light is quite excluded. They should be then exposed to a temperature of not less than from 130 to 140 deg. of Fahrenheit's thermometer for three or four hours, or until the leaves begin to shrivel. They are then to be turned in the same temperature, and the heat kept up for six or eight hours longer, when the operation is generally finished; which is known by the leaves crumbling without much difficulty in the hand. If the process has been in all its parts properly managed, the result will be, that the leaves retain a beautiful green colour, and also in a high degree the medical properties of the plant to which they belong.

To preserve them in this desirable state, oil jars, made perfectly clean and dry, are found to answer best. Place the leaves lightly in the jars, and hermetically seal them. The filled jars ought to be kept in a dry and warm situation.



**PULVIS AL' OES COMPOSITUS**, Lond. *Compound Powder of Aloes.*<sup>1</sup>

“Take of Extract of Spiked Aloes, *an ounce and a half*; Guaiacum Gum-resin, *an ounce*; Compound Powder of Cinnamon, *half an ounce*. Powder the Extract of Aloes and the Guaiacum separately; then mix them with the Compound Powder of Cinnamon.”

**PULVIS ALOES CUM GUAIACO**, Dub. *Powder of Aloes with Guaiac.*

“Take of Hepatic Aloes, *an ounce and a half*; Guaiacum Gum-resin, *an ounce*; Aromatic Powder, *half an ounce*. Rub the Aloes and the Guaiacum separately to powder; then mix them with the Aromatic Powder.”

Both the active substances in these powders are ill adapted for this form of preparation; and the addition of the aromatic is scarcely sufficient to cover the nauseous taste of the Aloes. They are warm sudorific cathartics, and may be given in doses of from gr. x. to ℥j. but are seldom ordered.

**PULVIS ALOES CUM CANELLA**, Dub. *Powder of Aloes with Canella.*

“Take of Hepatic Aloes, *a pound*; White Canella, *three ounces*. Rub them separately to powder, and then mix them.”

This powder is liable to the same objection as the former, although the Canella covers the taste better than the aromatic powder. It has been long known in the shops under the name of *Hiera Picra*; and is used as a domestic remedy, infused in wine or spirits. From grs. x to ℥j. may be given for a dose.

**PULVIS ASARI COMPOSITUS**, Edin. *Compound Powder of Asarabacca.*

“Take of the Leaves of Asarabacca, *three parts*; the Leaves of Marjoram, Flowers of Lavender, of each, *one part*. Rub them together to a powder.”

*Dublin.*

“Take of Dried Leaves of Asarabacca, *an ounce*; Lavender Flowers dried, *two drachms*. Rub them together to a powder.”

A few grains of this powder snuffed up the nostrils for several successive evenings at bed-time, excite sneezing and a copious discharge of mucus, which continues to flow on the succeeding day. It has been particularly used in toothach and chronic ophthalmia.

<sup>1</sup> *Pilulæ de Diambraë*, P. L. 1720. *Pulvis Aloes cum Guaiaco*, P. L. 1787.



**PULVIS CINNAMOMI COMPOSITUS**, Lond. *Compound Powder of Cinnamon.*<sup>1</sup>

“Take of Cinnamon Bark, *two ounces*; Cardamom Seeds, *an ounce and a half*; Ginger Root, *an ounce*; Long Pepper, *half an ounce*. Rub them together to a very fine powder.”

PULVIS AROMATICUS, Edin. *Aromatic Powder.*

“Take of Cinnamon Bark, Cardamom Seed, Ginger Root, of each, *equal parts*. Rub them to a very fine powder, which is to be preserved in a well-stopped phial.”

*Dublin.*

“Take of Cinnamon Bark, *an ounce*; Lesser Cardamom Seeds freed from the husks, Ginger, Long Pepper, of each, *an ounce*. Rub them together to a powder.”

*Syn.* Poudre aromatique (*F.*), Gewurpolver (*G.*), Polvere aromatica (*I.*).

These combinations of aromatics are stimulant and carminative, and may be used to promote digestion, and expel flatus in cold phlegmatic habits; but they are more generally employed to give warmth to other compositions. The dose is from gr. viij, to ʒj. given in the form of bolus, or diffused in water.

Officinal preparations. *Pulvis Alöes compositus*, L.D. *Electuarium aromaticum*, E. *Electuarium opiatum*, E.

**PULVIS CONTRAYERVÆ COMPOSITUS**, Lond. *Compound Powder of Contrayerva.*<sup>2</sup>

“Take of Contrayerva Root powdered, *five ounces*; Prepared Shells, *a pound and a half*. Mix them.”

This powder is stimulant and sudorific; and is given with advantage in typhoid fevers; the malignant exanthemata; the sinking stage of dysentery; and in atonic gout. The dose is from grs. x. to grs. xl. given either diffused in simple water, or rubbed up with mucilage and mint water.

**PULVIS CORNU CERVINI USTI**, Dub. *Powder of Burnt Hartshorn.*

“Let Pieces of Hartshorn be burnt until they become white, then reduce them to a very fine powder.”

**PULVIS CORNU USTI CUM OPIO**, Lond. *Powder of Burnt Hartshorn with Opium.*

“Take of Hard Opium powdered, *a drachm*; Hartshorn burnt and prepared, *an ounce*; Cochineal Powder, *a drachm*. Mix them.”

PULVIS OPIATUS<sup>3</sup>, Edin. *Opiate Powder.*

“Take of Opium, *one part*; Prepared Carbonate of Lime, *nine parts*. Rub them together to a fine powder.”

<sup>1</sup> Species Diambrae sine odoratis, P. L. 1720. Species aromaticæ, P. L. 1745. Pulvis aromaticus, P. L. 1787.

<sup>2</sup> Lapis Contrajervæ, P. L. 1720, so named from its having been made into balls.

<sup>3</sup> Pulvis opiatum, P. L. 1787.



*Syn.* Poudre opiate (*F.*), Opiumspulver (*G.*), Polvere oppiata (*I.*).

Ten grains of either of these powders contain one grain of Opium. They are intended chiefly for exhibiting Opium in very small doses. The substances used to divide the Opium are of no consequence as to the effect of the remedy; and, therefore, the burnt Hartshorn being more brittle than the Chalk, is better fitted for this purpose.

**PULVIS CRETÆ COMPOSITUS**, Lond. *Compound Powder of Chalk.*

“Take of Prepared Chalk, *half a pound*; Cinnamon Bark, *four ounces*; Tormentil Root, Acacia Gum, of each, *three ounces*; Long Pepper, *half an ounce*. Rub them separately to fine powder, then mix them.”

**PULVIS CARBONATIS CALCIS COMPOSITUS**, Edin. *Compound Powder of Carbonate of Lime.*

“Take of Prepared Carbonate of Lime, *four ounces*; Cinnamon Bark, *a drachm and a half*; Nutmegs, *half a drachm*. Rub them together to a powder.”

The London preparation, owing to the larger proportion of aromatics it contains, and the addition of the Tormentil root is better adapted for checking diarrhœa connected with acidity of the primæ viæ, than the Edinburgh powder, which may be regarded as a simple but grateful antacid. The dose is from grs. v. to ℥j. given generally in the form of mixture rubbed up with mucilage and some distilled water.

**PULVIS CRETÆ COMPOSITUS CUM OPIO**, Lond. *Compound Powder of Chalk with Opium.*

“Take of Compound Powder of Chalk, *six ounces and a half*; Hard Opium powdered, *four scruples*. Mix them.”

The addition of Opium to the compound powder of Chalk renders it more useful in diarrhœa; and from the minute division of the Opium, one grain only being contained in two scruples of the powder, it forms an useful opiate powder for children suffering under the irritative diarrhœa of teething. The dose is from ℥j. to ʒj. for adults.

**PULVIS JALAPÆ COMPOSITUS**, Edin. *Compound Powder of Jalap.*

“Take of Powder of Jalap Root, *one part*; Supertartrate of Potass, *two parts*. Rub them together to a fine powder.”

The addition of the Supertartrate, besides dividing the Jalap very minutely, modifies also its purgative operation. This powder is an useful purgative in habitual costiveness: it is also very serviceable to children with tumid bellies, in worm cases, and in dropsy. The dose is from ℥j. to ℥ij. for adults.

**PULVIS IPECACUANHÆ COMPOSITUS**, Lond. Dub. *Compound Powder of Ipecacuanha.*



“Take of Ipecacuanha Root powdered, Hard Opium powdered, of each, *a drachm*; Sulphate of Potass powdered, *an ounce*. Mix them.”

PULVIS IPECACUANHÆ ET OPII, Edin. *Powder of Ipecacuanha and Opium.*

“Take of Ipecacuanha Root powdered, Opium, of each, *one part*; Sulphate of Potass, *eight parts*. Rub them together to a fine powder.”

*Syn.* Poudre d'Ipecacuanha et d'opium (F.), Davers schmerzstillendes pulver (G.), Polvere d'Ipecacuanha ed oppio (I.).

In this powder the Sulphate of Potass is intended chiefly to divide the Opium mechanically; but it modifies also the action of the Opium and Ipecacuanha.<sup>1</sup> Compound Ipecacuanha powder operates as a powerful sudorific; and is very efficaciously given in all cases, whether inflammatory or not, in which full sweating is indicated. The dose is from grs. v. to ℥j. It may be given diffused in water, or in the form of bolus, and should be assisted in its action by plentiful dilution with tepid fluids; but these must not be drank immediately after taking the powder, as from such a circumstance it is very apt to be rejected by vomiting. Ten grains of this powder contain one grain of opium.

PULVIS KINO COMPOSITUS, Lond. *Compound Powder of Kino.*

“Take of Kino, *fifteen drachms*; Cinnamon Bark, *half an ounce*; Hard Opium, *a drachm*. Rub them separately to a very fine powder, and then mix.”

This is an astringent anodyne powder, which is now, for the first time, introduced into the Pharmacopœia. The dose is from grs. x. to ℥j. Twenty grains of the powder contain one grain of Opium.

PULVIS QUERCUS MARINÆ, Dub. *Powder of Yellow Bladder-wrack.*

<sup>1</sup> In the original *Dover's Powder*, the saline ingredient was procured by de-flagrating a mixture of equal parts of nitrate of potass, and sulphate of potass; and the nitre is still retained as an ingredient in the *Compound Powder of Ipecacuanha and Opium* of the French Codex. The Pharmacopœia Danica and the Pharmacopœia Austriaca, order sugar instead of any salt; but it is less calculated to assist in the pulverization of the opium, and is apt also to attract moisture, and form the powder into a solid mass. The following table, shows the proportion of opium in this powder, as ordered in the principal Pharmacopœias of Europe.

Lond.	Edin.	Gallica.	Swed.	Dan.	Boruss.	Austr.
$\frac{1}{16}$	$\frac{1}{16}$	$\frac{1}{11}$	$\frac{1}{80}$	$\frac{1}{10}$	$\frac{1}{16}$	$\frac{1}{10}$



“Take of Bladder-wrack in flower, *any quantity*. Let it be dried, and freed from the sordes, then exposed to heat in an iron vessel, or a crucible, to which a perforated lid is adapted, until, the vapours ceasing, it become obscurely red hot. Reduce the carbonaceous matter which remains to powder.”

This powder is a mixture of Soda, Iodine, and Charcoal. For its medicinal effects see *Fucus*, Part II.

PULVIS SALINUS COMPOSITUS, Edin. *Compound Saline Powder*.

“Take of pure Muriate of Soda, Sulphate of Magnesia, of each, *four parts*; Sulphate of Potass, *three parts*. Dry the salts with a gentle heat, then pulverize them separately, and afterwards rub them together. Preserve the powder in a well-stopped phial.”

This powder is a simple purgative, and does not appear to possess any peculiar advantages.

PULVIS SCAMMONIÆ COMPOSITUS, Lond. *Compound Powder of Scammony*.

“Take of Scammony, Hard Extract of Jalap, of each, *two ounces*; Ginger Root, *half an ounce*. Rub them separately to a very fine powder, and then mix them.”

PULVIS SCAMMONII COMPOSITUS, Edin. *Compound Powder of Scammony*.

“Take of Scammony, Supertartrate of Potass, of each, *equal parts*. Rub them together to a very fine powder.”

*Syn.* Poudre de Scammonée composée (F.).

These powders, although agreeing in name, differ very considerably in their nature. In the first, the activity and the stimulating quality of the Scammony are increased by the Jalap, while the griping effect of the mixture is in some degree obviated by the Ginger. In the second, the addition of the Supertartrate of Potass, detracts from the violence of the operation of the Scammony, and renders it less irritating; although, at the same time, it renders it more certain. The dose of the first is from grs. x. to grs. xv.; that of the second, from grs. x. to ʒss. They are chiefly used in hydropic and worm cases, and to remove mucous obstructions.

PULVIS SCILLÆ, Dub. *Powder of Squill*.

“Let Squill Roots (bulbs) freed from their membranous integuments and cut in transverse slices, be dried upon a sieve with a low degree of heat; and then reduce them to powder, which must be preserved in well-stopped glass phials.”

PULVIS SENNÆ COMPOSITUS, Lond. *Compound Powder of Senna*.

“Take of Senna Leaves, Supertartrate of Potass, of each, *two ounces*; Scammony, *half an ounce*; Ginger Root, *two*



*drachms.* Reduce to very fine powder, the Scammony by itself, and the other ingredients together; then mix the whole."

*Syn.* Poudre composée de Senna (*F.*), Polvere di Senna composta (*I.*).

This powder is hydragogue and cathartic; but it is an inconvenient form of preparation, owing to the bulk of the dose, which is very considerable, although from ℥j. to ʒj. only in weight.

**PULVIS SPONGIÆ USTÆ, Dub.** *Powder of burnt Sponge.*

"Let Sponge cut into small pieces be beaten so as to free it from little stones; then burn it in a covered iron vessel, until it become black and friable; finally, reduce it to powder."

**PULVIS ALUMINIS COMPOSITUS, Edin.** *Compound Powder of Alum.*

"Take of Sulphate of Alum, *four parts*; Kino, *one part*. Rub them together to a fine powder."

This is a powerful astringent powder, and is sometimes used internally in menorrhagia and diarrhœa; but is more generally employed as an external application. The dose is from grs. x. to grs. xv.; but it must be taken in the dry state, as the Kino is decomposed by the Alum, when a fluid vehicle is employed.

**PULVIS TRAGACANTHÆ COMPOSITUS, Lond.** *Compound Powder of Tragacanth.*

"Take of Tragacanth powdered, Acacia Gum powdered, Starch, of each, *an ounce and a half*; Refined Sugar, *three ounces*. Rub the Starch and the Sugar together to a powder; then add the Tragacanth and the Acacia Gum, and mix the whole together."

*Syn.* Poudre composée de Tragacanthé (*F.*), Tragacanthæ Gummi pulver (*G.*). Polvere di Tragacanta composta (*I.*).

In this composition the starch might well be omitted, as it is insoluble in cold water. This compound powder is efficaciously used as a demulcent in hectic fever, and to allay the tickling cough of catarrh: in gonorrhœa and strangury it is given combined with Nitre; and in dysentery, with Ipecacuanha powder. The dose is from ʒss. to ʒiij. mixed in water or any bland fluid.


## PILULÆ.

### PILLS.

PILLS are masses of a consistence sufficient to preserve a round form, yet not so hard as to be of too difficult solution in the stomach. This form of preparation is particularly adapted for medicines which have a very nauseous taste or



flavour, and such as operate in minute doses. Extracts, when not too hard, may be formed into pills without any addition; but more generally pills are composed of either vegetable, or earthy, or metallic powders, combined by means of syrup into a coherent mass. Salts also may be formed into pills, except such as are deliquescent; and when efflorescent salts are used, they should be first freed from the water of crystallization, for the pills formed with uneffloresced salts which are apt to effloresce, fall into powder as they dry. The masses, which are ordered to be kept prepared for the formation of pills, require to be preserved in covered pots, wrapped in bladders, and occasionally moistened. When they are to be formed into pills, a given portion of the mass is rolled into a cylinder, the length of which is regulated by the number of pills into which it is to be divided; and the division is effected either as equally as possible by the hand, or by a machine invented for the purpose. This machine consists of two pieces. The first *a* is divided into three compartments; 1. a vacant space to receive the divided mass, which is to be rolled into pills; 2. a grooved brass plate, which assists in dividing the mass into pills; and 3. a box for containing a powder for covering the pills after they are formed. The second, *b*, consists of a brass plate 1, grooved to match the plate 2 on piece *a*, and bounded at both ends by moveable projecting plates 2 2, containing two wheels under the ledge of the plate 2; and a wooden back 3, with two handles, 4 4, to which this plate is affixed. In using the machine the cylindrical mass is placed on 2 *a*; and divided by passing *b* over it, the wheels of *b* enabling it to run easily, being applied to a brass plate which forms the margin of *a*.



After the round form is given to each of the pills, by rolling the divided pieces between the fingers, they are dropped



into 3 *a*, and covered by some dried powder, as, for instance, subcarbonate of magnesia or starch, to prevent them from adhering. With the same intention pills were formerly gilded; but as simple dry powders answer all the purposes of this covering, it is now altogether laid aside.

PILULÆ ALOES COMPOSITÆ, Lond. *Compound Aloetic Pills.*

“Take of Extract of Spiked Aloes powdered, *an ounce*; Extract of Gentian, *half an ounce*; Oil of Carraway, *forty minims*; Syrup, *a sufficient quantity*. Beat them together until they combine into an uniform mass.”

PILULÆ ALOETICÆ, Edin. *Aloetic Pills.*

“Take of Socotorine Aloes in powder, Soap, of each, *equal parts*. Beat them with simple syrup, so as to make a mass fit for forming pills.”

PILULÆ ALOES CUM ZINGIBERE, Dub. *Pills of Aloes and Ginger.*

“Take of Hepatic Aloes, *an ounce*; Ginger Root in powder, *a drachm*; Soap, *half an ounce*; Essential Oil of Peppermint, *half a drachm*. Let the Aloes and the Ginger be rubbed together to a powder: then add the Soap and the Oil so as to form a mass.”

*Syn.* Pilule d'Aloë composée (*F.*), Pillole d'Aloë composta (*I.*).

In the London preparation, the quantity of extract of gentian ordered is too large; for, owing to its re-action on the Aloes, the mass becomes rather too soft to form into pills: at all events, no syrup is required in this instance. The soap ordered in the two other formulæ is well adapted for giving consistence and form to the Aloes. This is an useful pill, and is advantageously employed for obviating the habitual costiveness of the sedentary, and of leucophlegmatic habits. The dose is from grs. x. to grs. xv. or more.

PILULÆ AL'OEES ET ASSAFŒTIDÆ, Edin. *Pills of Aloes and Assafoetida.*

“Take of Socotrine Aloes in powder, Assafoetida, Soap, of each, *equal parts*. Beat them into a mass with mucilage of Gum-arabic.”

*Syn.* Pilules d'Aloë avec assafoetida (*F.*), Pillole Bloeteche con assafetida (*I.*).

These pills are anodyne and cathartic, allaying any irritability of the bowels, at the same time that they open them freely. They have been found extremely useful in dyspepsia attended with flatulence. The dose is grs. x. given twice a day.

PILULÆ ALOES CUM MYRRHA, Lond. *Pills of Aloes with Myrrh.*<sup>1</sup>

“Take of Extract of Spiked Aloes, *two ounces*; Saffron,

<sup>1</sup> Pilulæ Ruffi seu communes, P. L. 1720.



Myrrh, of each, *an ounce*; Syrup, *a sufficient quantity*. Rub separately to powder the Aloes and the Myrrh; then beat all the ingredients together until they form an uniform mass."

*Dublin.*

"Take of Hepatic Aloes, *an ounce*; Myrrh, *half an ounce*; Saffron, *two drachms*; Essential Oil of Carraway, *half a drachm*. Rub the Myrrh and the Aloes separately to powder, and beat the whole together into a mass.

PILULÆ ALOES ET MYRRHÆ, Edin. *Pills of Aloes and Myrrh.*

"Take of Socotorine Aloes, *four parts*; Myrrh, *two parts*; Saffron, *one part*. Beat them into a mass with simple syrup."

*Syn.* Pilules d'Aloë avec la Myrrh (*F.*), Pillole Aloetique con Mirra (*I.*).

These pills have been employed since before the time of Rhazes, to stimulate and open the bowels in chlorotic, hypochondriacal, and chachectic habits; and are not the less valuable because they are of very ancient origin. The dose is from grs. x. to ʒj. They may be given twice a day.

PILULÆ AMMONIARETI CUPRI, Edin. *Pills of Ammoniaret of Copper.*

"Take of Ammoniaret of Copper rubbed to fine powder, *sixteen grains*; Crumb of Bread, *four scruples*; Water of Carbonate of Ammonia, *a sufficient quantity*. Beat them into a mass, and divide it into thirty-two equal pills."

*Syn.* Pilules cuivreuses de Swédiaur (*F.*).

This is a convenient form for the exhibition of the Ammoniaret of Copper, half a grain of which is contained in each of the pills. They are given in epilepsy and other spasmodic diseases. One pill given at night and in the morning, is sufficient at first; but the number may be gradually increased till five be taken for a dose.

PILULÆ CAMBO'GIÆ COMPOSITÆ, Lond. Edin. *Compound Pills of Gamboge.*

"Take of Gamboge in powder, *a drachm* (Edin.); Extract of Spiked Aloes in powder, *a drachm and a half*; Ginger in powder, *half a drachm*; Hard Compound Powder of Cinnamon, *a drachm*; Soap, *two drachms*. Mix the powders together; then add the soap, and beat the whole into an uniform mass."

This is considerably more active than the aloetic pills. The dose is from grs. x. to ʒj. given at bed-time in obstinate costiveness.

PILULÆ COLOCYNTHIDIS COMPOSITÆ, Edin. *Compound Colocynth Pills.*

"Take Socotorine Aloes, Scammony, of each, *eight parts*; Colocynth Pulp, *four parts*; Sulphate of Potass, Oil of Cloves, of each, *one part*. Beat the Extract, Gum-resin, and Sulphate



together into powder, then, with the Colocynth Pulp rubbed to fine powder, mix them with the Oil, and, finally, beat the whole into a mass with Mucilage of Gum."

*Dublin.*

"Take of Colocynth Pith, *half an ounce*; Hepatic Aloes, Scammony, of each, *one ounce*; Castile Soap, *two drachms*; Oil of Cloves, *one drachm*. Pulverize separately the aloes, scammony, and colocynth, then triturate them with the soap and oil, and form them into a mass by means of syrup."

Both these are excellent forms of purgative pills. They are more powerful in their operation than the other aloetic pills; and do not so soon lose their power when taken for any considerable length of time in habitual costiveness.

PIL'ULÆ FER'RI COMPOSITÆ, Lond. *Pills of Iron with Myrrh.*

"Take of Myrrh in powder, *two drachms*; Subcarbonate of Soda, Sulphate of Iron, Sugar, of each, *a drachm*. Rub the Myrrh with the Subcarbonate of Soda; then, having added the Sulphate of Iron, rub again; and, lastly, beat the whole into an uniform mass."

This is an useful emmenagogue pill, similar in its properties to Griffith's mixture. The dose is from grs. x. to ℥j. and may be given twice or three times a day.

PIL'ULÆ GAL'BANI COMPOSITÆ, Lond. *Compound Pills of Galbanum.*<sup>1</sup>

"Take of Galbanum, *an ounce*; Myrrh, Sagapenum, of each, *an ounce and a half*; Assafoetida, *half an ounce*; Syrup, *a sufficient quantity*. Beat them together into an uniform mass."

PILULÆ ASSAFÆTIDÆ COMPOSITÆ, Edin. *Compound Assafoetida Pills.*

"Take of Assafoetida, Galbanum, Myrrh, of each, *eight parts*; Purified Oil of Amber, *one part*. Beat them into a mass with simple syrup."

PILULÆ MYRRHÆ COMPOSITÆ, Dub. *Compound Pills of Myrrh.*

"Take of Assafoetida, Myrrh in powder, Galbanum, of each, *an ounce*; Oil of Amber, *half a drachm*. Rub them together, and make them into a mass with simple syrup."

*Syn.* Pilules de Galbanum composée (*F.*), Pillole di Galbano composta (*I.*)

These preparations are useful antispasmodics and emmenagogues; and are given with advantage in chlorosis, hysteria, and hypochondriasis. The dose is from grs. x. to ℥j. and may be taken every night at bed-time.

Of the three appellations employed by the Pharmacopœias,

<sup>1</sup> Pilulæ gummosæ, P. L. 1720.



that of the Edinburgh is the least objectionable, the assafoetida being undoubtedly the most powerful article.

PILULÆ HYDRARGYRI, Lond. Dub. *Mercurial Pills.*

“Take of Purified Mercury, *two drachms*; Confection of Red Roses, *three drachms*; Liquorice Root in powder, *a drachm*. Rub the Mercury with the Confection until the globules disappear; then add the Liquorice Root, and beat the whole into an uniform mass.”

*Edinburgh.*

“Take of Purified Mercury, Conserve of the Red Rose, of each, *an ounce*; Starch, *two ounces*. Rub the mercury with the conserve in a glass mortar until the globules entirely disappear, adding, if necessary, a little mucilage of gum arabic; then add the starch, and with a little water beat the whole into a mass, which is to be directly divided into four hundred and eighty equal-sized pills.”

*Syn.* Pilules mercuriales (*F.*), Pillole mercuriale (*I.*).

One grain of Mercury is contained in three grains of the mass, made according to the London and Dublin formulæ, and in four grains according to the Edinburgh.

In these preparations the mercury is first minutely divided by the viscosity of the conserve, the substance with which it is triturated: and it is with much probability believed that this mechanical division is all that is effected by the trituration. It is, however, also supposed, that the metal is oxidized; and that the great extension of surface and, in some degree, the substance used in the trituration facilitate this effect. Further experiments are certainly requisite to determine this question. Syrup, honey, mucilage, soap, guaiacum and other matters, have been occasionally employed; but the Colleges have agreed in preferring Conserve of Roses; and it is not improbable that the operation is shortened by the weak Acid which the Conserve contains. On the continent the oil of eggs has been employed for dividing mercury<sup>1</sup>, and certainly no substance so rapidly assists in producing the desired effect as this oil, when it has been kept for some time. In the above preparations, therefore, the mercury is supposed to be nearly in the state of the black oxide, and on this combination of oxygen its activity as a remedy depends. The more assiduously the trituration is continued, the more perfect is the preparation. The sufficient division of the metal, or the oxidization of the whole of the globules, or the extinction or killing of the mercury in the common language of the laboratory, is known to be completed, when, on rubbing a small portion of

<sup>1</sup> Vide *Lond. Med. Repository*, vol. v. p. 166.



the mass with the point of the finger on a piece of clean paper, no metallic globules are perceptible. The mass must be then immediately formed into pills, as it very rapidly becomes too hard, if allowed to remain.

*Medical properties and uses.* — These pills are stimulant and antisyphilitic, and are the most common form of preparation under which mercury is exhibited for the cure of venereal affections, being much less liable to act on the bowels than any of the other forms. When they act on the bowels, they should be combined with Opium, or with a few grains of Rhubarb, which enable the bowels to resist the mercurial irritation. The common dose is grs. vi. to grs. viij. or two pills, given twice a day, until the mouth be affected. Larger doses are apt to excite purging.

PILULÆ HYDRARGYRI SUBMURIATIS COMPOSITÆ, Lond. Edin. *Pills of Submuriate of Mercury.*

“Take of Submuriate of Mercury (*calomel*), Precipitated Sulphuret of Antimony, of each, *a drachm*; Guaiacum Gum-resin, *two drachms*. Rub the Submuriate of Mercury with the precipitated Sulphuret of Antimony, then with the Guaiacum, and add a sufficient quantity of Mucilage of Gum to give the mass a proper consistence.”

This preparation was introduced into practice by Dr. Plummer, and admitted into the Edinburgh Pharmacopœia under the name of Plummer’s pill. It was, however, afterwards expunged; but as it continued to be much used in practice, the London College has given it a place in its Pharmacopœia. It is a very useful alterative in lepra, in secondary syphilis affecting the skin, and in other cutaneous diseases. The dose is from grs. v. to grs. x. given night and morning.

PILULÆ RHEI COMPOSITÆ, Edin. *Compound Rhubarb Pills.*

“Take of Rhubarb Root in powder, *one ounce*; Socotorine Aloes, *six drachms*; Myrrh, *half an ounce*; Volatile Oil of Peppermint, *half a drachm*. Beat them into a mass with Syrup of Orange-peel.”

*Syn.* Pilule de Rhubarbe composée (*F.*), Pillole di Rhabarbaro composta (*I.*).

This is a warm, stomachic, laxative pill, very useful for obviating costiveness, and at the same time giving tone to the bowels in dyspepsia and hypochondriasis. The dose is from grs. x. to ʒj. given twice a day.

PILULÆ SAPONIS CUM OPIO, Lond. *Pills of Soap and Opium.*

“Take of Hard Opium powdered, *half an ounce*; Hard Soap, *two ounces*. Beat them together into an uniform mass.” Five grains contain one grain of Opium.



PILULÆ OPIATÆ; olim, PILULÆ THEBAICÆ, Edin. *Opiate Pills*; formerly, *Thebaic Pills*.

“Take of Opium, *one part*; Extract of Liquorice, *seven parts*; Pimenta berries, *two parts*. Mix the Opium and the Extract, separately softened with diluted alcohol, and beat them into a pulp; then add the Jamaica Pepper rubbed to powder, and beat the whole to a mass.” Ten grains contain one grain of Opium.

PILULÆ E STYRACE, Dub. *Storax Pills*.

“Take of Purified Storax, *three drachms*; Soft Purified Opium Saffron, of each, *a drachm*. Mix them well together, by beating.” Five grains contain one grain of Opium.

*Syn.* Pilule d’Opium (*F.*), Storaxpillen (*G.*), Pillole d’Oppio (*I.*).

The substances with which the opium is combined in these pills do not interfere with its operation as an anodyne, but are intended chiefly to cover its odour and taste, in cases where the patient or his friends have an objection to opium; and as it is also sometimes necessary that it should not appear even in the prescription, the name adopted by the Dublin College is preferable to the others. The dose of the three preparations differs, and must be regulated by the quantity of opium contained in that one which is adopted.

PILULÆ SCILLÆ COMPOSITÆ, Lond. *Compound Squill Pills*.

“Take of Fresh Squill Root (bulb) dried and powdered, *one drachm*; Ginger Root powdered, Hard Soap, of each, *three drachms*; Ammoniacum powdered, *two drachms*. Mix the powders together; then beat them with the Soap, as much Syrup being added as will give them a proper consistence.”

PILULÆ SCILLITICÆ, Edin. *Squill Pills*.

“Take of Squill Root (bulb) dried, and rubbed to a fine powder, *one scruple*; Ammoniacum, Cardamom Seeds powdered, Extract of Liquorice, *one drachm*. Beat them with syrup into a mass.”

PILULÆ SCILLÆ CUM ZINGIBERE, Dub. *Pills of Squill with Ginger*.

“Take of Powder of Squill, *a drachm*; Ginger Root in powder, *two drachms*; Essential Oil of Anniseed, *ten drops*. Beat them together, and form them into a mass with jelly of Soap.”

*Syn.* Pilules de Scille (*F.*), Pillole Squilitiche (*I.*).

These pills are useful expectorants in chronic catarrh, dyspnoea, and asthma; and combined with calomel and digitalis in hydropic affections. They are liable, however, to the same objections as the Squill powder, the efficacy of the Squill being much injured by keeping in either form; and it is perhaps better that it should be always given under an extemporaneous



form, except when the tincture is used. The dose is from grs. iv. to xj. given three or four times a day.

PILULÆ SUBCARBONATIS SODÆ, Edin. *Pills of Subcarbonate of Soda.*

“Take of Exsiccated Subcarbonate of Soda, *four parts*; Hard Soap, *three parts*. Beat into a mass with simple syrup.”

This preparation was recommended by the late Dr. Beddoes; and has been found occasionally useful.

PILULÆ SULPHATIS FERRI COMPOSITÆ, Edin. *Compound Pills of Sulphate of Iron.*

“Take of Sulphate of Iron reduced to powder, *one ounce*; Extract of Chamomile flowers, *one ounce and a half*; Oil of Peppermint, *a drachm*. Beat into a mass with simple syrup.”

This is a useful tonic pill, and may be given with advantage in dyspepsia and other affections in which steel is indicated. A five grain pill will contain two grains and a half of the Sulphate of Iron.

## TROCHISCI.

### TROCHES.

THESE are little cakes or tablets composed of powders combined with sugar and mucilage. They are hard and dry, but readily dissolve in the mouth, for which purpose they are chiefly intended; and, therefore, such remedies only as are designed to produce a local effect are given in this form. They are of little importance as remedies; and the preparation of them ought to be left entirely to the confectioner. They should be altogether rejected from the Edinburgh Pharmacopœia, as has been done by the London and the Dublin Colleges.

TROCHISCI CARBONATIS CALCIS, Edin. *Troches of Carbonate of Lime.*

“Take of Prepared Carbonate of Lime, *four ounces*; Acacia Gum, *an ounce*; Nutmegs, *one drachm*; Refined Sugar, *six ounces*. Rub them to powder, and form them by means of water into a mass fit for making troches.”

“These troches are intended as antacids; but in the state of the stomach when it requires the use of these remedies, the effect of the carbonate of lime is counteracted by the sugar.”

TROCHISCI CARBONATIS MAGNESIÆ, Edin. *Troches of Carbonate of Magnesia.*

“Take of Carbonate of Magnesia, *six ounces*; Refined Sugar, *three ounces*; Nutmegs, *a scruple*. Beat them into



powder and form them into troches with mucilage of Tragacanth."

These are used in the same cases as the former; and are more serviceable when the bowels are confined.

**TROCHISCI GLYCYRRHIZÆ GLABRÆ**, Edin. *Troches of Liquorice.*

"Take of Extract of Liquorice, Gum Arabic, of each, *one part*; Refined Sugar, *two parts*; Boiling Water, *a sufficient quantity*. Dissolve and strain, then evaporate the solution, by a gentle heat, to a proper consistence for forming troches."

These troches are demulcent, and from the viscid nature of the extract of liquorice are well adapted for allaying the tickling irritation which induces coughing; but they are not more useful than the simple extract of liquorice refined by straining and inspissation, such as is found in the shops under the name of refined liquorice.

**TROCHISCI GLYCYRRHIZÆ CUM OPIO**, Edin. *Liquorice Troches with Opium.*

"Take of Opium, *two drachms*; Tincture of Balsam of Tolu, *half an ounce*; Simple Syrup, *eight ounces*; Extract of Liquorice, softened by hot water, Gum Arabic in powder, of each, *five ounces*. First rub the Opium well with the Tincture; then add gradually the Syrup and the Extract; afterwards sprinkle in the powdered Gum Arabic; lastly, dry the mass, and form it into troches, each weighing ten grains."

These troches are used for the same purposes as the former; and from the opium they contain are more efficacious in allaying tickling cough. Six troches contain one grain of opium; and from six to ten may be taken in twenty-four hours.

**TROCHISCI GUMMOSI**, Edin. *Gum Troches.*

"Take of Gum Arabic, *four parts*; Starch, *one part*; Refined Sugar, *twelve parts*. Rub the whole to powder, and form it into a mass with Rose Water fit for forming troches."

These troches are simple demulcents, and answer sufficiently well for allaying the tickling irritation of the fauces which excites coughing.

**TROCHISCI NITRATIS POTASSÆ**, Edin. *Troches of Nitrate of Potass.*

"Take of Nitrate of Potass, *one part*; Refined Sugar, *three parts*. Beat them to powder, and form them into a mass fit for forming troches by means of mucilage of Tragacanth."

These troches afford an agreeable form of taking Nitre in the dry state; and are useful for cooling the mouth in salivations, and in stopping the progress of inflammatory sore throat, when taken at its commencement. They may also be used as a general refrigerant in fevers, diluting largely during their use. The dose is one or two taken every second or third hour.



## PRÆPARATA EX ANIMALIBUS.

## PREPARATIONS FROM ANIMALS.

THE substances of this division are few in number, and are not remedies of much efficacy.

ADEPS PRÆPARATA, Lond. *Prepared Lard.*<sup>1</sup>

“Cut the Fat into small fragments; then melt it by a gentle heat, and press it through linen.”

ADEPS SUILLUS PRÆPARATUS, Dub. *Prepared Hog's Lard.*

“Let Fresh Lard, cut into small pieces, be melted by a moderate heat, and strained by pressing it through a linen cloth.

“Lard, which is prepared by the dealers, and is preserved with salt, is to be melted with twice its weight of boiling water, and the mixture well stirred: it is then to be allowed to cool, when the lard may be separated.”

SEVUM PRÆPARATUM, Lond. *Prepared Suet.*<sup>2</sup>

“Cut the Suet in pieces; then melt it by a gentle heat, and press it through linen.”

The properties of Lard and Suet have been already detailed. (*Part ii.*) The above processes are intended to purify them; but, in order to obtain them very pure, it is necessary that they be washed in water until the water come off colourless, before they be melted. Any water that may remain attached to the fat is evaporated during the melting; and that it is all evaporated, is known by throwing a little of the melted fat into the fire, when it will crackle if any water be present. The heat must not be raised above 97°, the melting point of fat; as otherwise the fat is decomposed, rendered acrid, and assumes a yellow colour. This purification is seldom attempted by the apothecary, as both kinds of fat can be procured very well purified from the dealers. To keep lard clean, and preserve it from the action of the air, it is generally run into bladders while in the liquid state.

CORNU USTUM, Lond. *Burnt Hartshorn.*<sup>3</sup>

“Burn pieces of Hart's horn in an open fire until they become thoroughly white; then powder them, and prepare them in the manner directed for the preparation of chalk.”

<sup>1</sup> Adeps Suillæ preparata, P. L. 1787.

<sup>2</sup> Sevum Ovillum preparatum, P. L. 1787.

<sup>3</sup> Cornu Cervi ustum, P. L. 1787.



From the Latin title given to this preparation, one might be led to suppose that any kind of horn would serve as a substitute for Hart's horn, which is intended to be designated: but the properties of Hart's horn are more similar to those of bone than of the horns of other animals, the chief difference being in the proportion of cartilage, which is greater in the Hart's horn than in bone.

In performing this operation the fire must not be too violent, as the horn is apt to suffer a species of vitrification of the surface, when exposed to a very strong heat, which prevents the internal parts from being completely burnt. The residue of 100 parts of Hart's horn consists, after the burning, of 57·5 of Phosphate of Lime, one of Carbonate of Lime, and a minute portion of Phosphate of Magnesia.

*Medical properties and uses.* — Phosphate of Lime is perfectly inert when taken into the stomach; and the analysis of burnt Hart's horn has clearly proved, that the former idea of its antacid properties was erroneous. It has been proposed as a remedy in rickets and mollities ossium; but we cannot easily conceive how it can be taken up by the absorbents, and thrown upon the bones; and experience has not yet confirmed the utility of burnt Hart's horn for any purpose as a remedy.

Officinal preparation. *Mistura Cornu usti*, L.

SPONGIA USTA, Lond. *Burnt Sponge*.

“Cut Sponge into small pieces and bruise it, in order to free it from any adhering extraneous substances; then burn it in a covered iron vessel, until it become black and friable; finally, rub it to a very fine powder.”

The properties of fresh sponge have been already noticed: when burnt, the residue consists of carbonate and phosphate of lime, subcarbonate of soda, iodine, and charcoal. The active ingredients are the subcarbonate of soda and the iodine.

*Medical properties and uses.* — Burnt Sponge is tonic, deobstruent, and antacid. It has been much recommended in bronchocele, scrophulous affections, and herpetic eruptions: and I have witnessed its efficacy in schirrhous testicle, when given in combination with Cinchona Bark. It seems to derive its efficacy from the Iodine it contains. (*See Appendix*, No. 2.) The dose is from ʒj. to ʒiij. mixed into the form of an electuary, with powdered cinnamon and honey. In bronchocele the patient is directed to swallow the portion of electuary very slowly, from a supposition that some local effect is produced.

TESTÆ PRÆPARATA, Lond. *Prepared Shells*.

“Wash the Shells with Boiling Water, having previously



freed them from extraneous matters, then prepare them in the manner directed for the preparation of chalk."

OSTREARUM TESTÆ PRÆPARATÆ, Dub. *Prepared Oyster Shells.*

"These are to be prepared in the same manner as chalk."

OVORUM TESTÆ PRÆPARATÆ, Dub. *Prepared Egg Shells.*

"These are to be prepared in the same manner as chalk."

Both in oyster and egg shells the predominating ingredient is carbonate of lime, and therefore these prepared shells do not differ from chalk, except in containing a small portion of gelatin or albumen. But as this does not in any degree affect their medicinal properties, which are exactly the same as those of chalk, they might well be spared from the list of preparations. The dose is from grs. x. to ʒij. or more.

## EMPLASTRA.

### PLASTERS.

THESE are solid, tenaceous compounds, adhesive in the ordinary heat of the human body. The base of the majority of plasters is a chemical combination of the semivitreous oxide of lead and oil; but some of them owe their consistence to wax and resin; and others contain no oily nor fatty matter whatsoever. Deyeux proposes<sup>1</sup> to confine the name of plasters to the combinations of metallic oxides with oils or fat; and to give those not containing oxides, the term solid ointments; but this definition would include among the plasters some of the ointments, and exclude many of the plasters.

Plasters should not adhere to the hand when cold; they should be easily spread when heated; and should remain tenacious and pliant after they are spread; but should not be so soft as to run when heated by the skin. All plasters become too consistent and brittle when long kept; but in this case, those which are unctuous may be re-melted by a gentle heat, and some oil added to them. They are usually formed into rolls, which are wrapped in paper; and when they are to be used, they are melted and spread on leather, calico, linen, or silk. Those which contain metallic oxides ought to be melted by boiling water, for in a greater degree of heat the fatty matter is apt to reduce the oxide.

<sup>1</sup> *Annales de Chimie*, xxxiii. 52.



Plasters are employed as local remedies to answer various indications. When the materials of which they are formed are soft and bland, they are used simply as coverings to sores and abraded surfaces, to protect them from the action of the air, and give support to the parts; but in many instances they contain acrid and stimulating substances, and operate as rubefacients, or as blisters.

**EMPLASTRUM AMMONIACI**, Lond. Edin. *Ammoniacum Plaster.*

“Take of Purified Ammoniacum, *five ounces*; Acetic Acid, (*distilled vinegar*,) *half a pint*. Dissolve the Ammoniacum in the Vinegar, then evaporate the solution in an iron vessel placed in a water-bath, constantly stirring until it acquire a proper consistence.”

This plaster is stimulant and resolvent. It is applied to scrophulous tumours and white swellings; and sometimes over the scalp, in tinea capitis.

**EMPLASTRUM AMMONIACI, CUM HYDRARGYRO**, Lond. *Ammoniac Plaster with Mercury.*

“Take of Purified Ammoniacum, *a pound*; Purified Mercury, *three ounces*; Sulphuretted Oil, *a fluid drachm*. Rub the Mercury with the Sulphuretted Oil until the globules disappear; then add gradually the Ammoniacum previously melted, and mix the whole together.”

*Dublin.*

“Take of pure Gum Ammoniacum, *a pound*; Purified Mercury, *three ounces*; Turpentine, *two drachms*. Rub the Mercury with the Turpentine until the globules disappear; then add gradually the Ammoniacum previously melted, and melt the whole together.”

In these plasters the Mercury is probably in the state of oxide, with a minimum of oxygen. They are discutients, and are applied to indurated glands, hydarthus, nodes, tophi and indolent tumours.

**EMPLASTRUM AROMATICUM**, Dub. *Aromatic Plaster.*

“Take of Frankincense, *three ounces*; Yellow Wax, *half an ounce*; Cinnamon Bark in powder, *six drachms*; Oil of Pimenta, Oil of Lemons, of each, *two drachms*. Melt the Frankincense and the Wax together, and strain the mixture; when it thickens by cooling, mix with it the Powder of Cinnamon previously rubbed with the oils, and form them into a plaster.”

This plaster, which is an elegant stimulant, is applied on the region of the stomach in dyspepsia, and increased irritability of that viscus, to allay pain and vomiting; and to expel flatus. As the oils are very volatile, it must be spread with the thumb



without being melted. It requires to be frequently renewed, and is consequently not very adhesive.

EMPLASTRUM ASSAFOETIDÆ, Edin. *Assafoetida Plaster.*

“Take of Plaster of Semivitreous Oxide of Lead, Assafoetida, of each, *two parts*; Galbanum, Yellow Wax, of each, *one part*.”

This plaster is sometimes applied over the umbilical region, in flatulence and hysteria.

EMPLASTRUM CALEFACIENS, Dub. *Warm Plaster.*

“Take of Plaster of Cantharides, *one part*; Burgundy Pitch, *seven parts*. Melt them together with a moderate heat, and mix them so as to form a plaster.”

This plaster is stimulant and rubefacient, and is applied with advantage in catarrh, hooping-cough, sciatica and local pains. The proportion of plaster of Cantharides is rather too great, as it seldom fails to blister or produce a running sore in the majority of persons who use it.

EMPLASTRUM CERÆ, Lond. *Wax Plaster.*

“Take of Yellow Wax, Prepared Suet, of each, *three pounds*; Yellow Resin, *a pound*. Melt them together, and strain.”

EMPLASTRUM SIMPLEX, Edin. *Simple Plaster.*

“Take of Yellow Wax, *three parts*; Mutton Suet, White Resin, of each, *two parts*.”

These plasters were originally intended for dressing blistered parts, with the view of promoting a discharge; but owing to the pain and irritation they induce, they are now seldom employed. They may be spread with a hot iron.

Officinal preparation. *Emplastrum Lyttæ*, L.

EMPLASTRUM CUMINI, Lond. *Cumin Plaster.*

“Take of Cumin Seeds, Carraway Seeds, Laurel Berries, of each, *three ounces*; Dried Pitch, *three pounds*; Yellow Wax, *three ounces*. Melt the Pitch and the Wax together, then add the other ingredients in powder, and mix.”

This plaster is stimulant and discutient. It is applied to the hypogastric region in flatulence and a cold feeling of the bowels, and to indolent tumours.

EMPLASTRUM GALBANI COMPOSITUM, Lond. *Compound Galbanum Plaster.*

“Take of Purified Galbanum, *eight ounces*; Plaster of Lead, *three pounds*; Common Turpentine, *ten drachms*; Resin of the Spruce-fir powdered, *three ounces*. Having melted the Galbanum and the Turpentine together, mix in first the Resin, and then the Plaster of Lead previously melted by a slow fire, and mix the whole together.”

EMPLASTRUM GUMMOSUM, Edin. *Gum Plaster.*

“Take of Plaster of Semivitreous Oxide of Lead, *eight*



*parts* ; Ammoniacum Gum-resin, Galbanum, Yellow Wax, of each, *one part*. Add the Gum-resins to the melted Plaster and Wax, and mix."

EMPLASTRUM GALBANI, Dub. *Plaster of Galbanum*.

"Take of Litharge Plaster, *two pounds* ; Galbanum, *half a pound* ; Yellow Wax sliced, *four ounces*. To the Galbanum, melted by heat add the Litharge Plaster and the Wax ; then melt the whole together by a gentle heat."

These plasters are stimulant and suppurative. They are applied with advantage to scrophulous tumours ; to joints which have been long affected with arthritic pains ; and to the loins in rickets. As a suppurative they are applied to indolent tumours, and to reduce the induration which often remains around abscesses, after they are discharged.

EMPLASTRUM HYDRARGYRI, Lond. *Mercurial Plaster*.

"Take of Purified Mercury, *three ounces* ; Sulphuretted Oil, *a fluid drachm* ; Plaster of Lead, *a pound*. Rub the Mercury with the Sulphuretted Oil until the globules disappear ; then add by degrees the Lead Plaster melted, and mix the whole."

*Edinburgh.*

"Take of Olive Oil, Resin, of each, *one part* ; Mercury, *three parts* ; Plaster of Semivitreous Oxide of Lead, *six parts*. Rub the Mercury with the Oil and the Resin previously melted together and cooled, until the globules disappear ; then add gradually the Plaster of Semivitreous Oxide of Lead melted, and let the whole be carefully mixed together."

*Syn. Quecksilberpflaster (G.).*

The Mercury in these plasters is supposed to be in the state of oxide, with a minimum of oxygen ; and the Sulphuretted Oil, ordered by the London College, is intended to diminish the labour required for this oxidizement of the metal. The plasters are powerful discutients, and are applied to buboes, venereal tumours, nodes, when they are not very painful to the touch, and indurations : they are also applied to joints affected with obstinate syphilitic pains.

EMPLASTRUM CANTHARIDIS, Lond. *Blistering Plaster*.

"Take of Blistering Flies reduced to a very fine powder, *a pound* ; Wax Plaster, *a pound and a half* ; Prepared Lard, *a pound*. Melt the Plaster and the Lard together, and having removed them from the fire, when the mixture is just about to become solid, sprinkle in the blistering flies, and mix the whole together."

EMPLASTRUM CANTHARIDIS VESICATORIÆ, Edin. *Blistering Plaster*.



“Take of Mutton Suet, Wax, White Resin, Blistering Flies reduced to a very fine powder, of each, *equal weights*. Mix the powder with the other articles previously melted together, and removed from the fire; then stir until the mixture stiffens in cooling.”

EMPLASTRUM CANTHARIDIS, Dub. *Blistering Plaster*.

“Take of Purified Yellow Wax, Mutton Suet, of each, *a pound*; Yellow Resin, *four ounces*; Blistering Flies in fine powder, *a pound*. Melt the Wax, the Suet, and the Resin together, and a little before they concrete in becoming cold, sprinkle in the Blistering Flies, and form the whole into a plaster.”

*Syn.* Emplâtre de Cantharides (*F.*), Kanthariden pflaster (*G.*), Emplastro di Cantarelle (*I.*).

These plasters are of a moderately soft consistence, so as to admit of being spread without the assistance of heat, which destroys the acrimony and epispastic property of the flies; but they seldom fail of raising a blister, if the flies be good, and have not been added when the other ingredients were too hot. When they are to be used, a piece of leather of a proper shape and size is first spread with adhesive plaster, and over this the blistering plaster is extended of a moderate degree of thickness, and as smooth as possible, by means of the thumb; a proper margin being left, so as to enable it to adhere closely to the skin; but they should not be bandaged down. There is, however, an evident waste of flies, as those flies only which are on the surface of the plaster, when it is spread, act on the skin; and it has been suggested by Parmentier<sup>1</sup>, that the same effect would be more economically produced by sprinkling the powdered flies on a piece of farinaceous paste, spread on linen or leather. Blistering plasters require to remain applied for twelve hours to raise a perfect blister; they are then to be removed, the vesicle to be cut at the most depending part, and without removing the cuticle, the vesicated part is to be dressed with simple cerate or spermaceti ointment spread on lint; and the old cuticle allowed to remain until a new one is formed under it, when it peels off, and the whole is healed in the course of a few days. The application of these plasters, however, is sometimes attended with strangury and bloody urine, which arise from the active principles of the insect being absorbed, and irritating the kidneys and urethra. This effect is very much increased if the blister be applied over an abraded surface, as, for example, on the head immediately after it has been shaved; and it also occurs if the plaster remain too long applied. To prevent strangury, camphor has been recom-

<sup>1</sup> *Annales de Chimie*, xlviij.



mended to be mixed with the blistering composition, but it has no good effect; and it is better obviated by copious dilution with milk, or mucilaginous fluids, and fomentations of warm milk and water to the blistered part, after the removal of the plaster. When the head is the part intended to be blistered, it should be shaved at least ten hours before the plaster is applied; and in all cases it is perhaps a good rule to interpose a thin piece of gauze between the vesicatory and the skin, wetted with vinegar, and applied smooth and very close over the plaster.

In persons of delicate habits, and those labouring under some diseases of irritation, particularly in children, the blistered part, instead of healing kindly, becomes a spreading sore: the cutis vera is destroyed, and the part cannot be healed until the irritability of habit which induced this unpleasant state is allayed. In such cases, the blistering plaster should not be allowed to remain on longer than is requisite to raise the cuticle: but if ulceration have commenced, the best local application is a warm emollient poultice; and bathing the denuded surface frequently with tepid milk and water; while at the same time Cinchona bark is internally administered.

**EMPLASTRUM CANTHARIDIS VESICATORIÆ COMPOSITUM**, Edin. *Compound Plaster of Spanish Flies.*

“Take of Venice Turpentine, *eighteen parts*; Burgundy Pitch, Blistering Flies, of each, *twelve parts*; Yellow Wax, *four parts*; Subacetate of Copper, *two parts*; White Mustard Seeds, Black Pepper, of each, *one part*. Melt the Burgundy Pitch and the Wax, and add to them the Turpentine. While these remain still warm, after being melted, sprinkle in the other ingredients reduced to fine powder, and mix them, stirring constantly, so as to form a plaster.”

This plaster is intended to raise a blister more quickly than the former; and thence is adapted for cases of rheumatism and cramps of the stomach, in which the effect of the blister is almost instantly produced. Its operation is accompanied with great pain, and a very pungent sense of heat; and it is apt to produce very unpleasant ulceration if allowed to remain too long applied.<sup>1</sup>

**EMPLASTRUM OPII**, Lond. Edin. *Plaster of Opium.*

“Take of hard Opium powdered, *half an ounce*; Resin of the Spruce-fir powdered, *three ounces*; Lead Plaster, *a pound*. Melt the Plaster and the Resin together, then add the Opium, and mix the whole.”

---

<sup>1</sup> Blisters are rapidly raised by digesting Cantharides in Acetic Ether, in the proportion of ʒj. of the powdered flies, to fʒj. of the Ether. They are still more quickly raised by the application of boiling water.



*Syn.* Opiumspflaster (*G*).

This plaster is anodyne, and supposed to be useful in relieving rheumatism and local pains: but although it is undoubtedly certain that opium, in that state of minute division in which it exists in the tincture, or when it is dissolved in oil, produces its specific effect on the system when externally applied, yet, we doubt whether the anodyne properties of this plaster are such as to sanction its employment.

**EMPLASTRUM OXIDI FERRI RUBRI,** Edin.  
*Plaster of Red Oxide of Iron.*

“Take of Plaster of Semivitreous Oxide of Lead, *twenty-four parts*; White Resin, *six parts*; Yellow Wax, Olive Oil, of each, *three parts*; Red-oxide of Iron, *eight parts*. Rub the Red-oxide of Iron with the Oil, and adding the other ingredients melted, mix the whole well together.”

**EMPLASTRUM THURIS,** Dub. *Plaster of Frankincense.*

“Take of Litharge Plaster, *two pounds*; Frankincense, *half a pound*; Red-oxide of Iron, *three ounces*. To the Plaster and Frankincense melted together, add the Oxide, stirring them together so as to form a Plaster.”

These plasters are supposed to be tonic, and are used in muscular relaxations, and weakness of the joints after sprains; but they act chiefly in affording mechanical support to the parts.

**EMPLASTRUM PICIS COMPOSITUM,** Lond.  
*Compound Pitch Plaster.*

“Take of Burgundy Pitch, *two pounds*; Resin of the Spruce-fir, *a pound*; Yellow Resin, Yellow Wax, of each, *four ounces*; expressed Oil of Nutmeg, *an ounce*; Olive Oil, Water, of each, *two fluid ounces*. To the Pitch, Resin, and Wax, melted together, add first the Resin of the Spruce-fir, then the Oil of Nutmeg, the Olive Oil, and the Water: mix the whole, and reduce to a proper consistence.”

This plaster is stimulant and rubefacient. It is used in catarrh, and other pulmonary affections, applied to the thorax; and in headach and chronic ophthalmia, applied to the temples. When a serous exudation takes place, the plaster should be frequently renewed.

**EMPLASTRUM PLUMBI,** Lond. *Lead Plaster.*

“Take of Semivitreous Oxide of Lead, rubbed to a very fine powder, *five pounds*; Olive Oil, *a gallon*; Water, *two pints*. Boil them together over a slow fire, stirring constantly until the Oil and Oxide of Lead cohere into the consistence of a plaster. It will be necessary, however, to add a little Boiling Water, if that which was employed in the beginning shall be consumed before the end of the process.”



EMPLASTRUM OXIDI PLUMBI SEMI-VITREI, Edin. *Plaster of Semivitreous Oxide of Lead.*

“Take of the Semivitreous Oxide of Lead, *one part*; Olive Oil, *two parts*; Water, *a sufficient quantity*. Boil them, stirring constantly, until the Oil and the Oxide unite into a Plaster.”

EMPLASTRUM LITHARGYRI, Dub. *Litharge Plaster.*

“Take of Litharge in fine powder, *five pounds*; Olive Oil, *nine pounds*; Boiling Water, *two pints*. Mix them at a high temperature, constantly stirring, until the Oil and the Litharge unite so as to form a Plaster; supplying occasionally any waste of water that may take place.”

*Syn.* Emplatre de diachylon (F.), Bleipflaster (G.).

The use of the water in the formation of these plasters is to moderate the heat of the mixture, until the oil and the oxide combine, by which means the reduction of the metal is prevented: a circumstance which is apt to take place from the strong attraction of the oil for oxygen when raised to a high temperature. By continuing the boiling, the water is dissipated; and the temperature can then be increased to a sufficient degree to give the plaster the necessary consistency. The water which is added should be previously made hot; as cold water is apt to produce an explosion. When long kept, these plasters change their colour, and lose most of their sensible properties. They are intended chiefly to defend excoriated surfaces from the action of the air; and to form the basis of some other plasters.

Officinal preparations. *Emplastrum Hydrargyri*, L. E. *Emplastrum Opii*, L. *Emplastrum Assafoetida*, E. *Emplastrum gummosum*, E. *Emplastrum Galbani*, D. *Emplastrum Galbani compositum*, L. *Emplastrum Oxidi Ferri rubri*, E. *Emplastrum Resinae*, L. E. D. *Emplastrum Saponis*, L. E. D. *Emplastrum Thuris*, D.

EMPLASTRUM RESINÆ, Lond. *Resin Plaster.*

“Take of Yellow Resin, *half a pound*; Lead Plaster, *three pounds*. Melt the Lead Plaster by a gentle heat, then add the Resin in powder, and mix.”

EMPLASTRUM RESINOSUM, Edin. *Resinous Plaster.*

“Take of Plaster of Semivitreous Oxide of Lead, *five parts*; Resin, *one part*. Melt them with a gentle heat; then continue stirring the mixture until it becomes stiff in cooling.”

EMPLASTRUM LITHARGYRI CUM RESINA, Dub. *Litharge Plaster with Resin.*

“Take of Litharge Plaster, *three pounds and a half*; Yellow Resin, *half a pound*. Melt the Litharge Plaster by a moderate heat, then add the Resin reduced to a very fine powder, that it may melt quickly, and form a Plaster.”

*Syn.* Harzigtes Bleipflaster (G.).



These plasters are defensive, adhesive, and gently stimulant. They are used for retaining together the lips of recent wounds, when it is wished to heal them by the first intention; to give support to ulcerated parts; and to assist their granulation and cicatrization, according to the excellent method of Mr. Baynton. The plaster, however, originally used by Mr. Baynton, contained less resin; 3vi. only being added to lb. j. of the litharge plaster; but this preparation answers the purpose equally well, except in very irritable habits. The best substance for spreading it on for the above purpose is calico; and it is of some importance to spread it equally and thin; to effect which the calico must be stretched, and the plaster melted and beginning to cool, must be poured on one end of it, and equally extended over the whole surface by means of a spatula, held horizontally, and one edge of the blade raised to an angle of 45 degrees: or it may be still more equally done by passing the calico, on which the fluid plaster has been poured, through a machine formed of a straight blade of steel, fixed by screws, at a proper distance from a polished plate of the same metal. It is sold ready spread.

EMPLASTRUM SAPONIS, Lond. Dub. *Soap Plaster.*

“Take of Hard Soap sliced, *half a pound*; Lead Plaster, *three pounds*. Mix the Soap with the melted Plaster; then boil it down to a proper consistence.”

EMPLASTRUM SAPONACEUM, Edin. *Soap Plaster.*

“Take of Semivitreous Oxide of Lead, *four parts*; Gum Plaster, *two parts*; Soap sliced, *one part*. Mix the Soap with the Plasters melted together; then boil them a little so as to form a Plaster.”

*Syn.* Seifenpflaster (G.).

Dr. Powell properly observes, that the Soap Plaster of the London College “must be formed into rolls when it begins to thicken, for afterwards, although it be still somewhat soft, it loses its tenacity, and will break to pieces.”<sup>1</sup>

Soap plaster is discutient; and is applied to lymphatic tumours: but it is much less useful than the mercurial plaster.

## CERATA.

### CERATES.

THESE are unctuous compositions possessing a certain degree of firmness, intermediate between that of plasters and

<sup>1</sup> *Powell's Translation of the London Pharmacopœia*, 2d. edit. 324.



that of ointments. Their consistence depends on the Wax they contain; and from it they derive their generic appellation. The most important circumstance to be attended to in their preparation is the freshness of the fat and the oils employed; and their preservation in this state.

**CERATUM SIMPLEX**, Lond. *Cerate.*

“Take of Olive Oil, *four fluid ounces*; Yellow Wax, *four ounces*. Add the Oil to the melted Wax, and mix.”

*Syn.* Cerat simple (F.), Cerotto semplice (I.).

This is an useful simple emollient dressing to excoriations and sores.

**CERATUM CALAMINÆ**, Lond. *Calamine Cerate.*

“Take of Prepared Calamine, Yellow Wax, of each, *half a pound*; Olive Oil, *a pint*. Mix the Oil with the melted Wax; then remove the mixture from the fire, and as soon as it begins to thicken, add the Calamine, stirring constantly until it be cold.”

**CERATUM CARBONATIS ZINCI IMPURI**, Edin. *Cerate of Impure Carbonate of Zinc.*

“Take of Simple Cerate, *five parts*; Prepared Impure Carbonate of Zinc, *one part*. Mix.”

**UNGUENTUM CALAMINARE**, Dub. *Calamine Ointment.*

“Take of Ointment of Yellow Wax, *five pounds*; Prepared Calamine, *a pound*. Make them into an ointment.”

These preparations are very useful dressings to excoriations and ulcers; and as they are in some degree desiccative, they are also applied to burns after the inflammation is abated; and to the eye-lids in ophthalmia tarsi. They have been long known in practice under the name of *Turner's Cerate*.

**CERATUM CETACEI**, Lond. *Spermaceti Cerate.*

“Take of Spermaceti, *half an ounce*; White Wax, *two ounces*; Olive Oil, *four fluid ounces*. Melt the Spermaceti and the Wax together, then add the Oil, and stir them until they be cold.”

**CERATUM SIMPLEX**, Edin. *Simple Cerate.*

“Take of Olive Oil, *six parts*; White Wax, *three parts*; Spermaceti, *one part*. Melt the Wax and Spermaceti in the Oil, with a gentle heat; then keep constantly stirring until the mixture stiffen in cooling.”

*Syn.* Cerat de blanc de balaine (F.), Cerotto di Spermaceti (I.).

These are soft cooling dressings.

Officinal preparations. *Ceratum Cantharidis*, L. *Ceratum Carbonatis Zinci impuri*, E.

**CERATUM CANTHARIDIS**, Lond. *Cerate of Blistering Flies.*

“Take of Spermaceti Cerate, *six drachms*; Blistering Flies reduced to a very fine powder, *a drachm*. Add the Blister-



ing Flies to the Cerate softened by the fire, and mix them together."

*Syn.* Cerat de Cantharides (*F.*), Cerotto di Cantarille (*I.*).

This Cerate is intended to promote a purulent discharge from a blistered surface; and in general it answers this intention without occasioning much irritation. In some habits, however, it occasions strangury, great pain of the part, swellings of the lymphatics, and so much general irritation as to produce œdematose swellings and erysipelas of the neighbouring parts.<sup>1</sup>

It may be proper to observe, that it is preferable to spread Cerates or Ointments, intended to keep open issues, on lint; and that the dressings should in all cases be renewed once in twenty-four hours.

**CERATUM PLUMBI ACETATIS, Lond.** *Cerate of Acetate of Lead.*

"Take of Acetate of Lead in powder, *two drachms*; White Wax, *two ounces*; Olive Oil, *half a pint*. Melt the Wax in seven fluid ounces of the Oil; then add, gradually, the Acetate of Lead separately rubbed down with the remaining oil, and stir with a wooden spatula, until they be thoroughly incorporated."

This is an excellent cooling Cerate for burns, excoriations, and other inflamed sores.

**CERATUM PLUMBI COMPOSITUM, Lond.** *Compound Cerate of Lead.*<sup>2</sup>

"Take of Solution of Acetate of Lead, *two fluid ounces and a half*; Yellow Wax, *four ounces*; Olive Oil, *nine fluid ounces*; Camphor, *half a drachm*. Melt the Wax and mix it with eight fluid ounces of the Oil; then remove them from the fire, and as soon as they begin to thicken, add gradually the solution of Acetate of Lead, and stir assiduously with a wooden spatula till they be cold. Finally, mix with these the Camphor dissolved in the remainder of the Oil."

This composition is similar to what was recommended by Goulard, as a mode of applying lead in the form of ointment, and long known under the name of *Goulard's Cerate*. It is applicable to the same cases as the former cerate.

**CERATUM RESINÆ, Lond.** *Resin Cerate.*

"Take of Yellow Resin, Yellow Wax, of each, *a pound*;

<sup>1</sup> In one case, which came under my observation, a blister on the scalp was dressed for four days with this cerate. On the fourth day the head swelled to an alarming size; and an œdematose erysipelas covered the scalp and face, and shut up the eyes; accompanied with a great degree of fever. On removing the acrid dressings, and employing emollient fomentations, with dressings of cetaceous ointment, these alarming symptoms soon subsided.

<sup>2</sup> This name is improper. It ought to have been *Ceratum Plumbi Acetatis comp.*; the virtue of the composition depending on the acetate of lead, not on lead.



Oil of Olive, *a pint*. Melt the Resin and the Wax together by a slow fire, then add the Oil, and strain the Cerate while it is hot through a linen cloth."

UNGUENTUM RESINOSUM, Edin. *Resinous Ointment*.

"Take of Hog's Lard, *eight parts*; Resin, *five parts*; Yellow Wax, *two parts*. Melt the whole with a gentle heat, and stir the mixture until it stiffen in cooling."

UNGUENTUM RESINÆ ALBÆ, Dub. *Ointment of White Resin*.

"Take of Yellow Wax, *a pound*; White Resin, *two pounds*; prepared Hog's Lard, *four pounds*. Make them into an ointment, which is to be strained, while it is hot, through a sieve."

*Syn.* Cerat Resineux (*F.*), Harzzerat (*G.*), Cerotto Resinoso (*I.*).

These ointments are stimulant, digestive, and cleansing; and, therefore, form an excellent dressing for foul and indolent ulcers.

Officinal preparation. *Linimentum Terebinthinæ*, L.

CERATUM SABINÆ, Lond. *Cerate of Savine*.

"Take of the Fresh Leaves of Savine bruised, *a pound*; Yellow Wax, *half a pound*; Prepared Lard, *two pounds*. Melt the Lard and the Wax together, and boil the Savine Leaves in the mixture; then strain through a linen cloth."

UNGUENTUM SABINÆ, Dub. *Savine Ointment*.

"Take Fresh Leaves of Savine freed from the stalks and bruised, *half a pound*; prepared Hog's Lard, *two pounds*; Yellow Wax, *half a pound*. Boil the Leaves with the Lard until they become crisp; then strain with expression: lastly, add the wax, and melt them together."

The preparation of this ointment is exceedingly difficult, as the acrid principle of the Savine, on which its efficacy depends, is much injured by long boiling, or too high a temperature. Might it not be better to express the acrid juice from the fresh leaves, and mix it with the ointment when it begins to thicken by cooling? If the fresh leaves cannot be procured, it may be prepared from the dry leaves reduced to a fine powder: but the acrimony of the savine is impaired by drying it. The ointment, when good, has a beautiful deep-green colour, and the odour of the fresh bruised herb. It should be kept in closely-covered pots, as it soon loses its virtue by exposure to the air.

Savine ointment, which was first described by Mr. Crowther<sup>1</sup>, is well calculated for keeping up a purulent discharge from a blistered surface; which it does as effectually, and with much less irritation than the ointment of blistering flies. A white coat is apt to form on the discharging surface, and

<sup>1</sup> *Observations on White Swelling.*



must be removed occasionally so as to allow the cerate to be applied to the sore.

**CERATUM SAPONIS, Lond.** *Cerate of Soap.*

“Take of Hard Soap, *eight ounces*; Yellow Wax, *ten ounces*; Semivitreous Oxide of Lead powdered, *a pound*; Olive Oil, *a pint*; Vinegar, *a gallon*. Boil the Vinegar on the Oxide of Lead over a slow fire, stirring diligently until they incorporate; then add the soap, and boil again in a similar manner, until the moisture be entirely evaporated: lastly, mix the wax previously melted with the oil.”

The efficacy of this cerate evidently depends on the acetate of lead which is formed in the first stage of the process, the soap answering scarcely any other purpose than to give consistence and adhesiveness. It is occasionally used as a cooling dressing.

## UNGUENTA.

### OINTMENTS.

THESE are unctuous substances of nearly the same nature as cerates, but having a consistence much less firm, scarcely exceeding that of butter.

**UNGUENTUM ACIDI NITROSI, Edin.** *Ointment of Nitrous Acid.*

“Take of Hog’s Lard, *one pound*; Nitrous Acid, *six drachms*. Mix the Acid gradually with the melted lard, and beat the mixture assiduously as it cools.”

*Dublin.*

“Take of Olive Oil, *a pound*; Prepared Hog’s Lard, *four ounces*; Nitrous Acid, *an ounce by weight*. Melt the Oil in a glass vessel, and add the Acid to it; let them be exposed to a medium heat in a water-bath for a quarter of an hour, then remove them from the bath, and stir them constantly with a glass rod until they become firm.”

In this process the Acid is partially decomposed, Nitrous gas is evolved, and the ointment is oxidized, assuming a yellow colour and a firm consistence. It was invented by Alyon, who found it useful in syphilitic and herpetic ulcers; and it has been occasionally used in this country for the same purposes; but it is less useful than the ointment of nitrate of mercury.

**UNGUENTUM CERÆ FLAVÆ, Dub.** *Ointment of Yellow Wax.*

“Take of purified Yellow Wax, *a pound*; Prepared Hog’s lard, *four pounds*. Form them into an ointment.”



UNGUENTUM CERÆ ALBÆ, Dub. *Ointment of White Wax.*

“This is to be prepared in the same manner as the former, with the substitution of White for Yellow Wax.”

These are useful dressings to benign ulcers and excoriations, and form the basis of the majority of the compound ointments of the Dublin Pharmacopœia.

UNGUENTUM CETA'CEI, Lond. *Spermaceti Ointment.*<sup>1</sup>

“Take of Spermaceti, *six drachms*; White Wax, *two drachms*; Olive Oil, *three fluid ounces*. Melt them together over a slow fire, and stir them constantly until they be cold.”

UNGUENTUM SPERMATIS CETI, Dub. *Ointment of Spermaceti.*

“Take of White Wax, *half a pound*; Spermaceti, *a pound*; Prepared Lard, *three pounds*. Make them into an ointment.”

These ointments form the ordinary dressings for healing blistered surfaces and excoriations.

UNGUENTUM ELEMI COMPOSITUM, Lond. *Compound Ointment of Elemi.*<sup>2</sup>

“Take of Elemi, *a pound*; Common Turpentine, *ten ounces*; Prepared Suet, *two pounds*; Olive Oil, *two fluid ounces*. Melt the Elemi with the Suet; then remove it from the fire, and mix in immediately the Turpentine and the Oil: lastly, strain the mixture through a linen cloth.”

UNGUENTUM ELEMI, Dub. *Ointment of Elemi.*

“Take of Elemi Resin, *a pound*; White Wax, *half a pound*; Prepared Hog's Lard, *four pounds*. Form them into an ointment, which is to be strained through a sieve while it is hot.”

*Syn.* Onguent d'Elemi et de Terebenthine (*F.*), Elemisalbe (*G.*), Unguento di Elemi e Trementina (*I.*).

These ointments are stimulant and digestive. They are used to keep open issues and setons; and as a dressing to ulcers which do not admit of the application of the adhesive straps.

UNGUENTUM HYDRAR'GYRI FOR'TIUS, Lond. *Strong Mercurial Ointment.*<sup>3</sup>

“Take of Purified Mercury, *two pounds*; Prepared Lard, *twenty-three ounces*; Prepared Suet, *an ounce*. First rub the Mercury with the Suet and a little of the Lard, until the globules disappear; then add the remainder of the fat, and mix.”

<sup>1</sup> Unguentum Spermaceti, P. L. 1787.

<sup>2</sup> Unguentum e Gummi Elemi, P. L. 1745.

<sup>3</sup> Ung. cæruleum fortius, P. L. 1745.



Two drachms of this ointment contain one drachm of mercury.

UNGUENTUM HYDRARGYRI, Edin. *Mercurial Ointment.*

“Take of Mercury, Mutton Suet, of each, *one part*; Hog’s Lard, *three parts*. Rub the Mercury diligently in a mortar with a little of the Hog’s Lard until the globules disappear; then add the remainder of the lard.”

“It may also be made with double or triple the quantity of Mercury.”

One drachm of this ointment contains twelve grains of Mercury.

*Dublin.*

“Take of Purified Mercury, Prepared Hog’s Lard, *equal weights*. Rub them together in a marble or an iron mortar until the globules disappear.”

*Syn.* Onguent. Mercuriale (*F.*), Quecksilbersalbe (*G.*), Unguento Mercuriale (*L.*).

One drachm of this ointment contains thirty grains of mercury.

UNGUENTUM HYDRAR’GYRI MITIUS<sup>1</sup>, Lond. Dub. *Milder Mercurial Ointment.*

“Take of the Stronger Mercurial Ointment, *a pound*; Prepared Lard, *two pounds*. Mix them.”

One drachm of this ointment contains ten grains of Mercury; but prepared according to the Dublin Pharmacopœia, with two parts of Lard to one of Mercury, one drachm contains a scruple of Mercury.

The preparation of the stronger Mercurial Ointments requires much labour, care, and patience. During the trituration the Mercury is mechanically divided into minute globules, which are prevented from running together again by the viscosity of the suet. They are by some supposed to be gradually oxidized during the trituration by attracting the oxygen of the atmosphere; the lard, the extension and the constant renewal of the surface exposed, favouring very much this effect. The fact of the oxidizement of the metal in this process, however, has been often questioned; and lately some experiments of M. Roux have thrown much light on the subject. That chemist triturated Mercury and Maltha, a species of pitch, in a vacuum, and produced the extinction of the metal as well as if the operation had been performed in the air; thence he concludes that the metal is not oxidized, but merely mechanically divided in the ointment. There are still, however, some difficulties in deciding this point. Whatever tends to favour oxidation, as for instance,

<sup>1</sup> Unguentum cæguleum mitius, P. L. 1745.



a slight degree of rancidity of the lard, or the oil of eggs, shortens the time, and lessens the labour required for the preparation of the ointment. It is not uncommon, however, to use other means, which are not admissible to facilitate the process, such as the use of sulphur or turpentine. The first may be detected by the very black colour of the ointment, which is produced by the sulphuret of mercury; and also by the sulphurous odour exhaled, when a paper covered with a little of it is held over the flame of a candle: and the turpentine is detected by its odour also, when the ointment containing it is treated in the same manner. When newly prepared, Mercurial ointment has a light gray or blueish colour; this has been ascribed to its containing some unoxidized metal, which separates in globules when it is liquefied by a gentle heat: when kept for some time the colour is much deepened, and less metallic mercury subsides, owing to the more complete oxidizement of the metal. It is probable, therefore, that long kept Mercurial ointment contains, besides the Oxide, a Sebate of mercury. As it is of great consequence to procure so important a preparation always of the same degree of strength, and as this can never be accomplished by the present method of preparing the ointment, Mr. Donovan has proposed to prepare it by rubbing together, lard and black oxide of mercury, at the temperature of  $350^{\circ}$  Fahrenheit, continuing the friction for two hours. By this method of proceeding, Mr. Donovan found that every ounce of lard dissolves and combines with twenty-one grains of oxide; that the ointment thus prepared can be introduced into the habit in one-third of the time required by the common ointment: and that it is equally efficient with the officinal preparation.

*Medical properties and uses.* — The Strong Mercurial Ointment rubbed upon the skin introduces a large quantity of mercury into the system. About 3j. is rubbed upon the inside of the thighs, or any other part of the body where the cuticle is thin, every night and morning, until the system is affected. The ointment is absorbed during the friction, and carried into the habit; where it produces the same effects as those which result from taking the remedy by the mouth, without the unpleasant affection of the bowels that very commonly follows the introduction of preparations of Mercury into the stomach. In order, however, to produce the full effect of the friction, it must be continued until every particle of the ointment disappear; and the operation should be performed by the patient himself. The Stronger Mercurial ointment is used in this form as an antisyphilitic, as a deobstruent in hepatic affections, and to excite the absorbents in hydrocephalus. The weaker ointment is used only as a topical dressing in



venereal sores. During a course of mercurials the patients should be kept in a moderately warm and dry, but airy chamber; and his diet should be chiefly weak broths, milk, and gruel.

The following Table shows at one view the quantity of mercury contained in each of the different ointments ordered by the British Colleges.

one drachm.	{	of the Lond.	{ stronger ointment contains of merc. 30 grs.	
			{ weaker ointment _____	10
	{	of the Edin.	common ointment _____	12
		of the Dub.	{ stronger ointment _____	30
			{ weaker ointment _____	20

UNGUENTUM OXIDI HYDRARGYRI CINEREI, Edin. *Ointment of gray Oxide of Mercury.*

“Take of Gray Oxide of Mercury, *one part*; Hog’s Lard, *three parts*. Mix.”

*Syn.* Graue Quecksilbersalbe (G.).

As the whole of the mercury in this ointment is oxidized, it might, *à priori*, be supposed that it would answer all the purposes of the mercurial ointment; but it cannot be so easily introduced by friction, the oxide remaining on the surface of the cuticle after the unctuous matter is absorbed. Dr. Paris justly remarks, that this is owing to its being a mechanical mixture instead of a chemical combination; an opinion, however, which is rendered doubtful by the experiments made to prove the non-oxidation of the mercury, in the preparation of the mercurial ointment. It has been too seldom employed to enable a correct judgment to be formed of its efficacy.

UNGUENTUM HYDRARGYRI NITRATIS, Lond. *Ointment of Nitrate of Mercury.*

“Take of Purified Mercury, *an ounce*; Nitric Acid, *eleven fluid drachms*; Prepared Lard, *six ounces*; Olive Oil, *four fluid ounces*. First dissolve the Mercury in the Acid; then mix the solution while it is hot, with the Lard and Oil melted together.”

UNGUENTUM NITRATIS HYDRARGYRI FORTIUS; vulgo, UNGUENTUM CITRINUM, Edin. *Stronger Ointment of Nitrate of Mercury.*

“Take of Purified Mercury, *one part*; Nitrous Acid, *two parts*; Olive Oil, *nine parts*; Hog’s Lard, *three parts*. Dissolve the Mercury in the Acid; then beat up the solution strongly with the Lard and Oil previously melted together and nearly cold, in a glass mortar, so as to form an ointment.”

UNGUENTUM SUPERNITRATIS HYDRARGYRI, Dub. *Ointment of Supernitrate of Mercury.*

“Take of Purified Mercury, *an ounce*; Nitrous Acid, *two*



*ounces by weight* ; Olive Oil, *one pint* ; Hog's Lard, *four ounces*. Dissolve the Mercury in the Acid ; then mix the solution with the Oil and Lard previously melted together, and form an Ointment in the same manner as the Ointment of Nitrous Acid."

*Syn.* Onguent citrin (*F.*), Gelbe Quecksilbersalbe (*G.*), Unguento Citrino (*I.*).

UNGUENTUM NITRATIS HYDRARGYRI MITIUS, Edin. *Milder Ointment of Nitrate of Mercury.*

"It is made in the same manner as the stronger ointment, with a triple proportion of Oil and Lard."

In all of these formulæ too large a proportion of lard is used : for the excess of acid in the metallic solution oxidizing the fatty matters, occasions them to become too hard and brittle after the ointment has been kept for some time, when more than one-sixth of lard is employed. The addition of the metallic solution to the melted mixture of lard and oil should be gradual, and made in a broad flat vessel, so as to expose a large surface to the action of a current of air ; while the stirring should be performed with a wooden spatula, and continued until the ointment be perfectly cold.

When prepared in the above manner, and with one-sixth part only of lard, this ointment has a beautiful golden colour, and is of the consistence of butter, which it retains if preserved in close pots ; but when made with a larger proportion of lard, it becomes hard, brittle, and of a pale dirty yellow hue, marbled with green blotches.

*Medical properties and uses.*— This ointment is stimulant and detergent. When moderately diluted with lard it is a local remedy of great efficacy in herpetic eruptions, psoriasis, porrigo, and other cutaneous eruptions. The weaker ointment may almost be regarded as a specific in psorophthalmia, in the purulent ophthalmia of infants producing ectropium, and in ulcerations of the tarsi. It is applied by taking a little on the finger, liquefying it by the fire or the flame of a candle, and applying it along the inner part of the eyelids.

UNGUENTUM GALLÆ, Edin. *Ointment of Galls.*

"Take of Galls, in fine powder, *one part* ; Lard, *eight parts*. Mix."

This is a very useful application in piles, and has long been employed, although it has never before obtained a place in the pharmacopœia.

UNGUENTUM HYDRAR'GYRI NIT'RICO OX'Y-DI, Lond. *Ointment of Nitric Oxide of Mercury.*

"Take of Nitric Oxide of Mercury, *an ounce* ; White Wax, *two ounces* ; Prepared Lard, *six ounces*. Melt together the Wax and Lard, then add to the mixture the Nitric Oxide of Mercury in very fine powder, and mix."



UNGUENTUM OXIDI HYDRARGYRI RUBRI, Edin. *Ointment of Red Oxide of Mercury.*

“Take of Red Oxide of Mercury by Nitric Acid in fine powder, *one part*; Hog’s Lard, *eight parts*. Mix.”

UNGUENTUM SUBNITRATIS HYDRARGYRI, Dub. *Ointment of Subnitrate of Mercury.*

“Take of Ointment of White Wax, *half a pound*; Subnitrate of Mercury, *half an ounce*. Form them into an ointment.”

*Syn.* Rothe Quecksilbersalbe (G.).

These are excellent stimulant ointments, well adapted for giving energy to indolent foul ulcers. They are also of great use in inflammation of the conjunctiva, with a thickening of the inner membrane of the palpebræ; and to specks of the cornea. They are to be applied in the same manner as the ointment of Nitrate of Mercury.

UNGUENTUM HYDRARGYRI PRÆCIPITATI ALBI<sup>1</sup>, Lond. *Ointment of White Precipitate of Mercury.*

“Take of White Precipitate of Mercury, *a drachm*; Prepared Lard, *an ounce and a half*. Add the Precipitated Mercury to the Lard previously melted by a gentle heat, and mix.”

UNGUENTUM SUBMURIATIS HYDRARGYRI AMMONIATI, Dub. *Ointment of Ammoniated Submuriate of Mercury.*

“Take of Ointment of White Wax, *a pound*; Ammoniated Submuriate of Mercury, *an ounce and a half*. Form them into an ointment.”

These ointments are stimulant and detergent. They are recommended by Werlhoff, and some other German authors, as a remedy for the itch. They may be safely used on infants.

UNGUENTUM CANTHARIDIS<sup>2</sup>, Lond. *Blistering Ointment.*

“Take of Blistering Flies finely powdered, *two ounces*; Distilled Water, *eight fluid ounces*; Resin Cerate, *eight ounces*. Boil the Water with the Blistering Flies to half its quantity, and strain. Mix the Cerate into the strained liquor, and evaporate it to a proper consistence.”

*Syn.* Kantharidensalbe (G.).

UNGUENTUM INFUSI CANTHARIDIS VESICATORIÆ, Edin. *Ointment of Infusion of Blistering Flies.*

“Take of Blistering Flies, Resin, Yellow Wax, of each, *one part*; Venice Turpentine, Hog’s Lard, of each, *two parts*; Boiling Water, *four parts*. Macerate the flies in the Water for a night, and strain the liquor, expressing it strongly; add the liquor to the fat, and boil until the water be evaporated;

<sup>1</sup> Unguentum e Mercurio præcipitato albo, P. L. 1745. Ung. Calcis Hydrargyri albi, P. L. 1787.

<sup>2</sup> Unguentum Cantharidis, P. L. 1787.



then add the Wax and the Resin, and when these are melted, remove the mixture from the fire; add the Venice Turpentine, and mix."

These ointments are sufficiently mild, but they do not always succeed in keeping open a blistered surface, the purpose for which they are designed. Little efficacy can be ascribed to the blistering flies, the acrimony of which is destroyed by the heat employed for the evaporation of the water.<sup>1</sup>

UNGUENTUM JUNIPERI SABINÆ, Edin. *Ointment of Savine.*

"Take of Fresh Leaves of Savine, *two parts*; Yellow Wax, *one part*; Lard, *four parts*. Melt the Wax and Lard together, then boil the Leaves in the mixture, and express through a cloth."

This ointment is intended for keeping a blistered surface discharging, and answers the purpose of the cerate of savine.

UNGUENTUM CARBONATIS PLUMBI, Edin. *Ointment of Carbonate of Lead.*

"Take of Simple Ointment, *five parts*; Carbonate of Lead, *one part*. Mix."

UNGUENTUM CERUSSÆ sive SUBACETATIS PLUMBI, Dub. *Ointment of Cerussa, or Subacetate of Lead.*

"Take of Ointment of White Wax, *a pound*; Cerussa reduced to a very fine powder, *two ounces*. Form them into an ointment."

*Syn.* Onguent blanc (*F.*), Bleiweissalbe (*G.*), Unguento bianco (*I.*).

These are useful, cooling, desiccative ointments.

UNGUENTUM OXIDI ZINCI IMPURI, Edin. *Ointment of Impure Oxide of Zinc.*

"Take of Simple Liniment, *five parts*; Prepared Impure Oxide of Zinc, *one part*. Mix."

UNGUENTUM TUTIÆ, Dub. *Ointment of Tutty.*

"Take of Ointment of White Wax, *ten ounces*; Prepared Tutty, *two ounces*. Form them into an ointment."

*Syn.* Onguent de Tuthie (*F.*), Unguento di Tuzia (*I.*).

These ointments were formerly much used in ophthalmia tarsi; but they are now seldom employed.

UNGUENTUM PICIS LIQUIDÆ<sup>2</sup>, Lond. *Tar Ointment.*

"Take of Tar, Prepared Suet, of each, *a pound*. Melt them together, and strain the mixture through a linen cloth."

UNGUENTUM PICIS LIQUIDÆ, Edin. *Tar Ointment.*

<sup>1</sup> Galen employed an ointment made by macerating the entire insect in melted lard for twenty-four hours, and then straining by expression. Boerhaave proposed to boil the flies in water, then to pour off the liquid, and make an ointment of the boiled insects with the addition of lard.

<sup>2</sup> Unguentum e Pice, P. L. 1745. Unguentum Picis, P. L. 1787.



“Take of Tar, *five parts*; Yellow Wax, *two parts*. Melt the Wax with a gentle heat: then add the Tar, and stir until the mixture stiffen in cooling.”

UNGUENTUM PICIS LIQUIDÆ, Dub. *Tar Ointment.*

“Take of Tar, Mutton Suet, of each, *half a pound*. Melt them together, and strain them through a sieve.”

Although the pitch and the tar ointments differ in their sensible qualities, yet they are both used with advantage as detergents in scabby, foul eruptions, and tinea capitis.

UNGUENTUM PICIS NIGRÆ, Lond. *Ointment of Black Pitch.*

“Take of Black Pitch, Yellow Resin, of each, *nine ounces*; Olive Oil, *one pint*. Melt them together, and strain through a linen cloth.”

This ointment is digestive and stimulant.

UNGUENTUM PIPERIS NIGRI, Dub. *Ointment of Black Pepper.*

“Take of Prepared Hog’s Lard, *a pound*; Black Pepper in powder, *four ounces*. Form them into an ointment.”

We are ignorant of the purpose for which this irritating ointment is designed.

UNGUENTUM PULVERIS CANTHARIDIS VESICATORIÆ, Edin. *Ointment of the Powder of Blistering Flies.*

“Take of Resinous Ointment, *seven parts*; Powdered Blistering Flies, *one part*. Sprinkle the powder into the melted ointment; and stir the mixture until it stiffen in cooling.”

UNGUENTUM CANTHARIDIS, Dub. *Ointment of Blistering Flies.*

“Take of Ointment of Yellow Wax, *half a pound*; Blistering Flies in powder, *one ounce*. Form them into an ointment.”

These ointments are intended for promoting a purulent discharge from blistered surfaces, and produce this effect sufficiently well when the irritation they excite can be supported, which, however cannot always be done. The flies should be reduced to a very fine powder, and very intimately mixed with the ointment.

UNGUENTUM RESINÆ NIGRÆ,<sup>1</sup> Lond. *Black Resin Ointment.*

“Take of Black Resin, Yellow Wax, Yellow Resin, of each, *nine ounces*, Olive Oil, *a pint*. Melt them together, and strain through a linen-cloth.”

UNGUENTUM SAMBUCI<sup>2</sup>, Lond. *Elder Ointment.*

“Take of Elder Flowers, Prepared Lard, of each, *two pounds*. Boil the Elder Flowers in the Lard, until they become crisp, then strain the ointment through a linen cloth.”

<sup>1</sup> Unguentum basilicum nigrum, vel tetra pharmacum, P. L. 1745.

<sup>2</sup> Unguentum sambucinum, P. L. 1720.



*Dublin.*

“Take of Fresh Elder Flowers, *three pounds*; Prepared Hog’s Lard, *four pounds*; Mutton Suet, *two pounds*. Make an ointment of these in the manner directed for the preparation of the Savine ointment.”

These ointments are simply emollient, and possess no advantages over simple ointment. They are vestiges of the redundant practice of former times.

UNGUENTUM SIMPLEX, Edin. *Simple Ointment.*

“Take of Olive Oil, *five parts*; White Wax, *two parts*. Melt the Wax in the Oil; then stir the mixture until it stiffen in cooling.”

An useful emollient ointment for softening the skin.

Officinal preparations. *Unguentum Oxidi Plumbi albi*, E. *Unguentum Acetatis Plumbi*, E.

UNGUENTUM SUBACETATIS CUPRI, Edin. *Ointment of Subacetate of Copper.*

“Take of Resinous Ointment, *fifteen parts*; Subacetate of Copper, *one part*. Sprinkle the Subacetate into the melted ointment, and stir until the mixture stiffen in cooling.”

UNGUENTUM ÆRUGINIS, Dub. *Ointment of Verdegris.*<sup>1</sup>

“Take of Ointment of White Wax, *a pound*; Prepared Verdegris, *half an ounce*. Make them into an ointment.”

*Syn.* Unguento Egiziaco (I.).

These ointments are escharotic and detergent. They are used as an occasional dressing to foul, flabby ulcers; and as an application to scrophulous ulcerations of the tarsi. They can scarcely be used in the undiluted state, unless to act as a caustic for taking down fungous flesh.

UNGUENTUM SULPHURIS, Lond. *Sulphur Ointment.*

“Take of Sublimed Sulphur, *three ounces*; Prepared Lard, *half a pound*. Mix them.”

*Edinburgh.*

“Take of Hog’s Lard, *four parts*; Sublimed Sulphur, *one part*. Mix.”

*Dublin.*

“Take of Prepared Hog’s Lard, *four pounds*; Sublimed Sulphur, *a pound*. Form them into an ointment.”

<sup>1</sup> *Syn.* Onguent soufré (F.), Schemfelsalbe (G.), Unguento Solforato (I.).

These ointments are specific in itch. They should be rubbed on the body every night until the disease be cured, but not more than one-fourth part of the body should be covered with it at a time.\*

\* As the smell of the sulphur ointments is objected to by many people, the following combination has been recommended.

“Take of Subcarbonate of Potass, *half an ounce*; Rose-water, *one ounce*; Red



UNGUENTUM SULPHURIS COMPOSITUM,  
Lond. *Compound Ointment of Sulphur.*

“Take of Sublimed Sulphur, *half a pound*; White Hellebore Root in powder, *two ounces*; Nitrate of Potass, *a drachm*; Soft Soap, *half a pound*; Prepared Lard, *a pound and a half*. Mix them.”

*Syn.* Onguent soufré composée (F.), Unguento solforato composto (I.).

This ointment is employed in the same cases as the simple ointment. It is supposed to derive more efficacy from the addition of the white hellebore, but it often excites too much irritation

UNGUENTUM VERATRI, Lond. *Ointment of White Hellebore.*<sup>1</sup>

“Take of White Hellebore Root powdered, *two ounces*; Prepared Lard, *eight ounces*; Oil of Lemon, *twenty minims*. Mix them.”

UNGUENTUM HELLEBORI ALBI, Dub. *Ointment of White Hellebore.*

“Take of Prepared Lard, *a pound*; White Hellebore Root in powder, *three ounces*. Make them into an ointment.”

These ointments are sometimes used for the cure of scabies, when the smell of the sulphur ointment is objected to; but they are less certain remedies.

UNGUENTUM ZINCI, Lond. UNGUENTUM OXIDI ZINCI, Edin. *Zinc Ointment.*

“Take of Oxide of Zinc, *an ounce*; Prepared Lard, *six ounces*. Mix them.”

UNGUENTUM OXIDI ZINCI, Edin. *Ointment of Oxide of Zinc.*

“Take of Simple Liniment, *six parts*; Oxide of Zinc, *one part*. Mix.”

*Dublin.*

“Take of Ointment of White Wax, *a pound*; Oxide of Zinc, *an ounce and a half*. Make them into an ointment.”

*Syn.* Onguent de Zinc (F.), Zinksalbe (G.), Unguento di Zinco (I.).

These ointments are moderately astringent and stimulant. They are generally applied in chronic inflammation of the eye, depending on a relaxed state of the vessels: we find them also of very considerable use in sore nipples; and for removing ring-worm, particularly when it attacks the scalp.

Sulphuret of Mercury, *one drachm*; Essential Oil of Bergamot, *half a fluid drachm*; Sublimed Sulphur, Hog's Lard, of each, *eleven ounces*. Mix them.”

*Bateman on Cutaneous Diseases, p. 200. note.*

<sup>1</sup> Unguentum Hellebori albi, P. L. 1787.



## LINIMENTA.

### LINIMENTS.

THESE are compositions which have the consistence of oil or balsam; so as to allow them to be easily rubbed upon the skin. They are in general more active remedies than cerates or ointments; and act as local stimulants, relieving deep-seated inflammations and pains.

**LINIMENTUM ÆRUGINIS**, Lond. *Liniment of Verdegris.*

“Take of Verdegris powdered, *an ounce*; Vinegar, *seven fluid ounces*; Clarified Honey, *fourteen ounces*. Dissolve the Verdegris in the Vinegar, and strain it through a linen cloth; then, having added the Honey, boil down the mixture to a proper consistence.”

**OXYMEL ÆRUGINIS**, Dub. *Oxymel of Verdegris.*

“Take of Prepared Verdegris, *one ounce*; Wine Vinegar, *seven fluid ounces*; Clarified Honey, *fourteen ounces*. Dissolve the Verdegris in the Vinegar, and strain it through a linen cloth; add the Honey, and boil the mixture to a proper thickness.”

This preparation, which is improperly named a liniment by the London College, is detergent and escharotic. In the above state it is used for taking down fungous flesh; and considerably diluted, is an useful stimulant to foul ulcers, which it clears and excites to a more healthy action. It has been employed as a gargle in venereal ulcerations of the mouth and fauces; but we cannot recommend it.

**LINIMENTUM AMMONIÆ FORTIUS**, Lond. *Stronger Liniment of Ammonia.*

“Take of Solution of Ammonia, *a fluid ounce*; Olive Oil, *two fluid ounces*. Shake them together until they unite.”

**OLEUM AMMONIATUM**, Edin. *Ammoniated Oil.*

“Take of Olive Oil, *eight parts*; Water of Ammonia, *one part*. Mix them.”

**LINIMENTUM AMMONIÆ**, Dub. *Liniment of Ammonia.*

“Take of Caustic Water of Ammonia, *two fluid drachms*; Olive Oil, *two fluid ounces*. Mix them.”

*Syn.* Liniment volatil (F.), Ammonium liniment (G.), Linamento volatile (I.).

In these preparations a chemical union takes place between the alkali and the fixed oil, and produces a white soap, which is kept fluid by the water of the solution of ammonia. It is an excellent rubefacient, and is efficaciously employed in cynanche tonsillaris, spread on a piece of flannel, and applied round the throat; and to relieve rheumatic pains, is rubbed upon the skin over the affected part, often with the addition of a little camphor, or, what is preferable, some Extract of Belladonna. We have found a medium proportion of solution of ammonia,



or half a fluid ounce to two fluid ounces of oil, form a preparation better fitted for general use than the above.

**LINIMENTUM AMMONIÆ SUBCARBONATIS**, Lond. *Liniment of Carbonate of Ammonia.*<sup>1</sup>

“Take of Solution of Subcarbonate of Ammonia, *a fluid ounce*; Olive Oil, *three fluid ounces*. Shake them together until they unite.”

This preparation is also a fluid soap, but the combination of the oil and alkali is prevented from being so perfect by the carbonic acid of the subcarbonate. It is also much less soluble in water, and after a little time, the soapy matter separates from the water. It is intended for the same purposes as the strong liniment, which can be readily rendered as mild by the addition of oil; and therefore this may be regarded as a superfluous preparation.

**LINIMENTUM AQUÆ CALCIS**; sive **OLEUM LINI CUM CALCE**, Edin. *Liniment of Lime-water.*

“Take of Linseed-oil, Lime-water, of each, *equal parts*. Mix them.”

**LINIMENTUM CALCIS**, Dub. *Liniment of Lime.*

“Take of Lime-water, Olive Oil, of each *three fluid ounces*. Mix them.”

These are solutions of earthy soaps, formed by the chemical union of the lime and the oil. They are thick, of a white colour, and devoid of acrimony, and are very advantageously applied to burns and scalds. The soapy matter separates from the water, when it is kept for a little time, and therefore it is always better to prepare it only when it is wanted.

**LINIMENTUM CAMPHORÆ**, Lond. *Liniment of Camphor.*

“Take of Camphor, *half an ounce*; Olive Oil, *two fluid ounces*. Dissolve the Camphor in the Oil.”

**OLEUM CAMPHORATUM**, Edin. *Camphorated Oil.*

“Take of Olive Oil, *four parts*; Camphor, *one part*. Mix them so as to dissolve the camphor.”

**OLEUM CAMPHORATUM**, Dub. *Camphorated Oil.*

“Take of Camphor, *half an ounce*; Olive Oil, *two fluid ounces*. Rub them together.”

These solutions of camphor in fixed oil are very useful embrocations to glandular swellings, sprains, bruises, and to joints affected with rheumatic pains. Mr. Ware recommends them with the addition of half an ounce of the solution of subcarbonate of potass, to be applied to the eyelids night and morning in incipient amaurosis.

**LINIMENTUM CAMPHORÆ COMPOSITUM**, Lond. *Compound Liniment of Camphor.*

<sup>1</sup> Linimentum volatile, P. L. 1745. Linimentum Ammoniae, P. L. 1787.



“Take of Camphor, *two ounces* ; Solution of Ammonia, *six fluid ounces* ; Spirit of Lavender, *a pint*. Mix the Solution of Ammonia with the Spirit ; then from a glass retort, by a gentle heat, distil a pint. Lastly, dissolve the Camphor in this distilled liquor.”

This is a very useful stimulant application to sprains, bruises, and rheumatic pains. It is also an excellent vehicle for introducing opium into the habit by means of friction. An embrocation composed of  $\text{f}\overline{\text{z}}$ jss. of this liniment, and  $\text{f}\overline{\text{z}}$ ss. of tincture of opium, warmed and rubbed over the surface of the abdomen, very quickly allays the pains of flatulent colic.

LINIMEN'TUM HYDRAR'GYRI, Lond. *Liniment of Mercury.*

“Take of the Stronger Mercurial Ointment, Prepared Lard, of each, *four ounces* ; Camphor, *an ounce* ; Rectified Spirit, *fifteen minims* ; Solution of Ammonia, *four fluid ounces*. First rub the Camphor with the Spirit, then with the Lard and Mercurial Ointment : lastly, drop in gradually the Solution of Ammonia, and mix the whole.”

This liniment is stimulant and discutient. It is employed as an embrocation to parts affected with chronic venereal pains, nodes, and tophi ; to indolent swellings, and to discuss morbid collections of fluid. One drachm should be rubbed on the affected part night and morning. When largely used, it salivates sooner than mercurial ointment.

LINIMEN'TUM SAPONIS COMPOSITUM, Lond. *Compound Soap Liniment.*<sup>1</sup>

“Take of Hard Soap, *three ounces* ; Camphor, *an ounce* ; Spirit of Rosemary, *a pint*. Dissolve the Camphor in the Spirit, then add the Soap, and macerate in the heat of a sand-bath, until they be dissolved.”

TINCTURA SAPONIS CAMPHORATA ; vulgo, LINIMENTUM SAPONACEUM, Edin. *Camphorated Tincture of Soap* ; commonly called, *Liniment of Soap*.

“Take of Hard Soap, sliced, *four ounces* ; Camphor, *two ounces* ; Volatile Oil of Rosemary, *half an ounce* ; Alcohol, *two pounds*. Digest the Soap in the Alcohol for three days, then add the Camphor and the Oil, frequently shaking the mixture.”

*Syn.* Kampferliniment (G.).

These preparations are stimulant and anodyne, and may be advantageously applied against local pains, and in bruises rubbed upon the parts.

TINCTURA SAPONIS ET OPII ; vulgo, LINIMENTUM ANODYNUM, Edin. *Tincture of Soap and Opium* ; commonly called *Anodyne Liniment*.

<sup>1</sup> Linimentum saponaceum, P. L. 1745.



“Take of Hard Soap sliced, *four ounces*; Opium, *one ounce*; Camphor, *two ounces*; Oil of Rosemary, *half an ounce*; Alcohol, *two pounds*. Digest the Soap in the Alcohol for three days, then to the strained solution add the Camphor and the Oil, frequently shaking the mixture.”

The addition of the opium to the soap liniment, renders it in many cases of rheumatism and local pains, more useful than the simple liniment: but if any effect is particularly desired from the external application of opium, it will be obtained with more certainty, by dissolving the opium in olive oil.

**LINIMENTUM TEREBIN'THINÆ**, Lond. *Turpentine Liniment.*

“Take of Cerate of Resin, *a pound*; Oil of Turpentine, *half a pint*. Melt the Cerate, then add to it the Oil of Turpentine, and mix them.”

This liniment was introduced into practice by Dr. Kentish, at that time a surgeon in Newcastle, as a dressing to burns immediately after they happen, and until the loosening of the eschars. Dr. Kentish's plan was first to bathe the parts with warm oil of turpentine, and then to apply over them plasters thickly spread, of this liniment; at the same time that he supported the strength with wine, opium, and cordials. After the life of the parts appeared to be restored, purges were given, the cordials omitted, and mild emollient dressings applied.<sup>1</sup> We have had several opportunities of witnessing the good effects of this plan of treatment.

---

## CATAPLASMATA.

### CATAPLASMS.

CATAPLASMS are in general extemporaneous preparations; but the two following formulæ are introduced into the Pharmacopœias to fix the proportions of the ingredients.

**CATAPLASMA FERMENTI**, Lond. *Yeast Cataplasm.*

“Take of Flour, *a pound*; Yeast of Beer, *half a pint*. Mix, and expose the mixture to a gentle heat, until it begin to swell.”

The inflation is produced by the extrication of carbonic acid gas, on which the efficacy of the cataplasm depends; and which is evolved by the heat applied to the mixture exciting the fermentative process. In this state it is applied to painful gangrenous, or foul ulcers; and soon corrects the foetor of the

---

<sup>1</sup> *Essays on Burns, &c.* by Edward Kentish, 1797 and 1800.



discharge, while at the same time it hastens the sloughing of the sores.

CATAPLASMA SINAPIS, Lond. *Cataplasma of Mustard.*

“Take of Mustard Seed, Linseed, of each in powder, *half a pound*; Hot Vinegar, *a sufficient quantity*. Mix them to the thickness of a cataplasm.”

CATAPLASMA SINAPEOS, Dub. *Mustard Cataplasm.*

“Take of Mustard Seed in powder, Crumb of Bread, of each, *half a pound*; Vinegar, *a sufficient quantity*. Mix them so as to make a cataplasm. This preparation may be rendered more acrid by adding two ounces of Horse-radish finely scraped.”

These cataplasms are powerful local stimulants, and rubefacients. They are to be spread on cloths to the thickness of about half an inch, and applied to the soles of the feet, in the low stage of typhus fever, particularly when stupor or delirium is present; and in apoplexy, coma, and other cases in which there is a great determination to the head. Their rubefacient effects are very quickly produced, and often so powerfully, as to raise blisters on the part.

## TABLE,

*Showing the Proportion in which Opium and certain Preparations of Iron, Antimony, Arsenic, and Mercury are contained in some compound Medicines.*

### *Opium.*

CONFECTIO OPII, Lond. *Confection of Opium.* Thirty-six grains contain one grain of opium.

ELECTUARIUM OPIATUM, Edin. *Opiate Electuary,* contains in each drachm about one grain and a half of opium.

ELECTUARIUM CATECHU COMPOSITUM, Edin. *Electuary of Catechu,* contains in each ounce about two grains and a half of opium; or one hundred and ninety-three grains contain one grain of opium.

ELECTUARIUM CATECHU COMPOSITUM, Dub. *Compound Electuary of Catechu,* contains in each ounce about two grains and a half of purified opium.

PILULÆ SAPONIS CUM OPIO, Lond. *Pills of Soap and of Opium.* Five grains contain one grain of opium.

PILULÆ OPIATÆ, Edin. *Opiate, formerly Thebaic, Pills.* Each drachm contains six grains of opium. A pill of five grains contains half a grain of opium.



PILULÆ STYRACIS, Dub. *Storax Pills* contain, in five grains of the mass, one grain of purified opium.

PULVIS CORNU USTI CUM OPIO, Lond. *Powder of Burnt Hartshorn with Opium*. Ten grains contain one grain of opium.

PULVIS CRETÆ COMPOSITUS CUM OPIO, Lond. *Compound Powder of Chalk with Opium*. Two scruples contain one grain of opium.

PULVIS IPECACUANHÆ COMPOSITUS, Lond. Dub. *Compound Powder of Ipecacuanha*. Ten grains contain one grain of opium.

PULVIS IPECACUANHÆ ET OPII, Edin. *Powder of Ipecacuanha and Opium*. Each drachm contains six grains of opium, or one grain in ten grains of the powder.

PULVIS KINO COMPOSITUS, Lond. *Compound Powder of Kino*. Each scruple contains one grain of opium.

SYRUPUS OPII, Dub. *Syrup of Opium*, contains in one fluid ounce about one grain of the watery extract of opium : for the liquor is more than doubled in bulk by the addition of the sugar.

TINCTURA OPII, Lond. *Tincture of Opium*. Nineteen minims contain about one grain of opium.

TINCTURA OPII, Edin. *Tincture of Opium* is made with two scruples of opium in each ounce of liquid, or each drachm should contain five grains. But one drachm of the tincture, when evaporated, yields only three grains and a half of opium.

TINCTURA OPII, Dub. *Tincture of Opium* contains in one fluid drachm about four grains and a half of purified opium.

TINCTURA CAMPHORÆ COMPOSITA, Lond. *Compound Tincture of Camphor*. TINCTURA OPII CAMPHORATA, Edin. Half a fluid ounce contains nearly one grain of opium.

TINCTURA OPII AMMONIATA, Edin. *Ammoniated Tincture of Opium*, is made with about eight grains of opium in each ounce of liquid ; or each drachm should contain nearly one grain of opium.

TINCTURA OPII CAMPHORATA, Dub. *Camphorated Tincture of Opium*. Four fluid drachms and a half contain nearly one grain of purified opium.

TINCTURA SAPONIS ET OPII, Edin. *Tincture of Soap and Opium*, is made with one scruple of opium in each ounce of the liquid.

TROCHISCI GLYCYRRHIZÆ CUM OPIO, Edin. *Troches of Liquorice with Opium*. Each drachm contains nearly one grain of opium.



## IRON.

TINCTURA ACETATIS FERRI CUM ALCOHOLE, Dub. *Tincture of Acetate of Iron with Alcohol.* Each fluid drachm contains about one grain of dry acetate of iron.

---

## ANTIMONY.

VINUM ANTIMONII TARTARIZATI, Lond. *Solution of Tartarized Antimony* contains in each fluid ounce two grains of Tartarized Antimony.

VINUM TARTRITIS ANTIMONII, Edin. *Wine of Tartrate of Antimony* contains in each ounce two grains of tartrate of antimony.

---

## MERCURY.

EMPLASTRUM HYDRARGYRI, Edin. *Mercurial Plaster.* Each drachm contains about sixteen grains of mercury, (fifteen grains, Lond.)

HYDRARGYRUS CUM MAGNESIA, Dub. *Mercury with Magnesia.* Three grains contain two of mercury.

HYDRARGYRUM CUM CRETA, Lond. *Mercury with Chalk.* Three grains contain one grain of mercury.

LIQUOR HYDRARGYRI OXYMURIATIS, Lond. *Solution of Oxymuriate of Mercury.* Two fluid ounces contain half a grain of oxymuriate of mercury.

LINIMENTUM HYDRARGYRI, Lond. *Mercurial Liniment.* Six drachms contain one drachm of mercury.

PILULÆ HYDRARGYRI, Lond. Dub. *Mercurial Pills.* Three grains contain one grain of mercury.

PILULÆ HYDRARGYRI, Edin. *Mercurial Pills.* Each drachm contains fifteen grains of mercury. Each five grain pill contains one and one-fourth grain of mercury.

PILULÆ HYDRARGYRI SUBMURIATIS COMPOSITÆ, Lond. Edin. *Pills of Submuriate of Mercury.* About four grains contain one grain of submuriate of mercury.

UNGUENTUM HYDRARGYRI FORTIUS, Lond. Dub. *Stronger Mercurial Ointment.* Two drachms contain one drachm of mercury.

UNGUENTUM HYDRARGYRI MITIUS, Lond. *Weaker Mercurial Ointment.* Six drachms contain one drachm of mercury.

UNGUENTUM HYDRARGYRI, Edin. *Mercurial Ointment.* Each drachm contains twelve grains of mercury; made with double the quantity of mercury, each drachm contains twenty grains.

UNGUENTUM NITRATIS HYDRARGYRI FORTIUS, Edin. *Stronger Ointment of Nitrate of Mercury.* Each drachm contains four grains of mercury.



UNGUENTUM NITRATIS HYDRARGYRI MITIUS, Edin. *Milder Ointment of Nitrate of Mercury.* Each scruple contains half a grain of mercury.

UNGUENTUM OXIDI HYDRARGYRI CINEREI, Edin. Each drachm contains fifteen grains of the oxide.

UNGUENTUM OXIDI HYDRARGYRI RUBRI, Edin. *Ointment of Red Oxide of Mercury.* Each drachm contains seven grains of the oxide.

---

ARSENIC.

LIQUOR ARSENICALIS, Lond. SOLUTIO ARSENICALIS, Edin. *Arsenical Solution.* One fluid ounce contains four grains of white sublimed arsenic.



1908-1909



TABLE  
OF  
NEW NAMES;

*Showing to what Name of the former London Pharmacopœia  
each belongs.*

NEW NAMES.	FORMER NAMES.
ABIETIS resina - - - - -	Thus.
Absinthium - - - - -	Absinthium vulgare.
Acaciæ Gummi - - - - -	Arabicum Gummi.
Acetosa - - - - -	Acetosa pratensis.
Acidum aceticum dilutum - -	Acidum aceticum.
———— arseniosum - - -	Arsenici oxydum album.
———— benzoicum - - -	Flores Benzoës.
———— nitricum - - -	Acidum nitrosum.
———— sulphuricum - - -	———— vitriolicum.
Alöes spicatæ extractum - -	Alöe Socotorina, <i>succus spissatus</i>
———— vulgaris extractum - -	—— Barbadensis, <i>succus spissatus</i>
Ammonia Murias - - - - -	Sal Ammoniacus.
———— Subcarbonas - - -	Ammonia præparata.
Anthemidis flores - - - - -	Chamæmelum, <i>flos simplex</i> .
Antimonii Sulphuretum - - -	Antimonium.
———— Sulphuretum præcipi- tatum - - - - -	Sulphur Antimonii præcipitatum.
Argenti Nitras - - - - -	
Armoraciæ radix - - - - -	Argentum nitratum.
Arsenicum album - - - - -	Raphanus rusticanus; <i>radix</i> .
———— ————— sublimatum -	Arsenici oxydum.
	———— ————— sublimatum.
	B.
Benzoinum - - - - -	Benzoe.
	C.
Calami radix - - - - -	Calamus aromaticus, <i>radix</i> .
Calamina - - - - -	Lapis Calaminaris.
Calumba - - - - -	Colomba, <i>radix</i> .
Cambogia - - - - -	Gambogia.
Canella cortex - - - - -	Canella alba, <i>cortex</i> .
Cantharis - - - - -	Lytta.
———— vesicatoria - - -	———— vesicatoria.
Capsici baccae - - - - -	Piper indicum, <i>capsula</i> .
Caryophylli - - - - -	{ Caryophyllus aromatica <i>pericar</i> <i>pium immaturum</i> .
Cassia pulpa - - - - -	
Castoreum - - - - -	Cassia fistularis, <i>fructus</i> .
	Castoreum Rossicum.
Ceratum Plumbi compositum	{ Ceratum Lithargyri acetati com positum.
———— ————— acetatis - -	
———— Cantharidis - - -	———— Plumbi Superacetatis
———— Resinæ - - - - -	———— Lyttæ.
Cetaceum - - - - -	Unguentum Resinæ flavæ.
Cinchonæ lancifoliæ cortex -	Sperma Ceti.
———— cordifoliæ cortex -	Cinchonæ cortex.
	<i>Vulgò cortex flavus.</i>



NEW NAMES.		FORMER NAMES.	
Cinchonæ oblongifoliæ cortex	-	<i>Vulgò</i> Cortex ruber.	
Coccus	- - - - -	Coccinella.	
Confectio Aurantii	- - - - -	Conserva Aurantii.	
———— Cassiæ	- - - - -	Electuarium Cassiæ.	
———— Opii	- - - - -	Confectio opiata.	
———— Rosæ caninæ	- - - - -	Conserva Cynosbati.	
———— Rosæ Gallicæ	- - - - -	———— Rosæ.	
———— Scammoneæ	- - - - -	Electuarium Scammonii.	
———— Sennæ	- - - - -	———— Sennæ.	
Conii folia	- - - - -	Cicuta, <i>herba</i> .	
Copaiba	- - - - -	Balsamum Copaivæ.	
Cucumis Colocynthis, <i>peponum</i>	}	Cucumis Colocynthis <i>pomarum</i>	
<i>pulpa</i>		<i>pulpa</i> .	
Cupri Sulphas	- - - - -	Vitriolum cæruleum.	
Cuspariæ cortex	- - - - -	<i>Vulgò</i> cortex Augusturæ.	
Cydoniæ semina	- - - - -	Cydonii Mali <i>semen</i> .	
D.			
Decoctum Cydoniæ	- - - - -	Mucilago seminis Cydonii Mali.	
———— Malvæ compositum	-	Decoctum pro enemate.	
———— Papaveris	- - - - -	———— fomento.	
E.			
Elaterii <i>pepones</i>	- - - - -	Elaterii <i>poma</i> .	
Emplastrum Cantharidis	- -	Emplastrum Lyttæ.	
———— Ceræ	- - - - -	———— Ceræ compositum.	
———— Galbani compositum	{	———— Lithargyri compo-	
———— Hydrargyri		situm.	
———— Picis compositum	{	———— Lithargyri cum hy-	
———— Lyttæ		drargyro.	
———— Plumbi	{	———— Picis Burgundicæ	
———— Resinæ		compositum.	
————	- - - - -	———— Cantharidis.	
————	- - - - -	———— Lithargyri. [ <i>sina</i> .	
————	- - - - -	———— Lithargyri cum re-	
F.			
Ferri Sulphas	- - - - -	Ferrum vitriolatum.	
Ferrum ammoniatum	- - - - -	———— ammoniacale.	
Fœniculi semina	- - - - -	Fœniculum dulce, <i>semen</i> .	
H.			
Hellebori fœtidi folia	- - - - -	Helleboraster, <i>folium</i> .	
Hydrargyri Nitrico-oxydum	-	Hydrargyrus nitratus ruber.	
———— Oxydum rubrum	-	———— calcinatus.	
———— Oxymurias	- - - - -	———— muriatus.	
———— Submurias	- - - - -	Calomelas.	
———— Sulphuretum rubrum	-	Hydrargyrus sulphuratus ruber.	
Hydrargyrum præcipitatum album	-	Calx Hydrargyri albi	
J.			
Jalapæ radix	- - - - -	Jalapium, <i>radix</i> .	
Infusum Lini compositum	- -	Infusum Lini.	
———— Rosæ compositum	- -	———— Rosæ.	
———— Sennæ compositum	-	———— Sennæ.	



## NEW NAMES.

## FORMER NAMES.

## L.

Linimentum Ammoniae Subcar-	}	Linimentum Ammoniae.
bonatus - - - - -		
Linimentum Æruginis - - -		Oxymel Æruginis.
Lini usitatissimi semina - - -		Linum, <i>semen</i> .
Liquor Aluminis compositus -		Aqua Aluminis composita.
—— Ammoniae - - - - -		—— Ammoniae puræ.
—— Ammoniae Acetatis - - -		—— Ammoniae acetatæ.
—— Antimonii tartarizati -		Vinum Antimonii tartarizati.
—— Calcis - - - - -		Aqua Calcis.
—— Cupri ammoniati - - -		—— Cupri ammoniati.
—— Plumbi subacetatis - - -		—— Lithargyri acetati. [sita.
—— Plumbi subacetatis dilutus		—— Lithargyri acetati compo-
—— Potassæ - - - - -		—— Kali puri.
Lytta - - - - -		Cantharis.

## M.

Magnesia - - - - -		Magnesia usta.
Magnesiae Subcarbonas - - -		—— carbonas.
—— Sulphas - - - - -		—— vitriolata.
Marmor Album - - - - -		Lapis calcarius.
Marrubium - - - - -		Marrubium album.
Matonia Cardamomum - - -		Elattaria Cardamomum.
Mentha piperita - - - - -		Mentha piperitis.
—— viridis - - - - -		—— sativa.
Menyanthes - - - - -		Trifolium paludosum.
Mistura Amygdalarum - - -		Lac Amygdalæ.
—— Ammoniacy - - - - -		—— Ammoniacy.
—— Assafœtidæ - - - - -		—— Assafœtidæ.
—— Camphoræ - - - - -		Mistura camphorata.
—— Cretæ - - - - -		—— cretacea.
—— Guaiacy - - - - -		Lac Guaiacy.
—— Moschi - - - - -		Mistura moschata.

## O.

Oleum Succini - - - - -		Oleum Succini rectificatum.
Oxymel simplex - - - - -		Mel acetatum.

## P.

Papaveris somniferi capsulæ -		Papaver album, <i>capsula</i> .
Pilulæ Saponis cum Opio - -		Pilulæ Opii.
—— Scillæ compositæ - - -		—— Scillæ.
Pix abietina - - - - -		Pix arida.
—— Nigra - - - - -		Resina Nigra.
Plumbi acetat - - - - -		Plumbi superacetat.
—— Subcarbonas - - - - -		Cerussa.
—— Oxydum semivitreum -		Lithargyrus.
Potassa cum Calce - - - - -		Calx cum Kali puro.
—— fusa - - - - -		Kali purum.
—— impura - - - - -		Cineres clavellati.
Potassæ Acetat - - - - -		Kali acetatum.
—— Nitras - - - - -		Nitrum.
—— Subcarbonas - - - - -		Kali præparatum.



NEW NAMES.	FORMER NAMES.
Potassæ Tartras - - - - -	Kali tartarizatum.
—— Sulphas - - - - -	—— vitriolatum.
—— Sulphuretum - - - - -	—— sulphuratum.
—— Supertartras - - - - -	Tartari Crystalli.
Pruna gallica - - - - -	Pruna.
Pterocarpi lignum - - - - -	Santalum rubrum, <i>lignum</i> .
Pulvis Alöes compositus - - -	Pulvis Alöes cum Guaiaco.
—— Cinnamomi compositus - -	—— aromaticus.
—— Cornu usti cum Opio - -	—— opiatum.
R.	
Rhœados Petala - - - - -	Papaver erraticum, <i>flos</i> .
Rhamni Baccæ - - - - -	Spina cervina, <i>bacca</i> .
Rhei Radix - - - - -	Rhabarbarum, <i>radix</i> .
Rosæ caninæ Pulpa - - - - -	Cynosbatus, <i>fructus</i> .
—— centifolæ Petala - - - -	Rosa Damascena, <i>petalum</i> .
—— Gallicæ Petala - - - - -	—— rubra, <i>petalum</i> .
S.	
Saccharum - - - - -	Saccharum non purificatum.
Scammoneæ Gummi resina - -	Scammonium, <i>gummi resina</i> .
Senegæ Radix - - - - -	Seneka, <i>radix</i> .
Serpentariæ Radix - - - - -	Serpentaria virginiana, <i>radix</i> .
Soda impura - - - - -	Barilla.
—— tartarizata - - - - -	Natron tartarizatum.
Sodæ Subboras - - - - -	Borax.
—— Murias - - - - -	Sal muriaticus.
—— Sulphas - - - - -	Natron vitriolatum.
—— Subcarbonas - - - - -	—— præparatum.
Spartii cacumina - - - - -	Genista <i>cacumen</i> .
Spiritus Camphoræ - - - - -	Spiritus camphoratus.
—— rectificatus - - - - -	—— vinosus rectificatus.
—— tenuior - - - - -	—— vinosus tenuior.
Sulphur lotum - - - - -	Flores Sulphuris loti.
—— sublimatum - - - - -	Sulphuris flores.
T.	
Terebinthina Canadensis - -	Balsamum Canadense.
Tinctura Camphoræ composita -	Tinctura Opii Camphorata.
—— Cantharidis - - - - -	—— Lyttæ.
—— Ferri Muriatis - - - - -	—— Ferri muriati.
V. & U.	
Veratri radix - - - - -	Helleborus albus, <i>radix</i> .
Vinum Antimonii tartarizati -	Liquor Antimonii tartarizati.
Unguentum Picis liquidæ - -	Unguentum Picis.
—— Cantharidis - - - - -	—— Lyttæ.
—— Picis nigræ - - - - -	—— Resinæ nigræ.
—— Cetacei - - - - -	—— Spermatidis Ceti.
Z.	
Zinci Oxydum - - - - -	Zincum calcinatum.
—— Sulphas - - - - -	—— vitriolatum.



TABLE  
OF  
FORMER NAMES;

*Showing to what Name of the present London Pharmacopœia  
each belongs.*

FORMER NAMES.	NEW NAMES.
Absinthium vulgare - - -	Absinthium.
Acetosa pratensis - - -	Acetosa.
Acidum aceticum - - -	Acidum aceticum dilutum.
Acidum nitrosum - - -	—— nitricum.
—— vitriolicum - - -	—— sulphuricum.
Alöe barbadensis - - -	Alöes vulgaris Extractum.
—— socotorina - - -	—— Spicatae Extractum.
Ammonia præparata - - -	Ammoniae Subcarbonas.
Antimonium - - -	Antimonii Sulphuretum.
Aqua Aluminis composita - -	Liquor Aluminis Compositus.
—— Ammoniae puræ - - -	—— Ammoniae.
—— Ammoniae acetatae - -	—— Ammoniae Acetatis.
—— Calcis - - -	—— Calcis.
—— Cupri ammoniati - - -	—— Cupri ammoniati.
—— Lythargyri acetati - -	—— Plumbi subacetatis.
—— Lythargyri acetati com- posita - - -	} —— Plumbi Subacetatis.
—— Kali puri - - -	
Arabicum Gummi - - -	—— Potassæ.
Argentum nitratum - - -	Acaciae Gummi.
Arsenici oxydum - - -	Argenti Nitras.
—— sublimatum - - -	Arsenicum album.
	—— sublimatum.
B.	
Balsamum canadense - - -	Terebinthina Canadensis.
—— Copaiva - - -	Copaiba.
Barilla - - -	Soda impura.
Benzöe - - -	Benzoinum.
Borax. - - -	Sodæ Subboras.
C.	
Calamus aromaticus, radix - -	Calami Radix.
Calumba radix - - -	Calumba.
Calomelas - - -	Hydrargyri Submurias.
Calx cum Kali puro - - -	Potassa cum Calce.
—— Hydrargyri alba - - -	{ Hydrargyrum præcipitatum album.
Canella alba, cortex - - -	
Cantharis - - -	Canellæ Cortex.
Caryophyllus aromatica, peri- carpium immaturum - - -	} Lytta.
Cassia fistularis, fructus - -	
Castoreum Rosicum - - -	Caryophillus.
Ceratum Lithargyri Acetati compositum - - -	{ Cassiæ Pulpa.
—— Lyttæ - - -	
Cerussa - - -	Castoreum.
	{ Ceratum Plumbi compositum.
	—— Cantharidis.
	Plumbi Subcarbonas.



FORMER NAMES.	NEW NAMES.
Chamæmelum, <i>flos simplex</i> . -	Anthemidis flores.
Cicuta, <i>herba</i> - - - - -	Conii folia.
Cinchona, <i>cortex</i> - - - - -	Cinchonæ lancifoliæ Cortex.
_____ <i>flavus</i> - - - - -	_____ cordifoliæ Cortex.
_____ <i>ruber</i> - - - - -	_____ oblongifoliæ Cortex.
Cineres clavellati - - - - -	Potassa impura.
Coccinella - - - - -	Coccus.
Colomba, <i>radix</i> - - - - -	Calumbæ radix.
Conserva Aurantii - - - - -	Confectio Aurantii.
_____ Cynosbati - - - - -	_____ Rosæ caninæ.
_____ Rosæ - - - - -	_____ Gallicæ.
Confectio opiata - - - - -	_____ Opii.
Cortex Angusturæ - - - - -	Cuspariæ cortex.
Cucumis Colocynthis, <i>pomorum</i>	Cucumis Colocynthis — <i>Pepo-</i>
<i>pulpa</i> - - - - -	<i>num pulpa</i> .
Cydonia Malus, <i>semen</i> - - - - -	Cydoniæ semina.
Cynosbatus, <i>fructus</i> - - - - -	Rosæ caninæ Pulpa.
D.	
Decoctum pro enemate - - - - -	Decoctum Malvæ compositum.
_____ fomento - - - - -	_____ Papaveris.
E.	
Elaterii poma - - - - -	Elaterii <i>pepones</i> .
Elettaria Cardamomum - - - - -	Matonia Cardamomum.
Electuarium Cassiæ - - - - -	Confectio Cassiæ.
_____ Scammonii - - - - -	_____ Scammoneæ.
_____ Sennæ - - - - -	_____ Sennæ.
Emplastrum Ceræ Compositum	Emplastrum Ceræ.
_____ Lithargyri - - - - -	_____ Plumbi.
_____ Lithargyri cum resina	_____ Resinæ.
_____ Lithargyri com- }	_____ Galbani compo-
positum - - - - - }	situm.
_____ Lithargyri cum }	_____ Hydrargyri.
Hydrargyro - - - - - }	
_____ Lyttæ - - - - -	_____ Cantharidis.
_____ Picis Burgundicæ }	_____ Picis compositum.
compositum - - - - - }	
F.	
Ferrum vitriolatum - - - - -	Ferri Sulphas.
_____ ammoniacale - - - - -	Ferrum ammoniatum.
Flores Benzoës - - - - -	Acidum benzoicum.
_____ Sulphuris loti - - - - -	Sulphur lotum.
Fœniculum dulce, <i>semen</i> - - - - -	Fœniculi Semina.
G.	
Gambogia - - - - -	Cambogia.
Genista, <i>cacumen</i> - - - - -	Spartii cacumina.
H.	
Helleboraster, <i>folium</i> - - - - -	Hellebori foetidi folia.
Helleborus albus. <i>radix</i> - - - - -	Veratri radix.
Hydrargyri calcinatus - - - - -	Hydrargyri Oxydum rubrum.



FORMER NAMES.	NEW NAMES.
Hydrargyrus muriatus - - -	Hydrargyri Oxymurias.
————— nitratus ruber - -	————— Nitrico-oxydum.
————— sulphuratus ruber -	————— Sulphuretum rubrum.
————— è sulphure - - -	————— Sulphuretum nigrum.

## J.

Jalapium, <i>radix</i> . - - - -	Jalapæ radix.
Infusum Lini - - - - -	Infusum Lini compositum.
———— Rosæ - - - - -	———— Rosæ compositum.
———— Sennæ - - - - -	———— Sennæ compositum.

## K.

Kali acetatum - - - - -	Potassæ Acetas.
———— purum - - - - -	Potassa fusa.
———— præparatum - - - - -	Potassæ Subcarbonas.
———— sulphuratum - - - - -	———— Sulphuretum.
———— tartarizatum - - - - -	———— Tartras.
———— vitriolatum - - - - -	———— Sulphas.

## L.

Lac Amygdalæ - - - - -	Mistura Amygdalarum.
———— Ammoniæ - - - - -	———— Ammoniæ.
———— Assafoetidæ - - - - -	———— Assafoetidæ.
———— Guaiaci - - - - -	———— Guaiaci.
Lapis calcarius - - - - -	Marmor album.
———— calaminaris - - - - -	Calamina. [bonatis.
Linimentum Ammoniæ - - -	Linimentum Ammoniæ Subcar-
Linum, <i>semen</i> - - - - -	Lini usitatissimi Semina.
Liquor Antimonii tartarizati	Vinum Antimonii tartarizati.
Lythargyrus - - - - -	Plumbi Oxydum semivitreum.
Lytta - - - - -	Cantharis.
———— vesicatoria - - - - -	———— vesicatoria.

## M.

Magnesia Carbonas - - - -	Magnesiae Subcarbonas.
———— usta - - - - -	Magnesia.
———— vitriolata - - - - -	Magnesiae Sulphas.
Marrubium album - - - - -	Marrubium.
Mel acetatum - - - - -	Oxymel.
Mentha piperitis - - - - -	Mentha Piperita.
———— sativa - - - - -	———— viridis.
Mistura camphorata - - - -	Mistura Camphoræ.
———— cretacea - - - - -	———— Cretæ.
———— moschata - - - - -	———— Moschi.
Mucilago seminis Cydonii mali	Decoctum Cydonii.

## N.

Natron præparatum - - - -	Sodæ Subcarbonas.
———— tartarizatum - - - -	Soda Tartarizata.
———— vitriolatum - - - - -	Sodæ Sulphas.
Nitrum - - - - -	Potassæ Nitras.

## O.

Oleum Succini rectificatum -	Oleum Succini.
Oxydum Arsenici album - -	Acidum Arseniosum.
Oxymel Æruginis - - - - -	Linimentum Æruginis.



FORMER NAMES.	NEW NAMES.
P.	
Papaver album, <i>capsula</i> - - -	Papaveris somniferi Capsulæ.
——— erraticum, <i>flos</i> - - -	Rhœados petala.
Pilulæ Opii - - - - -	Pilulæ Saponis cum Opio.
——— Scillæ - - - - -	——— Scillæ compositæ.
Pix arida - - - - -	Pix abietina.
Plumbi superacetas - - -	Plumbi acetas.
Pulvis Alöes cum Guaiaca - -	Pulvis Alöes compositus.
——— aromaticus - - - -	——— Cinnamomi compositus.
Pulvis opiatus - - - - -	——— Cornu usti cum Opio.
Pruna - - - - -	Pruna Gallica.
R.	
Raphanus rusticanus, <i>radix</i> -	Armoraciæ radix.
Rhabarbarum, <i>radix</i> - - -	Rhei radix.
Resina nigra - - - - -	Pix nigra.
Rosa Damascena, <i>petalum</i> - -	Rosæ centifoliæ petala.
——— rubra <i>petalum</i> - - -	——— Gallicæ petala.
S.	
Saccharum non purificatum -	Saccharum.
Sal Ammoniacus - - - - -	Ammonix Murias.
——— muriaticus - - - - -	Sodæ Murias.
Santalum rubrum - - - - -	Pterocarpi Lignum.
Scammonium, <i>gummi-resina</i> -	Scammoneæ Gummi-resina.
Seneka, <i>radix</i> - - - - -	Senegæ radix.
Serpentaria virginica, <i>radix</i> -	Cetaceum.
Sperma Ceti - - - - -	Rhamni Baccæ.
Spina cervina, <i>bacca</i> - - -	Rhamni Baccæ.
Spiritus camphoratus - - -	Spiritus Camphoræ.
——— vinosus rectificatus -	——— rectificatus.
——— vinosus tenuior - - -	——— tenuior.
Sulphur Antimonii præcipitatum {	Antimonii Sulphuretum præcipi-
Sulphuris Flores - - - - -	tatum.
	Sulphur sublimatum.
T.	
Tartari Crystalli - - - - -	Potassæ Supertartras.
Tinctura Lyttæ - - - - -	Tinctura Cantharidis.
——— Opii camphorata - - -	——— Camphoræ composita.
——— Ferri muriati - - - -	——— Ferri Muriatis.
Thus - - - - -	Abietis Resina.
Trifolium paludosum, <i>herba</i> -	Menyanthes.
V. & U.	
Liquor Antimonii tartarizati -	Vinum Antimonii tartarizati.
Vitriolum cæruleum - - - -	Cupri Sulphas.
Unguentum Lyttæ - - - - -	Unguentum Cantharidis.
——— Picis - - - - -	——— Picis liquidæ.
——— Resinæ Nigræ - - - -	——— Picis Nigræ.
——— Resinæ flavæ - - - -	Ceratum Resinæ.
——— Spermatidis Ceti - - -	Unguentum Cetacei.
Z.	
Zincum calcinatum - - - - -	Zinci Oxydum.
——— vitriolatum - - - - -	——— Sulphas.



# APPENDIX.

## No. I.

### OF WATER.

*Syn.* Aqua (*Lat.*), ὕδωρ (*Greek*), Eau (*F.*) Agua (*S.*), Acqua (*Ital.*), Aae (*Dan.*), A (*Swed.*), Ea, Eha (*old Saxon*), Awe (*Pers.*), Aw (*Welch*), Aa (*Low Germ. Islandic*), Ab (*Shans.*), Abh (*Gaelic*), Immek (*Esquimaux*), Tannee (*Tam.*), Panic (*Duk.*), Mah (*Arab.*), Watoora (*Cyng.*), Ayer (*Malay*), Panicum (*Sans.*), Hai (*Tonquinese*), A (*Gothic*).

**W**ATER is an agent of great importance, independent of the part it sustains in the magnificent operations of Nature. Its efficacy in the cure of diseases is indubitable; yet it is not admitted into the list of *materia medica* of any of the British Pharmacopœias, either in the state in which it is most commonly procured, or that in which it holds in solution substances from which it receives new properties, and is rendered capable of producing important changes in the animal œconomy. In the first state it is denominated **COMMON WATER**: in the second, **MINERAL WATER**; and under both of these forms, it is necessary that its qualities and effects should be known to the medical practitioner.

#### I. COMMON WATER.

The usual appearance of water is too well known to require description. It retains its fluidity under the ordinary pressure of the atmosphere, at any degree of temperature between  $32^{\circ}$  and  $212^{\circ}$ <sup>1</sup>, *Fah.*: but under  $32^{\circ}$  it crystallizes and becomes solid, or is changed into ice; and above  $212^{\circ}$ <sup>2</sup> assumes an aëriform character, or becomes steam, expanding to 1698 times its ordinary bulk. One cubic inch of pure water at  $60^{\circ}$ , and under a pressure of the atmosphere indicated by  $30^{\circ}$  of the barometer, weighs 252.42 grains, or nearly  $\frac{1}{15}$ th part of a grain less than two hundred and fifty-two grains and a half.

Although water is almost universally diffused over the surface of the earth, yet it is not found perfectly pure in any place; which is owing to its great solvent powers enabling it to take up a portion of many things with which it must come into contact in its natural state. These impregnations, however, are not sufficient in general to give it any very sensible taste or odour, or to render it unfit for the ordinary purposes of life; and it is in this state that common water is usually obtained. Common water varies considerably according to the source whence it is derived, and other circumstances; but all the varieties may be reduced under the three following heads:

1. Rain Water — *Aqua pluvialis*.
2. Spring Water — *Aqua fontana*.
3. River Water — *Aqua fluviatilis*.

1. **RAIN WATER** is the purest kind of natural water; but it, nevertheless, contains in solution, in every 100 cubic inches, about  $3\frac{1}{2}$  cubic inches of air, rather more oxygenous than common atmospherical air, and about one cubic inch of carbonic acid gas, besides minute portions of carbonate of lime and muriate of lime. Its specific gravity scarcely differs from that of distilled water; and after precipitating the muriate of lime, by dropping into it a little barytic water, and exposing it to the atmosphere until the precipitate be totally deposited, it is sufficiently pure for most pharmaceutical purposes.<sup>3</sup> When rain water, how-

<sup>1</sup> This degree varies according to the pressure of the atmosphere. Thus, in Gemmilaro's hut, on the side of Etna, about 11332 feet from the level of the sea, Dr. Irvine found that water boiled at  $191^{\circ}$ . Vide *Letters on Sicily*, 8vo. p. 153.

<sup>2</sup> Guy Lussac.

<sup>3</sup> Morveau, *Annales de Chimie*, xxiv. 320.



ever, is collected in towns, or from the roofs of houses, it contains a small portion of sulphate of lime, soot, and other impurities, and requires to be boiled and filtered before dropping in the barytic water.

Snow water, when newly melted, is destitute of air, which is the reason that fish cannot live in it; but, when allowed to remain for some time exposed to the atmosphere, it does not differ in its qualities from rain water.

2. SPRING WATER, if it have not filtered through a very soluble soil, is almost as pure as rain water. The best springs are those which rise through sand or gravel, at a small depth.<sup>1</sup> It generally contains, besides the ingredients which are found in rain water, a small portion of muriate of soda.

*Well or pump water*, which is spring water obtained by digging to a considerable depth, is by no means so pure. It is commonly distinguished by a property named hardness, implying an incapability of dissolving soap<sup>2</sup>; which is owing to its containing many earthy salts, the principle of which is sulphate of lime. It, also, contains more carbonic acid gas than common spring-water. Many of the foreign ingredients contained in hard water are simply suspended in it; for pump water is rendered softer and purer by only passing it through a filtering stone. The best mode of freeing hard water of its earthy salts, is first to boil it; then, after it has cooled, to drop into it an alkaline carbonate; and, lastly, to filter it. It cannot be employed for pharmaceutical purposes.

3. RIVER WATER, when the stream is rapid, and runs over a pebbly or a siliceous channel, is as pure as the softer spring water; but when the current is slow, and the bed clayey, it approaches nearer to the nature of well water, and frequently contains putrefied vegetable and animal matters, as is generally the case in the water of lakes and marshes.

Such are the foreign ingredients contained in *common water*. Boiling frees it from air and gases, and precipitates many of the earthy salts: but distillation in glass vessels frees it entirely from these ingredients, and it is obtained almost perfectly pure, transparent, colourless, insipid and inodorous.

The varieties of water enumerated above may be almost indiscriminately employed as diluents, the small proportion of foreign ingredients they contain occasioning no difference in their diluent properties. When the quantity of sulphate of lime and aluminous matter, however, is very considerable, as is the case in the water of many pumps, there is some reason for concluding that deleterious effects may arise from the use of the water; although it may be doubted whether the scrophulous and glandular swellings, peculiar to some populous towns, can be justly ascribed to this cause.<sup>3</sup> Even a few of the waters which are regarded as mineral waters owe more to the diluent property of the water for their efficacy, than to the impregnations they contain. This is particularly the case with the Malvern spring, which has been found to contain very little foreign matter. The diluting power of water is much modified by temperature; warm or tepid water being a much better diluent than cold water.

The medicinal properties of water as a diluent were well known to the ancients; and cold water, used as a drink in fevers, was the principal remedy of the father of physic in these complaints. The temperature of 60° is the proper degree, when it is intended that water should produce its diluent effects without the aid of heat. Under 45° it produces a sedative and astringent effect; above 60° and under 100°, it relaxes the fibres of the stomach, and is apt to induce nausea, particularly when bulk is added to this range of temperature; but at a higher temperature, the stimulus of heat, in the same manner as the addition of other stimulants, prevents that effect. Simple water may supersede the use of all other diluents; but

<sup>1</sup> The water conveyed to Hodsdon, in Hertfordshire, rises through a fine, white sand, and is so pure, that Dr. Hales affirms it left no incrustation in a boiler which had been in constant use for fifteen years. *Statical Essays*, ii. 242.

<sup>2</sup> Soap when agitated with hard water is decomposed; the alkali of the soap uniting with the acid of the earthy salts, while the oil and earths combine, and form new, nearly insoluble soaps, which swim in a curdy form on the surface of the water.

<sup>3</sup> Percival ascribes the glandular swellings common in Manchester to this cause. See *Essays*, i. 291.



animal and vegetable infusions are generally employed ; or toast and water (*infusum panis tostii*), which is more agreeable to most palates, and is an excellent diluent in fevers and inflammatory diseases. The temperature of water as a diluent should be regulated by the nature of the disease : in internal hæmorrhagies the temperature should not exceed  $45^{\circ}$ , but it may be  $60^{\circ}$  in fevers ; unless in the cold stage of the paroxysm of fever, when thirst should be allayed by tepid or warm water or other bland fluids ; and the same precaution is necessary when the sweat has become general and profuse. In cases in which there exists a morbid increase of bile disturbing the functions of the stomach and irritating the bowels, the temperature of the water used as drink may be from  $90^{\circ}$  to  $114^{\circ}$  ; and in some cases of dyspepsia, which are attended with the sensation of coldness at the stomach, and with cold extremities, a cup full of water, taken as hot as it can be drank, affords very considerable relief. In cases of redundant bile, by drinking half a pint of tepid water every morning before breakfast, and taking immediately afterwards moderate exercise, the acrid bile is diluted, and its passage through the bowels assisted, without the irritation which, in its undiluted state, it always excites ; and it produces the same benefit in cholera morbus in the commencement of the disease, the stomach being rendered by it more fit to receive opiates and other remedies. Some medicines, as sudorifics, diuretics, and emetics, scarcely produce their effects, unless their operation be assisted by copious dilution with water, or watery fluids.

Water is also an external remedy of great importance, but its effects are much modified by the degree of temperature at which it is applied.

COLD WATER, or of a temperature under  $70^{\circ}$ , gives the sensation of cold to the skin, and is applied under the form of *bath* and of *affusion*.

The cold Bath, (*balneum frigidum*), is water of any temperature, from  $42^{\circ}$  to  $85^{\circ}$  of Fahrenheit. When the body is immersed in it, it first induces the sensation of cold, excites shivering, renders the skin pale, and contracts it so as to produce the papillous appearance denominated goose skin (*cutis anserina*) ; the respiration at the same time is quickened and rendered irregular, producing sobbing ; and the pulse is diminished in force and velocity, but is also rendered firmer and more regular. If the immersion be not long continued, re-action takes place on coming out of the bath, a glow, or agreeable sensation of heat, is felt over the whole body, the tone and vigour of the muscles are increased, a buoyancy of spirit and aptitude for action succeed, and a sense of general refreshment is experienced by the bather. The protraction, however, of the immersion for a considerable space of time, particularly if the temperature of the bath be under  $50^{\circ}$ , is not followed by this re-action, but the cold water operates as a powerful sedative ; the action of the heart and arteries becomes languid, the pulse ceases at the wrist, the animal heat is rapidly diminished, and a sensation of coldness at the stomach is felt, which is succeeded by faintness, delirium, torpor, and sometimes death. These unpleasant effects are occasionally experienced in some degree, even when the immersion is not protracted, and the temperature of the bath is not under  $60^{\circ}$  ; in which case cold bathing proves always hurtful, and ought not to be repeated : but when the contrary effects are experienced it is found to be useful in many diseases of debility, particularly in scrophula, if the water be impregnated with salt, or sea-bathing be resorted to. The debilitated, however, in whom the use of sea-bathing produces these effects, when it is employed before breakfast, are not always affected in the same manner when it is used after breakfast, or when the stomach is full ; but, on the contrary, receive the same benefit from it as those with whom it agrees at all times. The use of cold water as a general bath is never employed with a view of producing its sedative effects : but for this purpose it is partially applied, either by the immersion of the affected parts, or by means of cloths dipped in very cold water, and laid over or near the parts. It is used as a remedy in active uterine hæmorrhagies, burns, and scalds, and in local inflammations, even when arising from general disease, as gout and acute rheumatism, when the surface of the pained part appears red and inflamed.

The cold affusion, or the suddenly pouring cold water over the whole surface of the body, operates as a powerful remedy, although its effects as such are of short duration. They are produced by the suddenness of the application affecting the nervous energy, and by the shock rousing the dormant sensibility, so as to



induce a new action, as it were, of the nervous system, dissolving the spasm on the extreme vessels of the surface, carrying off a large portion of morbid heat by general evaporation, and the remainder by insensible perspiration; thence restoring the healthy action of the exhalants and the capillaries. In typhus fever this mode of applying cold water has been productive of the best effects.<sup>1</sup> It should be applied in the first hot stage of the disease, if possible, and repeated every time the morbid heat returns. If the water can be impregnated with salt, so much the better; but when the disease is advanced, its temperature should not be more than  $26^{\circ}$ <sup>2</sup> under the heat of the body. It often stops suddenly the disease, if it be used during the three first days, and sometimes so late even as the fifth; but after this period it can be regarded as an useful auxiliary only even when properly employed. In tetanus, Currie affirms<sup>3</sup> that the cold affusion also proves useful, particularly when the shock is considerable, and applied during the convulsions. It is, however, in idiopathic tetanus only that it proves useful, no advantage being obtained from using it in tetanus arising from wounds.<sup>4</sup> Its utility has also been proved in many of the exanthemata; for instance, during the hot stage of the eruptive fever of small-pox; and we can bear ample testimony to its efficacy in scarlatina maligna, when the heat arises to above  $100^{\circ}$ .<sup>5</sup> This remedy, however, is productive of much mischief when misapplied; and therefore it is necessary to observe, that it is contra-indicated in the cold stage of fevers, and when a sense of chilliness is present, although the thermometer indicate the real heat to be more than natural. It is also said to be improper in fevers, when diarrhoea or dysentery are present; after the sweating stage in intermittents is formed; after the eruption is completely formed in confluent small-pox; and in symptomatic fever occasioned by great local inflammation: but Dr. A. Nicoll found it useful in India, in remittent and intermittent fevers accompanied with dysentery; when the heat of the surface has exceeded  $98^{\circ}$  Fahrenheit; for as in these cases the dysenteric symptoms seem to depend on the degree of febrile excitement, the cold water, by producing a solution of these, allays the griping and tenesmus, and natural stools follow. The affusion should always, in such cases, be preceded by bleeding and other depletory<sup>6</sup> means. The water should be dashed from a height of ten or twenty feet; and its temperature should be nearly that of the air at the time. The cold affusion, in the form of the shower-bath, is advantageously employed as a stimulant and tonic in diseases of general debility, when no fever or increased heat is present. I know of no remedy so generally useful in those affections which are known by the name of nervous complaints.

WARM WATER, or of a temperature from  $86^{\circ}$  to  $100^{\circ}$ , gives the sensation of warmth to the body, and is applied both locally and generally, in the form of vapour, fomentation, and bath. Water is found in a state of nature combined with different quantities of caloric within the above range of temperature. In the Buxton hot springs the temperature is about  $82^{\circ}$ ; at Bristol it is from  $76^{\circ}$  to  $84^{\circ}$ ; and at Bath the range is from  $110^{\circ}$  to  $114^{\circ}$ .<sup>7</sup> The necessary degree of temperature, however, is generally obtained by artificially heating the water.

The general application of warm water is obtained by means of baths. When the greater part of the entire body is immersed, the water constitutes properly a

<sup>1</sup> The cold affusion was employed by Antonius Musa, physician to Augustus, when that emperor was affected with a bowel complaint, which had resisted every other remedy. *Vide Q. Horatii F. Epistol. lib. xv. ad Num. Valam. C. Sueton. Tranq. Octavius Aug. ii. cap. 81. p. 104.* Cold affusions in the fevers of Asia are also prescribed by the Koran, and used in India by the Mahomedan and Hindoo physicians in various diseases.

<sup>2</sup> Currie — *Reports on Cold Water*, i. 31.      <sup>3</sup> *Ibid.* i. 138.      <sup>4</sup> *Ibid.* i. 159.

<sup>5</sup> Currie gives the following results of the affusion: — The heat of the body in fever, as indicated by the thermometer, being  $103^{\circ}$ , was by it reduced to  $98^{\circ}$  in half an hour; and the pulse from 112 to 80 beats, (vol. i. 22.); the heat  $101^{\circ}$  was reduced to  $99^{\circ}$ ; and the pulse from 112 to 98 in the same time. The heat  $106^{\circ}$  was reduced to  $98^{\circ}$ ; and the pulse from 130 to 90. Vol. i. 46.

<sup>6</sup> *Lond. Med Repository*, vol. ix. p. 123.

<sup>7</sup> The temperature of the Cross Bath pump is  $110^{\circ}$ ; the King's Bath  $112^{\circ}$ ; and the Hot Bath,  $114^{\circ}$ .



bath (*balneum*); but when half only is immersed, it is a half bath (*semicupium*). These may be either,

- a. The hot bath (*balneum calidum*), from 97° to 100°.
- b. The tepid bath (*balneum tepidum*), from 86° to 97°.
- c. The vapour bath (*balneum vaporis*), from 100° to 130°.

The two first differ in temperature only; but the last, from the water being applied in a very minutely divided state, acts with much greater effect than water in the liquid form. The operation of the first of these forms of applying water is stimulant; it augments the action of the heart and arteries, renders the skin red, quickens respiration, and produces a copious flow of sweat; but the others, although they excite the sensation of heat, yet lessen the frequency of the pulse, relax powerfully the skin and simple solids, and diminish generally increased excitement. It has been a question of some interest, whether water be ever taken into the body by the cutaneous absorbents? That it is taken in has been denied by many philosophers, and facts brought forward to support the opinion. Dr. Currie and Dr. Pearson, after half an hour's immersion in the Buxton bath, at 82°, found that the weight of the body was rather diminished than increased; and in a case of dysphagia, in which neither food nor drink was taken by the mouth for a considerable time, the patient, when put into the tepid bath, felt his thirst alleviated, and received much comfort, without his weight being at all increased. Dr. Currie supposed, that the abatement of thirst in this case arose from the relaxation of the exhalants of the surface produced by the bath, and those of the stomach sympathizing; and that although the exhalants terminate by open mouths, which pierce the epidermis, yet as the mouths of the absorbents terminate under it, and do not come into contact with the open air; so while the epidermis remains unirritated and entire, no absorption of fluid can therefore take place from the surface. Many experiments made by Seguin are also in favour of the opinion that no cutaneous absorption is effected in the bath. Among others, he immersed venereal patients in baths containing oxymuriate of mercury in solution, and found that while the cuticle remained entire, no salivation, nor other effect of the mercury on the system, was apparent; but the specific effect of the remedy always took place when the epidermis was injured or destroyed, as in itch. It must, however, be observed, that in the case of dysphagia the urine flowed as if drink had been taken by the mouth; a circumstance which Currie supposes to depend on the absorption from the atmosphere by the lungs. This, however, is an assumed position; the free exhalation from the lungs is evident, but it is by no means proved that any absorption takes place. It is true that the weight of the body in the above case was diminished; but from the sum of this loss we must abstract the cutaneous exhalation of the part of the body not immersed, the pulmonary exhalation, and the weight of the egesta; and were a supposition to be admitted as argument, it might be suggested that the relaxant power of the warm water acting on the epidermis as on inert matter, may open a way through it to the mouths of the absorbents. The question is still undecided, and, fortunately, it is not of much importance in a practical point of view.

Warm and vapour baths<sup>1</sup> are efficaciously employed in acute rheumatism, inflammation of the abdominal viscera, of the kidneys, bladder, and uterus; in suppression of urine, and in spasmodic affections, particularly those to which infants are liable, arising from dentition and other irritations. The general relaxation produced by their use has been taken advantage of, also, for assisting the reduction of strangulated hernia; for, although the effect be not topical as it regards the hernial tumour, yet the general relaxation produced gives a disposition

---

<sup>1</sup> A very simple and convenient vapour bath for military practice has been recommended by my friend Dr. A. Nicoll. It may be formed of a common beer or spirit barrel, with a false bottom, placed about a foot from the bottom of the cask, that is, the end on which it is standing, and perforated with numerous small holes. A gun barrel connected with the spout of a tea-kettle placed on a fire, is to be introduced into an opening in the side of the cask between the real and false bottom; and the patient, being seated on the false bottom, the steam or vapour from the boiling kettle soon surrounds him by rising through the holes. It must be prevented from escaping by means of a blanket, which should cover the open end of the cask, and apply closely round the neck of the patient.



to all the parts to regain their proper place. The tepid bath is found to be very useful in the rigidities which follow some acute diseases, as gout and rheumatism, nodosities of the joints<sup>1</sup>, and, according to some, the rigidities attendant on old age.<sup>2</sup> Its effects in promoting the natural excretions by the skin render it very serviceable in promoting the cure of herpetic eruptions; in slight cases of lepra the use of it with friction is all that is required; and in all cutaneous foulnesses it is a most important auxiliary. It has also been found very beneficial in cases of insanity. In general the period of immersion should not be less than twenty minutes, nor exceed one hour.<sup>3</sup>

The partial application of warm water as a remedy is made by means of

1. *a.* The foot bath (*pediluvium*);
- b.* The hip bath (*coxæluvium*); and
- c.* The hand bath (*manuluvium*).
2. *d.* Fomentations of vegetable decoctions; and
- e.* Flannel cloths wrung out of boiling water, by which the moisture is applied in a state of vapour.

These partial baths are useful in the same diseases for which the general baths are employed; but are better adapted for relieving the rigidity of single joints, and topical inflammation; and the hip bath has lately been found to be very beneficial in suppressed menstruation, and for relieving the pains of cancer in utero.

For fomentations it is the practice to employ vegetable decoctions; but the best of these can be regarded only as vehicles for retaining the heat and moisture. At all times, flannel cloths wrung out of boiling water are superior; both because the water is applied in the form of vapour, and also, while they continue as long warm, they do not wet the bed and linen of the patient. The flannel cloths should be each about two yards long, with the ends sewed together, so that by means of two sticks, one being at each end, turned in opposite directions, they may be wrung much dryer, when taken out of the boiling water, than could be effected by the hands. The principal circumstance to be attended to in the application of fomentations is the frequent renewal of them, in order that a steady and constant heat may be applied to the fomented part.

## No. II. MINERAL WATERS.<sup>4</sup>

It has been already noticed, that although no natural water is found in a state of absolute purity, yet that in general the quantity of foreign matters is not sufficient to give it any very sensible taste or odour. In some instances, however, these are so considerable, and of such a nature, as to prevent the water from forming a part of the nourishment of animals; in which case it is denominated a MINERAL WATER, and can be useful to mankind only in a medicinal point of view.

The substances found in mineral waters may be arranged under four heads.

### I. AIR and GASES:

1. Atmospheric Air, (*very common*.) It is generally in the proportion of  $\frac{1}{28}$  of the bulk of the water.
2. Oxygen Gas, (*rare*.)<sup>5</sup>
3. Azotic Gas, (*Buxton*<sup>6</sup>, *Harrowgate*<sup>7</sup>, *Lymington Priors*<sup>8</sup>, *Bath*.<sup>9</sup>)
4. Sulphureted Hydrogen Gas, (*Harrowgate*, *Moffat*.)

<sup>1</sup> Haygarth, *Clinical History of Diseases*, 8vo. Lond. 1805.

<sup>2</sup> Tepid bathing with friction is said by one author, “*vitam sæpe per plures menses, interdum etiam per aliquot annos, protraxisse.*” — Gregory, *Conspectus Med.* ii. 100.

<sup>3</sup> The Arabian physicians used the vapour bath in a singular mode, in scrophulous affections, which they denominated Bother, “*ponatur sub puero olla plena aquâ calidâ, in principio apparitionis pustularum, ut attrahat ab interioribus superfluum humorem ad corporis superficiem.*” *Rhazes de Morbis Infant.* cap. 19. by Willan, p. 31.

<sup>4</sup> The greater part of the chemical observations on mineral waters has been taken from Thomson’s *System of Chemistry*. <sup>5</sup> First detected by Scheele.

<sup>6</sup> Dr. Pearson.

<sup>7</sup> Dr. Garnet.

<sup>8</sup> Dr. Lambe.

<sup>9</sup> Dr. Priestley.



## II. ACIDS, in a free state:

1. Carbonic Acid, (*very common*).<sup>1</sup>
2. Sulphureous Acid, (*some hot springs in Italy.*)
3. Boracic Acid, (*some lakes in Italy.*)

## III. ALKALIES and EARTHS:

1. Soda, (*Geyzer and Rykum, hot springs in Iceland.*)<sup>2</sup>
2. Iodine, (*Salis, Piedmont.*)<sup>3</sup>
3. Silica, (*Geyzer, Rykum, Carlsbad, Pongues.*)<sup>4</sup>
4. Lime, (*doubtful.*)

## IV. COMPOUND SALTS:

1. Sulphate of Soda, (*very common.*)
2. ———— Ammonia, (*some volcanic springs.*)
3. ———— Lime, (*very common.*)<sup>5</sup>
4. ———— Magnesia, (*Epsom, and many other springs.*)<sup>6</sup>
5. ———— Alumina, (*very rare, but found at Ballycastle.*)<sup>7</sup>
6. ———— Iron, (*volcanic springs; also in some Danish waters, and those of Husley Green.*)<sup>8</sup>
7. ———— Copper, (*waters from copper mines.*)
8. Nitrate of Potash, (*some springs in Hungary, rare.*)
9. ———— Lime, (*some springs in Arabia.*)<sup>9</sup>
10. ———— Magnesia, (*rare.*)
11. Muriate of Potash, (*Uhleaborg, Sweden, rare.*)<sup>10</sup>
12. ———— Soda, (*very common.*)
13. ———— Ammonia, (*some springs in Italy and Siberia.*)
14. ———— Barytes, (*very uncommon.*)
15. ———— Lime, (*very common.*)
16. ———— Magnesia, (*very common.*)
17. ———— Alumina, (*uncommon.*)<sup>11</sup>
18. ———— Manganese, (*Lymington Priors, Tunbridge.*)<sup>12</sup>
19. ———— Carbonate of Potash, (*Berlin.*)<sup>13</sup>
20. ———— Soda, (*very common.*)
21. ———— Ammonia, (*rare.*)
22. ———— Iron, (*common.*)
23. Hydrosulphuret of Lime, } (*not uncommon in sulphureous springs.*)
24. ———— Soda, }
25. Sub-borate of Soda, (*lakes in Persia and Thibet.*)

These substances are not all contained in any mineral water, seldom more than five or six being present together; and they are generally in very minute quantity, the character and properties of the water depending on one or two ingredients which predominate. This allows mineral waters to be arranged into the four following classes: 1. ACIDULOUS WATERS; 2. CHALYBEATE WATERS; 3. SULPHUREOUS WATERS; 4. SALINE WATERS. We shall first give a sketch of the physical characters and medicinal properties of each of these classes; and then describe the method of determining the ingredients, and their proportions, contained in any mineral water.

1. ACIDULOUS WATERS owe their properties chiefly to carbonic acid. They sparkle when drawn from the spring, or when poured into a glass; have an acidulous taste, and become vapid when exposed to the air. Besides free carbonic acid, on the presence of which these qualities depend, acidulous waters contain generally also carbonates of soda, of lime, of magnesia, and of iron; and sometimes muriate of soda. They may be divided into *thermal* or *warm acidulous waters*, and *cold acidulous waters*; the temperature of the former, however, does not exceed 72°, while that of the latter is generally about 55°.

Of this kind are the springs of Cleves in Germany; Granshaw in Ireland; Languac, Upper Loire; Orston and Thorston, Nottinghamshire; Passi, near Paris; Stonefield, Lincolnshire; Tonstein in Germany; Wisbaden in Germany; and Bandola in Italy; but the most celebrated springs of this class are Pyrmont,

<sup>1</sup> Dr. Brownrigg.<sup>2</sup> Dr. Black.<sup>3</sup> M. L. Angelina of Voghera.<sup>4</sup> Dr. Bergman.<sup>5</sup> Dr. Lister, in 1682.<sup>6</sup> Dr. Grew, in 1610.<sup>7</sup> Dr. Garnet.<sup>8</sup> Dr. Rutty.<sup>9</sup> Dr. Home, in 1756.<sup>10</sup> Julin.<sup>11</sup> Dr. Withering.<sup>12</sup> Dr. Lambe, Dr. Scudamore.<sup>13</sup> Margraff



Seltzer, Spa, and Carlsbad. They are tonic and diuretic; and in large doses produce a sensible degree of exhilaration. They all afford a grateful and moderate stimulus to the stomach; but the Pyrmont and Spa, containing carbonate of iron, are especially useful in all cases of impaired digestion; while those which contain alkaline carbonates, as the Carlsbad and Seltzer, are more particularly employed as palliatives in calculous affections.

2. CHALYBEATE WATERS owe their properties to iron in combination generally with carbonic acid; and as this is usually in excess, they are often acidulous as well as chalybeate. The metal is found also in the form of a sulphate, but the instances of this are very rare.

Chalybeate waters have a styptic or inky taste; they are, when newly drawn, transparent, and strike a black with tincture of nut-galls; but an ochrey sediment soon falls, and the water loses its taste. If the iron be in the state of sulphate, however, no sediment falls; and the black colour is produced by the above test, even after the water has been boiled and filtered. Chalybeate springs are very numerous. On the Continent are those of Abcourt, St. Germain; Aumale and Forges near Rouen; Bologne; Buzot in Spain, (*a warm spring*); Caroline baths, Bohemia; Daswild, Bad, Germany; Driburgin, Westphalia; Naptha in Russia; Nizdeniee in Germany; Perekop, Russia; Sarepta, Russia; Scollensis, Switzerland; Suchalda, Hungary; and Vechy near Moulins. There are many chalybeates in Great Britain; as for instance, those of Arbroath and Peterhead in Scotland; Ashton in Wiltshire; Balemore, Worcestershire; Ballycastle, Antrim; Ballynahinch, Down; Ballyspellan, Kilkenny; Banden, Cork; Bromley, Kent; Brownstown, Kilkenny; Castlecomer, Kilkenny; Castleconnel, Limerick; Castlemain, Kerry; Coalcullen, Kilkenny; Corville, Tipperary; Coventry; Crowthorn, Waterford; Donemaile, Cork; Dunnard, Dublin; Galway; Garry-hill, Carlow; Haigh, Lancashire; Hampstead; Hartfell, Scotland; Islington; Kilcoran, Clare; Kilagee, Down; Kirby, Westmoreland; Lancaster; Llandridad, Wales; Luz, Essex; Listerlin, Kilkenny; Milltown, Mallay, Clare; Newtown Stewart, Tyrone; Oakfield, Cavan; Phoenix Park, Dublin; Scool, Clare; Shadwell, near London; Somersham, Huntingdonshire; and many others: but the most celebrated are Tunbridge, Brighton, and Peterhead: the Cheltenham spring also contains carbonate of iron; but on account of the large proportion of saline matter, and its strong purgative properties, it is not ranked in this class. The Spa springs also belong to this class.

Chalybeate waters are powerful tonics, and are employed in dyspepsia, scrophulous affections, cancer, amenorrhœa, chlorosis, and the other diseases of debility for which the artificial preparations of iron are used. Much of the benefit derived from the use of chalybeate waters depends on the extreme division of the metallic salts they contain, as well as the vehicle in which it is held in solution; while at the same time their operation is much modified by the carbonic acid by which the iron is suspended. When the water is a carbonated chalybeate, it should be drunk the moment it is drawn from the spring: but the same precaution is not necessary with a water containing sulphate of iron.

3. SULPHUREOUS WATERS derive their character chiefly from sulphureted hydrogen gas; which in some of them is uncombined, while in others it is united with lime or an alkali. They are transparent when newly drawn from the spring, and have the foetid odour of rotten eggs, which is gradually lost from exposure to the air and the water becomes turbid. When they are strongly impregnated with the gas, they redden infusion of litmus, and exhibit some other of the characteristics of acids; and, even in a weak state, they blacken silver and lead. Besides containing sulphureted hydrogen gas, they are not unfrequently, also, impregnated with carbonic acid. They generally contain muriate of magnesia or other saline matters, which modify their powers as a remedy.

The most important sulphureous springs in this island are those of Kilburn, Harrowgate, and Moffat; but the following are also of some note, Annaduff, Leitrim; Askeron in Yorkshire; Broughton in Yorkshire; Clonmell, Tipperary; Codsallwood, Staffordshire; Derrylister, Cavan; Drumasnane, Leitrim; Dudley, Worcestershire; Kedlistone, Derbyshire; Killashen, Fermanagh; Loansbury, Yorkshire; Maudley, Lancashire; Nottingham, Dorsetshire; Rippon, Yorkshire; Shapmoor, Westmoreland; Wardrew, Northumberland; and Wirksworth in Derbyshire. On the continent, Aix-la-Chapelle; Barege; Baden; Baia, Italy; Dux, Bayonne; Ems, Germany; Montmorency, near



Paris; Motte, near Grenoble; Viterbo in Italy; and St. Amands, near Valenciennes; which are resorted to chiefly for the cure of cutaneous eruptions, and are applied locally as well as drunk. They are slightly sudorific and diuretic, and are apt to occasion in some patients head-ach of short duration, directly after they are drunk. They are also employed for curing visceral and serophulous obstructions, torpor of the intestines, and some dyspeptic and hypochondriacal cases.

4. SALINE MINERAL WATERS owe their properties altogether to saline compounds. Those which predominate, and give their character to the waters of this class, are either,

1. Salts, the basis of which is lime;
2. Muricates of soda and magnesia;
3. Sulphate of magnesia;
4. Alkaline carbonates; particularly carbonate of soda.

They are mostly purgative, the powers of the salts they contain being very much increased by the large proportion of water in which they are exhibited. The most celebrated saline springs are those of Cheltenham, Leamington, Bristol, Kinalton, Pancras near London, Scarborough, Sydenham, and Thursk in Yorkshire, in England; Pitcaithly, in Scotland; and Sedlitz, on the continent. They are employed in diseases which require continued and moderate intestinal evacuations; such as dyspepsia, hypochondriasis, chronic hepatitis, jaundice, and strumous swellings. They are more grateful to the stomach when carbonic acid also is present; and when they contain iron, as in the case of the Cheltenham spring, their tonic powers, combined with their purgative qualities, render them still more useful in dyspeptic complaints and amenorrhœa.

To this class the water of the ocean belongs. The quantity of saline matter SEA WATER contains varies in different latitudes: thus, between  $10^{\circ}$  and  $20^{\circ}$  it is rather more than  $\frac{1}{24}$ th; at the equator it is  $\frac{1}{25}$ th; and at  $57^{\circ}$  north it is only  $\frac{1}{27}$ th. The saline ingredient, in 10,000 parts of sea water, according to the last analysis of Dr. Murray<sup>1</sup>, are, muriate of soda, 220.01; muriate of lime, 7.84; muriate of magnesia, 42.08; and sulphate of soda, 33.16. When brought up from a great depth, its taste is purely saline; but when taken from the surface it is disagreeably bitter, owing, perhaps, to the animal and vegetable matters suspended in it. Its specific gravity varies from 1.0269 to 1.0285; and it does not freeze until cooled down to  $28.5^{\circ}$  Fahrenheit. Its medicinal properties are the same as those of the saline purging waters, but more powerful; and as a bath, its efficacy is much superior to that of fresh water.

The general effects of mineral waters are modified by temperature, whether they be taken internally, or be externally applied. In some springs, as those of Bath, Matlock, Buxton, Wildbad, in Germany, Bagnols in France, Bonnes in the Lower Pyrennees, Borset near Aix-la-Chapelle, Canteries in the Upper Pyrennees, Digne in the Lower Alps, Klintschysela in Russia, Lucca in Italy, Plombieres, in Lorraine, and Pontgibault in France;<sup>2</sup> their virtues depend almost altogether on temperature; and in others, as Malvern, which has been found to contain scarcely any foreign matter, the simple diluent power of the pure water seems to produce the benefit that results from drinking them. Some of the good effects of all of them, however, must be allowed to proceed from change of scene, relaxation from business, amusement, temperance, and regular hours; and in these circumstances the drinking the waters at the springs possesses advantages which cannot be obtained from artificial waters, however excellent the imitations may be; nor even from the natural waters, when bottled and conveyed to a distance from the springs.

<sup>1</sup> *Edinburgh Transactions*, vol. viii. p. 205. The water was taken from the Frith of Forth, and was of the specific gravity 1.029.

<sup>2</sup> See, for a very full table of mineral waters, *Dr. Ryan's Essay on the Nat. Hist. of Water; Med. and Phys. Journ.* vol. liv. pp. 452—461.







Such are the known contents of the most celebrated mineral waters. Many more have been analysed, but it is unnecessary to introduce an account of them in this place; and I consider it to be of more importance to describe the method of determining the nature and proportion of the substances, or the analysis of mineral waters, one of the most difficult parts of practical chemistry.

### METHOD OF ANALYSING MINERAL WATERS.<sup>1</sup>

THE first circumstance to be attended to in the chemical examination of any mineral water, is to determine the gross weight of the substances held in solution. This is to be done by first ascertaining the specific gravity of the mineral water; then subtracting from it the specific gravity of distilled water (both expressed in whole numbers), multiplying the remainder by 14. The product is the gross saline contents, in a quantity of the water denoted by the number employed to indicate the specific gravity of distilled water.<sup>2</sup> Thus if the specific gravity of the mineral water be 1.079, as that of distilled water is 1.000, the remainder, after the subtraction of the latter from the former, in whole numbers, will be 79, which multiplied by 14 makes 1106; and therefore 110.6 is the sum of the saline contents of 1000 parts of the water; or 11.06 are contained in 100 parts. The next step is to ascertain the particular substances, and the proportion of each, contained in the water.

1. The AERIAL OR GASEOUS BODIES are to be first separated by boiling for a quarter of an hour, as much of the water as will fill two-thirds of a glass retort, connected with an inverted jar, divided into cubic inches and tenths, full of mercury, and placed in a mercurial trough. The air and gases will pass over into the jar, and depress the mercury; and when cool, after subtracting the air of the retort, the quantity of air expelled from the water may be easily determined.

The only gaseous bodies contained in water are atmospheric air, oxygen gas, azotic gas, carbonic acid gas, sulphureted hydrogen gas, and sulphurous acid: of which the following cannot exist together in the same water.

Oxygen gas and sulphureted hydrogen gas.

Sulphureted hydrogen gas and sulphurous acid.

If sulphureted hydrogen gas be present, it must be first separated; then the sulphurous acid, and carbonic acid gas; and, lastly, the oxygen and azotic gases.

a. *Sulphureted hydrogen gas* is known to be contained in water by its peculiar odour, by the water becoming turbid when exposed to the air and depositing sulphur, by its reddening the infusing of litmus fugaciously, blackening paper dipped in a solution of lead, and precipitating nitrate of silver black or brown. It may be separated from the air obtained from water during boiling, by carrying the jar into a tub of warm water and introducing nitric acid, which absorbs the sulphureted hydrogen. The bulk of this gas contained in any water is determined by filling a jar three-fourths with the water, inverting it in a water trough, and introducing nitrous gas at intervals, as long as red fumes appear, or the hepatic odour continues; when the jar is turned up and the air blown out. The nitrous gas in this operation mixing with the common air in the upper part of the jar, forms nitrous acid, which renders the water turbid, by decomposing the sulphureted hydrogen and precipitating sulphur. The bulk of hepatic gas is determined by the weight of the sulphur thrown down, one grain indicating the presence of 3.33 cubic inches of the gas.

b. *Sulphurous acid gas*, is ascertained by the same tests as discover the presence of sulphuric acid and water (*which see*).

c. *Carbonic acid gas* is detected by lime-water occasioning a precipitate soluble with effervescence in muriatic acid; by reddening fugaciously tincture of litmus, and losing this property when boiled.

To estimate the bulk of these gases, introduce into the air obtained by boiling the water, a solution of pure potass, and agitate the whole gently.

<sup>1</sup> The following observations on this important subject are chiefly extracted from the system of chemistry of Dr. Thomson.

<sup>2</sup> This useful formula was invented by Mr. Kirwan. See *Essay on Mineral Waters*, 145.



These acid gases will be absorbed, and any other gases left ; after which, the bulk of the residuum must be estimated, and subtracted from the bulk of the whole to obtain that of the acid gases absorbed. Evaporate next the potass slowly, nearly to dryness ; and by leaving it exposed to the atmosphere, sulphate of potass will be formed, which may be separated by dissolving the potass in diluted muriatic acid, and filtering the solution. 100 grains of sulphate of potass indicate 42.72 cubic inches of sulphureous acid gas, which being subtracted from the bulk of the gas absorbed by the potass, leaves the bulk of the carbonic acid gas.

- d. *Oxygen gas*, after the above gases are separated, may be examined by means of the solution of sulphate of iron saturated with nitrous gas.<sup>1</sup> A small graduated tube filled with the air to be examined is to be plunged into this solution, and moved backwards and forwards for a few minutes. The whole of the oxygen is rapidly absorbed, and by marking the greatest absorption, its bulk in a given quantity of the air is ascertained.
- e. *Azotic or nitrogen gas* is discovered by not being at all affected by eudiometrical processes.
2. **ALKALIES, and ALKALINE, EARTHY, and METALLIC CARBONATES.** Alkalies, even in minute quantities, are discovered in water by rendering infusion of turmeric, or paper stained with it, brown.<sup>2</sup> When the change is permanent, the fixed alkalies may be supposed to be present ; when fugacious, the alkali is ammonia. An infusion of Brazil-wood is rendered blue by the alkalies ; but this also is the case with the alkaline and earthy carbonates<sup>3</sup>, and the addition of sulphuric acid produces effervescence. Tincture of nutgalls discovers iron ; the colour is violet if alkaline carbonates or earthy salts be also present ; dark purple indicates other alkaline salts ; purplish red, sulphureted hydrogen gas ; and whitish and then black, sulphate of lime. Boiling the water precipitates the earthy and metallic carbonates.

The following substances of this class set down in the first column are incompatible, or cannot exist in mineral waters, with the salts placed in the opposite column.

Alkalies - - - - -	{	Fixed alkaline sulphates.
	{	Alum.
	{	Sulphate of magnesia.
	{	—— of iron.
Alkaline carbonates -	{	Muriate of barytes.
	{	—— of lime.
	{	—— of magnesia.
	{	Nitrate of lime.
Earthy carbonates -	{	Sulphate of iron.
	{	Muriate of barytes.
Carbonate of Magnesia	{	Sulphate of lime.
	{	Alum.
	{	Muriate of lime.

- a. *Alkalies and their carbonates* are ascertained to be present in mineral waters by the tests already mentioned ; and by the water, after being boiled, throwing down a precipitate on the addition of muriate of magnesia. The volatile nature of ammonia easily distinguishes it if present, which is very seldom ; and the best test for determining whether the fixed alkali be potass or soda is muriate of platina<sup>3</sup>, which forms an immediate precipitate with potass or any salt containing it, but is not at all affected by soda. The quantity of an alkali is determined by saturating it with sulphuric acid, and noting the quantity of real acid<sup>4</sup> necessary ; setting down, for every 100 grains of real acid used, 121.48 of potass, or 78.32 of soda ; and the loss of weight pro-

<sup>1</sup> Dr. Henry.

<sup>2</sup> This test is sufficiently delicate to detect soda when it amounts to  $\frac{1}{2217}$ th part only of the water.

<sup>3</sup> Sulphate of lime likewise produces the same effect.

<sup>4</sup> For a rule to determine the quantity of real acid in a diluted acid, see Introduction.



duced by the effervescence on dropping in the acid, being added to the above, shows the quantity of an alkaline carbonate.

b. *Iodine*, is detected by starch.

c. *Earthy carbonates*. If the water contains sulphureted hydrogen gas, this must be separated by exposing the water for a considerable time to the air, before the quantities of the earthy carbonates can be estimated. After thus exposing it, boil the water for fifteen minutes, filter it when cold; and treat what remains on the filter with muriatic acid, which will dissolve the carbonates of lime, of magnesia, and of iron. The residuum, which may contain carbonate of alumina, and perhaps sulphate of lime, is to be dried in a red heat, and its weight noted; and then boiled in a solution of carbonate of soda. The soda is next to be saturated with muriatic acid, and the mixture boiled for half an hour, which precipitates carbonate of lime and alumina. This precipitate being dried, the lime is to be separated by acetic acid, and the alumina that remains dried and weighed; so that by subtracting its weight from the original weight, the proportion of sulphate of lime is ascertained.

To estimate the contents of the muriatic solution, add to it ammonia as long as it throws down a reddish precipitate, which is the iron united with a portion of magnesia. The magnesia is to be separated by acetic acid, the precipitate being previously dried by exposure to the air, in a heat of  $200^{\circ}$ , and the solution added to the muriatic solution; and to determine the weight of the iron, it is to be redissolved in muriatic acid, then precipitated by an alkaline carbonate, and dried and weighed.

Sulphuric acid is now to be added to the muriatic solution; and the sulphate of lime, thus obtained, is to be heated to redness, and weighed; setting down for every 100 grains of it 74 of carbonate of lime. From the solution the magnesia is lastly to be separated by subcarbonate of soda, dried and weighed; then evaporate the remaining solution to dryness, and wash the residue with distilled water, so as to dissolve the muriate of soda. This residue is subcarbonate of magnesia, the weight of which, when dried, must be added to the former; which gives the entire weight of the subcarbonate of magnesia.

3. MINERAL ACIDS exist in mineral water, sometimes uncombined, but more generally combined with alkalies and earths, forming sulphates.

a. SULPHURIC ACID is readily detected by muriate of barytes, when it does not exceed the millionth part of the water. To render this test certain, however, the muriate must be diluted; the alkaline carbonates, if the water contain any, must be previously saturated with muriatic acid; and the precipitate must be insoluble in muriatic acid. The hydro-sulphurets are precipitated by muriate of barytes, but their presence is easily detected by their odour.

The proportion of sulphuric acid is easily estimated by saturating it with barytic water, and heating the precipitate to ignition; every 100 grains of this sulphate of barytes indicate 34 of real sulphuric acid.

b. The *Sulphates* contained in mineral waters are six in number, and are incompatible with the following salts placed in the opposite column.

Fixed alkaline sulphates	-				{ Nitrates of lime and of magnesia.
					{ Muriates of lime and of magnesia.
Sulphate of lime	-	-	-	-	{ Alkalies.
					{ Carbonate of magnesia.
					{ Muriate of barytes.
Alum	-	-	-	-	{ Alkalies.
					{ Muriate of barytes.
					{ Nitrate, muriate, carbonate of lime.
					{ Carbonate of magnesia.
Sulphate of magnesia	-	-			{ Alkalies.
					{ Muriate of barytes.
					{ Nitrate, muriate of lime.
Sulphate of iron	-	-			{ Alkalies.
					{ Muriate of barytes.
					{ Earthy carbonates.



- b. 1. *Sulphate of soda*. — To detect this salt, first evaporate the water to one-half, and add lime-water as long as any precipitate falls. This precipitates all the earths except sulphate of lime, which may be separated by evaporating the fluid till it become concentrated, then adding a little alcohol, and after filtration a little oxalic acid. If lime-water produces a precipitate in the water thus treated, immediately, or after a little alcohol be added, either sulphate of potass or of soda is present. To determine which, add acetate of barytes, which will precipitate sulphate of barytes; then filter and evaporate the filtered fluid to dryness, and dissolve the residue by digesting it in alcohol, and evaporate to dryness. If the sulphate be sulphate of potass, the dry salt thus obtained being acetate of potass, will deliquesce; but if it be sulphate of soda, the acetate will effloresce.

The proportion of the alkaline sulphates is found by precipitating their acid, by nitrate of barytes, from the water purified as above. If soda be the base of the salt contained in the water, for every 100 grains of this precipitate ignited, set down 61.2 grains of dried sulphate of soda; if potass be the base, for 100 grains of ignited precipitate set down 74.8 of dry sulphate of potass.

- b. 2. *Sulphate of lime* is detected by an immediate precipitate being formed by oxalic acid, or oxalate of potass, which, although a less sensible, is, nevertheless, a more accurate test. To determine its quantity, first saturate any earthy carbonates that may be present with nitric acid; then evaporate the fluid to a few ounces; and having precipitated the sulphate of lime by means of proof spirit, dry and weigh it.
- b. 3. *Alum* is detected by carbonate of magnesia, muriate of lime, muriate of magnesia, or succinate of ammonia. Twelve grains of alumina precipitated by carbonate of magnesia, heated to incandescence, indicate 100 grains of crystallized alum, or 49 of the dried salt.
- b. 4. *Sulphate of magnesia* may be detected in any water (previously freed from any alum or uncombined acids it might have contained) by hydro-sulphuret of strontian, which produces an immediate precipitate with this salt, and with no other. If no other earthy sulphate be present, the sulphuric acid may be separated by a barytic salt; every 100 grains of the ignited precipitate indicating 51 grains of dried sulphate of magnesia. If sulphate of iron be present, mix the water with a portion of argil, and expose it for some days to the air, during which time oxyd of iron and sulphate of alumina are precipitated, leaving the sulphate of magnesia alone in solution; which may be then estimated by the above method.
- b. 5. *Sulphate of iron* is detected by tincture of galls striking a black colour with the water after it has been boiled, and has cooled. Its quantity may be estimated by precipitating the iron by prussiate of potass.<sup>1</sup>
- c. *MURIATIC ACID*, either uncombined or combined, in mineral waters is detected by nitrate of silver, which forms with it a white precipitate, insoluble in nitric acid; but the alkaline carbonates, if any, must be first saturated by nitric acid; and any sulphuric acid removed by nitrate of barytes. The proportion of uncombined muriatic acid is ascertained by saturating it with barytic water, and then precipitating the barytes by sulphuric acid. For every 100 grains of the ignited precipitate set down 21 grains of real muriatic acid. Very minute portions of muriates may be detected by putting into some pure nitric acid contained in a porcelain capsule a minute quantity of finely divided gold, precipitated from its solution by sulphate of iron, and then adding the supposed muriate. If it be a muriate, a light tint will be gradually formed round the gold.<sup>2</sup>

<sup>1</sup> To make the calculation, the weight of a precipitate produced by the prussiate in a solution of a given weight of sulphate of iron in water must be previously determined.

<sup>2</sup> *Ann. de Chim.* xxviii. 26.



The *Muriates* contained in mineral waters are incompatible with the following  
d. articles in the second column.

Muriate of barytes	- -	{ Sulphates. Alkaline carbonates. Earthy carbonates.
Muriate of lime	- -	{ Sulphates, except of lime. Alkaline carbonates. Carbonate of magnesia.
Muriate of magnesia	-	{ Fixed alkaline sulphates. Alkaline carbonates.

- d. 1. *Muriates of soda and of potash* are detected in water by acetate of silver; but any earthy nitrates and muriates must first be decomposed by sulphuric acid, and the sulphates separated by alcohol and nitrate of barytes. To ascertain whether the precipitate be muriate of soda or of potass, evaporate to dryness, then dissolve the acetate in alcohol, and again evaporate to dryness. If it be acetate of potass the salt will deliquesce, but if acetate of soda, it will effloresce. To estimate the quantity of these salts, if they be unaccompanied by other salts, it is only necessary to dry and weigh the precipitate by nitrate of silver; setting down for every 100 grains of muriate of silver thus thrown down, 52 of muriate of potass; and 41 of muriate of soda. If alkaline carbonates be present, they must be first saturated with sulphuric acid, and sulphate of silver used to precipitate the muriatic acid.
- d. 2. *Muriate of Barytes* is detected by sulphuric acid. It is rarely found.
- d. 3. *Muriate of Lime*. To detect this salt the water must be first freed from the sulphates, then filtered, evaporated to dryness, the dry mass treated with alcohol, and the residue, after evaporating the alcohol, dissolved in water. If this solution yield a precipitate with acetate of silver, the water contained muriate of lime.
- d. 4. *Muriate of magnesia* is detected by separating the sulphates and proceeding as in the former case. If the aqueous solution of the dry mass treated with alcohol afford no precipitate with carbonate of lime; and if sulphuric acid and evaporation, with the addition of a little alcohol, occasion no precipitate, the solution contains only muriate of magnesia.
- d. 5. *Muriate of alumina* is detected by first saturating any alkali the water may contain with nitric acid, and separating any sulphuric acid by nitrate of barytes; and then adding carbonate of lime, which produces a precipitate if this salt be present. This process also precipitates muriate of iron and of manganese, if any be present.

To estimate the quantities of these muriates, which may all be contained in the same water, the earths, after separating any sulphates that may be present, are to be precipitated by barytes-water, and redissolved in muriatic acid. They are then to be separated by the rules already mentioned, and separately weighed. For every 50 grains of lime, set down 100 of dried muriate of lime; for 30 grains of magnesia, 100 of muriate of magnesia; and for 21.8 grains of alumina, 100 of muriate of alumina. The barytes of the muriate of barytes, which the addition of the barytes-water had formed in the mineral water by precipitating the earths, is now to be separated by sulphuric acid, and its muriatic acid expelled by heat; after which the muriate of soda, which the water originally contained, is to be obtained by evaporation.

- e. NITRIC ACID never exists in an uncombined state in mineral waters; and even the nitrates are comparatively of rare occurrence. The nitrates may be detected by an experiment, the inverse of that employed for detecting the muriates; that is by putting the fragment of gold into pure, colourless muriatic acid, and adding to it the suspected nitrate.

- f. The *nitrates* are incompatible with the salts, in the second column of the following table.

Nitrate of lime	- -	{ Alkaline carbonates. Sulphates, except of lime. Carbonates of magnesia and alumina.
Nitrate of magnesia	-	Fixed alkaline sulphates.



- f. 1. Nitrate of Potass* may occur in mineral waters in conjunction with sulphates and muriates; the former of which must be decomposed by acetate of barytes, and the latter by acetate of silver, before the nitrates can be estimated. After these previous steps, filter the water, then evaporate it to dryness, and treat the residue with alcohol; which dissolves the acetates, and leaves the nitre.
- f. 2. Nitrate of Lime* is detected by first concentrating the water, and separating the sulphates by alcohol; then filtering and distilling off the alcohol, and separating any muriatic acid by acetate of silver; afterwards, filtering again, evaporating to dryness, and dissolving the residue in alcohol, which must be also distilled off, and the dried residue dissolved in water. If oxalic acid detect lime in this solution, the mineral water contains nitrate of lime; the quantity of which may be estimated by precipitating with sulphuric acid, and calculating the quantity of lime contained in the sulphate; and for every 35 grains of lime setting down 100 grains of dry nitrate of lime.
- f. 3. Nitrate of Magnesia* is detected by nearly the same means; but to the last watery solution, instead of oxalic acid add potass, as long as any precipitate appears. Filter this solution; evaporate and treat the dry mass with alcohol. If a residue of nitre remains, the mineral water contained nitrate of magnesia.

Such is the general method of ascertaining the components of mineral waters, and the proportion of the ingredients contained in any particular water. To render the analysis complete, many minutiae must necessarily be attended to; but the detail of these would far exceed the limits which a work of this kind can admit of; and, after all, much must depend upon the ingenuity and expertness of the operator.

---

### No. III.

#### ON THE ART OF PRESCRIBING MEDICINES.

INDEPENDENT of the knowledge of diseases and the treatment of them, much of the success of the practitioner depends on circumstances connected altogether with the form in which the remedies are exhibited. In prescribing a medicine, even the best calculated to fulfil the object of the practitioner, it is necessary to consider the age, sex, temperament, habits, and idiosyncrasy of the patient, before the dose can be properly apportioned; and, as far as the medicine itself is regarded, the most convenient and agreeable form of exhibiting it, whether it should be given alone, or combined with other ingredients; and how far these are likely to impede, modify, or facilitate its operation. An attention to these circumstances is absolutely requisite to prevent the errors which too frequently occur in forming a prescription.

1. *Circumstances connected with the state of the patient. Age.* Here it must be observed, that the doses of the medicines described in the foregoing pages, are those adapted for an adult; but, as in the two extremes of life, childhood and old age, the body is weaker, and in early youth more susceptible of all impressions, these quantities cannot be administered with safety in every case; and thence the judgment of the prescriber must be exercised. Under ordinary circumstances, the following table, originally drawn up by Gaubius, may be considered as a sufficient guide for the young practitioner:



Ages.	Proportional quantities.	Doses.
For an Adult,	Suppose the dose to be - - - ONE	or 1 drachm.
Under 1 year	Will require only - - - $\frac{1}{12}$	— 5 grains.
2 years	_____ $\frac{1}{8}$	— 8 grains.
3 —	_____ $\frac{1}{6}$	— 10 grains.
4 —	_____ $\frac{1}{4}$	— 15 grains.
7 —	_____ $\frac{1}{3}$	— 1 scruple.
14 —	_____ $\frac{1}{2}$	— $\frac{1}{2}$ a drachm.
20 —	_____ $\frac{2}{3}$	— 2 scruples.
Above 21 —	The full dose - - - -	— 1 drachm.
65 —	The inverse gradation of the above.	_____

*Sex.* — Although some women possess as much bodily strength and vigour of constitution as the majority of men, yet the general greater delicacy and sensibility of the female frame, at every period of life, require not only caution in apportioning the doses of active medicines, which should be less than those ordered for men of the same age; but the medicines themselves should be such as are likely to fulfil the indications required, without much violence. The state of the uterine system likewise must not be overlooked in prescribing for a female. Thus the employment of aloetic and drastic purgatives, Cinchona bark, sulphuric acid, and astringents, should be suspended during the period of the catamenia.

*Temperament.* — It is undoubtedly true, that persons of different temperaments or original conformations of body are differently affected by the operation of medicines. Stimulants more readily affect those of a sanguine than those of a phlegmatic temperament; and, therefore, smaller doses are required. In the phlegmatic, also, the bowels are generally torpid, and require both a description of purgatives and such doses of them to excite the proper peristaltic motion, as would induce either visceral inflammation, or be followed by an alarming state of debility, were they administered to those of a sanguine temperament. Hence the necessity of attending to this circumstance in prescribing.

*Habits* — have a considerable influence in modifying the operation of medicines. Persons addicted to the use of spirits, narcotics, and other stimulants, are less easily excited both by medicinal stimulants and narcotics; and the knowledge of the habits of a patient, as far as the exhibition of purgatives is concerned, is absolutely necessary for the prescriber, many people being in the almost daily habit of taking this class of remedies without consulting a medical practitioner. In the first of these cases, larger doses of stimulants and narcotics are required to produce the ordinary effects of these remedies; but in the second a change of the purgative usually taken will generally be sufficient. In the employment of medicines, also, which require to be long continued, the beneficial effect is soon lost if the doses be not increased.

*Idiosyncrasy.* — Many persons have a peculiarity of disposition, unconnected with temperament, which renders them liable to be effected by substances, taken into the stomach, either in the form of food or of medicine, in a manner different from the majority of mankind. Such a state can be discovered only by accident or by time; but when it is known, it must be attended to by the practitioner. Instances in which opium proves deleterious in every form and dose, are not unfrequent. I knew a lady, in whom the smallest dose of squill excited an erythematic eruption over the whole body; spirit of turpentine, also, frequently produces a similar eruption; and many examples of the same kind might be quoted. But besides these guides in forming a prescription, the choice of the medicine must occasionally depend on the circumstance of the patient being more or less immediately under the eye of the prescriber. Thus, if the patient can be seen every day, or frequently, by the practitioner, the most active medicine which the nature of the case requires should be chosen; but if he cannot be frequently seen, or is not resident in the same place, the practitioner should choose a remedy of the same class, but less likely to have a sudden or violent effect. Thus in prescribing for ascites under such circumstances, Squill or Acetate of Potass, or Supertartrate of Potass, must be pre-



ferred to Elaterium; for intermittent fever, Cinchona bark to the Arsenical solution; and so in other cases, when the patient is not under the eye of the prescriber.

2. *Of the Form and Composition of extemporaneous Prescriptions.* — In every prescription simplicity should be kept in view, and when one medicine will answer the intention of the prescriber, it ought to be preferred. The nauseous taste, however, and the other qualities of the great majority of drugs require the addition of others to modify their action: but, although medicines are more generally prescribed in a compound form, yet the practice of accumulating a great variety of ingredients in one prescription must be avoided.

Medicines exhibited in the fluid form operate sooner, and with more certainty, than in the solid state; but in choosing the vehicle or solvent, the taste of the patient ought not to be overlooked. Thus, for those to whom peppermint-water is not disagreeable, the nauseous taste of Sulphate of Magnesia is more completely concealed by that vehicle than any other; if Cinchona bark in powder be ordered, milk effectually covers its taste, provided the dose be taken the moment it is mixed; and if aloes, the most nauseous article of the materia medica, be prescribed in a fluid form, a solution of extract of Liquorice renders it by no means unpalatable. Medicines which, when given alone, produce griping, require the addition of aromatics to correct that quality, and, when they operate with violence, mucilages and demulcents are sometimes necessary to obtund their acrimony, or narcotics to moderate their action. In prescribing purgatives, it is also necessary to consider the particular part of the alimentary canal on which they more immediately act. Thus Rhubarb acts chiefly on the pylorus and duodenum, Aloes on the rectum, and Calomel and Jalap on the larger intestines. Another reason for ordering medicines, in a compound form, is the necessity of producing two or more effects at one time. Thus the same dose may be required, in a case of colic for example, to allay pain and to open the bowels; or, in fever, to determine to the surface, to allay irritation, and to produce sleep. But in combining medicines, care must be taken not to bring together incompatibles, or substances that decompose each other, or chemically combine, and, consequently, alter the nature of the mixture, or render it inert; unless the resulting compound be the remedy on which the practitioner relies. Thus acids and alkalies are incompatible, unless the neutral salt they produce be the remedy required; and astringent vegetable infusions and decoctions destroy the emetic and diaphoretic property of tartar emetic. Hence the necessity of a knowledge of chemistry to the medical practitioner.

In writing a prescription, the first object is the principal or most active ingredient, which is called the *basis*; the next the *adjuvans*, or that which is designed to promote the action of the basis; the third, the *corrigens*, or that intended to correct or modify its action; and the last, the *vehiculum*, or that substance in which the more active ingredients are to be exhibited, and which, consequently, gives the formula its peculiar character. It has been usually regarded as a proper rule in writing a prescription to place the basis first, and the other articles in the form in which they have been enumerated; but this must depend on the mode best fitted for compounding the medicine. Thus salts and other soluble solids should be placed before the menstruum in which they are to be dissolved: and volatile substances should always be placed last, as they are necessarily the last ingredients added in the manipulation of the compound. Finally, the names of each ingredient should be written at full length, in a legible hand, and the symbols of the quantities distinctly marked; and no prescription should pass from the hand of the prescriber, without being deliberately read over, and its correctness ascertained.

To facilitate the art of prescribing to the young practitioner, I have added a Table of Incompatibles, and a few examples of the usual Forms of extemporaneous Prescriptions.



# LATIN INDEX.

[The initial letters, placed after the names of the articles, denote the Pharmacopœia in which they are found.]

	PAGE		PAGE
A.		Æther c. alcohole arom. E.	916
ABIETIS resina L.	478	Agrimonia D.	147
Abrotani folia D.	198	Alcohol L. E. D.	778
Absinthium L. D.	200	dilutus E.	575
Absinthii maritimi cacumina D.	ib.	ammoniatum E.	881
Acaciæ gummi L. E.	131	foetidum D.	882
Acetosæ folia L.	539	Alii radix L. E. D.	149
Acetosella L.	450	Alöes spicatae extractum L.	151
Acetas ferri D.	757	vulgaris extractum L.	153
hydrargyri E. D.	763	hepatica E. D.	ib.
plumbi E. D.	785	Altheæ folia et radix L. E.	155
kali D.	682	Alumen L. E. D.	156
potassæ E.	681	exsiccatum L. ustum D.	706
Acetum L. E. D.	133	Amomi repentis sem. E.	412
aromaticum E.	925	zingiberis radix E.	626
colchici L.	ib.	Ammonia sub-carbonas L.	667
distillatum D.	631	murias L.	159
scillæ L. D.	926	Ammoniacum L. E. D.	357
scilliticum E.	ib.	Ammoniaretum cupri E.	743
Acidum aceticum (dilutum) L. E.	631	Amygdala amara, dulcis L. E. D.	162
aceticum D.	632	Amylum L.	598
camphoratum D.	923	Amyridis gileadensis resina E.	165
forte E.	633	Anchusæ tinctoriæ radix E. D.	167
benzoicum L. E. D.	634	Angelica ; radix E.	169
citricum L.	636	Anethi semina L. E.	167
hydrocyanicum	656	Anthemidis flores L. E.	171
oxalicum	660	Angustura D.	216
muriaticum L. E. D.	639	Antimonii sulphuretum L.	174
dilutum D.	642	sulph. præcipitatum L.	727
nitricum L. E.	645	vitrum	176
dilutum L.	649	Anisi semen L. D.	473
nitrosum E. D.	645	Antimonium tartarizatum L.	728
dilutum E. D.	649	Aqua acetatis ammoniæ E. D.	672
succinicum D. E.	651	alkalina oxymuriatica D.	691
sulphuricum L. E. D.	137	ammoniæ E. caustica D.	670
dilutum L. E. D.	653	carbonatis D.	675
aromaticum E.	654	anethi L.	826
tartaricum	ib.	calcis E. D.	708
Aconiti folia L. E. D.	142	composita D.	841
Acori calami radix E. D.	144	carui L.	826
Adeps L.	589	cinnamomi L. D.	827
ovillus E.	449	citri aurantii E.	826
suillus D. E.	589	medicæ E.	827
Ærugo L. D.	307	cupri ammoniati D.	774
præparata D.	742	distillata L. E. D.	825
Æsculi hippocast. semen, cort. E. D.	145	fæniculi L. D.	827
Æther nitrosus D.	917	kali caustici D.	676
rectificatus L.	909	lauri cassiæ E.	827
sulphuricus L. E. D.	908	cinnamomi E.	ib.
cum alcohole E.	914	muriatis calcis D.	710
		menthæ piperitæ L. E. D.	827
		menthæ viridis L. E. sativæ D.	731



	PAGE		PAGE
Aqua oxymuriatica D.	642	Canceri paguri chelæ E. D.	225
picis liquidæ D.	841	Cancrorum lapilli præparati E.	ib.
pimentæ L. E. D.	827	Canellæ cortex L. E. D.	226
potassæ E.	676	Cantharis vesicatoria E. D.	227
pulegii L. D. E.	828	Capsici baccæ L. D. E.	232
rosæ L. E. D.	ib.	Carbo ligni L. E. D.	234
subcarbonatis kali D.	678	Carbonas ammoniæ E. D.	667
sulphureti ammoniæ D.	801	barytæ E.	210
sulphureti kali D.	799	calcis E.	223
supercarbonatis potassæ E.	694	præparatus E.	711
sodæ E.	703	ferri D. præcipitatus E.	748
Arbutus uvæ ursi, folia E.	178	magnesiæ E.	717
Ari radix recens D.	201	plumbi	491
Aristolochiæ serpentariæ, radix E.	182	potassæ E.	684
Aretii lappæ radix E.	180	purissimus E.	ib.
Argentum L. E. D.	ib.	sodæ E. D.	695
Argenti nitras L. D.	734	siccata D.	700
Armoraciæ radix L.	283	zinci impurus E.	624
Arnica montanæ herba E. D.	184	præparatus E.	790
Arsenicum album L. D.	186	Cardamines flores L. D. E.	236
sublimatum L.	738	Cardamomi semina L. D.	412
Arsenias kali D.	739	Cardui benedicti folia D.	244
Artemisiæ santonicæ cacumina E.	199	Caricæ fructus L. D.	337
absinthium E.	200	Carui semina L. E.	236
Asari folia L. D. E.	202	Caruon D.	ib.
Assafoetidæ gummi resini L. E. D.	335	Caryophylli L. E. D.	327
Astragalus tragacantha; gummi E.	205	oleum L. E. D.	ib.
Atropæ belladonnæ folia E.	206	Caryophyllum rubrum, flores D.	318
Aurantii baccæ, cortex L.	279	Cascarillæ cortex L. D.	299
Avenæ semina L. E.	209	Cassiæ pulpæ fistulæ fructus L. E.	238
		Cassia fistularis D.	ib.
B.		lignea cort. flor. nond. exp. D.	ib.
Balsamum copaibæ D.	293	sennæ foliæ E.	239
peruvianum L. D.	433	Castoreum L. E. D.	242
tolutanum L. D.	569	Cataplasma fermenti L.	989
canadensæ D.	477	sinapeos D.	990
Bardanæ radix D.	180	sinapis L.	ib.
Barilla D.	566	Catechu extractum L. E. D.	128
Beccabungæ herba D.	606	Centuriæ benedictæ herba E.	244
Belladonnæ folia L. D.	206	cacumina L.	256
Benzoinum L. Benzoe D.	583	Centaurium minus D.	ib.
Bismuthum L.	212	Cepæ radix D.	150
Bismuthi subnitras L.	740	Cera flava et alba L. E. D.	251
Bistorta L. D.	496	Ceratum L.	972
Bitumen petroleum E.	213	calaminæ L.	ib.
Boletus ignarius E.	215	cetacei L.	ib.
Borax D. Boras sodæ E.	564	cantharidis L.	972
Bonplandiæ cortex E.	216	plumbi acetatis L.	973
Bubon galbanum; gum. res. E.	221	compositum	ib.
		resinæ L.	973
C.		sabinæ L.	974
Cajaputi oleum L.	415	saponis L.	975
Calami radix L.	144	simplex E.	972
Calamina L.	624	Cerevisiæ fermentum L.	255
præparata L.	789	Cervi elaphi cornu E.	254
Calcis murias L.	709	Cerussa D. E.	491
Calumbæ radix L.	418	Cetaceum L.	471
Calx L. D.	222. 707	Chamæmelum; flores D.	171
Cambogia L.	578	Chamædrys herba D.	595
Camphora L. E. D.	325. 389	Chelæ cancrorum E.	225
Canceri astaci lapilli E.	225	Chironiæ centaurii semina E.	256
		Cicuta D.	288
		Cineres clavellati D.	501



	PAGE		PAGE
Cinchona cordifoliae cortex L.	265	Decoctum althæae officinalis E.	843
lancifoliae cortex L.	258	aloes compositum L.	ib.
oblongifoliae cortex L.	263	anthemidis nobilis E.	ib.
Cinnamomi cortex L. D.	384	chamæmeli comp. D.	ib.
oleum L. D.	ib.	cinchonæ L. E. D.	ib.
Citri aurantii cortex E. D.	279	cornu cervini D.	674
fructus immat. D.	ib.	cydoniæ L.	847
Citrus medicus; fructus, &c. E.	277	daphnes mezerei E.	898
Coccus L. cacti E.	281	digitalis D.	ib.
Coccinella D.	ib.	dulcamaræ L.	849
Cochleari armoraciæ radix E.	283	geoffrœae inermis E.	ib.
Coci butyraceæ oleum fixum E.	285	guaiaci compositum E.	ib.
Colchici radix L. D. E.	ib.	hordei L. E. D.	ib.
Colocynthis pulpa L.	302	compositum L. D.	850
fructus, medulla D.	ib.	lichenis L. D.	ib.
Colombæ radix E.	418	malvæ compositum L.	851
Colombo radix D.	ib.	papaveris L.	ib.
Confectio amygdalarum L.	939	polygalæ senegæ E.	853
aromatica L.	940	quercus L.	851
aurantiorum L. D.	ib.	sarsaparillæ L. D.	852
cassiæ L.	ib.	comp. L. E. D.	853
piperis nigra L.	941	senegæ L.	ib.
opii L.	ib.	ulmi L. D.	ib.
rosæ caninæ	942	veratri L.	854
gallicæ L.	ib.	Delphinium staphisagriæ sem. E.	317
rutæ L.	943	Dianthi caryophylli flores E.	ib.
scammoneæ L,	ib.	Digitalis folia L. E. D.	319
sennæ L.	ib.	Doliches, setæ leguminum D.	323
Conii maculati folia L. E.	288	Dolichi pubes L. E.	ib.
Conserva aurantii E. D.	940	Dorstenia contrayerva E.	324
rosæ caninæ E.	942	Dulcamaræ caulis L. D.	568
gallicæ E. D.	ib.	E.	
rosæ D.	ib.		
Contrayervæ radix L.	324		
Convulvi jalapæ, radix E.	292	Elaterium E.	864
Copaiba L. E.	293	Elaterii poma L. D.	424
Coriandri semina L. E. D.	295	Electuarium aromaticum E.	940
Cornua L.	254	cassiæ fistulæ E.	941
Cornu cervinum D.	ib.	cassiæ D.	ib.
Cornu ustum L.	961	catechu comp. D. E.	942
Creta L. D.	223	opiatum E.	941
præcipitata D.	712	scammonii D.	943
præparata L. D.	711	sennæ D. E.	944
Croci stigmata L. E. Crocus D.	296	Elemi L. D.	164
Croton eleutheriæ cortex E.	299	Emplastrum ammoniaci	964
Cubeba L.	485	cum hydrargyro L. D.	ib.
Cucumeris colocynthidis pulpa E.	302	aromaticum D.	ib.
Cummini semina L.	304	assafœtidæ E.	965
Cuprum E. D.	306	calefaciens D.	ib.
ammoniaturum L. D.	743	cantharidis L. E. D.	966
Cupri subacetas	307	cerae L.	965
Cupri sulphas L. E.	309	cumini L.	ib.
Cuspariæ cortex L.	216	galbani D.	966
Cydoniæ semina L.	512	galbani comp. L.	965
		gummosum E.	ib.
		hydrargyri L. E.	ib.
		lythargyri D.	970
		cum resina D.	ib.
Daphnes mezerei cortex E.	312	oxidi ferri rubri E.	969
Daturæ stramonii herba E.	314	plumbi semivit E.	970
Dauci radix L.	316	opii L.	968
carotæ semina E.	ib.	piceis compositum L.	969
silvestris semina D.	ib.		



	PAGE		PAGE
Mistura amygdalarum L.	872	Oleum olivæ L. D.	443
assafoetidæ L.	874	pimentæ L. D.	819
camphoræ L.	ib.	pini purissimum E.	822
camphorata D.	ib.	pulegii L. D.	819
cornu usti L.	875	ricini L.	812
cretæ L. D.	ib.	rosmarini L. D.	819
ferri composita L.	ib.	rutæ D.	820
guaiaici L.	876	sem. fœniculi dulcis D.	817
moschi L.	ib.	succini L. E.	821
Momordica elaterium, fruc. rec. E.	424	purissimum E.	ib.
Mori baccae L.	425	rectificatum D.	ib.
Moschus L. E. D.	426	sulphuratum L. E.	797
Mucilago acaciæ L. E.	842	terebinthinæ rectif. D.	822
amylæ L. E. D.	843	vol. pimpinellæ anisi E.	816
astragali tragacanthæ E.	ib.	juniperi communis E.	817
gummi arabici D.	842	lavandulæ spicæ E.	ib.
tragacanthæ D.	843	lauri sassafras E. D.	818
Murias ammoniæ E.	159	rorismarini officin. E.	819
et ferri E. D.	746	Olibanum L. D.	219
barytæ E.	713	Olivæ oleum L.	443
calcis L.	709	Opium L.	452
hydrar. corrosivus D. E.	786	succus concretus D.	ib.
sodæ E.	561	purificatum D.	808
siccatum E. D.	695	Opoponax L.	469
Myristicæ nuclei L.	430	Origanum L.	446
Myristici moschatæ nuclei involucrum,		marjoranæ herba E.	447
vulgo macis E.	ib.	Ovum L.	470
Myroxylæ peruif. balsamum E.	433	Oxidum arsenici E.	186
Myrrha L. E.	435	antim. cum phosphate calcis E.	732
gummi resina D.	ib.	ferri rubrum E. D.	752
Myrtæ pimentæ fructus E.	438	nigrum pur. E. D.	745
		hydrargyri D.	767
N.		rubrum per acid. nit. E.	761
Nicotianæ tabaci folia E. D.	439	cinerium E.	764
Nitrum D.	497	nitricum D.	761
Nitras argenti E. D.	735	sulphuricum D.	782
potassæ	497	plumbi rubrum E.	494
Nux moschatæ oleum, &c. D.	430	zinci E.	790
		impur. præp. E.	625
O.		Oxymel L. D.	928
Oleæ europææ oleum fixum E.	443	æruginis D.	986
Oleum ammoniatum E.	986	colchici D.	928
æthereum L.	913	scillæ D. E.	929
amygdalarum L. D.	810		
amygdalæ communis E.	ib.	P.	
anisi L. D.	816	Papaveris capsulæ L. E.	413
anthemidis L. D.	ib.	Papaver album; capsulæ D.	452
camphoratum E. D.	987	erraticum; petala D.	451
cajuput D.	415	som. capsulæ; opium E.	452
carui L. D.	816	Petroleum L. D.	214
cornu cervini rectif. D.	820	Phosphas sodæ E. D.	696
juniperi L. D.	817	Pistacia lentiscus; resina E.	488
sabinæ E. D.	820	Pilulæ aloeticæ E.	953
lavandulæ L. D.	817	alöes cum zingibere D.	ib.
lini L. D. E.	811	compositæ L.	ib.
menthæ piperitæ L. E. D.	818	cum assafoetida E.	ib.
viridis L.	ib.	alöes cum myrrha L. E. D.	953
sativæ D.	ib.	ammoniaretæ cupri E.	954
myrti pimentæ E.	819	assafoetidæ compositæ E.	955
origani L. E. D.	818	colocynthis compositæ E.	954



	PAGE		PAGE
Pilulæ galbani compositæ L.	955	Pulvis hydrargyri cinereus D.	765
gambogiæ compositæ L.	954	jalapæ compositus E.	948
ferri compositi L.	955	ipēcacuānhæ compositus L. D. ib.	
hydrargyri L. E. D.	956	et opii E.	949
hydrar. submuriatis comp. L.	957	kino compositus L.	949
opiatæ E.	958	opiatus E.	947
myrrhæ compositæ D.	955	salinus comp. E.	950
rhei compositæ E.	957	quercus marinæ D.	949
saponis cum opio L.	ib.	salinus comp. E.	930
storace D.	958	scammoneæ compositus L. E. ib.	
scillæ compositæ L.	ib.	sennæ compositus L.	ib.
scilliticæ E.	ib.	scillæ D.	ib.
scillæ cum zingibere D.	ib.	spongiæ ustæ D.	951
subcarbonatis sodæ	ib.	stanni D.	785
sulphatis ferri compositæ	ib.	aluminis comp. E.	951
Pimentæ baccæ L. D.	438	tragacanthæ comp. L.	ib.
Pimpinellæ semen E.	473	Punica granatum ; fructus, &c. E.	510
Piperis longi fructus L. E. D.	484	Pyrethri radix E. D.	172
Piper nigrum L. D.	482		
nigri fructus E.	484	Q.	
Pix abietina L.	478	Quassiæ lignum L. D. E.	515
burgundica D. E.	ib.	simaroubæ cort E.	513
liquida L. E. D.	474	Quercus cortex L. D. E.	519
Plumbum E.	489	marina, herba D.	341
Plumbi subcarbonas L.	491		
oxydum semivitreum L. E.	493	R.	
acetas L.	785	Raphani rustici radix D.	283
Polygalæ senegæ radix E.	495	Resina alba D. E.	432
Polygoni bistortæ radix E.	496	flava nigra L.	474
Polypodium filix mas ; radix E.	204	Ramni baccæ L. E. D.	521
Porri radix L.	148	Rhei radix L. D. E.	522
Potassa E.	679	Rhœados petala L.	451
impura L.	601	Rhodendri chrysanti, folia E.	529
fusa L.	679	Rhus toxicodendri, folia E.	530
cum calce L. E.	ib.	Ricini semina et oleum L. E.	531
Potassæ acetas L.	681	Ricinus ; oleum e seminibus exp. D. ib.	
impura carbonas L.	501	Rosæ caninæ pulpa L.	535
carbonas L.	684	Rosæ caninæ fructus recens E.	ib.
nitras L. E.	497	Rosæ centifoliæ petala L. E.	533
subcarbonas L.	685	Rosa damascena ; petala D.	ib.
sulphas L.	503. 688	Rosæ gallicæ petala L. E.	534
sulphuretum L. E.	796	Rosa rubra ; petala D.	ib.
supersulphas L.	691	Rosmarini cacumina L. E.	536
supertartras L.	529	Rosmarina herba D.	ib.
tartras L.	692	Rubiæ radix L. D.	537
Potio carbonatis calcis E.	875	Rubia tinctorum ; radix E.	ib.
Pruna L.	503	Rumicis acetosæ folia E.	539
Pruni gānciæ fructus E. D.	ib.	Rumex aquaticus radix D.	539
Pterocarpi lignum L.	504	Rutæ folia L. D.	540
Pulegium L. D.	256	Ruta graveolens ; herba E.	ib.
Pulvis alōes cum canella D.	946		
compositus L.	946	S.	
cum guaiaco D.	ib.	Sabinæ folia L. D.	376
antimonialis L. D.	732	Saccharum L. E.	541
aromaticus E. D.	947	rubrum D.	ib.
asari comp. D. E.	946	purificatum L. D.	ib.
carbonatis calcis comp. E.	948	Sagapenum ; gummi resina L. E. D.	544
cinnamomi comp. L.	947	Sal ammoniacum D.	26
contrayervæ comp. L.	ib.	commune D.	561
cornu cervini usti D.	ib.	Salicis cortex L. D.	545, 546
cornu usti c. opio L.	ib.		
cretæ comp. L.	948		
c. opio L.	ib.		



	PAGE		PAGE
Salvia officinalis ; folium E. D.	548	Spiritus menthæ viridis L.	886
Sambuci flores, baccæ, cort. L. E. D.	549	myristicæ L.	887
Santalum rubrum ; lignum E. D.	504	myrti pimentæ E.	ib.
Santonicæ semen D.	199	mucis moschatæ D.	ib.
Sapo durus L. E. D.	550	pimentæ L.	ib.
mollis L. E.	552	pimento D.	ib.
Sarsaparillæ radix L. D.	559	pulegii L.	ib.
Sassafras, lignum et radix L. E. D.	394	raphani comp. D.	883
Scammoniæ gummi resinæ L. E. D.	290	rectificatus L.	572
Scillæ radix L. D.	553	rosmarina L. D.	886
Scrophularia ; herba D.	555	officinalis E.	ib.
Senegæ radix L.	495	tenuior L.	575
Seneka radix D.	ib.	vinosus rectificatus D.	572
Sennæ folia L. D.	238	tenuior D.	575
Serpentariæ radix L. D.	182	Spongia L. E. D.	577
Sevum L. D.	449	usta L.	962
præparatum L.	961	Stannum L. E. D.	580
Simaroubæ cortex L. D.	513	Staphisagriæ semina L. D.	317
Sinapis albæ semina E. D.	556	Stramonium ; herba D.	314
semina L.	ib.	Styracis balsamum L. E.	582
Sium, herba D.	558	Styrax calamita ; resina D.	ib.
Smilax sarsaparilla ; radix E.	ib.	officinalis ; balsamum E.	ib.
Sodæ sub-boras L.	564	purificata D.	808
Soda impura L. (subcarbonas) E.	566	Subacetas cupri E.	307
Sodæ murias L. E.	561	Sub-boras sodæ D. L.	564
carbonas L. E.	695	Subcarbonas ferri E.	747
subcarbonas L.	698	kali D.	685
sulphas L. E. D.	565.700	potassæ impurus E.	501
tartarizata L.	703	Submurias hydrargyri mitis E.	775
Solani dulcamara caules	568	hydrarg. ammoniatum D.	773
Solutio acetatis zinci E.	794	hydrargyri præcip. E. D.	778
arsenicalis E.	738	hydrargyri sublimatum D.	775
muriatis barytæ E.	715	Subsulphas hydrargyri flavus E.	782
muriatis calcis L.	710	Succinum L. E. D.	584
sulphatis cupri comp. E.	745	Succus spis. atropæ belladonnæ E.	859
sulphatis zinci E.	794	cicutæ D.	863
Spartii cacumina L. E.	570	aconiti napelli E.	857
Spermaceti E. D.	471	spiss. conii maculati E.	863
Spigeliæ radix L. E. D.	571	Sambuci nigræ E.	870
Spiritus ætheris aromaticus L.	913	Sulphas aluminæ L.	156
comp. L.	916	barytæ E.	211
nitrici L.	914	ferri E. D.	750
nitrosi E.	ib.	exsiccatus E. D.	752
æthereus nitrosus D. E.	ib.	kali D.	688
ætheris sulphurici L.	916	magnesiae E. D.	406
comp. D.	ib.	potassæ cum sulphure E.	690
ammoniæ L. D.	831	potassæ E.	688
aromaticus L. D.	ib.	Sulphur L. E. D.	585
foetidus L. D.	832	antimoniatum fuscum D.	726
succinatus L.	ib.	lotum L.	798
colchici ammoniatus	884	præcipitatum L.	799
anisi L.	883	sublimatum L. E. D.	585
armoraciæ comp. L.	ib.	lotum E. D.	798
camphoræ L.	ib.	Sulphuretum antimonii E. D.	174
carui L. D.	884	ferri D.	800
cari carui E.	ib.	ant. præparatum E.	724
cinnamomi L. D.	ib.	præcipitatum E.	725
juniperus compositus L. D.	ib.	hydrargyri nigrum E.	779
lavandulæ L. D.	885	rubrum D.	780
lavandulæ spicæ E.	ib.	kali D.	797
lavandulæ comp. L. E. D.	885	Supersulphas aluminæ et potassæ E.	156
lauri cinnamomi E. D.	884	aluminæ et pot. exs. E.	706
menthæ piperitæ L. D.	886	Supertartras potassæ E.	587



	PAGE		PAGE
Swietenia febrifugæ cortex D.	591	Tinctura acetatis zinci D.	795
Syrupus L. D.	930	alöes L. D.	889
aceti E.	ib.	ætherea E.	890
allii D.	ib.	composita L. D.	ib.
althææ L.	931	et myrrhæ E.	ib.
officinalis E.	ib.	amomi repentis E.	983
ammomi zingiberis E.	938	angusturæ D.	982
aurantiorum L. D.	931	assafœtidæ L. D.	890
caryophylli rubri D.	932	ammoniatæ E.	832
citri aurantii E.	931	aurantii L. D.	891
medici E.	933	benzoini composita L. E.	ib.
colchici autumnalis E.	932	benzöes comp. D.	ib.
croci L.	ib.	bonplandia trif. E.	ib.
dianthi caryophylli E.	ib.	calumbæ L.	892
limonum L. D.	ib.	cantharidis E. D.	ib.
mori L.	933	colombæ E. D.	ib.
opii D.	ib.	camphoræ composita L.	ib.
papaveris L.	ib.	capsici L.	893
albi D.	934	cardamomi L. D.	ib.
somniferi E.	933	comp. L. D.	ib.
erratici D.	934	cascarillæ L. D.	894
rhæados L.	ib.	crotonis eleutheriæ E.	ib.
rhamni L.	935	castorei L. E. D.	ib.
cathartici E.	ib.	comp. L. E.	ib.
rosæ L.	ib.	catechu L. E. D.	895
centifoliæ E.	ib.	cinchonæ L. D.	ib.
gallicæ E.	936	lancifoliæ E.	ib.
sarsaparillæ L.	ib.	ammoniata L.	ib.
sennæ L. D.	ib.	cinnamomi L. D.	896
scillæ maritimæ E.	ib.	cinnam. comp. L. E. D.	ib.
tolutanus L.	937	conii maculati E.	897
toluiferæ balsami E.	ib.	convolvuli jalapæ E.	900
violæ odoratæ E.	ib.	croci D.	897
violæ D.	ib.	sativæ E.	ib.
zingiberis L. D.	938	digitalis L. D.	ib.
		purpureæ E.	ib.
T.		ferri ammoniati L.	756
Tabaci folia L.	439	muriatis L. E. D.	ib.
Tamarindi pulpa L.	592	furulæ assafœtidæ E.	890
indicæ fructus E. D.	ib.	galbani D.	897
Tanacetii vulgaris fol. flores E. D.	594	gallarum D.	ib.
Taraxaci radix L. (folia.) D.	396	gentianæ comp. L. E. D.	898
Tartarum	587	guaiaci L. D.	ib.
antimoniatum D.	728	officinalis E.	ib.
crystalli D.	529	ammoniata L. E. D.	ib.
ferri D.	753	hellebori nigri L. E. D.	899
Tartras kali D.	693	humuli L.	ib.
sodæ et kali D.	703	hyosciami L. D.	900
antimonii E.	641	nigri E.	ib.
potassæ E.	693	jalapæ L. D.	ib.
et ferri E.	753	kino L. E. D.	901
et sodæ	703	lauri cinnamomi E.	896
Testæ L.	448	mimosæ catechu E.	895
Terebinthina canadensis L.	477	moschi D.	901
resina liquida E. D.	476	muriatis ferri c. oxy. rubro D.	756
chia L.	487	myrrhæ L. E. D.	901
veneta D. E.	476	opii L. E. D.	902
vulgaris L.	474	camphorata E.	892
; resina D.	ib.	ammoniata E.	904
Tigllü oleum	301	quassia E. D.	903
Tinctura acetatis ferri D.	757	rhei L. E. D.	ib.
ferri c. alcohole D. ib.		composita L.	



	PAGE		PAGE
Tinctura rhei et alōes E.	904	Unguentum oxidi hydrarg. rubri E.	980
et gentianæ E.	ib.	præcip. albi L.	981
saponis E.	ib.	zinci impuri E.	982
et opii E.	988	zinci E. D.	985
scillæ L. D.	904	picis liquidæ L. D. E.	982
sennæ L.	ib.	nigræ, L.	983
composita E.	ib.	piperis nigri D.	ib.
serpentariæ L. D.	ib.	pulv. cantharidis E.	983
toluiferi balsami E.	905	resinæ nigræ E.	ib.
valerianæ L. D.	ib.	albæ D.	974
ammom. L.	906	sambuci L. D.	983
veratri albi E.	ib.	sabinæ D.	974
zingiberis L. D.	ib.	simplex E.	984
Toluiferæ balsami balsamum E.	596	spermatis ceti D.	976
Tormentillæ radix L. D. E.	597	subacetatis cupri E.	984
Toxicodendria folia L.	530	submur. hyd. ammon. D.	981
Tragacantha L. D.	205	subnitratis hydrarg. D.	ib.
Trifolium paludosum D.	423	sulphuris L. E. D.	984
Tritici farina amyllum D.	598	comp. L.	985
Trochisci carbonatis calcis F.	959	tutiae D.	982
magnesia	ib.	veratri L.	985
glycyrrhizæ glabræ E.	960	zinci L.	ib.
e. opio E.	ib.	Uvæ passæ L. E.	608
gummosi E.	ib.	sole siccata D.	ib.
nitratis potassæ E.	ib.	ursi folia L. D.	178
Tussilago L. D.	601	Valerianæ radix L. E. D.	603
farfara, folia, flores E.	ib.	Veratri radix L. E. D.	604
Tutia D.	625	Vinum L. E.	610
		alōes L. E. D.	920
U.		antimonii tartarizati L.	731
Ulmi cortex L. D.	621	colchici L.	921
campestris cortex E.	ib.	gentianæ comp. E.	922
Unguentum acidi nitrosi E. D.	975	ferri L. D.	759
acetatis plumbi E.	ib.	ipēcacuanhæ L. E. D.	922
æruginis D.	984	nicotianæ tabaci E.	923
calaminare D.	972	opii L.	ib.
cantharidis L.	982	rhei E.	924
carbonatis plumbi E. D.	981	tartratis antimonii E.	729
ceræ albæ D.	976	veratri L.	924
flavæ D.	975	Violæ flore L. D.	604
cerussæ D.	982	Virga aurea; flores, folia D.	564
cetacei L.	976		
elemi compositum L.	ib.	W.	
elemi L.	ib.	Winteræ aromaticæ cortex E.	62
gallæ E.	980		
hellebori albi D.	985	Z.	
hydrargyri L. E. D.	976		
fortius L.	ib.	Zedoaria; radix D.	3
mitius L. D.	977	Zincum L. E. D.	6
hydrarg. nitratis L. E.	979	Zinci oxidum E. D.	7
nitric. oxydi L.	980	oxydum L.	7
infusi cantharidis E.	981	sulphas L. E. D.	7
juniperi sabinæ E.	982	Zingiberis radix L. D.	6
nitratis hyd. mitius E.	980	Zingiber; radix condita D.	
oxydi hydrarg. cin. E.	979		

THE END.

LONDON:

Printed by A. & R. Spottiswoode,  
New-Street-Square.















